

Sensory and rheological properties of fortified coconut milk based chocolate drink as influenced by cocoa powder and sugar levels

Comment [HP1]: Perlu semaksemulapenulis antajukini.

ABSTRACT

Purpose: The sensory and rheological properties of fortified drinks produced from coconut milk, cocoa powder and sugar was evaluated.

Methodology: Drinks samples were formulated with different concentration of coconut milk, cocoa powder and sugar levels. Samples includes: K1 (Coconut milk 100 %), K2 (Coconut milk 99.8 % cocoa powder 0.2 %), K3 (Coconut milk 99.6 %, cocoa powder 0.4%), K4 (Coconut milk 98 %, sugar 2%), K5 (Coconut milk 97.8 %, cocoa powder 0.2%, sugar 2%), K6 (Coconut milk 97.6 %, cocoa powder 0.4%, sugar 2%), K7 (Coconut milk 96 %, sugar 4%), K8 (Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%), K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%). The samples were produced and subjected to physicochemical, proximate, vitamins and minerals analysis using standard methods.

Results: Sensory evaluation; appearance from 8.13(K9) to 7.00 (K2), Aroma from 7.93 (K9) to 6.53 (K1), Mouth feel from 7.83 (K8) to 6.63 (K1) and 6.63 (K3), Taste from 7.90(K9) to 6.27 (K1) and General Acceptability from 8.30 (K9) to 6.40 (K3). The rheological parameters using the power law; consistency index (m) 0.003 to 0.19 (Ns^n/m^2), flow behavior index (n) $n < 1$ for all drink samples. For the Arrhenius model, μ_0 (frequency factor) ranged from 1.9×10^{-5} to 5.1×10^{-8} . E_a (activation energy) ranged from 15.55 to 30.70 KJ/mol. The r squared for both power law model and Arrhenius model, ranges from 0.825 to 0.99 that is approximately 1. Cocoa powder and sugar levels improved sensory profile. The flow behaviour indices (n) of drinks were all less than 1 ($n < 1$) this implies pseudoplastic flow behaviour for drinks samples.

Comment [HP2]: Review the abstract writing

1.0 Introduction

Coconut is the fruit of the coconut palm (*Cocos nucifera* L.). It has a hard shell, edible white flesh and clear liquid (water), it is often used for its water, milk, oil, and tasty meat (Abdullahi & Yakubu, 2014). Coconut milk is an opaque, white liquid with a rich taste extracted from the meat of a mature coconut. Coconut contains sugar, dietary fiber, proteins, fat, antioxidants, vitamins and minerals which is essential for human nutrition (Belew et al., 2014).

Coconut milk is one of the most popular plant based milk alternatives to replace animal milk. The increasing demand for coconut milk worldwide is contributed by increasing veganism, increasing demand for lactose-free milk, and the discovery of the health benefits of coconut milk fats and vitamins (Matin et al., 2020).

Cocoa powder is derived from cocoa beans, seed of the cacao tree (*Theobroma cacao*), and the key ingredient in producing chocolate and chocolate drinks. A good source of high quality protein, vitamins A & D, B-12, riboflavin, and minerals (calcium phosphorus, magnesium, and zinc) (Hanks et al., 2014). Chocolate drinks are one of the cocoa-derived products that are in high demand among consumers

Food fortification is the practice of deliberately increasing the content of one or more micronutrients (vitamins and minerals) in a food or condiment to improve the nutritional quality of the food supply and provide a public health benefit with minimal risk to health (Regulations, 2021)

Many consumers are interested in reducing their consumption of animal products such as cow milk, goat milk etc due to health, environmental and ethical issues. Developing plant based alternatives such as coconut milk based chocolate drink could serve as an alternative with better nutritional profile for people with lactose intolerance, cow milk allergy and vegetarians. Producing a milk drink from coconut milk and cocoa powder will produce another variety of drink, thus diversification of food production (Aribah et al., 2020).

Most beverages available in the markets are energy drinks with high sugar content and little nutritional value which is a concern to diabetes patients and people with heart diseases. Producing a drink from coconut milk and cocoa powder with low sugar could be a good source of nourishment thus improved nutritional status (Putri & Sukma, 2021).

Also cow milk is becoming increasingly expensive, using coconut milk and cocoa powder as alternative could be of advantage since they are locally cultivated, readily available and affordable in the local markets thus reduce post-harvest losses of crop.

Most flavourant and colourants used in the industrial production of drink are synthetic. The use of natural cocoa powder from plant sources could be more beneficial to the consumer's health because they are sources of minerals, fibre and vitamins and bioactive compounds (Klis et al., 2023). The combination of coconut milk and cocoa powder could improve the

health benefits of milk drink thus improving the nutritional status and rendering it more functional non-alcoholic milk drink.

Comment [HP3]: Good introduction but the way of writing this introduction is not interesting. It is recommended that this introduction be rewritten to attract the reader's interest

2.0 MATERIAL AND METHODS

The material used for the processing of drink samples includes, coconut fruits, cocoa beans, sugar. These were purchased from modern market Makurdi Benue state Nigeria for processing.

Comment [HP4]: Set justify margin

2.1 Preparation of cocoa powder

Cocoa powder was produced from cocoa beans as shown on figure 1. The cocoa beans was sorted and cleaned to removed dirt and foreign particles. The beans was roasted at a temperature of 150-190 °C for 5 to 15 min. (Setiadi et al., 2021) . The roast beans was peeled and the shells removed by winnowing. The seeds was then crushed and ground for oil extraction. The cocoa mass / liquor was pressed using a hydraulic press to extract the cocoa butter. The mass of the roast beans after the cocoa butter was extracted was dried for the production of cocoa butter. As the pressing does not remove all the cocoa butter, so the particles will remain coated with a thin layer of cocoa butter with fat content of cocoa powder varies from 8% to 26%. The cake was then further dried at 60 °C for 24 h and then crushed and grind finely to very small particle sizes. The powder was then sieved with a sieve of particle size < 0.5 mm to obtain a fine powder. The cocoa powder was packaged and stored for further processing to chocolate drink.

Comment [HP5]: Figure 1

Comment [HP6]: Does this cocoa production method use the method used by Setiadi et al., 2021. If not, these processes should be written from where these methods are quoted


2.2 Preparation of coconut milk

Coconut milk was prepared from grated meat or endosperm by mechanically expressing the milk (figure 2).

Comment [HP7]: Figure 2

The husk of matured coconut used was removed, exposing the hard shell. The shell of the coconut was removed manually by cracking the shell open with a knife or hitting it against a hard surface. Paring was done by scraping away of the brown skin attached to the coconut

meat to give white coconut meat using a knife. The paired coconut was rinsed with clean water to remove specks of dirt and impurities attached to the coconut meat. The clean coconut meat was cut into smaller sizes with the use of a knife. The coconut cut was pulverized/ground into smaller particles using a blender with added potable water. Milk was extracted by pressing the ground coconut to extract the milk. A cheesecloth was used to squeeze out milk from the blended coconut. The extracted milk was filtered to remove any form of sediments. Extracted coconut milk was pasteurized at 70 °C for 15 min. Cooled, filled into plastics bottles, sealed and labeled and stored under refrigerated conditions (2-4) °C (Adhikari, 2018).







Figure 1: Flow chart for the production of cocoa powder

Comment [HP8]: Figure 1 should be placed before or after the explanation

Comment [HP9]: make sure this figure is on 1 page only






Figure 2: Flow chart for the production of coconut milk

Comment [HP10]: Figure 2 should be placed before or after the explanation

Comment [HP11]: make sure this figure is on 1 page only

2.3 Chocolate Drink formulation and sampling


Drink samples was formulated by varying the proportions of coconut milk, cocoa powder and sugar using 1 x 3 x 3 experimental design comprising 3 levels of cocoa powder (0 %, 0.2 %, 0.4 %) and 3 levels of sugar (0 %, 2 %, and 4 %) which yielded 9 experimental samples, where 100 % coconut milk was used as control. The drinks were subjected to nutritional quality analysis using standard methods (AOAC, 2012).

2.4 Production of coconut-based chocolate drink

Coconut based chocolate drink was produce by blending coconut milk and cocoa powder in different proportions to obtained the formulated sample that was subjected to analysis as seen

on figure 3. Coconut milk and cocoa powder was mixed during which sugar (Optional) was added. The blend was filter to get rid of any foreign particles that might have gained entering during the mixing. The mixture is homogenized to have uniform distribution of particles and also to achieve a uniform consistency. The drink was then be Pasteurization at the temperature of 85 -90 °C for 3-5min to get rid of pathogenic and food spoilage microorganisms. The drink is then be Cooled to 10-15 and packaged in air tight containers for storage.

Comment [HP12]: Figure 3






Figure 3: Flow chart for the production of coconut-based chocolate drink

Source: (Abadi et al., 2023)

Comment [HP13]:

Comment [HP14]: make sure this figure is on 1 page only

2.4.0 Analysis

2.4.1 Rheological analysis of drink samples

Apparent viscosity (Shear stress) was obtained using a Brookfield digital viscometer (Model DV-II, Spindle-0, Canada), using standard methods (Hymavathi et al., 2020). The experimental measurements was performed in triplicate using spindle 61 at different shear rates (5, 10, 20, 50 and 100 rpm) and at different temperatures, (10, 20, 30, 40 °C). The temperatures was defined in order to provide rheological data of the products aiming at a possible application on an industrial scale, based on its relevance in sizing of equipment

and piping (Silva et al., 2020). Apparent viscosity versus shear rate data was fitted with the power law model and Arrhenius equation as follows

2.4.1.1 Power law model

The rheological data were analyzed using Power law

$$\tau = m\dot{\gamma}^{n-1} \quad (1)$$

Linearizing gives

$$\ln \tau = \ln m + (n - 1) \ln \dot{\gamma} \quad (2)$$

Where,

μ = viscosity or shear stress of drink samples (Ns/M²)

$\dot{\gamma}$ = Share rate (rpm)

M = consistency index

n = flow behavior index

Plotting a graph of $\ln \mu$ versus $\ln \dot{\gamma}$ (ln of viscosity versus ln of shear rate) gives slope as n-1 and intercept as $\ln m$

2.4.1.2 Arrhenius equation

$$\tau = \tau_0 \cdot e^{-\frac{Ea}{RT}} \quad (3)$$

Linearizing this equation gives (3)

$$\ln \tau = \ln \tau_0 - \frac{Ea}{R} \left(\frac{1}{T} \right) \quad (4)$$

Where,

μ = viscosity or shear stress of drink samples (Ns/M²)

μ_0 = frequency factor (Arrhenius constant)

e=The base of natural logarithm (Eulers number)

Ea= Activation energy (KJ/mol)

R= The universal gas constant 0.008314 KJ/mol °C

T= absolute temperature associated with the reaction

Plotting a graph of $\ln \mu$ versus $\frac{1}{T}$ (ln of viscosity versus 1/temperature) gives

$$\text{Slope} = -\frac{Ea}{R} \quad \text{intercept} = \ln \tau_0$$

2.4.2 Sensory evaluation of drink samples

Sensory evaluation of the drink samples was carried out using 30 panelists. Panelists were required to evaluate the appearance, aroma, taste, mouth feel and general acceptability of the drink samples using a 9-point Hedonic scale with 1 = dislike extremely, 2 = dislike very much, 3 = dislike moderately, 4 = dislike slightly, 5 = neither like nor dislike, 6 = like slightly, 7 = like moderately, 8 = like very much, 9 = like extremely. Each panelist was provided with 9 different samples of drink at 10-12 °C in disposable plastic cups with coded labels. The panelist was required to taste the drink samples and fill the necessary information on the questionnaire. A cup of potable water was also provide to rinse the mouth after the analysis of each sample (Amadou et al., 2020; Perna et al., 2014; Steve & Olufemi, 2019).

2.4.2 Statistical analysis

Data were subjected to analysis of variance (one-way ANOVA) where it was appropriate and means separated by Duncan's Multiple Range test (DMRT) at 0.05 level of significance. Using the statistical package for social sciences SPSS version 28.(Matin et al., 2020)

3.0 Results and discussion

3.1 Sensory properties of fortified coconut milk based chocolate drinks

The mean sensory scores of fortified coconut milk based chocolate is presented in table 1

The appearance scores for fortified drinks samples ranged from 7.77 (like very much) to 8.70 (like extremely) for coconut milk 99.8 % cocoa powder 0.2 % and Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4% with significant difference at $p < 0.05$.

Aroma of fortified drink samples ranged from 7.40 (like moderately) to 8.57 (like for samples coconut milk 99.8 % cocoa powder 0.2 % and Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4% with significant difference at $p < 0.05$.

Mouth feel ranged from 7.20 (like moderately) to 8.37 (like very much) for Coconut milk 99.6 %, cocoa powder 0.4% and Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4% with significant difference at $p < 0.05$.

Comment [HP15]: Review the results and discussion writing. It should refer to the journal writing format

Comment [HP16]: Table 1

Taste of fortified drinks samples ranged from 7.20 (like moderately) to 8.37 (like very much) for samples coconut milk 99.6 %, cocoa powder 0.4% and Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4% with significant difference at $p < 0.05$

Overall acceptability of fortified drink samples ranged from 7.50 (like very much) to 8.40 (like very much) for samples coconut milk 99.6 %, cocoa powder 0.4% and coconut milk 95.6 %, cocoa powder 0.4%, sugar 4% with significant difference at $p < 0.05$

Table 1: Mean sensory scores for fortified coconut milk based chocolate drinks

Samples code	Appearance	Aroma	Mouth feel	Taste	Overall acceptability
524	7.87 ^{bc} ± 1.22	7.40 ^d ± 0.72	7.63 ^c ± 0.85	7.37 ^b ± 0.96	7.70 ^{bc} ± 0.92
382	7.77 ^c ± 0.86	7.47 ^d ± 1.19	7.53 ^c ± 0.89	7.37 ^b ± 0.99	7.77 ^{bc} ± 0.89
461	7.93 ^{bc} ± 0.83	7.63 ^{cd} ± 0.96	7.50 ^c ± 0.68	7.20 ^b ± 1.06	7.50 ^c ± 0.68

365	8.03 ^{bc} ±1.13	8.03 ^{bc} ±0.96	8.07 ^{ab} ±0.78	8.13 ^a ±0.68	8.03 ^{ab} ±0.72
412	8.13 ^{bc} ± 0.73	8.00 ^{bc} ±0.78	7.90 ^{bc} ±0.76	8.03 ^a ±0.85	8.03 ^{ab} ±0.93
354	8.23 ^a ±0.86	8.20 ^a ±0.99	7.83 ^{bc} ±0.75	7.97 ^a ±1.06	7.97 ^{ab} ±0.89
514	8.07 ^{bc} ± 0.91	7.97 ^{bc} ±0.85	8.03 ^{ab} ±0.63	8.00 ^a ±0.81	8.00 ^{ab} ±0.58
439	8.37 ^a ±0.62	8.57 ^a ± 0.86	8.43 ^a ±0.63	8.16 ^a ±0.79	8.37 ^a ±0.67
452	8.70 ^a ±0.54	8.37 ^a ±0.76	8.50 ^a ±0.66	8.37 ^a ±0.79	8.40 ^a ±0.72

Mean values within the same column having the same letter are not significantly different at $p > 0.05$

Data are means of 30 panelists response on a scale of 9 =like extremely and 1= dislike extremely

Key

The samples code represent fortified drink produced from:

524: Coconut milk 100 %

382: Coconut milk 99.8 % cocoa powder 0.2 %

461: Coconut milk 99.6 %, cocoa powder 0.4%

365: Coconut milk 98 %, sugar 2%

412: Coconut milk 97.8 %, cocoa powder 0.2%, sugar 2%

354: Coconut milk 97.6 %, cocoa powder 0.4%, sugar 2%

514: Coconut milk 96 %, sugar 4%

439: Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%

452: Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%

All drinks samples are fortified with 0.15 mg KI, 2.0 mg FeSO₄, and 1.6 mg Retinol palmitate per 100g of sample

3.1.1 Appearance of fortified coconut milk based chocolate drinks

For appearance, scores by the panelist ranged from 7.87 to 8.70 on a scale of 9 for sample K2 (Coconut milk 99.8 % cocoa powder 0.2 %), and K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%) respectively with significant different at $p > 0.05$. This means the score ranged from like very much to like extremely), this gave a positive impression about the appearance of the drink samples. Panelist appreciated the brown colouration imparted by cocoa powder to the drink.

3.1.2 Aroma of fortified coconut milk based chocolate drinks

In term of aroma, samples K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%) was preferred most and sample K1 (Coconut milk 100 %), was the least preferred. There was significant difference between the samples at $P < 0.05$. The scores ranged from 7.40 to 8.37 (like moderately to like extremely) for K1 and K9 respectively with significance difference between the samples at $p > 0.05$. This means the cocoa powder flavour in the drink samples was appreciated by the panelists.

3.1.3 Taste of fortified coconut milk based chocolate drinks

The taste of the drink samples, ranged from 7.37 to 8.37 (like moderately to like very much) for K1 (Coconut milk 100 %), and K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%) respectively Sample K9 was preferred most and sample K1 was the least preferred. There was significant difference between the samples at $P < 0.05$. Cocoa powder positively influenced the taste of the drink samples.

3.1.4 Mouth feel of fortified coconut milk based chocolate drinks

Mouth feel ranged from 7.53 (K1) to 8.50 (K9) there was significance difference between the samples. The lowest score was in K1 (Coconut milk 100 %) and K3 (Coconut milk 99.6 %, cocoa powder 0.4%) while highest was recorded for K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%). This means panelists like sample K9 extremely and sample K1 and K3 was liked moderately. K1 (Coconut milk 100 %), K2 (Coconut milk 99.8 % cocoa powder 0.2 %), K3 (Coconut milk 99.6 %, cocoa powder 0.4%), K4 (Coconut milk 98 %, sugar 2%), K5 (Coconut milk 97.8 %, cocoa powder 0.2%, sugar 2%), K6 (Coconut milk 97.6 %, cocoa powder 0.4%, sugar 2%), K7 (Coconut milk 96 %, sugar 4%), K8 (Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%), K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%).

3.1.5 General acceptability of fortified coconut milk based chocolate drinks

For overall acceptability, sample K9 (Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%) was the most preferred with the score of 8.30/9 (like very much) and the score was not significantly different from 8.03/9 for K8 (Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%), (like very much). Sample K7 (Coconut milk 96 %, sugar 4%), came third with the score of 7.77 (like very much) and it was not significantly different from samples K9 and K8. The

least preferred samples was sample K3 (Coconut milk 99.6 %, cocoa powder 0.4%), followed by K1 (Coconut milk 100 %), K2 (Coconut milk 99.8 % cocoa powder 0.2 %) with score of 6.40 /9, 6.53/9 (like slightly) and 6.60 (moderately). This was similar to like very much obtained by (Yakum et al., 2022) on the sensory properties of plant based yoghurt. This implies that drink from coconut milk based chocolate drink was generally accepted by the panelists.

3.2.0 Rheological properties of fortified coconut milk based chocolate drinks

The rheological properties of drink samples were studied using the power law and the Arrhenius equation. Results are presented on table 2 and table 3

3.2.1 Rheological properties of drink samples using the Power law model

The relationship between viscosity and shear rates at different temperatures of the drink samples were investigated using Power law model and results presented in table 1

The rheological parameters using the power law gave consistency index (m) as follows

At 10 °C, consistency index (m) ranged from 0.003 to 0.0032 Ns^n/m^2 while flow behavior index (n) were all $n < 1$. Coefficient of regression (r^2) ranged from 0.902 to 0.983 (approximately 1).

At 20 °C consistency index (m) ranged from 0.011 to 0.020 Ns^n/m^2 while flow behavior index (n) were all $n < 1$. Coefficient of regression (r^2) ranged from 0.833 to 0.973 (approximately 1).

At 30 °C consistency index (m) ranged from 0.009 to 0.019 Ns^n/m^2 while flow behavior index (n) were all $n < 1$. Coefficient of regression (r^2) ranged from 0.825 to 0.996 (approximately 1).

At 40 °C consistency index (m) ranged from 0.008 to 0.012 Ns^n/m^2 and flow behavior index (n) $n < 1$ for all drink samples. Coefficient of regression (r^2) ranged from 0.887 to 0.997 (approximately 1).

Table 2: Rheological properties of fortified drink samples using Power law model

T °C	Power law	Sample								
	Parameter	K1	K2	K3	K4	K5	K6	K7	K8	K9
	Intercept	-5.69	-5.754	-5.803	-5.795	-5.814	-5.789	-5.830	-5.748	-6.19
	$m(\text{Ns}^n/\text{m}^2)$	0.003	0.0032	0.0031	0.0030	0.0029	0.0031	0.0029	0.0032	0.003

10	Slope	-0.072	-0.104	-0.103	-0.099	-0.083	-0.086	-0.038	-0.077	-0.067
	N	0.928	0.896	0.897	0.901	0.917	0.914	0.9612	0.923	0.932
	r ²	0.930	0.949	0.9073	0.947	0.902	0.983	0.965	0.902	0.915
20	Intercept	-4.501	-4.200	-4.003	-3.891	-4.185	-3.937	-4.214	-4.158	-3.933
	m(Ns ⁿ /m ²)	0.011	0.014	0.018	0.012	0.015	0.019	0.015	0.016	0.020
	Slope	-0.091	-0.166	-0.248	-0.246	-0.311	-0.264	-0.347	-0.332	-0.273
30	N	0.908	0.834	0.752	0.753	0.688	0.736	0.653	0.668	0.727
	r ²	0.833	0.964	0.965	0.965	0.955	0.932	0.923	0.919	0.973
	Intercept	-4.757	-4.271	-4.173	-4.055	-4.080	-4.195	-4.247	-4.173	-4.056
40	m(Ns ⁿ /m ²)	0.009	0.014	0.015	0.016	0.017	0.018	0.016	0.017	0.019
	Slope	-0.123	-0.201	-0.190	-0.171	-0.175	-0.237	-0.253	-0.242	-0.249
	N	0.877	0.799	0.810	0.829	0.825	0.763	0.747	0.758	0.751
30	r ²	0.996	0.837	0.825	0.908	0.903	0.899	0.863	0.854	0.984
	Intercept	-5.329	-4.626	-4.647	-4.512	-4.932	-4.746	-4.747	-4.569	-4.401
	m(Ns ⁿ /m ²)	0.008	0.009	0.010	0.010	0.010	0.011	0.010	0.011	0.012
40	Slope	-0.308	-0.109	-0.100	-0.108	-0.216	-0.189	-0.182	-0.122	-0.103
	N	0.918	0.891	0.900	0.892	0.839	0.811	0.818	0.878	0.897
	r ²	0.934	0.948	0.889	0.960	0.887	0.973	0.914	0.997	0.974

Key

K1: Coconut milk 100 %

K2: Coconut milk 99.8 % cocoa powder 0.2 %

K3: Coconut milk 99.6 %, cocoa powder 0.4%

K4: Coconut milk 98 %, sugar 2%

K5: Coconut milk 97.8 %, cocoa powder 0.2%, sugar 2%

K6: Coconut milk 97.6 %, cocoa powder 0.4%, sugar 2%

K7: Coconut milk 96 %, sugar 4%

K8: Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%

K9: Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%

M = Consistency index (Nsⁿ/m²)

n = Flow behavior index

r² = Coefficient of regression

All drinks samples were fortified with 0.15 mg KI, 2.0 mg FeSO₄, and 1.6 mg Retinol palmitate per 100g of sample

3.2.1.1 Drinks behavior with Power law model

The changes in viscosity with shear rate for the fortified coconut milk based chocolate drink indicated that, viscosity decreased with shear rate and temperature in all the drink samples.

3.2.1.2 Consistency index (m) of drink samples

Apparent viscosity and consistency index of drink decreased with increasing temperatures.

Consistency index is an indication of the viscous nature of the drink (Alakali & Irtwange, 2015)

3.2.1.3 Flow behaviour index (n) of drink samples

It was observed that flow behaviour indices of drinks were all less than 1 ($n < 1$). Similar results for flow behavior index (n) ranging from 0.657 to 0.723 was obtained by Silva et al 2020 on Rheological behavior of plant-based beverages. The flow behavior index did not show major change with temperature and its value was generally below one, hence coconut milk based chocolate drink exhibit pseudoplastic behavior (Gupta et al., 2022). The increase in viscosity (shear stress) gives more than proportional increase in share rate. The curve is convex and begins at the origin. It is the most common type of non newtonian behavior. The viscosity is dependent on shear rate. And it obeys the power law equation (Silva et al., 2020) .

3.2.1.4 Coefficient of regression (r^2) for drink samples

The high r^2 values indicates that the power law model is appropriate for describing the rheological characteristics of fortified coconut milk based chocolate drink

Apparent viscosity decreases with increasing shear rate, indicating shear thinning behavior (Ashaver et al., 2023).

3.3.0 Rheological properties of drink samples using the Arrhenius equation model

The relationship between temperature and viscosity at different share rates of the drink samples was investigation using Arrhenius equation and results are presented in table 2

At rpm 5, frequency factor (μ_0) for drink samples ranged from 3.3×10^{-6} to 4.1×10^{-7} , Activation energy (Ea) in terms of energy per mole (KJ/mol) from 24.84 to 30.70 KJ/mol, Coefficient of regression (r^2) from 0.810 to 0.952 (Approximately 1).

At rpm 10, frequency factor (μ_0) for drink samples ranged from 1.9×10^{-5} to 1.7×10^{-7} , Activation energy (Ea) in terms of energy per mole (KJ/mol) from 22.25 to 26.81 KJ/mol, Coefficient of regression (r^2) from 0.810 to 0.993 (Approximately 1).

At rpm 20, frequency factor (μ_0) for drink samples ranged from 5.1×10^{-6} to 9.7×10^{-6} , Activation energy (Ea) in terms of energy per mole (KJ/mol) from 18.93 to 22.96 KJ/mol, Coefficient of regression (r^2) from 0.838 to 0.948 (Approximately 1).

At rpm 50, frequency factor (μ_0) for drink samples ranged from 1.1×10^{-5} to 9.7×10^{-6} , Activation energy (Ea) in terms of energy per mole (KJ/mol) from 17.68 to 19.82 KJ/mol, Coefficient of regression (r^2) from 0.803 to 0.961 (Approximately 1).

At rpm 100, frequency factor (μ_0) for drink samples ranged from 3.1×10^{-5} to 9.9×10^{-6} , Activation energy (Ea) in terms of energy per mole (KJ/mol) from 15.55 to 19.45 KJ/mol, Coefficient of regression (r^2) from 0.817 to 0.997 (Approximately 1).

Table 3 Rheological of fortified drink samples using Arrhenius equation model

Share rate	Arrhenius	Sample									
		Parameter	K1	K2	K3	K4	K5	K6	K7	K8	K9
5	Rpm										
	Intercept	-13.32	-14.39	-14.66	-15.46	-14.04	-14.93	-15.09	-15.66	-16.80	
	μ_0	1.6E-6	5.6E-7	4.3E-7	1.9E-7	7.9E-7	3.3E-7	2.7E-7	1.6E-7	5.1E-8	
	Slope	2987.7	3237.9	3387.1	3317.3	3473.7	3517.0	3589.2	3627.6	3692.6	
	Ea(KJ/mol)	24.84	26.92	28.16	27.58	28.88	29.24	29.84	30.16	30.70	
10	r^2	0.938	0.952	0.929	0.915	0.845	0.859	0.939	0.876	0.929	
	Intercept	-10.85	-12.59	-14.11	-12.17	-13.23	-14.30	-12.97	-14.49	-15.59	
	μ_0	1.9E-5	3.4E-06	1.0E-07	7.4E-7	1.8E-6	6.2E-7	2.2E-6	5.1E-7	1.7E-7	
	Slope	2676.2	2969.7	3115.2	2927.6	3013.0	3133.3	3128.5	3193.4	3224.7	
	Ea(KJ/mol)	22.25	24.69	25.90	24.34	25.05	26.05	26.01	26.55	26.81	
20	r^2	0.983	0.993	0.953	0.947	0.837	0.912	0.810	0.972	0.915	
	Intercept	-12.18	-14.70	-12.460	-11.30	-13.05	-13.14	-10.274	-11.55	-12.17	
	μ_0	5.1E-6	4.1E-6	3.9E-6	1.2E-6	2.2E-6	2.0E-6	3.5E-6	9.7E-6	5.2E-6	
	Slope	2276.9	2487.4	2504.2	2666.6	2694.3	2724.3	2681.0	2720.7	2761.6	
	Ea(KJ/mol)	18.93	20.68	20.82	22.17	22.40	22.65	22.29	22.62	22.96	
50	r^2	0.858	0.941	0.867	0.892	0.838	0.862	0.944	0.933	0.948	
	Intercept	-13.80	-13.44	-11.41	-9.43	-12.23	-12.791	-11.55	-11.96	-11.95	
	μ_0	1.0E-6	1.4E-6	1.1E-5	8.0E-5	4.9E-6	2.8E-06	9.7E-6	6.3E-6	6E-6	
	Slope	2126.5	2297.3	2355.1	2201.1	2351.5	2373.1	2284.1	2341.8	2383.9	
	Ea(KJ/mol)	17.68	19.10	19.58	18.30	19.55	19.73	18.99	19.47	19.82	
100	r^2	0.816	0.923	0.831	0.820	0.934	0.904	0.803	0.901	0.961	
	Intercept	-13.54	-12.58	-10.87	-10.38	-12.97	-12.87	-11.52	-11.85	-11.93	
	μ_0	1.3E-6	3.4E-06	1.9E-05	3.1E-5	2.3E-6	2.6E-6	9.9E-6	7.1E-6	6.6E-6	
	Slope	1870.3	1896.8	1975.0	1910.0	1967.8	2079.6	2127.7	2270.9	2339.4	
	Ea(KJ/mol)	15.55	15.77	16.42	15.88	16.36	17.29	17.69	18.88	19.45	
100	r^2	0.867	0.917	0.890	0.817	0.875	0.828	0.894	0.889	0.997	

KEY

- K1: Coconut milk 100 %
- K2: Coconut milk 99.8 % cocoa powder 0.2 %
- K3: Coconut milk 99.6 %, cocoa powder 0.4%
- K4: Coconut milk 98 %, sugar 2%
- K5: Coconut milk 97.8 %, cocoa powder 0.2%, sugar 2%
- K6: Coconut milk 97.6 %, cocoa powder 0.4%, sugar 2%
- K7: Coconut milk 96 %, sugar 4%
- K8: Coconut milk 95.8 %, cocoa powder 0.2%, sugar 4%
- K9: Coconut milk 95.6 %, cocoa powder 0.4%, sugar 4%
- μ_0 = frequency factor (Arrhenius constant)

E_a = Activation energy in terms of energy per mole (KJ/mol)
 r^2 = Coefficient of regression

All drinks samples are fortified with 0.15 mg KI, 2.0 mg FeSO₄, and 1.6 mg Retinol palmitate per 100g of sample

3.3.1 Drinks behavior with Arrhenius equation

There was an inverse relationship between viscosity and temperature hence viscosity decreases as temperature increases and thus exhibits Arrhenius type relationship. Arrhenius equation describes the relationship between the rate of reaction and temperatures for many physical and chemical reactions(Correa et al., 2018).

3.3.2 Frequency factor (μ_0) of fortified drink samples

There was an inverse relationship between viscosity and temperature hence viscosity decreases as temperature increases and thus exhibits Arrhenius type relationship

3.3.3. Activation energy (E_a) of drink samples

Activation energy (E_a) is defined as minimum energy required which overcomes the energy barrier before the elementary flow can occur (Azeddine et al., 2023).

The viscous flow occurs as a sequence of events which are shift of particles in the direction of shear force action from one equilibrium position to another position by overcoming a potential energy barrier. The barrier height determines the free activation energy of viscous flow. Higher activation energy values indicate a greater influence of temperature on the viscosity, i.e. more rapid change in viscosity with temperature. When temperature increased, the thermal energy of the molecules and intermolecular spacing increased significantly, which lead to decrease in the magnitude of viscosity

3.3.4 Coefficient of regression (r^2) of drink samples using the Arrhenius equation

The high r^2 values indicates that the Arrhenius equation is appropriate for describing the rheological characteristics of fortified coconut milk based chocolate drink since the values were all approximately 1.

Arrhenius equation describes the relationship between the rate of reaction and temperatures for many physical and chemical reactions(Correa et al., 2018)

4.0 CONCLUSION

The production of fortified coconut milk based chocolate drinks is possible and looking at its sensory profile it was generally accepted by the panelist

The flow behaviour indices (n) of drinks did not show major change with temperature and its value were all less than 1 ($n < 1$). This implies coconut milk based chocolate drink exhibit pseudoplastic behavior thus a non-Newtonian fluid. There is an inverse relationship between viscosity and temperature hence viscosity decreases as temperature increases and thus exhibits Arrhenius type relationship. Cocoa powder and sugar levels improved the sensory properties of the drink samples since the overall best sample contains 0.4 % cocoa powder and 4 % sugar content.

REFERENCES

- Abadl, M. M. T., Marzlan, A. A., Sulaiman, R., Abas, F., & Meor Hussin, A. S. (2023). Optimization of Coconut Milk Kefir Beverage by RSM and Screening of Its Metabolites and Peptides. *Fermentation*, 9(5), 1–17. <https://doi.org/10.3390/fermentation9050430>
- Abdullahi, M. K., & Yakubu, A. A. (2014). Determinants of Non Alcoholic Beverages (NAB) Consumption in North-Western Nigeria: A study of Sokoto Metropolis. *Nigerian Journal of Basic and Applied Sciences*, 21(4), 273. <https://doi.org/10.4314/njbas.v21i4.5>
- Adhikari, B. and N. G. (2018). Preparation and Quality Evaluation of coconut milk incorporated yoghurt. *Institute of Science and Technology Journal*, 3(5), 15–22.
- Alakali, J. S., & Irtwange, S. V. (2015). Rheological characteristics of food gum (*Cissus populnea*). *African Journal of Food Science*, 3(9), 237–242.
- Amadou, N. M., Richard, E. A., Jules-roger, K., & Dung, M. S. (2020). Advances in Nutrition and Food Science ISSN : 2641-6816 Minerals , Antioxidant Activity and Consumer Preference of Ginger Spiced Yoghurt Minerals , Antioxidant Activity and Consumer Preference of. *Www.Kosmospublishers.Com Contact@kosmospublishers.Com Research*, 2020(01).
- AOAC. (2012). *AOAC Official Method 2012.09 Vitamins A and E in Infant Formula and Adult/Pediatric Nutritional Formula*. 9.
- Aribah, S. Al, Sanjaya, A. P., Muhammad, D. R. A., & Praseptiangga, D. (2020). Sensorial and physical properties of chocolate beverage prepared using low fat cocoa powder. *AIP*

- Conference Proceedings*, 2219(May). <https://doi.org/10.1063/5.0003435>
- Ashaver, V. A., Ariaahu, C. C., Yusufu, M. I., Ariaahu, E. C., & Gbuusu, B. (2023). Formulation of Aqueous Drinks Using Tigernut and Moringa Seeds : Effect of Adding Chemical Hurdles and Fortificants on Physico-Chemical and Rheological Properties. *Asian Journal of Food Research and Nutrition Volume*, 2(4), 847–862.
- Azeddine, F., Sergio, P. A., Khadir, L. El, Ali, I., & Houssayne, B. El. (2023). *clean technologies Rheological Behavior and Characterization of Drinking Water Treatment Sludge from Morocco*. 259–273.
- Belewu, M. A., Muhammed-lawal, A., & Abdulsalam, K. (2014). Date-Coconut Drink : Physico-Chemical and Sensory Qualities D Ate -C Oconut D Rink. *Daffodil International University Journal of Science and Technology*, 9(2), 1–7.
- Correa, D. A., Margarita, P., & Castillo, M. (2018). *Rheology of a Traditional Dairy Beverage from Colombian Caribbean*. 11(23), 1103–1110.
- Gupta, M. K., Torrico, D. D., Ong, L., Gras, S. L., Dunshea, F. R., & Cottrell, J. J. (2022). Plant and Dairy-Based Yogurts : A Comparison of Consumer Sensory Acceptability Linked to Textural Analysis. *Foods*, 11(453), 1–18.
- Hanks, A. S., Just, D. R., & Wansink, B. (2014). Chocolate milk consequences: A pilot study evaluating the consequences of banning chocolate milk in school cafeterias. *PLoS ONE*, 9(4), 1–7. <https://doi.org/10.1371/journal.pone.0091022>
- Hymavathi, D. T., Roberts, T. P., Jyothsna, E., & Sri, V. T. (2020). Proximate and mineral content of ready to use minor millets. *International Journal of Chemical Studies*, 8(2), 2120–2123. <https://doi.org/10.22271/chemi.2020.v8.i2af.9065>
- Klis, V., Pühn, E., Jerschow, J. J., Fraatz, M. A., & Zorn, H. (2023). Fermentation of Cocoa (Theobroma cacao L.) Pulp by *Laetiporus persicinus* Yields a Novel Beverage with Tropical Aroma. *Fermentation*, 9(6), 1–13. <https://doi.org/10.3390/fermentation9060533>
- Matin, A., Rahman, N., Islam, T., & Ahmed, F. B. H. (2020). Effect of adding coconut milk on the physicochemical, proximate, microbial and sensory attributes of «Dahi». *Ukrainian Journal of Food Science*, 8(1), 49–57. <https://doi.org/10.24263/2310-1008-2020-8-1-6>
- Perna, A., Intaglietta, I., Simonetti, A., & Gambacorta, E. (2014). Antioxidant activity of yogurt made from milk characterized by different casein haplotypes and fortified with chestnut and sulla honeys. *Journal of Dairy Science*, 97(11), 6662–6670. <https://doi.org/10.3168/jds.2013-7843>
- Putri, N. T., & Sukma, M. (2021). Development of Chocolate Beverage Products by Considering Consumer Preferences. *Proceedings of the Second Asia Pacific International Conference on Industrial Engineering and Operations Management*, 1782–1793.
- Regulations, F. F. (2021). *Federal Republic of Nigeria Official Gazette*. 135.
- Setiadi, P. A., Saputro, A. D., Nurkholisa, Z., Hardiyanto, Y. F., Bintoro, N., & Karyadi, J. N. W. (2021). Physical characteristic of instanised chocolate beverage powder produced with palm sugar and sucrose as sweeteners. *IOP Conference Series: Earth and Environmental Science*, 653(1), 1–8. <https://doi.org/10.1088/1755-1315/653/1/012038>

Silva, K., Machado, A., Cardoso, C., Silva, F., & Freitas, F. (2020). Rheological behavior of plant-based beverages. *Food Science and Technology*, 2061(June), 258–263.

Steve, O., & Olufemi, O. (2019). Optimization and Evaluation of Some Physico- Chemical and Nutritional Properties of Cereal-Based Soya-Fortified Flours for Dough Meal. *Research Gate*, 6(1), 40–59.

Yakum, K. N., Amove, J., Peter, A., Forwukeh, V. H., & Kemjie, L. (2022). Physicochemical and sensory properties of yoghurt from cow milk , soy bean milk and tiger nut milk blends spiced with ginger powder. *International Journal of Food Science and Nutrition*, 7(3), 42–51.

UNDER PEER REVIEW