

**Original Research Article**  
**A Polyherbal Formulation's (Storg B) *In vitro***  
**Antioxidant, *In vivo* Bioavailability and**  
**Immunomodulatory Effects on Experimental**  
**Animal Model Induced by Cyclophosphamide**

---

**ABSTRACT**

**Introduction:** The immunomodulatory properties of plants are being explored extensively these days as they contain a wide range of bioactive compounds that can treat immunomodulatory disorders. The aim of present investigation was to determine the efficacy of the standardised polyherbal formulation Storg-B. The polyherbal formulation combining *Psidium guajava*, *Mangifera indica*, and *Citrus limon*.

**Methods:** The aqueous extract of polyherbal formulation (Storg B) investigated for DPPH radical scavenging activity to determine antioxidant activity. SD rats were selected for the evaluation of immunomodulatory activity of Storg-B. Haemagglutination and Hematology method was used to evaluate the immunomodulatory studies. For the pharmacokinetic studies LC-MS/MS technique is employed to quantify the Storg B.

**Results:** The percentage inhibition of DPPH radial IC<sub>50</sub> values of standard BHT and Storg-B were 60.52 and 68.358µg/ml, respectively. In the *in vivo* animal studies, Storg-B showed significant immunomodulatory activity by increasing the Primary and secondary antibody titer value by 6.58 and 8.25 mg/mL respectively, to the SRBC antigen, at the dose of 100mg/kg. There's also increase in the neutrophil adhesion percentage in the testing doses. Results of hematological parameters revealed that polyherbal extract (Storg-B) restored all the levels of blood components in 100 mg/kg dose. Polyherbal formulation (Storg-B) containing B vitamins showed greater potency in the bioavailability studies.

**Conclusion:** Thus, polyherbal formulation Storg-B may be used as an efficient and secure formulation in the treatment of immune deficiency by improving immune response.

*Keywords: Polyherbal formulation (PHF), Antioxidant, Immunomodulatory, Bioavailability, B vitamins, In vitro and In vivo*

## 1. INTRODUCTION

The majority of the world's population has recently witnessed a shift from synthetic to natural medicine. It incorporates the use of natural herbal elements to replenish the body's vital nutrients and fundamentals for leading a healthy lifestyle [1]. Currently available chemical based immunomodulators, such as levamisole, glycans, tellerones and L-fucose, continue to

cause immunosuppression leading to adverse health effects in humans, particularly flu-like symptoms, fatigue, nausea, neutropenia, thrombocytopenia, depression and can also cause other diseases. Due to the emergence of side effects associated with chemical drugs, natural immunomodulators are considered as potential tools that can replace chemical agents in treatment programs [2].

The prevention and treatment of diseases with herbal medicines can be traced throughout human history, in all cultures and throughout the ages for all kinds of ailments. According to the World Health Organization (WHO), approximately 80% of the world's population mainly uses herbal medicines in their primary health care [3,4].

In comparison to a single drug, herbal combinations frequently show strong potential in the treatment of diseases. Combination of herbals may act on multiple targets at the same time to provide a thorough relief [5]. The PHF (polyherbal formulation) removes the need for patients to take many distinct medicines at once, it improves patient convenience and indirectly improves therapeutic impact and compliance. When compared to a single herbal formulation, PHF's popularity in the market has been promoted by all these benefits [6].

Oxidative stress has been determined to be a major cause of many chronic and degenerative diseases, including atherosclerosis, ischemic heart disease, aging, diabetes, cancer, immunosuppression, neurodegenerative diseases, etc [7]. The most effective way to eliminate and reduce the activity of free radicals that cause oxidative stress is the antioxidant defense system. Antioxidants are substances that have the ability to break the chain reactions of free radicals. Recently, there has been increased interest in medicinal plants with curative potential as reactive antioxidants to reduce tissue damage caused by oxidative stress [8].

The immune system's primary function is to differentiate between self and non-self. This non-self might be a body's own cell that can be misinterpreted as a foreign object, an infectious organism, or an organ that has been transplanted [9]. Immunomodulators are biological or synthetic compounds that have the ability to activate, inhibit, or modify any part of the immune system, including the innate and adaptive systems. Extensive study and innovation are needed on natural products, plants, and their extracts to understand their immunomodulatory efficacy. It might be an essential commodity to develop a novel herbal formulation to supplement existing chemotherapies.

Pharmacokinetics research is an important part of innovation in herbal medicine. Pharmacokinetics deals with the absorption, distribution, metabolism, and absorption of compounds in the body. Bioavailability is the part of a drug dose that is absorbed from the place of administration and reaches the systemic circulation unchanged. The drug, its route of administration and the composition determine the amount of the administered dose that is absorbed into the bloodstream. Knowing the absorption and metabolism properties of the drug makes it easy to prepare and administer medicines for maximum benefit and minimum risk.

Our product, PHF (Storg B), is an enriched B vitamins supplementation from the three plants that have been used for decades for traditional medicinal and nutraceutical uses, namely *Psidium guajava*, *Mangifera indica*, and *Citrus limon*. We used an animal model of cyclophosphamide-induced immunosuppression in SD rats for the immunomodulatory studies.

## 2. MATERIAL AND METHODS

### 2.1 Preparation of sample

Storg B is manufactured and registered by Star Hi Herbs PVT Ltd Jigani, Bangalore, Karnataka, India.

### 2.2 Chemicals

All standards were purchased from Sigma-Aldrich. The Standards were stored in the original packages at -18°C prior to use. Methanol (JT baker for LC-MS), formic acid (Optima from Fisher Chemicals, LC-MS grade), Milli-Q-water from Sartorius ARIUM MINI.

### 2.3 Antioxidant activity of Storg-B

PHF Storg-B's ability to scavenge free radicals was evaluated using the DPPH radical scavenging assay, as described by Blois [10]. A solution of 0.1 mM DPPH in methanol was prepared, and 2.4 mL of this solution was mixed with 1.6 mL of Storg-B in methanol at different concentrations (20, 40, 60, 80, and 100 µg/mL). The sample was vortexed thoroughly and left in the dark at RT for 30 min. The absorbance of the sample was measured spectrophotometrically at 517 nm. BHT was used as a reference drug at different concentrations (20, 40, 60, 80, and 100). The percentage of DPPH radical scavenging activity was calculated by the following equation:

$$\% \text{ of DPPH radical scavenging activity} = (A_0 - A_1)/A_0 \times 100$$

Where  $A_0$  is the absorbance of the control and  $A_1$  is the absorbance of the Storg-B. Then the percent of inhibition was plotted against concentration, and the  $IC_{50}$  was calculated from the graph. The experiment was repeated three times at each concentration.

### 2.4 *In Vivo* Immunomodulatory activity Storg B

#### 2.4.1 Animals and experimental protocol

The Institutional Animal Ethics Committee (IAEC) granted its approval for the studies' ethical conduct prior to the experiment (BCP/IAEC/06/2022). In this investigation, SD rats weighing between 150g and 250g were employed. They were supplied by the animal house, Bharathi College of Pharmacy, Mandya, Karnataka, India. The rats were kept in standard laboratory settings (temperature  $24 \pm 1$  °C with a 12/12 h dark/light cycle) in polypropylene cages (38 x 23 x 10 cm) with a maximum of six animals per cage. Before the trial, all of the animals were acclimatized to laboratory conditions for a week.

#### 2.4.2 Grouping of Animals

The SD rats were divided into 5 groups each group consisting of 8 animals

Group I: Normal control received only normal saline

Group II: 20 mg/kg of Cyclophosphamide alone treated animals

Group III: Cyclophosphamide and 50 mg/kg of Levamisole treated animals

Group IV: Cyclophosphamide and 50 mg/kg of Storg B treated animals

Group V: Cyclophosphamide and 100 mg/kg of Storg B treated animals

#### 2.4.3 Sheep red blood cell (SRBC) preparation and standardization

Blood from the sheep was drawn out via the external jugular vein using a newly made 1:1 ratio of Alsever's solution. Centrifuging the blood at 2500 rpm for 10 minutes allowed the

SRBCs to be separated, and the collected blood was then rinsed with pyrogen-free normal saline (0.9% w/v). After that, the concentration was adjusted to  $1 \times 10^8$  SRBC in 1 mL of sheep blood [11].

#### **2.4.5 Hemagglutination Assay**

On days 7 and 14, all the groups except control were administered with 0.1 ml of 20% SRBC. On the 9<sup>th</sup> and 16<sup>th</sup> days, all the groups except control were administered cyclophosphamide to suppress immunity. This is also called the immunization of animals. Blood was withdrawn on the 14<sup>th</sup> and 21<sup>st</sup> days from the retro orbital plexus under mild ether anesthesia from all the antigenically sensitized and challenged rats, and serum was separated for antibody titer checks. 20 $\mu$ L of serum was serially diluted with 20 $\mu$ L of saline and filled into a microtiter plate. 20 $\mu$ L SRBC was added to each of these dilutions, and the plates were incubated at 37°C for one hour and then observed for hemagglutination and antibody reactions.

#### **2.4.6 Neutrophil adhesion test**

On the 21<sup>st</sup> day of drug treatment, blood samples were collected by puncturing the retro orbital plexus into heparinized vial and analyzed for total leucocyte counts (TLC) and differential leucocyte counts (DLC) by fixing blood smears and staining with Field stain I and II- Leishman's stain. After initial counts, blood samples were incubated with 80mg/ ml of nylon fibers for 15 min. at 37° C. The incubated blood samples were once again analyzed for TLC. The product of TLC and % neutrophil gives neutrophil index (NI) of blood sample. The percent neutrophil adhesion was calculated as follows:

$$\% \text{ Neutrophil Adherence} = \frac{NI_u - NI_t}{NI_u} \times 100$$

Where,  $NI_u$  is neutrophil index of untreated blood,  $NI_t$  is neutrophil index of fibre treated blood.

$$\text{Neutrophil Index(NI)} = \text{TLC} \times \% \text{neutrophil}$$

#### **2.4.7 Hematological Analysis**

The blood was collected from each group by retro orbital plexus into heparinized collecting tubes. The hematological analysis was carried out to determine the RBC, hemoglobin, total platelet count and DLC by using auto-analyzer.

### **2.5 *In Vivo* Bioavailability Study of Storg B**

#### **2.5.1 Animals used**

Healthy SD rats ( $n = 6$ ) weighing 150-200g and 8 to 12 weeks aged were selected for pharmacokinetic and bioavailability study. All rats were kept for acclimatization for seven days prior to start pharmacokinetic study. Body weight variation among the animals did not exceed 20% of the mean body weight. The room was environmentally-controlled at a temperature of 22° C with relative humidity of 60%. The photoperiod was 12 h light and 12 h darkness. Prescribed feed and ad libitum drinking water were provided during experimental period. The experiment was carried out in accordance with the guidelines of Institutional Animal Ethics Committee.

#### **2.5.2 Experimental design**

Study design six SD rats weighing about 150-200 g were selected as the animal model. The age of the rat was 8–12 weeks. The study involved rats had no medication for two weeks food was withdrawn 12 hours before drug administration and water was available throughout the study. The Storg B dose of 200 mg/kg was administered to rats using a gavage needle.

The study was approved by Institutional Ethical committee (BCP/IAEC/06/2022). The blood samples were taken at the time intervals of 0.5, 1, 2, 4, 8, and 24 hours from the retro orbital plexus of the rats and centrifuged at 6000 rpm for 10 minutes. EDTA disodium salt was used as an anticoagulant. Blood samples were collected into labeled heparin tubes, and stored in  $-20^{\circ}\text{C}$  freezer until use. The concentration of plasma, after PHF Storg-B consumption were determined by LC-MS/MS method. The value of  $C_{\text{max}}$ ,  $T_{\text{max}}$  and  $t_{1/2}$  of Storg-B containing vitamin B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>5</sub>, B<sub>6</sub> and B<sub>9</sub> were obtained by plotting the graph between plasma concentration and time.

### **2.5.3 LC-MS/MS analysis**

Liquid chromatography and Mass Spectrometry was performed by a Shimadzu LCMS-8050, high-end model of its UFMS (Ultra-Fast Mass Spectrometry) with Nexera X3 UHPLC series. Binary Analytical system was used which consisted of a Solvent Delivery Unit LC-40 Series with Mobile Phase Monitor MPM-40, Autosampler SIL-40, Column Oven CTO-40 Series. Heated ESI source which improves desolvation and escalate ionization efficiency with the addition of a heated gas used in combination with the nebulizer gas. UF sweeper® III collision cell increases CID efficiency by optimizing the collision cell pressure and a PC with Lab solution and Insight software.

The LC-MS/MS system was enabled with a chromatographic column Shim pack GIS C18-120 (100 x 4.6 mm, 3 $\mu\text{m}$ ) maintained at  $40^{\circ}\text{C}$ . Mobile phase A was prepared by adding 1mL of formic acid to 1000mL of water (0.1% formic acid in water). Mobile phase B used was 100% methanol. All mobile phases were prepared daily, filtered through a 0.2 $\mu\text{m}$  nylon filter and degassed before use.

Chromatographic partition was achieved at a flow rate 0.5mL/min using isocratic condition of mobile phase A and B as (60:40). Injection volume was 10 $\mu\text{L}$ .

Mass spectrometry analysis was achieved by multiple reactions monitoring (MRM) in ESI positive mode using the following optimized parameters:

Nebulizing gas flow (Nitrogen) = 3 mL/min, Heating gas flow (Zero Air) = 5 mL/min, Interface Temperature =  $300^{\circ}\text{C}$ , DL Temperature =  $180^{\circ}\text{C}$ , Heat Block Temperature =  $350^{\circ}\text{C}$ , Drying Gas Flow (Nitrogen) =  $10^{\circ}\text{C}$ , Interface voltage = 4kV, CID gas (Argon) = 270kPa.

### **2.5.4 Standard preparation**

Standard stock solutions were prepared individually by dissolving an appropriate amount of each Standard in Water (Vitamin B<sub>1</sub>, B<sub>3</sub>, B<sub>5</sub> and B<sub>6</sub>). While preparing Vitamin B<sub>2</sub>, and B<sub>9</sub>, add approximately 1mL of 0.1N NaOH in water for better solubility. Prepare a mixture of all vitamins where the concentration was 20000 ng/mL. All standard stock solutions were stored at  $-18^{\circ}\text{C}$ . The standard mixture solution 20000 ng/mL was prepared daily in an amber colored glass volumetric flask (5 mL) by diluting the calculated amount of each standard stock solution with water.

The calibration solutions were prepared daily in brown glass HPLC vials (1.8 mL) diluting the standard mixture solution 20000 ng/mL. Six different concentrations of stock standard solution were prepared 3 ng/mL to 300 ng/mL. Total volume of each calibration solution was 1mL for stock Standard solution prepared.

### **2.5.5 Sample preparation**

For the sample evaluation, a 2 mL eppendorf tube containing 40 mL of serum sample followed with 160  $\mu\text{L}$  of methanol. After being vortexed for 3 min, the mixture was centrifuged at 6000 rpm for 10 min at  $4^{\circ}\text{C}$ . After centrifugation, 100  $\mu\text{L}$  of the supernatant and 200  $\mu\text{L}$  of water were transferred to the brown glass HPLC vial (1.8 mL) for additional analysis. Vortex for 2 min and then inject.

For the spiked sample evaluation, 40  $\mu\text{L}$  of serum was taken in a 2 mL eppendorf tube, to which 160  $\mu\text{L}$  of methanol and 10  $\mu\text{L}$  of the appropriate standard solution were added. The mixture was vortexed for 3 minutes at  $4^{\circ}\text{C}$ , followed by a 10-minute centrifugation run at

6000 rpm. After centrifugation, 100  $\mu$ L of the cleared supernatant and 200  $\mu$ L of water were transferred to the brown glass HPLC crimp vial (1.8 mL) for further evaluation. Vortex for 2 min, then inject.

## 2.6 Statistical analysis

The statistical analysis of the experimental data was carried out using one-way ANOVA, followed by Tukey's multiple comparison tests, and the results were expressed as Mean  $\pm$  SEM.

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of Storg B on antioxidant Activity

The analysis specification of Storg B is shown in Table No.1. The PHF (Storg-B) antioxidant activity was compared with that of standard butylated hydroxyl toluene (BHT). The ability of the PHF (Storg-B) to donate hydrogen atoms was determined by decolorizing a methanol solution with 2, 2-diphenyl-1-picrylhydrazyl (DPPH). DPPH produced violet color in methanol solutions and faded to shades of yellow in the presence of antioxidants. The free radical antioxidant scavenging activity of standard BHT and Polyherbal extract Storg-B was found to have IC<sub>50</sub> values of 60.52 and 68.358 $\mu$ g/ml, respectively as shown in Fig. 1. The results obtained in the present study indicate that PHF Storg-B exhibits significant free radical scavenging activity as compared with the standard drug BHT.

**Table 1. Analysis specification of Storg-B**

TEST	SPECIFICATION	PROTOCOL
<b>PHYSICAL</b>		
<b>Description</b>	Light brown hygroscopic powder	Organoleptic
<b>Identification</b>	To comply by TLC	TLC
<b>Solubility</b>	Soluble in water Not less than 90% w/w	USP
<b>Loss on drying</b>	Not more than 6.00% w/w (dried at 105 <sup>0</sup> C)	USP<731>
<b>Sieve test ( passes through)-40#</b>	Not less than 98% w/w	USP<786>
<b>CHEMICAL ANALYSIS</b>		
<b>Each 1 gram contains (Assay of actives)</b>		
Natural Vitamin B1	Not less than 3.0 mg (0.30%)	HPLC
Natural Vitamin B2	Not less than 3.33 mg (0.33%)	HPLC
Natural Vitamin B3	Not less than 35mg (3.4%)	HPLC
Natural Vitamin B5	Not less than 17mg (1.7%)	HPLC
Natural Vitamin B6	Not less than 4.2mg (0.42%)	HPLC
Natural Vitamin B9	Not less than 1300 mcg (0.13%)	HPLC
<b>HEAVY METALS</b>		
<b>Lead</b>	Not more than 3.0 ppm	ICP-MS
<b>Arsenic</b>	Not more than 1.0 ppm	ICP-MS
<b>Cadmium</b>	Not more than 1.0 ppm	ICP-MS
<b>Mercury</b>	Not more than 0.1 ppm	ICP-MS
<b>PESTICIDES RESIDUE</b>	Should be Absent	USP
<b>MICROBIOLOGICAL PROFILE</b>		
<b>Total Plate count</b>	Not more than 10000 cfu/g	USP<2021>

<b>Total Yeast and Mould count</b>	Not more than 1000 cfu/g	USP<2021>
<b>Salmonella</b>	Negative/10g	USP<2022>
<b>E. Coli</b>	Negative/10g	USP<2022>
<b>Staph. Aureus</b>	Negative/10g	USP<2022>
<b>P. auriginosa</b>	Negative/10g	USP<62>
<b>Coli forms</b>	Negative/10g	USP<62>
<b>ADDITIONAL INFORMATION</b>		
<b>Sanitizing treatment</b>	Non-irradiated and Not-treated with ETO	
<b>Certification Status</b>	Kosher and Halal certified	
<b>Genetic Modification status</b>	GMO free	
<b>BSE/TSE status</b>	BSE/TSE free	
<b>Country of Origin</b>	India	
<b>Cultivated or wild crafted</b>	Cultivated	
<b>Manufactured by</b>	Star Hi Herbs Pvt Ltd, Plot No 50,3 <sup>rd</sup> Road,1 <sup>st</sup> phase ,KIADB Ind Area, Jigani ,Bangalore 560105	
<b>Shelf Life</b>	3 years	




Fig. 1.DPPH free radical scavenging activity and IC<sub>50</sub> of Storg-B and Standard BHT

### 3.2 Effect of Storg B on Humoral immune response

The hemagglutination test was used to assess humoral immune response. Cyclophosphamide, an immunosuppressant agent significantly decreases the HA titer value compared to control group. After administration of test doses PHF (Storg B) 100 mg/kg



Normal	6.55	5.32	23.72	18.47	155.3	98.26	36.72
Cyclophosphamide 20mg/kg	3.35	2.09	17.47	22.12	58.52	46.23	21.00
Cyclo + 50mg/kg Levamisole	5.20	4.47	21.6	17.53	112.32	78.35	30.24
Cyclo + Storg-B 50mg/kg	4.97	2.97	19.53	23.53	97.06	69.88	28.03
Cyclo + Storg-B 100mg/kg	5.18	4.51	22.31	18.10	115.56	81.63	29.36

\*UTB= Untreated Blood, NFTB = Nylon Fibre Treated blood

### 3.4 Effect of Storg B on hematological parameters

The harmful effects of cyclophosphamide on the animals disrupted their normal blood component levels. In comparison to the positive control group, the treatment group PHF Storg-B 100 mg/kg was effective in improving the normal level of blood parameters as shown in Table No.3. The outcomes of the study suggest that Storg- B has beneficial hematological effects.

**Table 3. Effect of Storg-B on Hematological parameters**

Groups	WBC (10 <sup>9</sup> /L)	RBC (10 <sup>12</sup> /L)	Hb (g/dL)	PLT (10 <sup>9</sup> /L)	DLC		
					Lympho cytes %	Monocy tes %	Granulocyt es %
Normal	6.55±0.1	5.3±0.13	12.6±0.32	15.9±1.2	62.72	16.48	25.25
Cyclophosphamide 20 mg/kg	3.35±0.3	4.1±0.43	12.42±0.65	9.8±1.3	43.98	10.28	30.93
Cyclo+Levamisole 50 mg/kg	5.20±1.7	4.8±0.14	12.54±0.17	12±1.02	56.33	15.47	24.8
Cyclo + Storg-B 50 mg/kg	4.97±0.88	4.74±0.12	12.56±0.32	11.8±1.36	54.88	14.88	26.71
Cyclo+ Storg-B 100 mg/kg	5.18±1.27	4.82±0.18	12.62±0.50	12.4±1.02	57.03	16.13	28.49

### 3.5 Effect of Storg B on Bioavailability

LC-MS/MS is the technique that is employed to quantify the Bioavailability study of Storg B. Single dosage of Storg B 200 mg/kg was orally administered to the SD rat. The main pharmacokinetic parameters  $C_{max}$ ,  $t_{max}$ , AUC and  $t_{1/2}$  are calculated and the values are shown in Table No.4 and Fig.3. The peak concentration ( $C_{max}$ ) and time to reach the peak concentration ( $T_{max}$ ) of B vitamin in the plasma were obtained from the concentration-time curve. In the graph the plasma concentration versus time curve represents the total amount of the drug reaching the circulatory system. PHF Storg B showed the greater potency.

**Table 4. Bioavailability results of Storg-B**

Parameters	$C_{max}$ (ng/mL)	$T_{max}$	$t_{1/2}$	AUC (ng h/mL)
Vitamin B1	28.74	1 hour	5 hour 15 minutes	269.3

Vitamin B2	23.44	1 hour	13 hour	318.3
Vitamin B3	148.59	1 hour	7 hour	1430
Vitamin B5	87.5	1 hour	5 hour 30 min	759.4
Vitamin B6	23.4	30 min	4 hour 30 min	200.4
Vitamin B9	31.8	30 min	3 hour 45 min	216.8




Fig. 3. Plasma concentration of Storg B time profile curve

#### 4. CONCLUSION

PHFs are becoming widely popular due to their effectiveness in treating a variety of diseases and infectious disorders, particularly via altering the host defense mechanism [12]. PHF has been used all around the earth due to its medicinal and therapeutic application. In comparison to a single herbal formulation, PHF offers more benefits [13]. Storg B is a standardized polyherbal supplement made from the extracts of the medicinal herbs *Psidium guajava*, *Mangifera indica*, and *Citrus limon* that contains enriched natural B vitamins and co-nutrients such as carotenoids and polyphenols. B-vitamin-mediated immunological regulation is specific to different immune cells and immune responses: that is, different B vitamins are required for different immune responses [14]. The B vitamins which are present in PHF Storg-B are identified as follows: thiamine (B<sub>1</sub>), riboflavin (B<sub>2</sub>), niacin (B<sub>3</sub>), pantothenic acid (B<sub>5</sub>), pyridoxine (B<sub>6</sub>), and folate (B<sub>9</sub>). B vitamins play essential role in

catabolic and anabolic metabolism and act as coenzymes in numerous enzymatic processes that support each aspect of biological functioning, including major functions within the brain and nerve system. Any B vitamin deficiency may adversely impact mitochondrial metabolism of amino acids, glucose, and fatty acids through the citric acid cycle and electron transport chain [15]. By playing a variety of roles in the immune system, B vitamins serve as essentials in the normal body's defense mechanism. From the recent investigations it has been recognized that vitamin B is enhancing a significant function in boosting the immune responses [16].

In this study, herbal immunomodulatory blend was prepared by synergized formulation of herbal extracts, scientifically quantified for their bioactive metabolites, tested *in vitro* and *in vivo* evaluated in CP-induced immunosuppression model in SD rat.

Antioxidants are considered to have an impact on DPPH due to their ability to donate hydrogen [17]. Radical scavenging activity is very essential in order to prevent free radicals from playing a harmful role in a variety of disorders, including cancer. The polyphenols primarily act as free radical scavengers, the majority of plant polyphenols have strong antioxidant qualities [18]. Since these plant polyphenols are co-nutrients in our PHF, they may be contributing to Storg B's antioxidant activity, which is demonstrated by its 68.358µg/ml free radical antioxidant scavenging activity in comparison to the standard BHT. Immunomodulatory agents of plant origin improve the immune responsiveness of an organism against a pathogen by nonspecifically activating the immune system [19]. The humoral immune system includes the interaction of B cells with the antigens, which replicate and differentiate into plasma cells that secrete antibodies [20,21]. In the study PHF treated group produced increased Primary and secondary antibody titres against SRBCs. The antibodies produced were intended to bind to the antigen and neutralized or facilitate recognition by phagocytes this leads to ultimate elimination of the antigen [22,23]. Polyherbal extract Storg B enhanced the humoral immune response to SRBCs in rats, as evidenced by an increase in antibody titer. This shows that the T and B lymphocyte subsets became more responsive, which are vital for antibody formation [24].

An animal's hematological characteristics can be employed to observe and quantify one of the immune responses. The total leukocyte count for both the control group and the groups that received PHF Storg-B medication were determined based on these characteristics. When non-self-components invade the body, blood cell reacts first. Any immunological material with immunomodulatory properties would first cause a shift in the leukocyte count [25]. The study group V, which was administered the testing medication, exhibited the highest leukocyte count of  $5.18 \times 10^3/\text{mm}^3$ , indicating the early activation of blood cells in order to develop a potent immune response. The findings indicate that higher concentrations of the medication PHF Storg-B can stimulate a significant immunological response. In the neutrophil adhesion test, adherence to nylon fibres was increased. This may be due to the up regulation of  $\beta$ -2 integrins and also to a decreased corticosterone level. The results are further supported with the highest percentage of neutrophils circulated among the group. Finally Result revealed that PHF Storg B has the potential humoral immunity and it can be a potential therapeutic candidate for several immune suppressed conditions.

Any molecule's bioavailability (BA) is determined by the interaction of its pharmacodynamics (PDs) and pharmacokinetics (PKs) factors [26]. PKs (ADME) and PDs are two aspects that influence the therapeutic efficacy of drugs. The water soluble vitamins biotin, folate, niacin, pantothenic acid, pyridoxine, riboflavin, and thiamin are absorbed through the intestinal tract by particular carrier-mediated pathways [27]. The estimations of  $C_{\text{max}}$  and area under the curve (AUC) parameters were included in the pharmacokinetic dataset. The time of peak concentration was not significantly different but it was significantly different in terms of AUC. The area under the curve is the most important parameter to evaluate the bioavailability of

the drug in the dosage form because it indicates the extent of absorption [28]. From the results the AUC value of the B vitamins is followed as B3 > B5 > B2 > B1 > B9 > B6. Where B3 (1430 ng h/mL) entered the systemic circulation for distribution to target cells and tissues maybe with the mild alteration compared to other B vitamins present in the PHF (Storg B). Vitamin B1 (269.3 ng h/mL) B9 (216.8 ng h/mL) and B6 (200.4 ng h/mL) are acidity and temperature sensitive. Since the pH of small intestine is greater than that of stomach phase, reductions in the pH sensitive vitamins may generally takes place [29]. Hence they are having slightly less absorption rate. While Vitamin B5 (759.4 ng h/mL) and B2 (318.3 ng h/mL) are showing moderate absorption rate. Since there aren't many bioavailability studies currently available for B vitamins, our study adds to the growing literature on Vitamin B by providing much-needed data on the treatment of these biological compounds and filling this knowledge gap.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by the Bharathi College of Pharmacy ethics committee(BCP/IAEC/06/2022).

## REFERENCES

1. Bhargava S, Bhargava P, Saraf S, Pandey R, Shukla SS. Evaluation of antipyretic activity of Sudarshan churna: An ayurvedic formulation. J Res Educ Indian Med. 2008.
2. Jantan I, Haque MA, Ilangkovan M, Arshad L. An insight into the modulatory effects and mechanisms of action of *Phyllanthus* species and their bioactive metabolites on the immune system. Front Pharmacol. 2019;10:1-19. DOI: [10.3389/fphar.2019.00878](https://doi.org/10.3389/fphar.2019.00878); PMID: [31440162](https://pubmed.ncbi.nlm.nih.gov/31440162/)
3. Makare N, Bodhankar S, Rangari V. Immunomodulatory activity of alcoholic extract of *Mangifera indica* L in mice. J Ethnopharmacol. 2001;78(2-3):133-7. DOI: [10.1016/s0378-8741\(01\)00326-9](https://doi.org/10.1016/s0378-8741(01)00326-9); PMID: 11694357
4. Parveen A, Parveen B, Parveen R, Ahmad S. Challenges and guidelines for clinical trial of herbal drugs. J Pharm Bioallied Sci. 2015;7(4):329-3. DOI: [10.4103/0975-7406.168035](https://doi.org/10.4103/0975-7406.168035); PMID: [26681895](https://pubmed.ncbi.nlm.nih.gov/26681895/)
5. Chorghade MS. Drug Discovery and Development: Drug Development. 2nd volume. John Wiley & Sons. New Jersey 2007; pp 36.
6. Parasuraman S, Thing GS, Dhanaraj SA. Polyherbal formulation: Concept of ayurveda. Pharmacogn Rev. 2014;8(16):73-80. DOI: [10.4103/0973-7847.134229](https://doi.org/10.4103/0973-7847.134229); PMID: 25125878
7. Young IS, Woodside JV. Antioxidants in health and disease. J Clin Pathol. 2001;54:176-86. DOI: [10.1136/jcp.54.3.176](https://doi.org/10.1136/jcp.54.3.176); PMID: 11253127
8. Pourmorad F, Hosseinimehr SJ, Shahabimajd N. Antioxidant activity, phenols, flavanoid contents of selected Iranian medicinal plants. S Afr J Biotechnol. 2006;5:1142-5.

9. Sharma P. RasayanaChikitsa in Charaka Samhita Chikitasasthana. Chaukhamba Orientalia Publication. Varanasi, India 1983; pp 54.
10. Blois MS. Antioxidant determinations by the use of a stable free radical. Nature. 1958; 181:1199–1200.  
[DOI: 10.1038/1811199a0](https://doi.org/10.1038/1811199a0)
11. Abebe D, Karim A, Bitew H, Periasamy G. *In-vivo* evaluation of immunomodulatory activity of crude extract and solvent fractions of *Cyphostemmaadenocaula* (Steud. ex A.Rich). Heliyon. 2022;8(12):e12377.  
[DOI: 10.1016/j.heliyon.2022.e12377](https://doi.org/10.1016/j.heliyon.2022.e12377); PMID: [36590494](https://pubmed.ncbi.nlm.nih.gov/36590494/)
12. Rajendra BM, Krishna RRV, Annapurna A, Ravi KBD. Immunostimulant profile of a polyherbal formulation RV08. Indian J Pharmacol. 2001;33(6):454-5.
13. Karole S, Shrivastava S, Thomas S, Soni B, Khan S, et al. Polyherbal formulation concept for synergic action: A review. J drug delivther. 2019; 9(1-s):453-66.  
[DOI: 10.22270/jddt.v9i1-s.2339](https://doi.org/10.22270/jddt.v9i1-s.2339)
14. Yoshii K, Hosomi K, Sawane K, Kunisawa J. Metabolism of dietary and microbial vitamin B family in the regulation of host immunity. Front Nutr. 2019;6:48.  
[DOI: 10.3389/fnut.2019.00048](https://doi.org/10.3389/fnut.2019.00048); PMID: 31058161
15. Kennedy DO. B Vitamins and the Brain: Mechanisms, Dose and Efficacy-A Review. Nutrients. 2016;8(2):68.  
[DOI: 10.3390/nu8020068](https://doi.org/10.3390/nu8020068); PMID: [26828517](https://pubmed.ncbi.nlm.nih.gov/26828517/)
16. Mitra S, Paul S, Roy S, Sutradhar H, Bin Emran T, Nainu F, et al. Exploring the immune-boosting functions of vitamins and minerals as nutritional food bioactive compounds: a comprehensive review. Molecules. 2022;27(2):555.  
[DOI: 10.3390/molecules27020555](https://doi.org/10.3390/molecules27020555); PMID: 35056870
17. Baumann J, Wurm G, von Bruchhausen F. Prostaglandin synthetase inhibition by flavonoids and phenolic compounds in relation to their O<sub>2</sub>--scavenging properties (author's transl). Arch Pharm (Weinheim). 1980;313(4):330-37.  
[DOI: 10.1002/ardp.19803130409](https://doi.org/10.1002/ardp.19803130409); PMID: 6772138
18. Dewick PM, Medicinal Natural Products; John Wiley & Sons: New York, USA, 1997
19. Rema R, Sutirtha R. Study of the immunomodulatory activity of a polyherbal drug. Pharmacologyonline. 2008;3:336-45.
20. Ose EE, Muenster OA: A method for evaluation of vaccines containing Pasteurella multocida. Am J Vet Res. 1968;29:1863–6.  
PMID: 5673520.
21. Niphade SR, Asad M, Chandrakala GK, Toppo E, Deshmukh P. Immunomodulatory activity of *Cinnamomum zeylanicum* bark, Pharm Biol. 2009;47(12):1168-73.  
[DOI: 10.3109/13880200903019234](https://doi.org/10.3109/13880200903019234)
22. Gokhale AB, Damre AS, Saraf MN. Investigations into the immunomodulatory activity of *Argyrea speciosa*. J Ethnopharmacol. 2003;84:109-114.  
[DOI: 10.1016/s0378-8741\(02\)00168-x](https://doi.org/10.1016/s0378-8741(02)00168-x); PMID: 12499085
23. Fulzele SV, Satturwar PM, Joshi SB, Dorle AK. Study of the immunomodulatory activity of Haridradighrita in rats. Indian J Pharmacol. 2003;35:51–4.
24. Benacerraf B, A hypothesis to relate the specificity of T lymphocytes and the activity of I region-specific Ir genes in macrophages and B lymphocytes, J Immunol.1978;120(6):1809-12.
25. Rama Bhat P. A Review on Immuno modulatory effects of plant extracts. Virol Immunol J. 2018;2(6).  
[DOI: 10.23880/VIJ-16000167](https://doi.org/10.23880/VIJ-16000167)
26. Mittal S, Sawarkar S, Doshi G, Pimple P, Shah J, Turabali B. Chapter twenty one- Pharmacokinetics and bioavailability of nutraceuticals, Industrial Application of Functional Foods, Ingredients and Nutraceuticals. 2023;725-83.  
[DOI: 10.1016/B978-0-12-824312-1.00021-2](https://doi.org/10.1016/B978-0-12-824312-1.00021-2)

27. Said HM. Intestinal absorption of water-soluble vitamins in health and disease. *Biochem J.* 2011;437(3):357–72.  
DOI: [10.1042/BJ20110326](https://doi.org/10.1042/BJ20110326); PMID: 21749321
28. Nepal S, Banskota S, Marasini N, Gupta B, Lohani SP, Basnet S, et al. Bioequivalence study of two brands of Phenytoin sodium 100mg formulations in healthy adult male rabbits. *Int J Pharm Sci Res.* 2013;4(1); 276-81.  
DOI: [10.13040/IJPSR.0975-8232.4\(1\).276-81](https://doi.org/10.13040/IJPSR.0975-8232.4(1).276-81)
29. Yaman M, Çatak J, Uğur H, Gürbüz M, Belli İ, Tanyildiz SN, et al. The bioaccessibility of water-soluble vitamins: A review, *Trends Food Sci Technol.* 2021;109:552-63.  
DOI: [10.1016/j.tifs.2021.01.056](https://doi.org/10.1016/j.tifs.2021.01.056)

UNDER PEER REVIEW