

# Short Term Effect of Vermicompost on Soil Chemical Properties Under Maize (*Zea mays* L.) field in Northern Ethiopia

## *Abstract*

**Aim:**The application of vermicompost (VC) to improve soil chemical properties in area that have low soil fertility status is crucial. Thus, the present study was initiated to determine impact of vermicompost on selected soil physicochemical properties.

**Study Design:** The experiment had 7 treatments arranged in a Randomized Complete Block Design (RCBD) with five replications.

**Place and Duration of the study:** Hence, a field experiment was carried out in 2018 main cropping season at farmer's field at Adiha village in Kolla Tembien district, Tigray, Ethiopia.

**Methodology:**The treatments were 5 levels of VC (0, 2.5, 5, 7.5 and 10 t ha<sup>-1</sup>), the recommended rate of N and P (46 N and 46 P<sub>2</sub>O<sub>5</sub> kg ha<sup>-1</sup>), compost at rate of 10 t ha<sup>-1</sup> and control. Surface soil samples were collected before planting and after maize harvesting at a depth of 0-30 cm to analyze selected soil chemical properties such as pH, extractable electric conductivity (ECe), cation exchange capacity (CEC), total nitrogen (TN), available P and exchangeable bases (K, Mg, Ca, and Na). Likewise, vermicompost and compost nutrient contents were also analyzed to know their nutrient composition.

**Results:**Soil analysis results before sowing revealed that most of the soil chemical properties were rated as low. However, vermicompost application significantly ( $P \leq 0.05$ ) improved the soil chemical properties. The studied soil chemical properties such as organic carbon (OC)(1.33%), TN (0.13%), CEC(17.2 Cmol(+)kg<sup>-1</sup>), and available P(6 .38 mg kg<sup>-1</sup>) showed an improvement at VC 10 t ha<sup>-1</sup>rate as compared to the control which was respectively 0.65%, 0.04%, 7.4 Cmol(+)kg<sup>-1</sup> and 2.45 mg kg<sup>-1</sup>.

**Conclusion:** A significant ( $P \leq 0.05$ ) increase in OC, TN and available P was observed with increased vermicompost rate.

**Key words:** CEC, Compost, Maize, Organic carbon, and RCBD

## INTRODUCTION

Land degradation, including soil erosion, deforestation and desertification, represents another major obstacle to agricultural production. Unsustainable agricultural practices, excessive grazing and inadequate land management lead to the degradation of fertile soils, reducing their productivity and agricultural potential [1]. Furthermore, deforestation to expand agriculture further exacerbates the loss of valuable land resources [2].

The soils of most African countries are regularly degraded by erosion. Loss of about 50 tons of soil per hectare per year, corresponding to 20 billion tons of N, 2 billion tons of P and 41 billion tons of K per year [3]. Soil fertility in sub-Saharan Africa is decreasing due to land degradation due to population growth, lack of attention to integrated soil fertility management (ISFM), inadequate plant nutrients, poor agricultural practices and long-term drought, and irregular rainfall patterns which are also leading to a decline in soil fertility in southern Africa the Sahara with low yields [4-7].

“In Ethiopia, agricultural cultivation involves the removal of nutrients (N, P, K, etc.) from the soil through agricultural products (food, fiber and wood) and crop residues. Consequently, nutrient deficiency leads to a decline in soil fertility when the supply of inorganic or organic nutrients is insufficient” [8–10]. “Likewise, declining soil fertility is a major constraint to yield limitation of smallholder farmers in Ethiopia due to shorter fallow periods, lower levels and the imbalanced use of organic and inorganic fertilizers, removal of crop residues and lack of adequate soil conservation practices” [11 – 14].

“The soil-level bottleneck such as organic matter depletion, soil biota depletion, nutrient depletion, limited biomass cover, soil compaction, soil erosion, soil salinity/sodicity, acidity, water logging, low moisture availability and physical soil degradation have a negative impact on physical, chemical and biological conditions of the soil and its fertility are common in Ethiopia” [15, 16]. “Possible measures are therefore soil fertility management, such as the application of organic and mineral fertilizers, crop rotation, catch crops, crop residue management, and physical protection of soil and water” [17].

“Vermicompost, made from livestock manure, potato waste, sugarcane residue and sunflower residue, resulted in significantly higher plant yields. Therefore, the nutrient content of vermicompost was always higher than that of normal compost” [18]. Many studies have shown

that the use of manure provides yields comparable or even superior to those of inorganic fertilizers [19].

“Application of different organic fertilizers favorably influences the soil physical, chemical and biological environment such as available nitrogen over the inorganic alone applied plots. Among different organic N sources, the application 75 per cent vermicompost was found superior in improving soil health over the treatments with inorganic fertilizers”[20].

“The organic carbon content of vermicompost (15.2%) was much lower than that of FYM (37.4%), while the N content of vermicompost (1.40%) was much higher than that of FYM (0.9%)” [21]. Furthermore, [22] reported that “vermicompost prepared from mustard residues and sugarcane waste using *Eiseniafetida* resulted in a significant increase in mineral nitrogen content and microbial activity”. “The use of vermicompost alone or with chemical fertilizers reduces bulk density, increases soil porosity and water holding capacity, but increases soil pH and organic carbon content” [23]. “Likewise, higher levels of available soil N were observed in plots that received 5 t/ha of vermicompost. It also improved the pH of soil towards neutrality”[24]. “The use of organic fertilizers such as VC and FYM progressively reduces the bulk density of the soil” [25, 26].

After applying vermicompost in two consecutive growing seasons, there was a significant decrease in soil density and a significant increase in pH and total organic carbon. Taken together, these changes in soil properties improve air and water availability, thereby promoting seedling emergence and root growth [27]. Thus, Mekelle University, with the help of AgriFoSe2030, recently introduced VC as an option for soil fertility management in Kolla Tembien District, located in the central part of Tigray Regional State. However, no study has been conducted so far on the influence of VC on the chemical properties of soils at the field level, especially in the study area. Therefore, this study was initiated to determine the effects of VC on soil properties under maize field.

## 2. Materials and Methods

### 2.1. Description of the Study Area

The experimental study was conducted at Adiha village located in Kolla Tembein district, central zone of Tigray, Ethiopia (Figure 1). The village is geographically located at 30 kilometers North east of Abyi - Adi town, and 95 kilometers north of Mekelle city (capital of Tigray Region) (Office of Agriculture and Rural Development (OoARD) [28], at a GPS location of 13°45'0.4"N and 38° 6' 5.8"E.




Figure 1. Location map of the study area

Most of the population lives in rural areas and earns their living through agricultural production. The main cereals grown in the study area are maize (*Zea mays L.*), sorghum

(*Sorghum bicolor L.*), teff (*Eragrostis tef (Zucc.)*) and millet (*Eleusine coracana L.*). Likewise, farmers in the district are involved in rearing animals such as oxen, goats, sheep, donkeys, mules, camels, bee colonies and others [28]. The geology of the study area is Adigrat sandstone [29]. Soil type of the experimental trial has a Reference Soil Group (RSG) of Petric Calcisols [30].




Figure 2: Soil type of Kolla Tembien District [30].

## 2.2. Experimental Design

The experimental study has seven (7) treatments with five replications namely: five levels of vermicompost (0, 2.5 t ha<sup>-1</sup>, 5 t ha<sup>-1</sup>, 7.5 t ha<sup>-1</sup> and 10 t ha<sup>-1</sup>), recommended rate of NP (100kg ha<sup>-1</sup> urea and 100 kg ha<sup>-1</sup> TSP) used by the farmers in the study area and compost at rate of 10 t ha<sup>-1</sup>. The seven treatments were laid out in randomized complete block design. The total area used for

the trial was 35 m×25.75 m (901.25 m<sup>2</sup>), whereas the plot size was 3.75 m × 4 m (15 m<sup>2</sup>). The spacing between plots and blocks were 1 m and 1.5 m, respectively.

### 2.3. Soil, Compost and Vermicompost Sampling and Analysis

Before planting, a composite soil sample (disturbed, for analysis of chemical properties) was collected from the entire study area to a depth of 0–30 cm. After harvest, 35 composite (disturbed) soil samples were collected from each subplot and pooled with the respective plots. The composite soil samples were air-dried, ground, passed through a 2 mm sieve, and prepared for analysis of soil chemical properties. One composite vermi-compost and compost produced by farmer's using locally available organic waste materials (cow dung, straw of crops, food waste like Lettuce and cabbage) were also taken for the same purpose.

Table 1. Soil parameter, method analysis and reference used in the experiment

Soil parameters	Method of analysis	Reference
pH	1:2.5 soil to water suspension	[31]
EC	1:2.5 soil to water suspension	[32]
Organic carbon	Wet oxidation method	[33]
Total nitrogen	Kjedahl method	[34]
Available P	Olsen method	[35]
CEC	Ammonium Acetate method	[36]
Exchangeable K and Na	Flame photometer	[37]
Exchangeable Ca and Mg	Titration method	[38]

Table 2. Initial chemical properties of vermicompost, compost and soil

Parameters	Rating			Reference
	VC	C	Soil	
pH	7.31	7.1	7.32	Slightly acidic to neutral [39,40]

ECe(ds m <sup>-1</sup> )		1.28	1.33	1.7	Low salinity	[41]
Organic Carbon (%)		23.01	22.14	0.65	Low	[40]
Total N (%)		1.64	1.04	0.041	Low	[42]
Total P (mg kg <sup>-1</sup> )		0.83	0.54	-	-	[35]
Total K (mg kg <sup>-1</sup> )		1.08	1.24	-	-	
CEC (cmol(+) kg <sup>-1</sup> )		-	-	7.61	Low	[43]
Ava. P(mg kg <sup>-1</sup> )		-	-	2.45	Low	[35]
Exchangeable Ca (cmol(+) kg <sup>-1</sup> )		-	-	2.8	Low	[44]
Exchangeable Mg (cmol(+) kg <sup>-1</sup> )		-	-	1.0	Low	[44]
Exchangeable K (cmol(+) kg <sup>-1</sup> )		-	-	0.12	Low	[44]
Exchangeable Na (cmol(+) kg <sup>-1</sup> )		-	-	0.17	Low	[44]

## 2.4. Data Analysis

The collected after harvesting soil data were subjected to statistical analysis like analysis of variance. Analysis of variance (ANOVA) was performed using SAS version 9.1 [45]. Significant difference between and among treatments means were assessed using the least significant difference (LSD) at 0.05 level of probability [46].

## 3. RESULTS AND DISCUSSION

### 3.1. Effect of Vermicompost on Selected Soil Chemical properties

#### 3.1.1. Soil reaction

The pH values were not significantly affected by VC dose and fertilizer type ( $P > 0.05$ ) (Table 3). However, the result varied numerically between treatments, namely the maximum pH value (7.63) was obtained from plots treated with a high dose of vermicompost, while the minimum pH (7.3) was obtained from plots treated with 10 t ha<sup>-1</sup> compost (Table 3). This could

be due to the addition of basic cations to the soil, which hinders the action of exchangeable acids at the site of soil exchange [26, 47].

Table 3. Effect of vermicompost on selected soil physico-chemical properties

Treatment	pH	E <sub>Ce</sub> (mmhos/cm)	OC (%)	TN (%)
Control	7.40	1.71	0.66 <sup>c</sup>	0.042 <sup>ed</sup>
46 N kg ha <sup>-1</sup> + 46 P <sub>2</sub> O <sub>5</sub> kg ha <sup>-1</sup>	7.35	1.72	0.63 <sup>c</sup>	0.038 <sup>e</sup>
10 t ha <sup>-1</sup> Compost	7.32	1.69	1.22 <sup>ab</sup>	0.072 <sup>bc</sup>
2.5 t ha <sup>-1</sup> VC	7.38	1.69	0.99 <sup>b</sup>	0.06 <sup>cd</sup>
5 t ha <sup>-1</sup> VC	7.44	1.67	1.01 <sup>b</sup>	0.08 <sup>bc</sup>
7.5 t ha <sup>-1</sup> VC	7.57	1.67	1.27 <sup>a</sup>	0.09 <sup>b</sup>
10 t ha <sup>-1</sup> VC	7.63	1.66	1.33 <sup>a</sup>	0.13 <sup>a</sup>
Mean	7.44	1.69	1.01	0.07
LSD (P≤0.05)	NS	NS	0.25	0.02
CV	3.99	11.57	12.01	14.71

Means followed by the same letters are not significantly different ( $P \leq 0.05$ ) according to Tukey Test; B.D= bulk density, SMC = Soil moisture content, pH= Power of hydrogen, E<sub>Ce</sub> = Extractable electrical conductivity, OC = Organic carbon, TN = Total nitrogen, N = nitrogen; P<sub>2</sub>O<sub>5</sub> = Di phosphate PentaOxide t= tone; ha= hectare; CV= Coefficient of variation;

### 3.1.2. Extractable electrical conductivity

E<sub>Ce</sub> values were not significantly ( $P > 0.05$ ) affected by the rates of VC and fertilizer types (Table 3). However, they had numerical difference within the treatments i.e. the maximum E<sub>Ce</sub> value (1.7 ds m<sup>-1</sup>) was obtained from plots treated with recommended NP and no fertilizer; while the minimum E<sub>Ce</sub> value (1.6ds m<sup>-1</sup>) was from the plot treated by 10 t ha<sup>-1</sup> VC (Table 3). This is in accordance with the findings of [48] which revealed that application of compost and vermicompost could lower E<sub>Ce</sub> than inorganic fertilizer. “This is possibly due to unlike chemical fertilizers, organic fertilizers release nutrients more gradually as a result of degradation process and their cheated effect”[49].

### 3.1.3. Organic carbon

Soil organic carbon (SOC) content was significantly ( $P \leq 0.05$ ) affected by VC doses and fertilizer type (Table 3). The SOC value increased as the VC rate increased. Therefore, the highest SOC content (1.33%) was obtained with the highest VC dose and with the lowest dose (0.66%) came from the recommended NP-treated plots (Table 3). This is consistent with the results of [50], who reported that the use of vermicompost at a dose of 10 t ha<sup>-1</sup> significantly increased the organic carbon content of the soil. However, the high dose (10 t ha<sup>-1</sup>) at a dose of 7.5 t ha<sup>-1</sup> VC and composting at a dose of 10 t ha<sup>-1</sup> were not significant.

### 3.1.4. Total N

Total N values were significantly ( $P \leq 0.05$ ) affected by the rates of VC (Table 3). Total N increased as the rate of VC increased. Hence, the maximum total N content of the soil (0.13%) was obtained from the high VC dose, whereas the lowest (0.042%) was from the recommended NP and unfertilized plots. This is in orthodoxy with the findings of [51-52] that revealed application of vermicompost to soils resulted in higher total N concentration in the soil and plant compared to the control. The increment in total N with addition of organic amendments (VC) to soil is caused by high release of total N content of vermicompost in the high dose of VC than control [53].

### 3.1.5. Cation Exchange Capacity (CEC)

The CEC was significantly ( $P \leq 0.05$ ) influenced by the rates of VC (Table 4). CEC showed a slight increase as the rate of VC increased. Thus, the highest CEC of the soil (17.2 cmol (+) kg<sup>-1</sup>) was obtained from the plots treated with high dose of Vermicompost, whereas the lowest (7.6 cmol (+) kg<sup>-1</sup>) was from unfertilized plots. This might have resulted from the humic acids contained in both vermicompost and compost which they bind cations as they contain carboxylic acid groups, which can in turn bind positively, charged multivalent ions [54].

Table 4. Effect of vermicompost on selected soil chemical properties.

Treatment	Exchangeable base (cmol(+) kg <sup>-1</sup> )					
	CEC (cmol(+) kg <sup>-1</sup> )	Ca	Mg	Na	K	AvaP (mg kg <sup>-1</sup> )
Control	7.60 <sup>d</sup>	2.60 <sup>e</sup>	1.00 <sup>d</sup>	0.14 <sup>f</sup>	0.10 <sup>e</sup>	2.43 <sup>e</sup>
46 N kg ha <sup>-1</sup> + 46 P <sub>2</sub> O <sub>5</sub> kg ha <sup>-1</sup>	7.90 <sup>d</sup>	3.16 <sup>d</sup>	1.10 <sup>d</sup>	0.21 <sup>e</sup>	0.21 <sup>d</sup>	2.46 <sup>e</sup>
10 t ha <sup>-1</sup> C	11.52 <sup>c</sup>	4.40 <sup>c</sup>	1.80 <sup>c</sup>	0.33 <sup>d</sup>	0.31 <sup>c</sup>	3.28 <sup>d</sup>
2.5 t ha <sup>-1</sup> VC	10.50 <sup>c</sup>	4.08 <sup>c</sup>	1.28 <sup>d</sup>	0.22 <sup>e</sup>	0.24 <sup>d</sup>	3.08 <sup>d</sup>
5 t ha <sup>-1</sup> VC	13.50 <sup>b</sup>	5.80 <sup>b</sup>	2.78 <sup>b</sup>	0.41 <sup>c</sup>	0.41 <sup>b</sup>	3.92 <sup>c</sup>
7.5 t ha <sup>-1</sup> VC	16.60 <sup>a</sup>	7.58 <sup>a</sup>	3.24 <sup>a</sup>	0.45 <sup>b</sup>	0.42 <sup>ab</sup>	5.61 <sup>b</sup>
10 t ha <sup>-1</sup> VC	17.20 <sup>a</sup>	8.10 <sup>a</sup>	3.46 <sup>a</sup>	0.47 <sup>a</sup>	0.46 <sup>a</sup>	6.38 <sup>a</sup>
Mean	12.12	5.11	2.10	0.32	0.31	3.88
LSD (P≤0.05)	1.97	0.55	0.36	0.02	0.04	0.48
CV	8.09	5.33	8.47	2.33	6.89	6.17

Means followed by the same letters are not significantly different ( $P \leq 0.05$ ) according to Tukey Test; Av. P = Available phosphorus, CEC = Cation exchange capacity, Ca= calcium, Mg= magnesium, K= potassium, Na= sodium, P<sub>2</sub>O<sub>5</sub> = Di phosphate PentaOxide t= tone; ha= hectare; CV= Coefficient of variation;

### 3.1.6. Exchangeable bases

Exchangeable bases showed significant difference ( $P \leq 0.05$ ) upon application of different rates of VC (Table 4). The highest (8.1 cmol (+) kg<sup>-1</sup>) exchangeable Ca was obtained from plots treated by 10 t ha<sup>-1</sup> VC, whereas the lowest (2.6 cmol (+) kg<sup>-1</sup>) was from unfertilized plots. An increase in exchangeable Ca content of soil might have happened due to organic amendments improve soil cation exchange capacity (CEC) through humus formation and increased concentrations of Ca. Similarly, the highest (3.5 cmol (+) kg<sup>-1</sup>) exchangeable Mg was obtained from plots which

received  $10 \text{ t ha}^{-1}$  VC, while the lowest ( $1.0 \text{ cmol (+) kg}^{-1}$ ) was from control. This could be due to storage and availability of Mg closely related to CEC [51, 55].

The maximum value of exchangeable Na ( $0.47 \text{ cmol (+) kg}^{-1}$ ) and K ( $0.46 \text{ cmol (+) kg}^{-1}$ ) were obtained from plots which received the highest dose of VC; whereas, minimum values ( $0.14 \text{ cmol (+) kg}^{-1}$ ) and ( $0.10 \text{ cmol (+) kg}^{-1}$ ) for the respective nutrients were from unfertilized plots. This might occur when organic material like VC are added to soils at high dose so that it improves organic matter content; they also supply soils with nutrients including exchangeable K and Na [55-57]. In harmony to this result, [50] revealed that application of VC at high dose significantly increased exchangeable base of post-harvest soil than the lowest rate and nil treatment.

### **3.1.7. Soil available P**

Soil available P of the post-harvest soil was significantly ( $P \leq 0.05$ ) influenced by the rates of VC (Table 4). Available P showed a slight increment as the rate of VC increased. The highest ( $6.38 \text{ mg kg}^{-1}$ ) available P was obtained from plots treated with  $10 \text{ t ha}^{-1}$  VC, while the lowest ( $2.43 \text{ mg kg}^{-1}$ ) was from control/check. This corresponds with the findings of [51-52] who revealed that application of VC at high dose significantly increased available P. This might have happened when organic material like VC add to soils at high dose can improve organic matter status, there by supply soil with nutrient of available P [56-57].

## **4. CONCLUSIONS**

The findings from this study showed that the initial soil reaction ranged from slightly acidic to neutral; E<sub>c</sub> value is rated as low soil salinity. Most of the chemical properties like OC content, TN, av.P, CEC and exchangeable bases were also rated as low. However, application of vermicompost at different rates significantly affected most of the studied soil parameters. Hence, VC improves the degraded soil and that have initially low soil nutrient through improving the structure of the soil, porosity of soil and improving soil nutrient. Vermicompost had also significantly affected soil chemical properties. A significant increase in OC, TN and available P was observed with increased vermicompost rate. Thus, the highest CEC, OC, TN and available P were recorded from plots which received the highest VC dose; while the lowest values were recorded from unfertilized plots and recommended NP.

### **Data Availability**

Data used to support the findings of this study are available from the corresponding author upon reasonable request of the publisher or other stakeholders.

## REFERENCES

1. UNCCD. (2018). The Global Land Outlook. United Nations Convention to Combat Desertification. Retrieved from <http://www.unccd.int/glo>
2. FAO. (2020). The State of the World's Forests 2020. Food and Agriculture Organization. Retrieved from <http://www.fao.org/state-of-forests/en/>
3. Bationo, A. (2009). Soil Fertility Paradigm Shift through Collective Action. <http://knowledge.cta.int>.
4. Cakir, R. (2004). Effect of water stress at different development stages on vegetative and reproductive growth of corn. *Field Crops Research*. 89:1-16. Doi.org/10.1016/j.fcr.2004.01.005.
5. Mando, A., Ouattara, B., Somado, A. E., Wwopereis, M. C. S., Stoosnijder, L. and Breman, H. (2005). Long term effects of fallow, tillage and manure application on soil organic matter and nitrogen fractions and on sorghum yield under Sudano-Sahelian conditions. *Soil Use Manage*, 21: 25-31.
6. Danso G, Drechsel P, Fialor S, Giordano M (2006). Estimating the demand for municipal waste compost via farmers' willingness-to-pay in Ghana. *Waste Manage*. 26(12):1400–1409. <http://dx.doi.org/10.1016/j.wasman.2005.09.021> PMID:16356706
7. Nederlof E. Suzanne and Dangbegnon. Constant (2007). Lessons for farmer-oriented research: Experiences from a West African soil fertility management project. *Agriculture and Human Values* 24:369–387.
8. Asgelil Dibabe, 2000. Effect of fertilizer on the yield and nodulation pattern of Faba bean on a Nitosol of Adet North Western Ethiopia. *Ethiopian J. Natural Resources*, 2: 237-244.
9. Sanchez, P.A., (2002). Soil fertility and hunger in Africa. *Sci.*, 295(5562): 2019-2020.
10. Taye Belachew and Yifru Abera. (2010). Assessment of Soil Fertility Status with Depth in Wheat Growing Highlands of Southeast Ethiopia. *World Journal of Agricultural Sciences* 6 (5): 525-531.

11. Eyasu, E. (2002). Farmers' Perception of Soil Fertility Change and Management, Institute for Sustainable Development and SOS Sahel International (UK), Addis Ababa.
12. Khanif, Y.M., (2010). Improvement of Soil Carrying Capacity for Better Living Department of Land Management, Faculty of Agriculture University Putra Malaysia, 43400, Serdang Selangor Malaysia.
13. Abebayehu, A. Eyasu E. and J. Diels. 2011. Comparative Analysis of Soil Nutrient Balance at Farm Level: A Case Study in Jimma Zone, Ethiopia. *In. J of soil sci* 6(4):259-266.
14. Khosro M. and Yousef S. (2012). Bacterial Bio fertilizers for Sustainable Crop Production: A Review. ARPN (Asian Research Publishing Network) *Journal of Agricultural and Biological Science* 7 (5):307-316.
15. Gete Z., Getachew A., Dejene A. and Shahid R. (2010). A Report on Fertilizer and Soil Fertility.
16. Agricultural Transformation Agency (ATA), 2013. Status of Soil Resources in Ethiopia and Priorities for Sustainable Management. GSP for Eastern and Southern Africa Mar 25-27, 2013, Nairobi, Kenya.
17. Abebe Bobo, Eyasu Elias and Gezaghegn Berecha. 2017. Assessment of Soil Fertility Management Practices Employed by Farmers in Selected Villages of Jimma Zone, South Western Ethiopia. *Journal of Biology, Agriculture and Healthcare* 7 (9):2224-320.
18. Singh, R., Gupta, R.K., Patil, R.T., Sharma, R.R., Asrey, R., Kumar, A. and Jangra, K.K., (2010). Sequential foliar application of vermicompost leachates improves marketable fruit yield and quality of strawberry (*Fragaria ananassa* Duch.). *Scientia Horticulturae*, 124, 34-39.
19. Butler TJ, Han KJ, Muir JP, Weindorf DC, and Lastly L. (2008). Dairy manure compost effects on corn silage production and soil properties. *Agronomic Journal*. 100:1541–1545.
20. Kanwar K, Gupta CS, and Paliyal S.S. 2005. Effect of chemical fertilizers vis-à-vis organic manures (vermicompost and FYM) on wheat yield and soil health. *Himachal Journal of Agriculture Research*. 31(2): 47-52.

21. Srikanth, K., Srinivasamurthy, C.A., Siddaramappa, R. and Ramakrishna Parama, V.R., (2000). Direct and residual effect of enriched composts, FYM, vermicompost and fertilizer on properties of an Alfisol. *J. Indian Soc. Soil Sci.*, 48(3): 496-499.
22. Bansal S, Kapoor KK (2000). Vermicomposting of crop residues and cattle dung with *Eisenia foetida*. *Bioresource Technology* 73, 95-98.
23. Maheswarappa H. P., Nanjappa H.V., and Hegde M. R. (1999). Influence of planting material, plant population and organic manures on galangal (*Kaempferia galanga* L.) grown as intercrop in coconut (*Cocos nucifera* L.) garden. *Journal of Spices and Aromatic Crops*. 8(1): 35-40.
24. Nethra NN, Jayaprasad KV and Kale RD. (1999). China aster [*Callistephus chinensis* (L)] cultivation using vermicompost as organic amendment. *Crop Research, Hisar* 17(2): 209–215.
25. Choudhari, V. K. and Kumar, S. P., (2013). Maize production, economics and soil productivity under different organic source of nutrients in eastern himalayan region, India. *Int. J. Plant Product*. 7 (2): 167-186.
26. Sujata, M. G., Longaraju, B. S., Palled, Y. B. and Shalatha, K. V., (2008). Importance of integrated nutrient management practices in maize under rainfed condition. *Karnataka J. Agric. Sci.*, 21 (3): 334-338.
27. Gopinath, K.A., Supradip, S., Mina, B.L., Pande, H., Kundu, S. and Gupta, H.S. (2008). Influence of organic amendments on growth, yield and quality of wheat and on soil properties during transition to organic production. *Nutrient Cycling in Agro ecosystems*, 82 (1), 51-60.
28. OoARD (Bureau of Agriculture and Rural Development), (2019). Annual report of Bureau of the Agricultural and Rural Development of Kolla Tembien, Woreda.
29. Andrea Sembroni, Paola Molin, Francesco Dramis & Bekele Abebe (2017). Geology of the Tekeze River Basin (Northern Ethiopia), *Journal of Maps*, 13:2, 621-631, DOI:10.1080/17445647.2017.1351907.
30. FAO, 1984a. Assistance to Land Use Planning Ethiopia: Geomorphology and Soils. Field Document AG DP/ETH/78/003. The United Nations Development Programmed and Food and Agriculture Organization. FAO, Rome.

31. Rhoades, J.D. (1982). In *Methods of Soil Analysis, Part 2. Second Edition* (A.L. Page, Miller and D.R. Keeney, Eds.), *American Society of Agronomy*. Madison, USA.
32. Jackson, M.L., (1967). *Soil chemical analysis*. Practice Hall of India. New Delhi.
33. Walkley, A., and Black, I.A. (1934). An examination of the method for determining soil organic matter and proposed chromic acid titration method. *Soil Science*, (37):29-38.
34. Bremner, J.M. and Mulvaney, C.S. (1982). Nitrogen - Total. In A.L. Page, R.H. Miller and D.R.Keeney (eds.). *Methods of soil analysis. Part 2-Chemical and microbiological properties. Agronomy*, 9 (2): 595-624. *American Society of Agronomy*. Madison, Wis.
35. Olsen, R., Cole, S., Watanabe, F., and Dean, L. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *United States Department of Agriculture Circ.* 939.
36. Chapman, H.D. (1965). Cation-exchange capacity. In: C. A. Black (ed.) *Methods of soil analysis - Chemical and microbiological properties*. *Agronomy* 9: 891-901.
37. Gupta, P.K. (2000). *Soil, Plant, Water and Fertilizer Analysis*. AGROBIOS Publisher, India.
38. Tucker. B. B. and Kurtz, L. T. (1961). Calcium and magnesium determinations by EDTA titrations. *Soil sci. soc. Am. Proc.* 25:
39. Murphy, H.F. (1968). Reports on fertility status of some soils and other data on soils of Ethiopia College of Agriculture HSIU, Addis Ababa, Ethiopia; Experiment Station Bulletin No.44. 551p.
40. Tekalign Tadese. (1991). *Soil, plant, water, fertilizer, animal manure and compost analysis*. Working Document No. 13. International Livestock Research Center for Africa, Addis Ababa, Ethiopia. and Subtropics. Longman Scientific and Technical, Essex, New York. 474p.
41. Lamond, R. and Whitney. D.A. (1992). *Management of saline and sodic soils*. Kansas State University, Department of Agronomy MF-1022.
42. Berhanu Debele. 1980. *The physical criteria and their rating proposed for land evaluation inthe highland region of Ethiopia*. Land Use Planning and Regulatory Department, Ministry of Agriculture, Addis Ababa, Ethiopia.

43. Landon, J.R. (1991). Booker Tropical Soil Manual: A hand book for soil survey and Agricultural Land Evaluation in the Tropics and Subtropics. Longman Scientific and Technical, Essex, New York. 474p. OR John Wiley & Sons Inc., New York.
44. FAO (Food and Agriculture Organization). (2006). Plant nutrition for food security: A guide for integrated nutrient management. FAO, Fertilizer and Plant nutrition Bulletin 16, Rome.
45. SAS. (2002). SAS/STAT User's Guide, Version 9.1.3. SAS institute Inc., Cary, NC.
46. Gomez, and Gomez H. (1984). Statistical analysis for agricultural research. John Willy and Sons Inc. pp.120-155.
47. Hubbard, R. K., Bosch D.D., Marshall, L.K., Strickland, T. C., Rowland, D., Griffin, T.S., Honeycutt, C.W., Alebrecht, S.L., Sistani, K.R., K.R., Torbert, H.A., Wienhold, B.J., Woodbury, B.L., and Powell, J.M., (2008). Nitrogen mineralization from broiler litter applied to southeastern Coastal plain soils. *Journal of Soil and Water Conservation* 63, 182-192.
48. Aung Naing OO. C. B. Iwai, and P. Saenjan. 2015. Soil Properties and Maize Growth in Saline and Non saline Soils Using Cassava-Industrial Waste Compost and Vermicompost with or without Earthworms. *Land Degradation and Development* 26(3): 300–310.
49. Sharma RC, Banik P. (2014). Vermicompost and fertilizer application: Effect on productivity and profitability of baby corn (*Zea mays* L.) and soil health. *Compost Science and Utilization*. 22:83-92.
50. Zaman, M.M., Chowdhury, M.A.H.; Islam, M.R.; Uddin, M.R. (2015). Effects of vermicompost on growth and leaf biomass yield of stevia and post-harvest fertility status of soil. *J. Bangladesh Agric. Univ.* 12, 169–174.
51. Azarmi R, Giglou MT, Taleshmikail. RD (2008). Influence of vermicompost on soil chemical and physical properties in tomato (*Lycopersicum esculentum*) field. *African Journal of Biotechnology* 7:2397-2401.
52. Mawiyah Mahmud, Rosazlin Abdullah and Jamilah Syafawati Yaacob. (2018). Effect of Vermicompost Amendment on Nutritional Status of Sandy Loam Soil, Growth Performance, and Yield of Pineapple (*Ananas comosus* var. MD2) under Field Conditions. [www.mdpi.com/journal/agronomy](http://www.mdpi.com/journal/agronomy). Doi: 10.3390/agronomy8090183.

53. Angelova V. R.; Akova V. I.; Artinova N. S. and Ivanov K. I. (2013). The Effect of Organic Amendments on Soil Chemical Characteristics. *Bulgarian Journal of Agricultural Science*, 19 (5), 958-971.
54. Pedra, F., Plaza, C., Garcia-Gill, J.C., and Polo, A. (2008). Effects of municipal waste compost and sewage sludge on proton binding behavior of humic acids from Portuguese sandy and clay loam soils. *Bio resource Technology*. 99, 2141-2147.
55. Uz Ilker, Sahriye Sonmez, Ismail Emrah Tavali, Sedat Citak, Dilek Saadet Uras, and Sevil Citak. (2016). Effect of Vermicompost on Chemical and Biological Properties of an Alkaline Soil with High Lime Content during Celery (*Apium graveolens* L.) Production. *Not Bot Horti Agrobi*. 44(1):280-290. DOI: 10.15835/nbha44110157.
56. Tiwari RC, Sharma PK, Khandelwal SK (2004). Effect of green-manuring through *Sesbania cannabina* and *Sesbania rostrata* and nitrogen application through urea to maize (*Zea mays* L.) in maize-wheat (*Triticumaestivum*) cropping system. *Indian Journal of Agronomy*. 49:15-1.
57. Zakir H.M., Sultana M.N., and Saha K.C. (2012). Influence of commercially available organic vs inorganic fertilizers on growth yield and quality of carrot. *Journal of Environmental Science and Natural Resources*. 5:39-45.