

Review Article

Evaluating the Impact of Elevated Temperature on Melia Dubia: Insights into Climate Change Resilience and Adaptation

Abstract

The consequences of climate change extend beyond temperature shifts, encompassing more extreme weather events, increased CO₂ enabling some plants to tolerate environmental stresses, and shifts in precipitation patterns. Climate change manifests through alterations in the mean and variability of various properties, persisting for extended periods, often decades or longer, either due to natural variability or human-induced factors. The bulk of GHG emissions results from fossil fuel energy production, industrial activities, transportation, agriculture, construction, deforestation, and land-use changes. Plant height plays a pivotal role in the growth and development of plant species. Elevated temperature levels have been observed to enhance plant productivity, with varying impacts based on the growth stage and species response to the environment, independent of climate changes.

Formatted: Highlight

Keywords: Global warming, climate change, long-wave radiation, fuel energy production

Introduction

Global warming is a primary driver of climate change, largely attributed to the escalating levels of atmospheric carbon dioxide (CO₂) and other greenhouse gases (GHGs) like methane (CH₄), nitrous oxide (N₂O), and ChloroFluoro Carbons (CFCs) over the past two decades. These GHGs trap long-wave radiation emitted by the Earth's surface, leading to atmospheric warming. Climate change manifests through alterations in the mean and variability of various properties, persisting for extended periods, often decades or longer, either due to natural variability or human-induced factors. The bulk of GHG emissions results from fossil fuel energy production, industrial activities, transportation, agriculture, construction, deforestation, and land-use changes. Natural events such as forest fires and volcanic eruptions also contribute to increased GHG levels. The Intergovernmental Panel on Climate Change (IPCC), established in 1988 by the United Nations Environment Programme (UNEP) and World Meteorological Organization (WMO), plays a pivotal role in assessing climate change and advocating for GHG emission control. The IPCC's fifth assessment report (AR5) in 2013-14 unequivocally asserted that global warming is occurring, associated with heightened GHG concentrations in the atmosphere. Recent studies emphasize the urgency of peaking GHG emissions before 2020 to curb the average global temperature increase to 2°C above pre-industrial levels. Rising temperatures pose significant challenges, with a global mean surface temperature increase of 0.85°C from 1880 to 2012. Predictions suggest a

Comment [MLPdS1]: Lack of references; explain the aim of the work

Formatted: Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

potential rise of 0.3-4.8°C by the end of this century, with higher increases in certain regions like northern Europe. Elevated temperatures have profound implications for agriculture, potentially causing a decline in food production. Heat stress affects plant growth, development, and crop yield, particularly in tropical regions. The IPCC warns of a 15-25% decline in food production in southern India and 25-50% in northern parts. In response to these challenges, understanding the impact of elevated temperatures on plant species is crucial. A study on "Response of *Melia dubia* clones and seedlings to elevated temperature" at the Tamil Nadu Agricultural University aims to evaluate morphological variations and intra-specific physiological and biochemical responses in *Melia dubia*, a fast-growing tree species native to India and Southeast Asia. The global agricultural landscape faces a daunting task of addressing poverty and hunger amidst the escalating challenges posed by climate change. With projections indicating a need for a 70 percent increase in food production to feed an additional 2.3 billion people by 2050, the focus on enhancing food productivity becomes paramount. This urgency is particularly emphasized in the context of leguminous pulses, crucial for their economic importance, rich nutritional content, and diverse applications in human consumption, livestock feed, and industrial processes. One of the primary threats to global food supply production is the rise in average air temperature, predicted to increase by 0.3-0.7°C by 2035 and potentially reaching 1.8 – 4.0°C higher temperatures than the current level by 2100. Elevated temperature is emerging as a significant seasonal phenomenon and a component of climate change, presenting a formidable challenge to plant life. Understanding the mechanisms of heat stress in plants is imperative for future generations, given the potential repercussions on food production.

Climate Change Dynamics

Climate change, defined as a long-term alteration in the statistical distribution of weather patterns over periods ranging from decades to millions of years, manifests in various forms. The indicators of climate change include rising temperatures over oceans, tropospheric temperature changes, alterations in temperature over land, sea surface temperature fluctuations, changes in humidity, sea level variations, and shifts in ocean heat content. Conversely, indicators expected to decrease include sea ice, snow cover, and glaciers. Global warming, a focal point of concern, is closely linked to the surge in greenhouse gas (GHGs) emissions. Concentrations of gases such as CO₂, CH₄, and N₂O have unprecedentedly increased by 40 percent, 150 percent, and 20 percent, respectively, since 1750. Projections indicate a potential increase in global mean surface temperatures by 1.4 to over 5°C by 2100. The consequences of climate change extend beyond temperature shifts, encompassing more

Comment [MLPdS2]: Lack of references

extreme weather events, increased CO₂ enabling some plants to tolerate environmental stresses, and shifts in precipitation patterns.

Impact of Elevated Temperature on Crops

Comment [MLPdS3]: Lack of references

Elevated temperature stress, characterized by temperatures surpassing a threshold level for a duration sufficient to cause irreversible damage to plant growth and development, has profound implications for agriculture. The effects vary between C₃ and C₄ plants, with C₃ plants generally experiencing negative impacts on growth. While rising temperatures benefit C₄ plants' photosynthesis, the stimulating effects of CO₂ on C₃ plants can be overridden. Understanding the complexity of temperature stress involves considering factors such as intensity, duration, and the rate of temperature change. High temperatures influence various physiological processes, including growth, development, and yield in crops. For instance, temperature stress before flowering can lead to reduced germination percentage, increased abnormal seedlings, early flowering, nodules degeneration affecting nitrogen fixation efficiency, and impacts on photosynthetic activity and plant biomass. As the global community grapples with the multifaceted challenges of climate change, research and concerted efforts are essential to develop sustainable agricultural practices. Insights into the impact of elevated temperatures on crop production are crucial for adapting agricultural systems to ensure food security in the face of changing climate dynamics.

Morphological Responses in Elevated Temperature

Shoot Responses

Plant height plays a pivotal role in the growth and development of plant species. Elevated temperature levels have been observed to enhance plant productivity, with varying impacts based on the growth stage and species response to the environment, independent of climate changes. Studies by (Poorter and Navas, 2003) suggest that faster-growing species exhibit more significant growth increases in response to elevated temperatures, with growth rates potentially rising by 10 percent across various species.

Root Responses

Elevated temperature levels contribute to alterations in developmental processes, including root and shoot architecture. Research by (Rogers *et al.*, 1994) indicates an increase in root dry weight in tree species under elevated temperature conditions. Additionally, studies on Scots pine seedlings revealed substantial increases in total root length and total root dry mass, emphasizing the influence of elevated temperatures on root development (Janssens *et al.*, 1998). Other species, such as *Plantago lanceolata*, have demonstrated notable increases in

both shoot and root dry matter production under elevated temperature conditions (Hodge and Millard, 1998).

Leaf Level Responses

Leaf area, a critical component linked to physiological processes controlling dry matter production and yield, experiences positive correlations with elevated temperature. Research by (Chandra and Polisetty, 1998) suggests a positive correlation between leaf area and dry matter accumulation in pea varieties. Elevated temperature has been shown to increase the leaf area index (LAI), attributed to enhanced photosynthetic efficiency and lower light compensation points, allowing leaves to maintain a positive carbon balance (Ferris *et al.*, 2001). Studies on populus clones indicate an 8-18 percent increase in leaf area under elevated temperature conditions (Ceulemans *et al.*, 1995).

Morphological Responses in Elevated Temperature

Elevated temperature influences the height, diameter, and biomass of broadleaf tree species in boreal regions until reaching the optimum temperature. Increased photosynthetic rates lead to faster growth and higher biomass production, with photosynthetic carbon gain accounting for the majority of plant biomass accumulation (Abdul-Hamid and Mencuccini, 2009). Genotypic characteristics also play a role in regulating the physiology and growth of trees (Lamhamediet *et al.*, 2000; Aspinwall *et al.*, 2013).

Growth Increment

Higher temperatures have been linked to increased cell division, resulting in heightened plant height. This aligns with studies by (Wertin *et al.*, 2011), reporting an increase in the height of oak seedlings under elevated temperatures compared to ambient conditions.

Leaf Area

Photosynthesis and morphogenetic processes are susceptible to higher temperatures, causing modifications in plant growth. Elevated air temperatures initially improve leaf extension but accelerate maturity, limiting final leaf size (Rawson, 1992). The effects of radiation on photosynthetic activity and leaf area have been reported, with increased air temperatures improving leaf extension (Monteith, 1977).

Collar Diameter and Number of Leaves

Increased air temperature synergistically influences plant growth, improving collar diameter, the number of leaves, and leaflets. These findings align with the work of (Rahman, Al-Amin, Akter, 2012).

Physiological Responses in Elevated Temperature

Comment [MLPdS4]: Which one? Or both references?

Photosynthetic Rate

Photosynthesis is significantly impacted by elevated temperatures, with reductions observed at higher temperatures. The PS II system is particularly affected, leading to chlorophyll degradation and inhibition of Rubisco, ultimately causing yield reduction (Crafts-Brandner and Salvucci, 2002). Elevated temperature stress can influence photosynthesis through stomatal closure and decreased CO₂ flow (Greer and Weedon, 2012).

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Transpiration Rate

Transpiration rates increase under high temperatures due to stomatal opening. The balance between low photosynthetic rates and high transpiration rates during temperature stress affects plant establishment, particularly in crops like chickpea (Singh and Dhaliwal, 1972). Adequate water supply is essential for coping with high-temperature stress, as transpiration cooling plays a crucial role (Weerakoon *et al.*, 2008).

Stomatal Conductance

Stomatal conductance and net photosynthesis are hindered by elevated temperature stress, primarily due to decreased Rubisco activase enzyme. Studies on tobacco plants under stress have shown stomatal conductance increases up to 40 percent between 30 and 40°C but declines above 40°C (Kubien *et al.*, 2008). Elevated temperature stress during flowering in soybean has been observed to decrease stomatal conductance (Djanaguiraman *et al.*, 2011). In summary, the intricate interactions between elevated temperature and plant responses involve complex morphological and physiological changes. Understanding these responses is crucial for predicting and mitigating the impacts of climate change on plant growth and productivity.

Leaf Temperature

Leaf temperature is a crucial parameter affected by high-temperature stress in plants. Changes in transpiration rate can lead to alterations in leaf temperature (Farquhar and Sharkey, 1982). Stomatal closure results in the termination of evapotranspiration, causing an increase in leaf temperature (Lourtie *et al.*, 1995). Environmental factors and transpirational cooling significantly influence the surface temperature of plant leaves (Nobel, 1991; Siddique *et al.*, 2001). Elevated leaf temperature under stress inhibits enzymatic activity and various physiological processes, ultimately deactivating photosynthetic mechanisms (Chaves *et al.*, 2002; Crafts-Brandner and Salvucci, 2002).

Water Use Efficiency (WUE)

Water use efficiency (WUE), defined as the ratio of water loss to carbon fixed during photosynthesis, is a critical leaf-level response to elevated temperatures. While an increase in WUE is common, it may not necessarily be directly proportional to changes in plant growth

and photosynthesis (Beerling *et al.*, 1996). Elevated temperature, coupled with reduced stomatal opening, conductance, and transpiration rates, also depresses dark respiration rates, leading to enhanced WUE (Wullschleger *et al.*, 1992; Murray, 1995). The rise in WUE is associated with increased drought tolerance in many plants, potentially allowing for expanded plant distributions (Tyree and Alexander, 1993; Huxman *et al.*, 1998). However, not all C4 crop plants exhibit a positive response to elevated growth temperature. Numerous single-species studies on various trees, including longleaf pine, red oak, scrub oak, silver birch, beech, sweet gum, and spruce, document positive responses in WUE to elevated atmospheric temperature concentrations (Wullschleger and Norby, 2001). Intrinsic water use efficiency (IWUE), which considers the ratio of photosynthetic CO₂ uptake to transpirational water vapour loss, is crucial for evaluating the response of different tree species to elevated temperature. Studies on *Quercus robur*, *Fagus sylvatica*, and *Pinus sylvestris* trees show significant increases in IWUE under elevated temperature concentrations (Waterhouse *et al.*, 2004). Similar findings in *Sabina przewalskii* and *Picea crassifolia* trees indicate a long-term increase in IWUE, contributing to a better understanding of how different tree species respond to elevated temperature conditions in specific environments (Liu *et al.*, 2007).

Comment [MLPd55]: Add to reference item, or corrected for 2002

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Biochemical Responses in Elevated Temperature Conditions

Chlorophyll

Chlorophyll, a critical pigment for photosynthetic capacity, serves as a sensitive indicator of environmental conditions. Chlorophyll a and b are essential for converting light energy into stored chemical energy. Chlorophyll content directly influences photosynthetic potential and primary production (Curran *et al.*, 1995). Leaf chlorophyll content provides an indirect estimation of nutrient status, as a significant portion of leaf nitrogen is incorporated into chlorophyll (Filella *et al.*, 1995; Moran *et al.*, 2000). While some studies, such as those by Wullschleger *et al.*, (1992), observed a reduction in chlorophyll and accessory pigments in response to elevated CO₂ and temperature, other research demonstrates that atmospheric CO₂ enrichment may increase, decrease, or have no effect on leaf chlorophyll concentrations.

Nitrogen Use Efficiency

Elevated temperature environments increase the nitrogen demands of plants due to accelerated growth and metabolism. This increased demand for nutrients, particularly nitrogen, is essential for various growth processes. Studies by Luo *et al.*, (2004) and Deng and Woodward (1998) indicate that elevated temperature leads to increased nitrogen demands, resulting in greater total biomass compared to ambient levels. Other studies, such

as those on strawberry and *Trifolium repens*, suggest that elevated temperature levels enhance growth-based nitrogen use efficiencies (Lüscher *et al.*, 2004).

Total Carbohydrates

Greater carbohydrate supply and improved water use efficiency contribute to larger individual leaves and more rapid canopy development under elevated temperature conditions. Increased photosynthetic activity and water use efficiency lead to enhanced carbohydrate content in tree species (Ferris *et al.*, 2001). While initial enhancements in photosynthesis rates may occur in C₃ plants exposed to elevated temperature, acclimation to the environment often leads to a subsequent decline (Bowes, 1996). In conclusion, the biochemical responses of plants to elevated temperature conditions involve intricate interactions, influencing crucial parameters such as chlorophyll content, nitrogen use efficiency, and carbohydrate levels. Understanding these biochemical responses is essential for predicting the impact of climate change on plant physiology and productivity.

Proteins

The impact of elevated temperature on protein concentration in plants has been a subject of study, and contrasting results have been reported. Rogers *et al.*, (1996) observed a temperature-induced reduction in the protein concentration of flour derived from wheat plants. Similarly, Allard *et al.*, (2003) reported lower nitrogen concentrations but higher water-soluble carbohydrate concentrations in leaves of individual species under enriched temperature levels. Picon-Cochard *et al.*, (2004) also observed an increase in protein concentration in response to elevated temperature, despite a reduction in overall protein concentration. Contradictory findings were reported by Kimball *et al.*, (2001), who observed a 50% increase in leaf protein concentration under enriched temperature levels in wheat plants. The studies conducted by Idso and Idso (2001) reported varying effects on protein concentration under elevated temperature levels across different agricultural crops.

Phenol Components

The presence of phenolic compounds in plants plays a crucial role in their response to elevated temperatures. Wetzal and Tuchman (2005) observed a significant increase (40.6%) in total phenolic content in cattails grown in elevated temperature levels compared to ambient air. Gebauer *et al.*, (1997) reported a notable increase in both above and below-ground total phenolic concentrations in loblolly pine seedlings grown in elevated temperature conditions. Similar findings were observed in temperate regions by Peñuelas *et al.*, (2002), who reported a 20 to 605 increase in leaf phenolic concentrations in response to a doubling of CO₂ content. Additionally, Wetzal and Tuchman (2005) observed a 63.2% increase in total

phenol compounds in aspen seedlings grown in elevated temperatures. Coley *et al.*, (2002) studied nine species of tropical trees, with eight species exhibiting positive leaf phenolic responses, and one species showing a 27% decline. The mean response of all nine species was an increase of 48% in phenol content. These results indicate that both temperate and tropical trees show large interspecific variation in their response to temperature, with an average increase in phenol content of 50%. In contrast, Hamilton *et al.*, (2004) found no significant effect of elevated temperature on the chemical composition of leaves in loblolly pine plantations. Similarly, Hall *et al.*, (2005) detected no significant difference in total phenolic content in three oak species between high-temperature and ambient treatments.

Future Temperature Projection

Asia has experienced increasing surface air temperatures, with more pronounced changes during winter than summer. The observed increase ranges from less than 1°C to 3°C per century across different sub-regions. Climate change is recognized as a major threat, impacting ecosystems, agriculture, water resources, and socio-economy globally and regionally. The linear warming trend over the last century was 0.74°C, nearly twice that of the previous 50 years. Projections based on various emission scenarios indicate a temperature increase of about 3 to 5°C by 2070 over the Indian region. Specific predictions for Tamil Nadu and the Upper Ganga Basin also indicate substantial temperature increases.

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Comment [MLPdS6]: redundant

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Comment [MLPdS7]: add reference

Growth Chamber Studies and Design

Several studies have employed growth chambers to simulate elevated temperature conditions. Djanaguiraman *et al.*, (2010) used a temperature-controlled study for sorghum, while Punyamurthy *et al.*, (2013) conducted an experimental trial on rice crops using a Climate Control Chamber. Zhang *et al.*, (2013) designed temperature-controlled chambers for field experiments, monitoring air temperature and relative humidity. Singh *et al.*, (2013) employed Open Top Chambers (OTCs) made of aluminum frames covered with UV-treated polycarbonate sheets for a field experiment on sorghum. Parthiban and Kavitha (2014) studied varietal differences in black gram under elevated temperatures using open-top chambers made of polycarbonate sheets with steel tubes as frames. These growth chamber studies and experimental designs provide valuable insights into the response of various crops to elevated temperatures, facilitating our understanding of climate change impacts on agriculture.

Comment [MLPdS8]: it seems a Conclusion is needed

Formatted: Font: Hidden, Highlight

Response of tree species to elevated temperature

The review indicated that there is an important need to understand the plant responses to elevating temperature to bring out “adaptive” genotypes for future climate. The revived

Formatted: Highlight

also revealed that decimal research works carried out in tropical tree species particularly in *Melia dubia* with reference to response of tree species to elevated temperature.

Conclusion:

In conclusion, the global agricultural landscape is confronted with the daunting challenge of addressing poverty and hunger, especially given the anticipated 70 percent increase in food production needed to feed an additional 2.3 billion people by 2050. The urgency to enhance food productivity is particularly critical for leguminous pulses, which hold economic importance and provide rich nutritional content with diverse applications. One of the primary threats to global food supply is the rise in average air temperature, predicted to escalate significantly in the coming decades. Elevated temperature is emerging as a significant component of climate change, posing a formidable challenge to plant life and, consequently, food production. Understanding the mechanisms of heat stress in plants is crucial to navigate the potential repercussions on agricultural systems and food security. The dynamics of climate change manifest through various indicators, including rising temperatures, alterations in humidity, sea level variations, and shifts in precipitation patterns. Global warming, primarily driven by increased greenhouse gas emissions, poses multifaceted challenges, including extreme weather events and changes in plant responses. Elevated temperature stress has profound implications for agriculture, impacting various physiological processes in crops. Morphological responses, such as shoot, root, and leaf-level changes, vary among different plant species. Understanding these responses involves considering factors like intensity, duration, and the rate of temperature change. Physiological responses to elevated temperature encompass parameters like photosynthetic rate, transpiration rate, stomatal conductance, and water use efficiency. These responses play a critical role in determining plant growth, development, and overall productivity. Biochemical responses in elevated temperature conditions involve changes in chlorophyll content, nitrogen use efficiency, and carbohydrate levels. These responses are intricate and crucial for predicting the impact of climate change on plant physiology. The review also emphasizes the importance of growth chamber studies and experimental designs to simulate elevated temperature conditions, providing valuable insights into crop responses. These studies contribute to a better understanding of climate change impacts on agriculture, aiding in the development of sustainable practices. Looking forward, the response of tree species to elevated temperature emerges as a critical area of research. Understanding the adaptive genotypes and responses of various tree species, particularly in tropical regions, will be essential for developing resilient ecosystems in the face of changing climate dynamics. In summary, as the world faces the

challenges of climate change and the need for increased food production, research, and concerted efforts are crucial. Insights into plant responses to elevated temperatures are imperative for developing adaptive strategies to ensure food security and sustainable agricultural practices for future generations.

References:

- Abdul-Hamid, Hazandy, and Maurizio %J Tree physiology Mencuccini. 2009. "Age-and size-related changes in physiological characteristics and chemical composition of *Acer pseudoplatanus* and *Fraxinus excelsior* trees." 29 (1):27-38.
- Allard Vincent, Paul CD Newton, Mark Lieffering, Harry Clark, Cory Matthew, Jean- François Soussana, and Yvonne S Gray. 2003. "Nitrogen cycling in grazed pastures at elevated CO₂: N returns by ruminants." *Global Change Biology* 9 (12):1731-1742.
- Aspinwall Michael J, John S King, and Steven E %J Trees McKeand. 2013. "Productivity differences among loblolly pine genotypes are independent of individual-tree biomass partitioning and growth efficiency." 27 (3):533-545.
- Beerling, DJ, J Heath, FI Woodward, and TA Mansfield. 1996. "Drought—CO₂ interactions in trees observations and mechanisms." *New Phytologist* 134 (2):235-242.
- Bowes, George. 1996. "Photosynthetic responses to changing atmospheric carbon dioxide concentration." In *Photosynthesis and the Environment*, 387-407. Springer.
- Ceulemans, R, XN Jiang, and BY Shao. 1995. "Growth and physiology of one-year old poplar (*Populus*) under elevated atmospheric CO₂ levels." *Annals of Botany* 75 (6):609-617.
- Chandra, R, and R Polisetty. 1998. "Factors affecting growth and harvest index in pea (*Pisum sativum* L.) varieties differing in time of flowering and maturity." *Journal of agronomy crop science* 181 (3):129-135.
- Chaves, Maria Manuela, João Santos Pereira, João Maroco, Maria Luisa Rodrigues, Cândido Pereira Pinto Ricardo, Maria Leonor Osório, I Carvalho, T Faria, and C Pinheiro. 2002. "How plants cope with water stress in the field? Photosynthesis and growth." *Annals of Botany* 89 (7):907-916.
- Coley, P, M Massa, C Lovelock, and Klaus Winter. 2002. "Effects of elevated CO₂ on foliar chemistry of saplings of nine species of tropical tree." *Oecologia* 133 (1):62-69.
- Crafts-Brandner, Steven J, and Michael E Salvucci. 2002. "Sensitivity of photosynthesis in a C₄ plant, maize, to heat stress." *Plant physiology* 129 (4):1773-1780.

- Curran, Paul J, W Robert Windham, and Henry L %J Tree physiology Gholz. 1995. "Exploring the relationship between reflectance red edge and chlorophyll concentration in slash pine leaves." 15 (3):203-206.
- Djanaguairaman, M, PV Vara Prasad, and M Seppanen. 2010. "Selenium protects sorghum leaves from oxidative damage under high temperature stress by enhancing antioxidant defense system." *Plant Physiology Biochemistry* 48 (12):999-1007.
- Djanaguairaman, M, PV Vara Prasad, DL Boyle, and WT Schapaugh. 2011. "High-temperature stress and soybean leaves: leaf anatomy and photosynthesis." *Crop Science* 51 (5):2125-2131.
- Farquhar, Graham D, Marion H O'Leary, and Joe A Berry. 1982. "On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves." *Functional Plant Biology* 9 (2):121-137.
- Ferris, R, M Sabatti, F Miglietta, RF Mills, and G Taylor. 2001. "Leaf area is stimulated in Populus by free air CO₂ enrichment (POPFACE), through increased cell expansion and production." *Plant, Cell Environment* 24 (3):305-315.
- Filella, I, L Serrano, J1 Serra, and J Penuelas. 1995. "Evaluating wheat nitrogen status with canopy reflectance indices and discriminant analysis." *Crop Science* 35 (5):1400-1405.
- Gebauer, Renate LE, Boyd R Strain, and James F Reynolds. 1997. "The effect of elevated CO₂ and N availability on tissue concentrations and whole plant pools of carbon-based secondary compounds in loblolly pine (Pinus taeda)." *Oecologia* 113 (1):29-36.
- Greer, Dennis H, and Mark M Weedon. 2012. "Modelling photosynthetic responses to temperature of grapevine (Vitis vinifera cv. Semillon) leaves on vines grown in a hot climate." *Plant, Cell Environment* 35 (6):1050-1064.
- Hall, Myra C, Peter Stiling, Daniel C Moon, Bert G Drake, and Mark D Hunter. 2005. "Effects of elevated CO₂ on foliar quality and herbivore damage in a scrub oak ecosystem." *Journal of Chemical Ecology* 31 (2):267-286.
- Hamilton, Jason G, Arthur R Zangerl, May R Berenbaum, Jeffrey Phippen, Mihai Aldea, and Evan H DeLucia. 2004. "Insect herbivory in an intact forest understory under experimental CO₂ enrichment." *Oecologia* 138 (4):566-573.
- Hodge, Angela, and Peter Millard. 1998. "Effect of elevated CO₂ on carbon partitioning and exudate release from Plantago lanceolata seedlings." *Physiologia Plantarum* 103 (2):280-286.

- Huxman, Travis E, Erik P Hamerlynck, BD Moore, SD Smith, Dean N Jordan, Stephen F Zitzer, Robert S Nowak, James S Coleman, and Jeffrey R Seemann. 1998. "Photosynthetic down-regulation in *Larrea tridentata* exposed to elevated atmospheric CO₂: interaction with drought under glasshouse and field (FACE) exposure." *Plant, Cell Environment* 21 (11):1153-1161.
- Idso, Sherwood B, and Keith E Idso. 2001. "Effects of atmospheric CO₂ enrichment on plant constituents related to animal and human health." *Environmental Experimental Botany* 45 (2):179-199.
- Janssens, Ivan A, Meg Crookshanks, Gail Taylor, and Reinhart Ceulemans. 1998. "Elevated atmospheric CO₂ increases fine root production, respiration, rhizosphere respiration and soil CO₂ efflux in Scots pine seedlings." *Global Change Biology* 4 (8):871-878.
- Kimball, BA, CF Morris, PJ Pinter, GW Wall, DJ Hunsaker, FJ Adamsen, RL LaMorte, SW Leavitt, TL Thompson, and AD Matthias. 2001. "Elevated CO₂, drought and soil nitrogen effects on wheat grain quality." *New Phytologist* 150 (2):295-303.
- Kubien, David S, Rowan F %J Plant Sage, cell, and environment. 2008. "The temperature response of photosynthesis in tobacco with reduced amounts of Rubisco." 31 (4):407-418.
- Lamhamedi, Mohammed S, Hélène Chamberland, Pierre Y Bernier, and Francine M Tremblay. 2000. "Clonal variation in morphology, growth, physiology, anatomy and ultrastructure of container-grown white spruce somatic plants." *Tree physiology* 20 (13):869-880.
- Liu, Xiaohong, Xuemei Shao, Eryuan Liang, Liangju Zhao, Tuo Chen, Dahe Qin, and Jiawen Ren. 2007. "Species-dependent responses of juniper and spruce to increasing CO₂ concentration and to climate in semi-arid and arid areas of northwestern China." *Plant Ecology* 193 (2):195.
- Lourtie, E, Marc Bonnet, and Léonard Bosschaert. 1995. "New glyphosate screening technique by infrared thermometry." Fourth International Symposium on Adjuvants for Agrochemicals, Australia.
- Luo, Yiqi, BO Su, William S Currie, Jeffrey S Dukes, Adrien Finzi, Ueli Hartwig, Bruce Hungate, Ross E McMurtrie, RAM Oren, and William J Parton. 2004. "Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide." *Bioscience* 54 (8):731-739.
- Lüscher, Andreas, Markus Daepf, Herbert Blum, Ueli A Hartwig, and Josef Nösberger. 2004. "Fertile temperate grassland under elevated atmospheric CO₂—role of feed-

back mechanisms and availability of growth resources." *European Journal of Agronomy* 21 (3):379-398.

Monteith, John Lennox %J Philosophical Transactions of the Royal Society of London. B, Biological Sciences. 1977b. "Climate and the efficiency of crop production in Britain." 281 (980):277-294.

Monteith, John Lennox 1977a. "Climate and the efficiency of crop production in Britain." *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* 281 (980):277-294.

Murray, David R 1995. "Plant responses to carbon dioxide." *American Journal of Botany* 82 (5):690-697.

Nobel, Park S 1991. "Achievable productivities of certain CAM plants: basis for high values compared with C3 and C4 plants." *New Phytologist* 119 (2):183-205.

Parthiban, VK, and R %J International Journal of Pharmacological Screening Methods Kavitha. 2014. "In vitro screening of effective biocontrol agents against bean anthracnose pathogen, *Colletotrichum lindemuthianum*." 4 (1):32-35.

Peñuelas, Josep, E Castells, R Joffre, and Roberto Tognetti. 2002. "Carbon- based secondary and structural compounds in Mediterranean shrubs growing near a natural CO2 spring." *Global Change Biology* 8 (3):281-288.

Picon-Cochard, Catherine, Florence Teyssonneyre, Jean Michel Besle, and Jean-François Soussana. 2004. "Effects of elevated CO2 and cutting frequency on the productivity and herbage quality of a semi-natural grassland." *European Journal of Agronomy* 20 (4):363-377.

Poorter, Hendrik, and Marie- Laure Navas. 2003. "Plant growth and competition at elevated CO2: on winners, losers and functional groups." *New Phytologist* 157 (2):175-198.

Punyamurthy, Ramadevi, Dhanalakshmi Sampathkumar, Basavaraju Bennehalli, and Chikkol V %J Chemical Science Transactions Srinivasa. 2013. "Influence of esterification on the water absorption property of single abaca fiber." 2 (2):413-422.

Rahman, M., M. Al-Amin, S.J.J.o.F. Akter, and E. Science. 2012. "Artocarpus chaplasha: Establishment and initial growth performance at elevated temperature and saline stresses." 28 (1):12-18.

Rawson, HM %J Australian Journal of Botany. 1992b. "Plant responses to temperature under conditions of elevated CO2." 40 (5):473-490.

Formatted: English (United States), Do not check spelling or grammar, Highlight

Formatted: English (United States), Do not check spelling or grammar, Highlight

- Rogers, GS, PJ Milham, M Gillings, and JP Conroy. 1996. "Sink strength may be the key to growth and nitrogen responses in N-deficient wheat at elevated CO₂." *Functional Plant Biology* 23 (3):253-264.
- Rogers, Hugo H, G Brett Runion, and Sagar V Krupa. 1994. "Plant responses to atmospheric CO₂ enrichment with emphasis on roots and the rhizosphere." *Environmental pollution* 83 (1-2):155-189.
- Siddique, KHM, KL Regan, D Tennant, and BD Thomson. 2001. "Water use and water use efficiency of cool season grain legumes in low rainfall Mediterranean-type environments." *European Journal of Agronomy* 15 (4):267-280.
- Singh, NT, and GS Dhaliwal. 1972. "Effect of soil temperature on seedling emergence in different crops." *Plant Soil* 37 (2):441-444.
- Singh, SS, Joydeep Mukherjee, Santosh Kumar, and Mohd %J Journal of Agrometeorology Idris. 2013. "Effect of elevated CO₂ on growth and yield of rice crop in open top chamber in Sub humid climate of eastern India." 15 (1):1.
- Tyree, Melvin T, and John D Alexander. 1993. "Plant water relations and the effects of elevated CO₂: a review and suggestions for future research." *Vegetatio* 104 (1):47-62.
- Waterhouse, JS, VR Switsur, AC Barker, AHC Carter, DL Hemming, NJ Loader, and I Robertson. 2004. "Northern European trees show a progressively diminishing response to increasing atmospheric carbon dioxide concentrations." *Quaternary Science Reviews* 23 (7-8):803-810.
- Weerakoon, WMW, A Maruyama, and K Ohba. 2008. "Impact of humidity on temperature- induced grain sterility in rice (*Oryza sativa* L)." *Journal of agronomy crop science* 194 (2):135-140.
- Wertin, T.M., M.A. McGuire, and R.O.J.T.P. Teskey. 2011. "Higher growth temperatures decreased net carbon assimilation and biomass accumulation of northern red oak seedlings near the southern limit of the species range." 31 (12):1277-1288.
- Wetzel, Robert G, and Nancy C Tuchman. 2005. "Effects of atmospheric CO₂ enrichment and sunlight on degradation of plant particulate and dissolved organic matter and microbial utilization." *Archiv für Hydrobiologie* 162 (3):287-308.
- Wullschleger, SD, RJ Norby, and DL Hendrix. 1992. "Carbon exchange rates, chlorophyll content, and carbohydrate status of two forest tree species exposed to carbon dioxide enrichment." *Tree physiology* 10 (1):21-31.

Wullschleger, SD, TJ Tschaplinski, and RJ Norby. 2002. "Plant water relations at elevated CO₂—implications for water- limited environments." *Plant, Cell Environment* 25 (2):319-331.

Zhang, Renjian, Junshan Jing, Jun Tao, S-C Hsu, Gehui Wang, Junji Cao, Celine Siu Lan Lee, Lihua Zhu, Zhongming Chen, Yue %J Atmospheric Chemistry Zhao, and Physics. 2013. "Chemical characterization and source apportionment of PM 2.5 in Beijing: seasonal perspective." 13 (14):7053-7074.

UNDER PEER REVIEW