

## Review Article

# Analysis of the ecological value of pioneer plants in the Yellow River Basin

---

### ABSTRACT

Pioneer plants are plant species that appear early in the community succession or show strong adaptability, which can improve the local ecological environment and create conditions for the succession and recovery of other plants and animals. Based on the analysis of the general situation of the Yellow River Basin, this paper explores the ecological value of pioneer plants to the treatment and restoration of the fragile ecological areas such as Pisha sandstone area, coastal saline-alkali land and abandoned mine area in the Yellow River Basin, in order to provide a theoretical basis for pioneer plants to further reflect the ecological functions in the Yellow River Basin.

*Keywords:* Yellow River Basin; pioneer plant; ecological value

### 1. INTRODUCTION

The Yellow River originates in mountainous western China and flows through nine provinces and autonomous regions, namely Qinghai, Sichuan, Gansu, Ningxia, Inner Mongolia, Shaanxi, Shanxi, Henan and Shandong, before finally discharging into the Bohai Sea at Dongying, Shandong. The Yellow River traverses four major geomorphic units, namely the Qinghai-Tibet Plateau, the Inner Mongolia Plateau, the Loess Plateau, and the Huang-Huai-Hai Plain, in a west-to-east direction. The basin exhibits significant variation in altitude differences, as well as longitudinal and latitudinal spans. The origin is perpetually obscured by ice and snow; The central part consists primarily of plateaus and loess formations, experiencing severe soil erosion due to the high cement sand content from the Yellow River; The eastern part, however, mainly consists of Yellow River alluvial plains and is characterized by flat terrain.

The source region of the Yellow River is high-altitude, rugged, cold, with frozen soil and glacial landforms, low precipitation and low vegetation coverage, making the ecological environment fragile. The middle reaches of the Yellow River flow through the Loess Plateau, which has a loess topography, sparse vegetation, frequent and concentrated summer storms, and severe soil erosion. The lower reaches of the Yellow River, due to their own soil erosion, combined with sediment brought from the middle and upper reaches, have resulted in significant development of the overland flow.

**Comment [G1]:** A short intro and objective were observed in the abstract. What have you found out? Conclusion? Add some components in your abstract.

**Comment [G2]:** At least 3-5 keywords

The Yellow River Basin is home to a diverse range of pioneer plants that play a crucial ecological role in various areas along the upper, middle and lower reaches of the river due to their exceptional adaptability to different vegetation conditions.

## 2. OVERVIEW OF PIONEER PLANTS AND MAJOR TYPES OF PIONEER PLANTS IN THE YELLOW RIVER BASIN

Pioneer plants are those that appear earliest in the succession of communities. They can survive in areas where soil nutrients and water are severely lacking, such as deserts, polar regions, and other harsh environments. Pioneer plants are not only able to withstand drought and barrenness, but also grow quickly, produce large seeds, and have a strong ability to spread. Pioneer plants have the ability to fix nutrients in the soil, enhance soil quality, attract fauna, and gradually enhance the ecological environment of the area. They play a crucial role in the initial stages of community succession.

Pioneer plants include not only lichens, algae and mosses, but also many herbs, shrubs and trees such as *Suaeda salsa*, *Tamarix chinensis* Lour. and *Eucommia ulmoides* Oliv. The common pioneer plant species distributed in the Yellow River Basin are presented in Table 1.

**Table 1.** Common pioneer plants of the Yellow River Basin

Name	Family	Genus	Life type	Main distribution area
<i>Artemisia blepharolepis</i> Bge.	Compositae	Artemisia	Annual herb	Inner Mongolia, Shaanxi (north), Ningxia
<i>Poa cynosuroides</i> (Hook. f.) Woodson	Apocynaceae	Poa cynosuroides	Perennial herb	Qinghai, Gansu, etc
<i>Cynanchum chinense</i> R. Br.	Asclepiadaceae	Cynanchum	Perennial herb	Henan, Shandong, Shanxi, Shaanxi, Ningxia, Gansu
<i>Suaeda salsa</i>	Amaranthaceae	Suaeda	Annual herb	Henan, Shandong, Shanxi, Shaanxi, Ningxia, Gansu, Qinghai
<i>Pyracantha fortuneana</i> (Maxim.) Li	Rosaceae	Pyracantha	Evergreen shrub	Shaanxi, Henan
<i>Hippophae rhamnoides</i>	Elaeagnaceae	Hippophae	Deciduous shrubs or trees	Inner Mongolia, Shaanxi, Shaanxi, Qinghai
<i>Tamarix chinensis</i> Lour.	Tamaricaceae	Tamarix	Deciduous shrubs or trees	Henan, Shandong
<i>Lonicera japonica</i> Thunb.	Caprifoliaceae	Lonicera	Perennial evergreen shrub	Shandong, Henan, Shaanxi, Shaanxi, Sichuan
<i>Robinia pseudoacacia</i> Linn.	Leguminosae	Robinia	Deciduous trees	Qinghai, Gansu, Ningxia, Inner Mongolia, Shaanxi, Shanxi, Henan, Shandong, Sichuan
<i>Zanthoxylum bungeanum</i> Maxim.	Rutaceae	Zanthoxylum	Small deciduous perennial tree	Qinghai, Gansu, Ningxia, Inner Mongolia, Shaanxi, Shanxi, Henan,

**Comment [G3]:** How did you arrange the common pioneer plants? It is not alphabetical. Is it random? Make it organize. If it is organized in a certain way, please mention it on a paper or maybe provide a map of distribution if that will work.

**Comment [G4]:** Some of the pioneer plants listed here are not found in the article. So the purpose of listing all of the common species is for awareness only?

Shandong, Sichuan

<i>Eucommia ulmoides</i> Oliv.	Eucommiaceae	Eucommia	Deciduous trees	Henan, Shaanxi, Gansu, Sichuan
<i>Broussonetia papyrifera</i>	Moraceae	Broussonetia	Deciduous trees	Qinghai, Gansu, Ningxia, Inner Mongolia, Shaanxi, Shanxi, Henan, Shandong
<i>Populus euphratica</i> Oliv.	Salicaceae	Populus	Deciduous trees	Inner Mongolia (west), Gansu, Qinghai

### 3. ECOLOGICAL EFFECTS OF PIONEER PLANT

#### 3.1 Treatment of saline-alkali land

Many pioneer plants are salt-tolerant and salt-resistant, and can be divided into salt-accumulating plants and salt-secreting plants.

Salt-accumulating plants exhibit robust salt resistance and can efficiently accumulate salt within their tissues without sustaining damage. As a representative example of such plants, *Suaeda salsa* demonstrates the ability to significantly reduce soil salinity levels, enhance microbial species diversity and abundance, as well as improve the structural complexity of microbial communities<sup>[1]</sup>. Furthermore, its root, stem, and leaf tissues harbor numerous halophilic microorganisms that actively contribute to the restoration process of saline-alkali soils<sup>[2]</sup>.

Saline-secreting plants have the ability to excrete salt from their bodies, allowing them to survive in high-salt soils. Examples of these plants include *Tamarix chinensis* Lour, a typical salt-secreting plant that can absorb and accumulate soil salts and then secrete them through salt glands in the form of crystals from leaves and branches to regulate internal salt balance, maintain osmotic pressure, and grow in high-salt environments. Therefore, in the treatment of saline-alkali land, the leaves of *Tamarix chinensis* Lour can be transferred by defoliation and manual removal, so as to realize the transfer of soil salinity. At the same time, plant growth can increase surface cover by forming mulch or increasing the canopy to protect the soil surface and reduce evaporation and wind erosion<sup>[3]</sup>.

#### 3.2 WINDPROOF SAND FIXATION, SOIL EROSION IMPROVEMENT

The root system of plants is a polymeric organism, and its cellulose molecular bonds give it a certain tensile resistance in the axial direction, and when applied it exhibits a certain elastic or viscoelasticity to accommodate the axial tension. When plants are attacked by strong external forces, such as strong winds or heavy rain, their roots and the soil form a root-soil complex, which has strong adhesion and friction and is better able to resist disturbance than pure soil<sup>[5]</sup>. The roots of pioneer plants are generally longer, faster growing, more numerous, and produce greater traction on the soil than those of common plants. As a result, pioneer plants have excellent soil consolidation, preventing water erosion from carrying away the soil and improving soil erosion to a certain extent. *Robinia pseudoacacia* Linn., for example, has strong roots, excellent soil fixation properties, wide adaptability, rapid growth rate and high survival rate, and can grow normally and thrive on hills, barren mountains, sunny slopes and

**Comment [G5]:** Exposition of ecological effects is good but requires more depth. The author just describes the plants and give example.

sandy waste lands. The roots of *Robinia pseudoacacia* are interwoven in a network structure, with lateral and fibrous roots closely connected to the soil, effectively preventing surface soil loss<sup>[8]</sup>.  
Error! Reference source not found.

### 3.3 IMPROVEMENT OF SOIL QUALITY

When plants root in the soil, they loosen the soil, promote aeration and permeability, and improve soil structure. The roots have the ability to absorb nutrients while secreting root material, and are symbiotic with rhizospheric microorganisms that can activate and fix various nutrients, thereby increasing the soil nutrient content<sup>[8]</sup>. Pioneer plants, which normally have faster growth rates and more developed roots, are excellent at absorbing water and nutrients deep into the soil layer, helping to improve soil structure, improve aeration and permeability<sup>[9]</sup>, and reduce ionic concentration and salinity vulnerability, thus enhancing soil fertility<sup>[10]</sup>. For example, *Hippophae rhamnoides* has shown greater ability than other legumes in nitrogen fixation by forming nodules through symbiosis with actinobacteria and mycobacterium, which can significantly increase the amount of organic matter and nitrogen in the soil, thereby increasing soil strength and water storage capacity and enhancing soil fertility<sup>[11]</sup>.

### 3.4 PROMOTING BIODIVERSITY AND COMMUNITY SUCCESSION

Pioneer plants play an important role in community succession. With their rapid growth and well-developed roots, pioneer plants can effectively improve the soil environment, which increases the survival chances of more plants, attracts animals, increases biodiversity, and promotes the improvement of the local ecological environment and community succession. In arid and semi-arid areas, pioneer plants are often the leading species in the restoration of vegetation, providing the necessary ecological conditions for other plants during their growth<sup>[14]</sup>. For example, plants such as *Psammochloa villosa* (Trin.) Bor and *Artemisia ordosica* can stabilize dunes and provide soil and water for subsequent vegetation.

**Comment [G6]:** Not on the list for common pioneer

## 4 ANALYSIS OF TYPICAL CASES OF ECOLOGICAL VALUE OF PIONEER PLANTS IN THE YELLOW RIVER BASIN

### 4.1 HIPPOPHAE RHAMNOIDES ECOLOGICAL SAND REDUCTION PROJECT IN SHANXI-SHAANXI-INNER MONGOLIA PISHA SANDSTONE AREA

In the northern part of the Loess Plateau, bordering Shanxi, Shaanxi, and Inner Mongolia, there are large patches of Pisha sandstone area, a species peculiar to China. The structure of the Pisha sandstone is loose, shrinking slowly in water and turning to sand in the wind. Once there is a heavy rain, the crops growing on it are washed away by the roots, resulting in a crop-less harvest. Moreover, the Yellow River, which flows through the Pisha sandstone area, carries away a large amount of Pisha sandstone, which are deposited when the river flows through the middle and lower reaches, forming overland river water and endangering other areas<sup>[15]</sup>.

**Comment [G7]:** Proper format

**Comment [G8]:** Please make this more in depth.

The roots of *Hippophaerhamnoides* are strong and numerous, and each plant has an area of 70 - 80 m<sup>2</sup> for sand and soil consolidation. First-order lateral root nodules are abundant and can fix up to 180 kg of nitrogen per square kilometre of *Hippophaerhamnoides* woodland. In addition, a canopy forest can be formed in 3-5 years with low time cost<sup>[16]</sup>. Therefore, planting *Hippophaerhamnoides* soils-water conserving forests in this area can effectively reduce the damage caused by coarse sand loss.

After more than 20 years of ecological construction of *Hippophaerhamnoides*, soil erosion has been effectively curbed, and the effect of sand retention is outstanding. The once barren Pisha sandstone area now presents a landscape of vegetation covering the wasteland and has achieved good ecological, economic and social benefits.

#### 4.2 SALINE ALKALI LAND MANAGEMENT IN THE YELLOW RIVER BASIN

Saline-alkali land is a type of soil where the accumulated salt content exceeds the level of normal cultivated soil, affecting the normal growth of crops. According to incomplete statistics from UNESCO and FAO, the total area of saline-alkali land in China amounts to 9.9\*10<sup>5</sup>km<sup>2</sup><sup>[17]</sup>. The distribution of saline-alkali land is found in the upper, middle and lower reaches of the Yellow River Basin. The Hetao region in Ningxia and Inner Mongolia, the river valley plains in Shaanxi and Shanxi, and the Huang-Huai-Hai plains in Shandong are all very well-known saline-alkali land distribution areas. However, saline-alkali land is an important reserve land resource with great potential for development.

The properties and degree of salinization of different types of saline-alkali soils in the Yellow River Basin vary widely. In the treatment of saline-alkali lands, the principles of specific analysis should be followed. Salt-alkali land in coastal Shandong Province has high groundwater levels, high soil salt content, hard soil structure, poor ventilation and permeability, low vegetation coverage and a relatively fragile ecological environment. In the long process of species selection, salt-tolerant plants such as *TamarixChinensis* Lour., *Salix psammophila*, *Nitrariatangutorum*Bobr., *Phragmites australis*, *Suaeda salsa* (L.) Pall., *Apocynumvenetum* Linn. and *Cynodondactylon*(L.) Pers. have been naturally selected to survive here. *TamarixChinensis*Lour. is an excellent tree species for windbreak and sand fixation and salt-alkali matrix<sup>Error! Reference source not found.</sup>, and its communities also have ecological functions such as resisting coastal erosion, preventing soil erosion, and conserving water, which can enhance ecosystem productivity, nutrient cycling, soil and water conservation, and soil carbon sequestration.<sup>[20]</sup>

#### 4.3 ECOLOGICAL RESTORATION OF ABANDONED MINE

Abandoned Mine refers to serious damage such as surface collapse and pollution caused by land excavation and mineral waste stacking during mineral extraction, which has deprived the land of its original use value<sup>Error! Reference source not found.</sup>. As a result of the large amount of ore residues, waste and pollutants, soil, water and air quality are severely polluted, and the ecological environment is usually severely damaged.

In vegetation restoration in abandoned mines, it is necessary to achieve vegetation coverage as soon as possible and play a role in soil consolidation, so pioneer plants that grow quickly should be selected in the initial stages of restoration. For example, in the abandoned

**Comment [G9]:** This is what I am talking about.

Please do the same exposition on other parts of the paper where you need an in depth analysis or writeup.

mine restoration work in Shanxi Province, *Populus simonii*Carr., *Ulmus pumila* Linn., *Cupressus funebris*, *Robinia pseudoacacia*Linn. and other plantations were planted according to the criteria of barrenness tolerance, salinity tolerance, ability to enrich heavy metals and resistance<sup>[26]</sup>.

In addition, vegetation litter has good water permeability and water retention capacity, and is good at intercepting precipitation, conserving water and reducing soil erosion<sup>[27]</sup>. For example, *Robinia pseudoacacia* has strong roots, strong soil and water retention, and is highly adaptable to soil. *Robinia pseudoacacia*Linn. not only survives in acidic, alkaline, sandy soils, but also plays a good role in ecological restoration<sup>[29]</sup>.

## 5 CONCLUSION

Based on an analysis of the ecological characteristics of pioneer plants in the Yellow River Basin, this paper summarizes the many ecological values of pioneer plants, such as salt-alkali land management, protection from wind and sand, reduction of soil erosion, and promotion of community succession, and performs a case-by-case analysis. The Yellow River Basin covers a wide area and includes harsh cold, arid, barren, saline, and other adverse environments, so there are many pioneer plant species with widespread distribution. The environment of many ecologically fragile areas in the basin, such as the Shanxi-Shaanxi-Inner Mongolia Pisha sandstone region, coastal saline-alkali lands, and abandoned mining areas, has been improved by the action of pioneer plants. In light of this, it is necessary to strengthen research and analysis of the ecological value of pioneer plants and make rational use of them so that they can play a greater role in community succession, ecological restoration and other aspects of the Yellow River Basin and provide more ecological value.

## REFERENCES

- [1] Liu Y. Study on the mechanism of *lyciumbarbarum's* salt tolerance and microbial-fertilizer salt regulation based on osmotic-photosynthetic feedback under brackish water in Hetao irrigation area [D]. Inner Mongolia Agricultural University, 2023. DOI:10.27229/d.cnki.gnmnu.2023.000016. China.
- [2] Zhang Xiaoling. Study on the application of TiO<sub>2</sub> NPs and ZnO NPs improving salt tolerance of rapeseed [D]. Huazhong Agricultural University, 2023. DOI:10.27158/d.cnki.ghznu.2023.001907. China.
- [3] Hu Guangde, Zhou Jiqiang, Da Junshan, Lei Longjv, Zhao Changxing, Li Chaonan, et al. Relationship between understory plant diversity and soil properties of different forests in sandy land of Jinta [J]. Acta Agrestia Sinica, 2023, 31(06):1834-1841. DOI:10.11733/j.issn.1007-0435.2023.06.027 China.
- [4] Qixin D, Zhihui Z, Baoshan W, Min C. Recent Progress on the Salt Tolerance Mechanisms and Application of *Tamarisk* [J]. International Journal of Molecular Sciences, 2022, 23(6):3325-3325.

**Comment [G10]:** Please organize your thoughts in here. Search for a proper format on how to do conclusion.

Also, recommendation at the end is very general and vague. Please make it more useful and should be in line with your analysis.

**Comment [G11]:** You mentioned this ecological values but some are not present on the projects. Make a proper recommendation or maybe a mention on how some pioneer plants can be used for restoration on some parts of Yellow River Basin.

- [5] Yi Bole. Study on the mechanical properties of soil fixation and erosion resistance of the root system of the pioneer sand fixation plant *Astragalus adsurgens* [D]. Inner Mongolia Agricultural University, 2023. DOI:10.27229/d.cnki.gnmnu.2023.000111. China.
- [6] Yang Baobao, Li Jie. Forest ecological service function and its value evaluation in the loess hilly and gully regions of Longdong [J]. Journal of Northwest Forestry University, 2023, 38(06):282-288. DOI:10.3969/j.issn.1001-7461.2023.06.37. China.
- [7] Jin Siyu, Peng Zuodeng. Changes in response of carbon and water physiological parameters of *Robinia pseudoacacia* seedlings to long-term drought and rehydration. [J]. Journal of Beijing Forestry University, 2023, 45(08):43-56. DOI:10.12171/j.1000-1522.20220096. China.
- [8] Xiang Yinchun, Peng Chunhua, Lin Lijin, Song Panhui, Zhou Jiayun, Li Xinyu, et al. Effects of amino acid water-soluble fertilizer on loquat seedling growth and nutrient uptake [J/OL]. Journal of Sichuan Agricultural University, 2024-01-16:1-15. DOI:10.16036/j.issn.1000-2650.202302299. China.
- [9] Li Chun, Liu Jingjing, Zhang Yingbin. Progress of phytoremediation in rare earth mining area [J]. Ecological Science, 2022, 41(03):264-272. DOI:10.14108/j.cnki.1008-8873.2022.03.031. China.
- [10] Zheng Zhilin, Luo Youfa, Zhou Jiajia, Qiu Jing, Wu Xingyu, Gao Bingting, et al. The fraction characteristics of phosphorus in the rhizosphere of four pioneer restoration plants in lead-zinc waste slag yeeds. [J]. Research of Soil and Water Conservation, 2019, 26(03):269-278. DOI:10.13869/j.cnki.rswc.2019.03.040. China.
- [11] Li Linshan, Wang Ziyu, Bai Huihui, Zhang Kaiyu, Liu Mili, Shi Jianguo, et al. Structure and diversity of bacteria communities in rhizosphere soil of four plant species in Mu Us Sandy Land [J]. Journal of Arid Land Resources and Environment, 2024, 38(02):142-149. DOI:10.13448/j.cnki.jalre.2024.038.
- [12] Cheng Yanjun, Zhu Yanfeng, Ma Jing, You Yunnan, Dong Wenxue, Chen Fu. Effects of plant restoration on the functional genes of soil nitrogen cycle in open-pit mine dump [J]. Chinese Journal of Soil Science, 2023, 54(06):1409-1417. DOI:10.19336/j.cnki.trtb.2022082801. China.
- [13] Qian Ling. Study on the release characteristics of heavy metal leaching from gold tailings and the combination of substrate improvement and biological remediation [D]. University of Science and Technology Beijing, 2023. DOI:10.26945/d.cnki.gbjku.2023.000256. China.
- [14] Juan Chen, Yuhu Lin, Ling Fang, Jinfang Li, Suju Han, Yudong Li, et al. Sex-related ecophysiological responses of *hippophaerhamnoides* saplings to simulate sand burial treatment in desertification areas [J]. Forests, 2023, 14(1):101-101.
- [15] Li Huahua, Zhang Maosheng, Feng Li, Zheng Shimei, Du Zhen. Research status and prospect of soil and water loss mechanism and prevention measures in pisha sandstone area [J]. Northwestern Geology, 2023, 56(03):109-120. DOI: 10.12401/j.nwg.2023090. China.
- [16] Yao Nana, Che Fengbin, Li Yonghai, Zhang Ting, Zhang Hui. Nutritional Value and Comprehensive Development and Utilization of *Hipopphaerhamnoides* [J]. Storage and Process, 2020, 20(02):226-232. DOI:10.3969/j.issn.1009-6221.2020.02.038. China.

- [17] Feng Yuqian, Mi Junzhen, Zhao Baoping, Xu Zhongshan, Chen Xiaojing, Li Yinghao, et al. Effect of straw combined with microbial fertilizer on salt content of soil and crops in saline alkali land [J].*Acta Agriculturae Boreali-Sinica*,2023,38(06):101-107.DOI : 10.7668/hbxb.20193788. China.
- [18] Zhao Haodi.Research on ecological basis and resistant plant resources in the process of "three types of natural disasters" in Lankao [D].The University of Henan,2020.DOI:10.27114/d.cnki.ghnau.2020.000753. China.
- [19] Kang Jiapeng, Ma Yingying, Ma Shuqin, Xue Zhengwei, Yang Lili, Han Lu, et al.Dynamic changes of spatial pattern and structure of the *Tamarix ramosissima* population at the desert-oasis ecotone of the Tarim Basin [J].*Acta Ecologica Sinica*,2019,39(01):265-276.DOI: 10. 5846 /stxb201711262112. China.
- [20] Yu Qianjun. Regulation of ecological function of wetland production in the Yellow River Delta: A case study of *T. chinensis* and rice [D].Ludong University,2023.DOI:10.27216/d.cnki.gysfc.2023.000573. China.
- [21] Miri Abbas, Robin Davidson-Arnott. The effectiveness of a single *Tamarix* tree in reducing aeolian erosion in an arid region [J]. *Agricultural and Forest Meteorology*, 2021, 300.
- [22] Fanghan Q, Xinjian H, Xiangmiao S, Yanyu B. Responses of microbial communities and metabolic profiles to the rhizosphere of *Tamarix ramosissima* in soils contaminated by multiple heavy metals [J]. *J Hazard Mater*, 2022, 438(12): 9469.
- [23] Shi Xinxin, He Yuting, Wang Rui, Wang Zhibo, Liu Zhujun, Gao Caiqiu, et al..*Tamarix hispidula* ThEIL1 improves salt tolerance by adjusting osmotic potential and increasing reactive oxygen species scavenging capability [J].*Environmental and Experimental Botany*, 2022, 194(10): 4707.
- [24] Tao Baoxian, Chen Qinghai, Wang Jingdong, Zhang Baohuan, Yuan Haiyan, Chen Yongjin. "Fertile island" of *Tamarix Chinensis* accelerated the carbon decomposition in the coastal wetlands of the Yellow River Delta, China [J].*Catena*, 2022, 211.
- [25] Zhai Quande.Study on the selection and effect of sewage sludge used as ecological restoration matrix in mine wasteland [D].Guizhou University,2020.DOI:10.27047/d.cnki.ggudu.2020.000818. China.
- [26] Hu Landi.Construction and research of Yuanling County ecological corridor based on comprehensive evaluation method [D].Central South University of Forestry and Technology,2023.DOI:10.27662/d.cnki.gznlc.2022.000496. China.
- [27] Zhang Jianhua, Zhang Kun, Liu Yong, Zhang Hong, Zhang Kaiquan, Zhou Xiaoyang, et al.Study on the water-holding capacity of litters from typical artificial forests in reclaimed regions of the opencast coal mine in Shanxi Province [J/OL].*Arid Zone Research*:1-11[2023-11-15].DOI: 10.13866/j.azr.2023.12.16. China.
- [28] Zou Yiqiao, Sun Ouwen, Liu Haiying, Cai Renyue, Lin Song, Ge Hongli, et al. Hydrological characteristics of litters and soils of different forest types in Tiantai County of Zhejiang province [J].*Bulletin of Soil and Water Conservation*,2020,40(03):170-174.DOI:10.13961/j.cnki.stbctb.2020.03.024. China.
- [29] Zhang Kaiquan. Effects of harvesting measures on growth and spatial patterns of multi-rotation *Robinia pseudoacacia* coppice forests [D].Beijing Forestry University,2021.DOI:10.26949/d.cnki.gblyu.2021.000853. China.

[30] FanjingB ,Yuefeng G ,Wei Q .Effects of different soil types on gas exchange parameters and fruits of *Hippophaerhamnoides* ssp. *mongolica* “Ulanshalin” plants[J].PeerJ,2023,11e15264-e15264.

UNDER PEER REVIEW