

**INFLUENCE OF DIFFERENT SPACINGS UNDER VARYING FERTILITY LEVELS ON
PROTEIN CONTENT, PROTEIN YIELD AND AVAILABLE NPK IN SOIL OF SUMMER
GREENGRAM (*Vignaradiata* L.)**

ABSTRACT

A field experiment on “Influence of different spacings under varying fertility levels on protein content, protein yield and available nPK in soil of summer greengram (*Vigna radiata* L.)” was carried out at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar during summer 2020 on loamy sand soil. Twelve treatment combinations comprising of four spacings S₁ (30 × 10 cm), S₂ (30 × 20 cm), S₃ (45 × 10 cm) and S₄ (45 × 20 cm) and three fertility levels viz., F₁ (75% RDF), F₂ (100% RDF) and F₃ (125% RDF) were evaluated in factorial randomized block design with three replications. The results revealed that significantly higher protein yield (256.10 kg/ha) under 30 × 10 cm spacing as compared to other treatments. Among the fertility levels application of 100% RDF (20-40-00 N-P₂O₅-K₂O kg/ha) recorded significantly higher protein content (23.10%), protein yield (241.10 kg/ha). Higher value of available N and P₂O₅ status in soil after harvest was observed in treatment (F₃) 125% RDF of summer greengram (RESULTS not conclusion).

Keywords: -Greengram, Protein content, Yield, Spacing.

1. Introduction

Pulses are wonder present of nature to the living universe and are the real gateway of sustainable agriculture. Food legumes constitute an important dietary ingredient of Indian diet as they supply protein and essential amino acids and play significant role in Indian farming (CITATION). Pulses are the cheapest source of quality protein for the human beings (CIT). The protein hunger is common problem in India, where majority of the people have vegetative diet. Greengram native is India and central Asia (CIT). Greengram (*Vignaradiata*) is cultivated in the countries of India, Burma, Srilanka, Pakistan, China, Fiji, Queens land and Africa. Greengram (*Vignaradiata*) is commonly known as moog, goldengram, mung. It's belonging to family leguminosae. It is third important pulse crop after chickpea and pigeonpea, cultivated throughout India for its multipurpose uses as vegetable, pulse, fodder and green manure crop (CIT)... Its seed is more palatable, nutritive, digestible and non-flatulent than other pulses grown in the world. (CIT).. Spacing plays an important role in contributing to the high yield because thick plant population will not get proper light for photosynthesis and high infestation of diseases (CIT)... On other hand, very low plant population will also reduce the yield (CIT). Due to this reason normal plant population is essential for high yield. Advantage of optimum spacing under irrigated conditions is due to reduced competition for light because when the moisture is lacking, light is no longer (ONLY) limiting factor and advantage of uniform spacing is lost (Ihsanullah et al. 2002). It is the most important non-monetary input, which can be manipulated to attain the maximum production per unit area (Jain and Chauhan, 1988 (recent citation). plays important role for obtaining crop production. For greengram, 20 kg nitrogen (N), 40 kg phosphorus should be given at sowing time (CIT). It is advisable to use fertilizers on the basis of soil test and recommendations, normally 100 kg DAP/ha is sufficient for one hectare the fertilizer should be applied by drilling either at the time of sowing or just before sowing in such

way that they are placed about 2-3 cm below the seed(CIT).. Where is the hypothesis and objective of the paper?

2. Methodology

The field experiment was conducted at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar during summer season of year 2020. The experiment field was fairly leveled and uniform. The soil of the experimental plot was loamy sand in texture and slightly alkaline(soil pH? –soil chemical analysis?). The soil was low in organic carbon and available (N)nitrogen, medium in available phosphorus and high in potassium status(Based on the?). Twelve treatments were laid out in Randomized Block Design with factorial concept with three replications. The four levels of spacings (S_1 -30× 10 cm, S_2 -30 × 20 cm, S_3 - 45 × 10 cm and S_4 -45 × 20 cm) and three levels of fertility levels (F_1 -75% RDF(explain), F_2 - 100% RDF and F_3 -125% RDF). The greengram variety GM 6 was sown on 4th march 2020. (Irrigated?)

3. Results and Discussion

The results obtained from the present investigation as well as relevant discussion have been summarized under following heads.

3.1. Effect of spacings on quality parameters and nutrient studies

3.1.1. Protein content and Protein yield

The data presented in Table 1 revealed that incase of protein content different spacings found non-significant effect on protein content in seed of greengram. However numerically(incorrect based on estastistics)higher protein content (23.33 %) was found in 45 × 20 cm spacing (S_4) followed by 30 × 20 cm spacing (S_2 : 23.18 %). The lowest protein content (22.22 %) was noted with 30 cm × 10 cm spacing (S_1). These might be due to(more emphatic)more competition between plants for growth resources like nutrient, moisture, light under narrow spacing that reduced protein content in seed. Similar results observed by Panchal (2015), Patel (2012), Patel (2017) and Patel *et al.* (2019) in mothbean and incase of protein yield in Table 1 showed that 30 × 10 cm (S_1) spacing gave significantly higher protein yield (256.10 kg/ha) but it was remained statistically at par with 45 × 10 cm (S_3) and 30 × 20 cm (S_2). This might be due to increased seed yield under 30 × 10 cm (S_1) spacing. The lowest protein yield (189.54 kg/ha) was recorded under 45× 20 cm spacing (S_4). Similar results were observed by Panchal (2015) and Patel (2017).

3.1.2. Available NPK in soil after harvest

The data presented in Table 2 indicated that available N, P_2O_5 and K_2O (none application)in soil after harvest did not differ significantly. Though marginally higher available N, P_2O_5 and K_2O (167.56, 41.19 and 261.90 kg/ha) was recorded under 45 × 20 cm (S_4) spacing. This might be due to less competition for nutrient in wider spacing which ultimately lead more nutrient left over in soil. The lowest available nitrogen of 161.19 kg/ha was observed under the spacing of 30 × 20 cm while available P_2O_5 and K_2O less under spacing 30 × 10 cm. Similar findings were reported by Patel (2017) and Ezunget *al.* (2020).

3.2. Effect of fertility levels on quality parameters and nutrient studies

3.2.1. Protein content and protein yield

A perusal of data presented in Table 1 indicated that application of 125% RDF (F_3)registered significantly higher protein content (24.17 %) and protein yield (260.27 kg/ha) in seed of greengrambut it was statistically at par with 100% RDF (F_2). This mainly because of nitrogen is primary component of amino acids which constitute the basis of protein that higher nitrogen in seed is directly responsible for higher protein content in seed of greengram(CIT). The

lowest protein content (20.99 %) and protein yield of 168.77 kg/ha registered with application of 75% RDF (F₁). These results are also close conformity with the finding of Singh (2012) and Meena *et al.* (2020).

3.2.2. Available NPK in soil after harvest

The beneficial effect of varying fertility levels on available N and P₂O₅ in soil after harvest (**lost phrase**). Application of 125% RDF (F₃) resulted Significantly higher available N and P₂O₅ in soil 179.97 and 42.15 kg/ha) and was statistically at par with 100% RDF (F₂). **This might be due** to higher rate of fertilizer application and nitrogen fixation by plants, while non-significant effect of varying fertility levels on available K₂O in soil after harvest of greengram. Similar results were also observed by khavithet *al.* (2019) in cowpea and Jat *et al.* (2012) in greengram.

Table 1: Effect of spacings and fertility levels on protein content in seed and protein yield of summer greengram

Treatments	Seed yield (kg/ha)	Protein content (%) in seed	Protein yield (kg/ha)
Spacings (S)			
S ₁ : 30cm × 10cm	1133	22.22	256.10
S ₂ : 30cm × 20cm	927	23.18	216.01
S ₃ : 45cm × 10cm	1037	22.33	231.86
S ₄ : 45cm × 20cm	812	23.33	189.54
S.Em. ±	107.04	1.22	22.45
C.D. (P=0.05)	314	NS	65.84
Fertility levels (F)			
F ₁ : 75% RDF	808	20.99	168.77
F ₂ : 100% RDF	1044	23.13	241.10
F ₃ : 125% RDF	1081	24.17	260.27
S.Em. ±	92.70	1.05	19.44
C.D. (P=0.05)	272	3.093	57.02
Interaction(S × F)			
S.Em. ±	61.80	0.70	12.96
C.D. (P=0.05)	NS	NS	NS
C.V.%	10.95	5.35	10.05

Use letters to difference the statistics results

The change to graphic designs will valorize the paper

Table 2: Effect of spacings and fertility levels on available N, P₂O₅, and K₂O content in soil after harvest of summer greengram

Treatments	Available nutrient(kg/ha)		
	N	P ₂ O ₅	K ₂ O
Spacings (S)			
S ₁ : 30 cm × 10 cm	163.48	38.16	259.96
S ₂ : 30 cm × 20 cm	161.19	39.93	261.43

S ₃ : 45 cm × 10 cm	162.38	39.30	260.74
S ₄ : 45 cm × 20 cm	167.56	41.19	261.90
S.Em. ±	12.95	2.26	17.15
C.D. (P=0.05)	NS	NS	NS
Fertility levels (F)			
F ₁ : 75% RDF	146.04	36.27	256.69
F ₂ : 100% RDF	164.95	40.51	258.45
F ₃ : 125% RDF	179.97	42.15	267.88
S.Em. ±	11.22	1.96	14.85
C.D. (P=0.05)	32.9	5.742	NS
Interaction(S × F)			
S.Em. ±	7.48	1.31	9.90
C.D. (P=0.05)	NS	NS	NS
C.V.%	7.91	5.70	6.57

Use letters to difference the statistics results

The change to graphic designs will valorize the paper

Conclusion

It is concluded that that higher protein yield (256.10 kg/ha) under 30 cm × 10 cm spacing. Among the fertility levels application of 100% RDF (20-40-00 N-P₂O₅-K₂O kg/ha) recorded higher protein content (23.13%), protein yield (241.10 kg/ha). Higher value of available N and P₂O₅ status in soil after harvest was observed in 125% RDF (This is the results, not a conclusion).

Reference

- Ezung, N. K., Yanthan M., Rajkhowa, D. J. and Jamir T. Performance of greengram as influenced by different levels of vermicompost + rhizobium and spacing on growth, yield and economics under upland terrace condition of north east India. *Research Journal of Agricultural Sciences*.2020;11(2):349-354.
- Ihsanullah, F. H., Akbar, H., Basir, A. and Ullah, N. Effect of row spacing on agronomic traits and yield of mungbean (*Vigna radiata* L. Wilczec). *Asian Journal of Plant Sciences*. 2002;1(4):328-329.
- *Jain, V. K. and Chauhan, Y. S. Performance of greengram cultivars under different row spacings. *Indian Journal of Agronomy*. 1988;33(3): 300-302.
- Jat, R. A., Arvadia, M. K., Tndel, B., Patel, T. U. and Mehta, R. S. Response of saline water irrigated greengram (*Vignaradiata* L.) to land configuration, fertilizer and farm yard manure in Tapi command area of south Gujarat. *Indian Journal of Agronomy*.2012;57(3):270-274.
- Kavitha, M. P., Balakumbahan, R. and Prabukumar, G. Effect of foliar spray and fertilizer levels on growth and yield of vegetable cowpea [*Vigna unguiculata* (L.) Walp.]. *Indian Journal Agricultural Research*. 2019;53(6):745-748.
- Meena, R. S., Verma, T., Verma, S., Singh, A., Kumar, S. and Gurjar, D. S. Influence of organic and inorganic sources of nutrients on growth, yield and quality of mungbean (*Vignaradiata*). *Indian Journal of Agricultural Sciences*. 2020;90(11):2233-2236.
- Panchal, C. C. (2015). Response of summer greengram (*Vignaradiata* L.) to different row spacing and weed management practices under south Gujarat conditions. Unpublished M.Sc. (Agri.) Thesis submitted to Navsari Agricultural University, Navsari (Gujarat).

- Patel, B. J., Patel, H. H. and Gavit, V. C. Effect of spacing and weed management on yield, quality and nutrient uptake of moth bean (*Vigna conitifolia*). *Journal of Pharmacognosy and Phytochemistry*. 2019;8(5):948-952.
- Patel, B. R. (2017). Performance of summer greengram (*Vigna radiata*) cultivars to plant densities under north Gujarat conditions. Unpublished M. Sc. (Agri.) Thesis submitted to Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar (Gujarat).
- Patel, D. K. (2012). Influence of irrigation based on IW/CPE ratio and spacing on growth, yield attributes and yield of summer greengram (*Vigna radiata* L.) cv. Meha under middle Gujarat condition. Unpublished M.Sc (Agri) Thesis submitted to Anand Agricultural University, Anand (Gujarat).
- Singh. Effect of organic and inorganic fertilizers on growth, yield and quality of different varieties of soybean (*Glycine max*). 2012; Accessed 10 August 2015. Available: <https://www.researchgate.net/publication/280829958>