

# **Natural Bioactive Compounds on *Schistosomahaematobium* and *Schistosoma mansoni***

---

## **ABSTRACT**

17

This documentary review aimed to make a synthesis of bioactive plants or natural compounds on schistosomes. We carried out a review from 2000 to 2022, oriented towards the plants in order to search natural compounds bioactive on schistosomes which can be a great contribution for new drugs discovering.

Terms such as “schistosomiasis and medicinal plants”, “natural anti-schistosome compounds”, “phytochemical screening”, “bioactive compounds on schistosomes” were used for the online literature search through the following sites: PubMed (<https://pubmed.ncbi.nlm.nih.gov/>); Google Scholar (<https://scholar.google.com>); ScienceDirect (<http://www.sciencedirect.com/>) and Hinari (<https://extranet.who.int/hinari/fr/journals.php>).

There are many plants already used for the traditional treatment of schistosomiasis. This review found nearly 72 families of plants belonging to 228 botanical species have been listed. Some of these have been studied *in vitro* / *in vivo* and preliminary results have shown their schistosomicidal properties. Moreover, the results of qualitative phytochemical screening and toxicity testing of these natural compounds provide hope for new drug discoveries.

Even if plants efficacy is proven, we need more in-depth investigations to determine their chemical components in order to minimize the undesirable effects and to guarantee their safety by assessing their toxicity.

*18Keywords: Schistosomiasis; Schistosoma; Praziquantel; natural compounds; bioactive*

19

20

## 1. INTRODUCTION

Schistosomiasis or bilharzia, also called “snail fever” is one of the most commonly encountered neglected tropical diseases (NTDs) with high prevalence very often, in low-income countries, especially among schools where hygiene conditions are quite precarious [1, 2]. It is a parasitic illness caused by flatworms of the trematode genus. Three species are generally implicated in human pathology. These include: *Schistosoma mansoni*, which is found in the large intestine, *Schistosoma japonicum* which has a preference for the small intestine and *Schistosoma haematobium* with the bladder environment as its preferred zone [3]. Schistosomiasis is the second endemic parasitosis worldwide after malaria [4, 5]. It affected 230 million people worldwide in 2020 with 80 to 90% of cases in Africa area. This parasitological disease linked both to fecal peril and to certain water activities such as fishing, swimming and agriculture [2], [6–8].

Prevention of schistosomiasis through mass chemotherapy, and therapy are the only alternative currently, as there is no known effective vaccine for large-scale use[9]. Some drugs such as metrifonate (introduced in 1960), Amoscanate and Oxamniquine have been used and abandoned either for their side effects, or for their low efficacy or for their hepatotoxicity [10, 11].

The management of schistosomiasis cases generally uses praziquantel (PZQ), the anthelmintic which is preferentially used to support schistosomiasis cases [12–14]. After nearly a century of research into schistosomiasis drugs, it was not until 1970 that PZQ was developed [15]. This anthelmintic, produced by Merck and used since 1970 [5] is known as Biltricide® [15,16]. It is the only drug approved by the world health organization (WHO) to treat schistosomiasis [17–19]. Its repeated and frequent use could induce cases of resistance or therapeutic failures [19–22]. Although effective, taking PZQ is often difficult in children with swallowing difficulties, while chewing is contraindicated according to recommendations [5]. This parasitic disease remains so far a heavy burden, with high prevalence especially for developing countries despite mass treatments as a preventive measure [1], [12], [20], [23–26]. In rural areas, where living conditions are precarious, the use of PZQ is not systematic due to poverty[19]. People generally use plants to treat schistosomiasis [19], [27, 28]. Chemoprevention is an alternative to get away or to eradicate this NTD, but the cost of this operation constitutes a difficulty for its implementation [29, 30].

Also, the success of this operation must integrate vector control through the use of molluscicides as recommended by the WHO control strategy [31].

The search for new perspectives for the management or even the eradication of this parasitic disease is an imperative for the researchers. Moreover, one of the major challenges of the WHO is to eradicate schistosomiasis by 2030[32,33,34]. Indeed one of the WHO objectives in its traditional medicine strategy 2014-2023 is to promote traditional medicine which plays a complementarity role with conventional medicine [35]. For this purpose, research should be directed towards new horizons by involving the plant environment in the search for new medicines as an alternative to the current use of the treatment of schistosomiasis [36, 37]. A literature review will allow us to have an overview of some studies conducted on plants or natural bioactive substances that can be potential sources for the development of new drugs against schistosomiasis or others diseases[38]. Phytotherapy can be an effective, accessible and inexpensive solution to effectively control schistosomiasis in addition to water treatment and health education [39,40]. Some advocate that we could modify, or even create a new formulation of PZQ in order to make it more bioactive and bioavailable [41]. Developing a new generation of antischistosomal drugs is a necessity for searchers.

However, the lack of information or scientific data on these plants, compound extracts or chemical groups constitutes an obstacle for their use [30], [42–44].

## 2. MATERIAL AND METHODS

We carried out a systematic search for articles deadline from 2000 to 2022 with anti-schistosomal plants, extracts or chemical compounds through the following sites: PubMed (<https://pubmed.ncbi.nlm.nih.gov/>); Google Scholar (<https://scholar.google.com/>); ScienceDirect (<http://www.sciencedirect.com/>) and Hinari (<https://extranet.who.int/hinari/fr/journals.php>)

Terms such as “schistosomiasis and medicinal plants”, “natural anti-schistosome compounds”, “phytochemical screening”, “bioactive compounds on schistosomes” were used for the online literature search. These search terms allowed us to have publications and an overview related to our research topic. The analysis of this documentation allowed us to identify plants and natural compounds active on schistosomes. We noted that *in vitro* and *in vivo* tests were carried out in purebred mice with the different extracts of plants belonging to several families according to the botanical nomenclature.

Our literature review focused on articles published from January 2001 to December 2022 and we sorted articles that best met our selection criteria. The documentation collection period is from September 2022 to April 2023. We also searched the libraries of the WHO and the West African Health Organization (WAHO) on plants with schistosomicidal properties.

The selected articles had to meet one of the following criteria: deal with plants or natural bioactive compounds effective on schistosomiasis; *in vitro* and *in vivo* tests on schistosomes involving natural substances; the phytochemical profile of plant extracts against schistosomiasis; toxicity testing of plant extracts effective against schistosomiasis. We proceeded to a full reading of each article in order to take stock of the research work already done and to establish the link with our research topic.

### 3. REVIEW

#### 3.1. YEARS OF PUBLICATIONS

Between 2001 and 2022, more than two hundred articles were submitted to us based on the keywords and expressions used with the aim of taking stock of the existing plants repertory may possibly be explored in the search for bioactive substances on schistosomes.

Among the documentation proposed through our research, around one hundred articles met our selection criteria (see Figure 1).




Figure 1 Number of publications exploited and interval of year of appearance

The highest rate of publication of the documentation was recorded between 2019 and 2020, followed by the last two years with a slight decline. This could be explained by a motivation of researchers in view of emergence of NTDs in recent years. It could be due to the funding allocated by WHO to researchers in achieving its considerable reduction objectives for burden of morbidity and mortality, or even eradication of NTDs by 2030 [32, 33].

### 3.2. SCHISTOSOMICIDAL PLANTS

The plant environment is increasingly explored in search of new alternatives in the search for solutions against schistosomiasis which has become a real challenge in the response to NTDs [45].

This approach is in line with the objectives of the WHO in its program for NTDs eradication by 2030 [33].

The documentary review allowed us to know that there are a lot of schistosomicidal plants belonging to different botanical families. This plant environment is a potential source of natural substances as an alternative which can contribute to the development of new antischistosomal drugs, and which will strengthen the capacity to treat this parasitosis on a global scale [46]. The plant material listed is the result of ethnobotanical surveys that have made it possible to inventory plants used in traditional medicine and with anthelmintic potential in general and antischistosomal particularly [47, 48]. The listed plants belong to 72 families containing a total of 228 botanical species (see tables 1 and 2). The majority of plants have been listed through ethnobotanical surveys [48], on the shelves of the WAHO [47] and the WHO libraries.

Table1: The different families of plants encountered in the literature

Botanic family	Botanic family	Botanic family
<i>Agavaceae</i>	<i>Dryopteridaceae</i>	<i>Ochnaceae</i>
<i>Amaranthaceae</i>	<i>Ebenaceae</i>	<i>Olacaceae</i>
<i>Amaryllidaceae</i>	<i>Euphorbiaceae</i>	<i>Papaveraceae</i>
<i>Anacardiaceae</i>	<i>Fabaceae</i>	<i>Passifloraceae</i>
<i>Annonaceae</i>	<i>Gentianaceae</i>	<i>Phyllanthaceae</i>
<i>Apiaceae</i>	<i>Humiriaceae</i>	<i>Phytolaccaceae</i>
<i>Apocynaceae</i>	<i>Hypericaceae</i>	<i>Pinaceae</i>
<i>Asclepiadaceae</i>	<i>Iridaceae</i>	<i>Piperaceae</i>
<i>Asparagaceae</i>	<i>Ixonanthaceae</i>	<i>Poaceae</i>
<i>Asphodelaceae</i>	<i>Lamiaceae</i>	<i>Polygalaceae</i>

Botanic family	Botanic family	Botanic family
<i>Asteraceae</i>	<i>Lauraceae</i>	<i>Punicaceae</i>
<i>Bignoniaceae</i>	<i>Loganiaceae</i>	<i>Renonculaceae</i>
<i>Boraginaceae</i>	<i>Loranthaceae</i>	<i>Rhamnaceae</i>
<i>Brassicaceae</i>	<i>Lumiaceae</i>	<i>Rosaceae</i>
<i>Burseraceae</i>	<i>Lythraceae</i>	<i>Rubiaceae</i>
<i>Capparaceae</i>	<i>Malpighiaceae</i>	<i>Rutaceae</i>
<i>Caricaceae</i>	<i>Malvaceae</i>	<i>Sapindaceae</i>
<i>Celastraceae</i>	<i>Melastomataceae</i>	<i>Sapotaceae</i>
<i>Clusiaceae</i>	<i>Meliaceae</i>	<i>Scrophulariaceae</i>
<i>Cochlospermaceae</i>	<i>Menispermaceae</i>	<i>Simaroubaceae</i>
<i>Combretaceae</i>	<i>Moraceae</i>	<i>Solanaceae</i>
<i>Cucubitaceae</i>	<i>Moringaceae</i>	<i>Verbenaceae</i>
<i>Cupressaceae</i>	<i>Myrtaceae</i>	<i>Zingiberaceae</i>
<i>Dipterocarpaceae</i>		<i>Zygophyllaceae</i>

Table 2: Directory of botanical families and species bioactive on schistosomes

<b>Agavaceae family</b>	<b>Asteraceae family</b>	<b>Ebenaceae family</b>
<i>Agave americana</i>	<i>Ageratum conyzoides</i>	<i>Eucleadivinatorum</i>
<i>Agave lophantha</i>	<i>Ageratum conyzoides L</i>	<i>Eucleanatalensis</i>
<i>Asclepiassinaica</i>	<i>Ambrosiamaritima</i>	<b>Euphorbiaceae family</b>
<i>Asiminatriloba</i>	<i>Arteisiaannua</i>	<i>Alchorneacordifolia,</i>
<i>Solanum nigrum</i>	<i>Artemisia absinthium</i>	<i>Antidesmavenosum</i>
<i>Solanum nigrum (L)</i>	<i>Artemisia afra</i>	<i>Euphorbiamauritana</i>
<b>Amaranthaceae family</b>	<i>Baccharis dracunculifolia</i>	<i>Euphorbiaeploides (A)</i>
<i>Chenopodium album (L)</i>	<i>Baccharis trimera</i>	<i>Euphorbiaroyleana</i>
<i>Dysphaniaambrosioides</i>	<i>Berkheyaspeciosa</i>	<i>Euphorbiatirucalli (L)</i>
<b>Amaryllidaceae family</b>	<i>dracunculifolia DC</i>	<i>Jatropha curcas</i>
<i>Allium cepa</i>	<i>Echinopskebericho</i>	<b>Fabaceae family</b>
<i>Allium cepa (L)</i>	<i>Taraxacum officinale</i>	<i>Abrus precatorius</i>
<i>Allium sativum</i>	<i>Vernonia amygdalina</i>	<i>Abrus precatorius (L)</i>
<i>Allium sativum (L)</i>	<i>Vernonia colorata</i>	<i>Acacia nilotica</i>
<b>Anacardiaceae family</b>	<i>Vernonia shirensis</i>	<i>Acacia polyacantha</i>
<i>Lanneabarberi</i>	<b>Bignoniaceae family</b>	<i>Albizia adianthifolia</i>
<i>Lanneachimperi</i>	<i>Kigeliaaethiopum</i>	<i>Albizia antuneziana</i>
<i>Mangiferaindica</i>	<b>Boraginaceae family</b>	<i>Albizia versicolor</i>
<i>Orozoapulcherrima</i>	<i>Alkana orientalis</i>	<i>Bauhinia reticulata</i>
<i>Schinopsisbrasiliensis</i>	<b>Brassicaceae family</b>	<i>Bauhinia variegata L</i>
<i>Scierocaryabirrea</i>	<i>Zillaspinosa L</i>	<i>Bobgunniamadagascarien</i>
<i>Searsialongipes</i>	<b>Burseraceae family</b>	<i>Cassia petersiana</i>
<b>Annonaceae family</b>	<i>Commiphoramolmol</i>	<i>Cassia sieberiana</i>

---

<i>Annona senegalensis</i>	<b>Capparaceae</b> family	<i>Cratylia mollis</i>
<i>Anonidiummannii</i>	<i>Cleomedroserifolia</i>	<i>Dichrostachys cinerea</i> (L)
<i>Asiminatriloba</i>	<b>Caricaceae</b> family	<i>Emilia javanica</i>
<i>Xylopiiiaethiopica</i>	<i>Caricapapaya</i> (L)	<i>Eminiapoladenia</i>
<i>Xylopiiaochrantha</i>	<i>Caricapapaya</i>	<i>Eriosemagriseum</i>
<b>Apiaceae</b> family	<b>Celastraceae</b> family	<i>Erythrinaabyssinica</i>
<i>Apiumgraveolens</i> (L)	<i>Maytenus senegalensis</i>	<i>Erythrina senegalensis</i>
<i>Coriandrumsativum</i>	<b>Clusiaceae</b> family	<i>Isoberliniaangolensis</i>
<i>Foeniculumvulgare</i>	<i>Garcinia huilensis</i>	<i>Piliostigmathonningii</i>
<b>Apocynaceae</b> family	<b>Cochlospermaceae</b> family	<i>Pterocarpus angolensis</i>
<i>Alstonia boonei</i>	<i>Cochlospermumtinctorium</i>	<i>Pterodonpubescens</i>
<i>Calotropisprocera</i>	<b>Combretaceae</b> family	<i>Rhynchosiainsignis</i>
<i>Cryptostegiagrandidflora</i>	<i>Anogeissusleiocarpus</i>	<i>Tetrapleuratetraptera</i>
<i>Diplorynchuscondlocarpoum</i>	<i>Combretumangustifolium</i>	<b>Gentianaceae</b> family
<i>Landolphia kirkii</i>	<i>Combretumglutinosum</i>	<i>Anthocleistanobilis</i>
<i>Neriumoleander</i>	<i>Combretummucronatum</i>	<b>Humiriaceae</b> family
<i>Neriumoleander</i> (L)	<i>Combretummucronatum</i>	<i>Sacoglottisgabonensis</i>
<i>Picralimanitida</i>	<i>Combretumsp</i>	<b>Hypericaceae</b> family
<i>Rauwolfia vomitoria</i>	<i>Terminalia mollis</i>	<i>Harunganamadagascariensis</i>
<b>Asclepiadaceae</b> family	<b>Cucubitaceae</b> family	<b>Iridaceae</b> family
<i>Pergulariatomentosa</i>	<i>Cucurbitapepo</i>	<i>Iris germanica</i> (L)
<b>Asparagaceae</b> family	<i>Cucurbitapepo</i> L	<b>Ixonanthaceae</b> family
<i>Asparagus stipularis</i> (R)	<b>Cucurbitaceae</b> family	<i>Phyllocosmuslemaireanus</i>
<i>Furcraeaselloa</i>	<i>Momordicacharantia</i>	<b>Lamiaceae</b> family
<i>Furcraeaselloa</i> (L)	<b>Cupressaceae</b> family	<i>Coleuskilimandscharica</i>
<i>Ledebouriaovatifolia</i>	<i>Juniperusvirginiana</i> (L)	<i>Ocimumamericanum</i>
<i>Scillanatalensis</i>	<b>Dipterocarpaceae</b> family	<i>Origanummajorana</i>
<b>Asphodelaceae</b> family	<i>Monoteskatangensis</i>	<i>Plectranthusneochilus</i>
<i>Aloe vera</i>	<i>Monoteskerstingii</i>	<i>Plectranthustenuiflorus</i>
	<b>Dryopteridaceae</b> family	<i>Tetradenia riparia</i>
<b>Lamiaceae</b> family	<i>Dryopterisgenus</i>	
<i>Coleuskilimandscharica</i>	<b>Myrtaceae</b> family	<b>Rutaceae</b> family
<i>Ocimumamericanum</i>	<i>Melaleucaarmillaris</i>	<i>Cissusquadrangularis</i> (L)
<i>Origanummajorana</i>	<i>Melaleucaleucadendron</i>	<i>Citrus limon</i> (L)
<i>Plectranthusneochilus</i>	<i>Pimenta racemosa</i>	<i>Citrus reticulata</i>
<i>Plectranthustenuiflorus</i>	<i>Syzygium aromaticum</i>	<i>Citrus sinensis</i> (L)
<i>Tetradenia riparia</i>	<i>Syzygium guineense</i>	<i>Pilocarpusmicrophyllus</i>
<b>Lauraceae</b> family	<b>Ochnaceae</b> family	<i>Zanthoxylumleprierii</i>
<i>Cinnanomoumcamphora</i>	<i>Ochnaschweinfurthiana</i>	<i>Zanthoxylumnaranjillo</i>
<b>Loganiaceae</b> family	<b>Olacaceae</b> family	<b>Sapindaceae</b> family
<i>Buddleia lindleyana</i>	<i>Olaxsubscorpioidea</i>	<i>Dodonaeaviscosa</i>
<i>Strychnos inocua</i>	<i>Ximenia americana</i>	<i>Paulina pinnata</i>

---

---

<i>Strychnos spinosa</i>	<b>Papaveraceae</b> family	<b>Sapotaceae</b> family
<i>Strychnos stuhlmannii</i>	<i>Fumaria officinalis</i>	<i>Manilkara subsericea</i>
<b>Loranthaceae</b> family	<b>Passifloraceae</b> family	<b>Scrophulariaceae</b> family
<i>Tapinanthus bangwensis</i>	<i>Smeathmanniasp</i>	<i>Verbascum sinuatum</i>
<b>Lumiaceae</b> family	<b>Phyllanthaceae</b> family	<b>Simaroubaceae</b> family
<i>Salvia fruticosa</i>	<i>Phyllanthus muellerianus</i>	<i>Harrisonia abyssinica</i>
<b>Lythraceae</b> family	<i>Phyllanthus niruri</i>	<b>Solanaceae</b> family
<i>Punicagranatum</i>	<i>Phyllanthus amarus</i>	<i>Brugmansia unguiculata</i>
<b>Malpighiaceae</b> family	<b>Phyllanthaceae</b> family	<i>Solanum elaeagnifolium</i>
<i>Flabellaria paniculata</i>	<i>Hymenocardia acida</i>	<i>Solanum incanum</i> (L)
<i>Malvaceae</i>	<b>Phytolaccaceae</b> family	<i>Solanum lycocarpum</i>
<i>Brachychiton rupestris</i>	<i>Phytolacca dodecandra</i>	<i>Solanum nigrum</i>
<i>Sida pilosa</i>	<b>Pinaceae</b> family	<i>Solanum solanagine</i> (F)
<b>Melastomataceae</b> family	<i>Pinus canariensis</i>	<i>Solanum tuberosum</i>
<i>Miconia willdenowii</i>	<b>Piperaceae</b> family	<b>Verbenaceae</b> family
<b>Meliaceae</b> family	<i>Piper nigrum</i>	<i>Cleome droserifolia</i>
<i>Azadirachta indica</i>	<i>Piper tuberculatum</i>	<i>Clerodendrum umbellatum</i>
<i>Khaya grandifoliola</i>	<b>Poaceae</b> family	<i>Stachytarpheta cayennensis</i>
<i>Khaya senegalensis</i>	<i>Cymbopogon densiflorus</i>	<b>Zingiberaceae</b> family
<i>Swietenia mahogani</i>	<i>Imperata cylindrica</i> (L)	<i>Aframomum albobolale</i>
<i>Trichilimon adelpha</i>	<b>Polygalaceae</b> family	<i>Curcuma longa</i>
<i>Menispermaceae</i>	<i>Securidaca longipedunculata</i>	<i>Curcuma longa</i> (L)
<i>Tinospora bakis</i>	<i>Eriogonum umbellatum</i>	<i>Zingiber officinale</i>
<b>Moraceae</b> family	<b>Punicaceae</b> family	<b>Zygophyllaceae</b> family
<i>Chlorophora excelsa</i>	<i>Prunicagranatum</i>	<i>Balanites aegyptiaca</i>
<i>Ficus carica</i>	<b>Renonculaceae</b> family	<i>Balanites aegyptiaca</i> (L)
<b>Moringaceae</b> family	<i>Nigella sativa</i>	<i>Fagonia mollis</i> Delile
<i>Moringa oleifera</i>	<i>Pulsatilla chinensis</i>	<b>Rubiaceae</b> family
<b>Myrtaceae</b> family	<b>Rhamnaceae</b> family	<i>Craterispermum caudatum</i>
<i>Callistemon umbellatum</i>	<i>Ziziphus spina-christi</i>	<i>Crossopteryx febrifuga</i>
<i>Callistemon rigidus</i>	<i>Ziziphus spina-christi</i> (L)	<i>Gardenia jovis-tonantis</i>
<i>Callistemon speciosus</i>	<i>Ziziphus mauritiana</i>	<i>Mitracarpus frigidus</i>
<i>Cadabaglandulosa</i>		<i>Mitragynastipulosa</i>
<i>Cajanus cajan</i> (L)	<b>Rosaceae</b> family	<i>Morinda lucida</i>
<i>Callistemon citrinus</i>	<i>Malus domestica</i>	<i>Nauclea latifolia</i>
<i>Callistemon viminalis</i>		<i>Pavetta owariensis</i>
<i>Callistemon viridiflorus</i> (L)		<i>Permacoche verticillata</i>
<i>Eucalyptus citriodora</i>		
<i>Eucalyptus rostrata</i>		
<i>Eucalyptus species</i>		
<i>Eugenia edulis</i>		
<i>Melaleuca styphelioides</i>		

---

### 3.3. PREPARATION OF PLANT MATERIAL

In search of plant extracts bioactive on schistosomiasis, the researchers proceeded to collect plants that had been previously identified by botanist experts. All harvested plants were fully dried in the shade for at least two weeks before being sprayed for the various extractions with different solvents. Virtually all plant parts have been used to search for bioactive substances on schistosomes: leaves, barks, roots, fruits, stems, branches, bulbs, pods, rhizomes, or even the whole plant.

The solvents used for the different extractions were: distilled water, hydro-alcohol, ethanol, methanol (in varying percentages), acetone, chloroform, ethyl-acetate, acetonitrile, ether, methane, olive oil and hexane [49].

### 3.4. ANTISCHISTOSOMAL ACTIVITY OF NATURAL SUBSTANCES

We noticed that there is a varied range of plants with antischistosomal and antimolluscicidal properties [47], [50] and therefore, the ability to neutralize the parasite [51].

All the strains of schistosomes used for the different tests were acquired through research institutions, and *S. mansoni* has been the subject of most studies unlike other species of schistosomes [52, 53]. Obtaining cercariae for *in vitro* and *in vivo* testing required collection of intermediate host snails that were maintained and infected in the laboratory for their excretion of cercariae.

### 3.5. *IN VITRO* STUDY OF THE ANTISCHISTOSOMAL ACTIVITY OF PLANTS

Several researchers have carried out work on the *in vitro* anti schistosome activity of plants. Most authors focused on one or at most three plants and priority was given to *in vitro* tests. *In vivo* testing remained a challenge [54]. Others, on the other hand, have shown the effectiveness of natural substances by taking these two aspects into account [52], [55].

For most authors *in vitro* activity was evaluated in the culture medium RPMI-1640 medium. This medium has been used either in combination with serum or with antibiotics such as Gentamicin, Streptomycin, Penicillin and many others as supplements [56]. All *in vitro* tests were incubated for 24 or 48 hours at 37°C with 5% of CO<sub>2</sub>. The concentrations of the different extracts tested were between 62.25µg/ml and 500µg/ml in search of lethal concentrations of the extract. The lethal concentrations that we observed

varied between 125µg/ml to 500µg/ml [57]. Those concentrations below 125µg/ml had a reduced activity on the parasites because they let perceive the mobility of certain parasites with microscopy [57].

The *in vitro* test was the most common and all results were promising, as they all showed efficacy of crude plant extracts on at least one stage of the parasite and even on the intermediate host snail [52], [54], [58].

The antischistosomal activity of a plant extracts may be linked to a chemical substance or to a combination of chemical groups which should be identified [59, 60].

According to many authors, the lethal dose LC<sub>50</sub> (the dose of the plant substance at which 50% of the parasites die) can vary according to the part of the plant used for the same incubation period: *Pterocarpus angolensis* DC; LC<sub>50</sub> (T=1 hour): leaves 102mg/mL; stem 51.3 mg/mL and bark 33.8mg/mL [53]. Also, the effectiveness of the active substance may depend on the solvent used: *Calotropis procera* (at 25mg/mL): the ethanolic extract caused the death of all adult forms of *S.mansoni* in ½ hour while the aqueous extract didn't become effective until 4 hours later [61, 62] ; *Clerodendrum umbellatum*: LC<sub>50</sub> of the aqueous extract= 805.21µg/mL against 343.10 µg/mL for the methanolic extract [28].

### **3.6. IN VIVO STUDY OF THE ANTISCHISTOSOMAL ACTIVITY OF PLANTS**

As far as *in vivo* tests are concerned, some research conducted has shown some shortcomings in their effectiveness. Some natural compounds with high *in vitro* mortality, on the other hand, exhibited low mortality rates in mice [57]. This is the case of *Allicin* which has been tested *in vitro* and *in vivo* [52]. Others, on the other hand have given interesting results *in vitro* and *in vivo* [63].

Many authors have carried out *in vivo* tests by taking PZQ as control. In the majority of cases for the results obtained, this drug had the greatest parasiticidal activity, compared to the different crude extracts [64–70]. Furthermore, it could also be noted that there was no statistically significant difference between PZQ and other plant extracts in terms of efficacy on schistosomes [71]. Many studies have shown that PZQ, in combination with plant extracts gave very good results in the treatment of schistosomiasis [62], [72]. Others, on the other hand, had as a control group infected and untreated mice in comparison with infected mice and treated with crude plant extracts to research the lethal activity of the extract [52].

Many authors recommend better exploring the *in vivo* activity of plant extracts that have shown conclusive results *in vivo* [54], [73–76]. Others still advocate conducting studies on the biology of intermediate hosts and considering breaking the chain of transmission of schistosomiasis [77]. Also, it would be interesting to

conduct more in-depth studies on plants that have shown their effectiveness on schistosomes using a variety of solvents at concentrations. In addition, the schistosomicidal activity of a plant or a bioactive substance is undoubtedly linked to a chemical substance or group that is important to identify, isolate and quantity ; which also makes it possible to better understand the mechanism of action of these plants or bioactive substance [78, 79].

It should also be noted that some authors have carried out *in vitro* / *in vivo* tests on mice, with other drug groups such as antimalarial drugs [15]. Some molecules have shown higher mortality rates on adults than cercariae: Artemeter at 300mg/kg/day for 2 days caused the death of 85.4-98.3% of *S. mansoni* adults and 70% cercariae. Others, on the other hand, were less effective on adult forms: Chlorambucil at 25 mg/kg for 5 days caused the death of 22.7% of adults against 75.8% of cercariae [37].

### 3.7. PHYTOCHEMICAL SCREENING

We noticed that the compounds of the natural substances were detected qualitatively. Among plants with potential schistosomicidal activity, their active ingredients are mostly unknown. The technic of spectrophotometry through gas chromatography for the detection of chemical compounds of crude extracts was used [51], [71], [80–82].




Figure 2 Proportion in percentages of chemical compounds

It should be noted that some chemical groups are more present than others in terms of percentage as we notice at the figure 2. We have alkaloids and terpenes groups with almost equal proportion, followed by flavonoids and saponins groups with the same trends. The chemical compounds present in high

proportion could be explained by the fact that some authors have sought them especially [51], [71], [80–82]. These include alkaloids, flavonoids, saponins, tannins, terpenoids, steroids and quinones.

Indeed, plants are a huge source of chemical compounds belonging to several chemical groups with a lot of medicinal properties such as anti-oxidant, anti-inflammatory, antimicrobial etc. [83].

### **3.8. CYTOTOXICITY TESTS**

The search for bioactive substances in the plant environment must also take into account their harmful effects or their toxicity. The toxicity test is also an essential step in the process of researching new drugs in general. There are similarities between toxicity test and *in vivo* test. The toxicity test is also carried out in RPMI-1640 medium, supplemented with fetal bovine serum and/or antibiotics. The extracts to be tested are brought into contact with the “Vero cells”. Incubation is also done with 5% CO<sub>2</sub> at 37°C for 24 to 48 hours [55]. Several studies have shown promising results as the toxicity of the herbal extracts used was extremely low or non-existent [84]. Others, however, have shown their effectiveness, but their toxicity remains to be investigated.

The table below (picture 1) is a summary of the work done on some authors on some plants with their antischistosomal properties. We could also note the different parts of plants and solvents used for the extractions process.

## **4. CONCLUSION**

According to many studies, plants could be an alternative in the treatment of schistosomiasis. Several plants are already used by traditional healers, but very few have been analyzed in the laboratory. It is also necessary to study the phytochemical profiles in a qualitative and quantitative approach of these natural compounds for the implementation of new molecules against schistosomiasis through clinical research trials.

## REFERENCES

- 1 Bagayan M, Zongo D, Oueda A, Sorgho H, Savadogo B, Drabo F, et al. Prevalence of schistosomiasis and soil-transmitted helminth infections among schoolchildren in Burkina Faso. 2016;(July). doi:10.1684/mst.2016.0570
2. Angora Etienne K, Menan H, Rey Olivier, Karim Tuo, Andre O Tour, Jean T. Coulibaly. Prevalence and Risk Factors for Schistosomiasis among Schoolchildren in two Settings of Côte d'Ivoire. Published online 2019.
3. Basha H, Mamo H. The Activity of Plant Crude Extracts against *Schistosoma mansoni*. 2021
4. Ly B, Yaro AS, Sodio B. Persistence of urinary schistosomiasis in endemic areas subjected to repeated mass treatments in Mali. 2019;13(February):369-381. French.
5. Amara R, Saadawi S. Praziquantel As The Gold Standard Of Schistosomiasis Control Drug Delivery And Nanotechnological Strategies. 2022;(April).
6. Professor Pierre Aubry. Doctor Bernard-Alex Gaüzère, Schistosomiasis or bilharzia. Published online 2021:1-10.
7. World Health Organization. Weekly epidemiological record. Weekly epidemiological report. 2021;(48):585-596. French.
8. High Authority of Health. Update of medical biology procedures relating to the diagnosis of schistosomiasis (bilharzia). Published online 2017. French.
9. Nicolas BA, Anthropologies S, Isad DD. Antibilharzia Control in Rural Areas in Ivory Coast: Between Modern, Traditional Treatment and Medical Syncretism. Published online 2019:181-190. French.
10. Abaza S. Treatment of Schistosomiasis: From Praziquantel to Development of New Drug Targets Treatment of Schistosomiasis: From Praziquantel to Development of New Drug Targets. 2015;(December).
11. Ferrari MLA, Coelho PMZ, Antunes CMF, Tavares CAP, Cunha AS. Efficacy of oxamniquine and praziquantel in the treatment of *Schistosoma mansoni* infection: a controlled trial. 2003;81(02):190-196.
12. Egie E Enabulele, Roy N Platt, Ehizogie Adeyemi, Esther Agbosua, 1016/j.actatropica.2021.105916.Urogenital
13. Ndegwa FK, Kondam C, Aboagye SY. Traditional Kenyan herbal medicine: exploring natural products' therapeutics against schistosomiasis. 2022;(March). doi:10.1017/S0022149X22000074

14. Corrêa SDAP, Oliveira RN De, Mendes TMF, Rodrigues K. In vitro and in vivo evaluation of six artemisinin derivatives against *Schistosoma mansoni*. Published online 2019.
15. Rodriguez-morales EAJ. Current topics in tropical medicine, 2012
16. Rosimeire Nunes De O Oliveira, Sierpe Jeraldo Veronica De Lourdes, Tarsila Fernaz Frezza, Claudia Melo. Epidemiology, Chemotherapy and New Therapeutics.2014
17. El-nour Mohamed F Abdoul. Antischistosomal Activity of Zingiber officinale, Piper nigrum, and Coriandrum sativum Aqueous Plant Extracts on Hamster Infected with *Schistosoma mansoni*. 2021.
18. Abdal Y and Fadladdin J. Antischistosomal Activity of Origanum majorana, Ziziphus spina-christi, and Salvia fruticosa Plant Extracts on Hamster Infected with *Schistosoma haematobium*. 2021.
19. Ibezim A, Olubiyi OO, Mbah C, Nwodo NJ. Current Computer-Aided Drug Design. 2017;(January). doi:10.2174/1573409913666170119114859
20. In DEM, Bassin LE, Fleuve DU, Abdellahi M, Ndir O, Niang S. Assessment of the prevalence of bilharzia among children aged 5 to 14 years after several years of mass treatment in the Senegal River basin Assessment of schistosomiasis prevalence among children 5 to 14 years old after several years of m. Published online 2016. doi:10.3917/spub.164.0535
21. Balahbib A, Omari N El, Ghchime R, Hakkour M, Bouyahya A. Anti-Schistosomal Activity of Medicinal Plants: 2021;10(3):2352-2360.
22. Ali SA. Natural Products as Therapeutic Agents for Schistosomiasis. doi:10.3923/rjmp.2011.1.20
23. Atalabi TE, Lawal U, Ipinlaye SJ. Prevalence and intensity of genito-urinary schistosomiasis and associated risk factors among junior high school students in two local government areas around Zobe Dam in Katsina State, Nigeria. Parasite Vectors. Published online 2016:1-12. doi:10.1186/s13071-016-1672-5
24. Nyasa L, Mazigo H.D, Uisso C, Kazyoba P, Nshala A. Prevalence, Infection intensity and geographical distribution of schistosomiasis among pre-school and school aged children in surrounding villages. SciRep. 2021;(0123456789):1-11. doi:10.1038/s41598-020-80317-x
25. Joof E, Sanyang AM, Camara Y, Sey AP. Prevalence and risk factors of schistosomiasis among primary school children in four selected regions of The Gambia. Published online 2021:1-15. doi:10.1371/journal.pntd.0009380
26. Natalie G Exum, Simon PS Kibira, Ssenyonga Ronald, Julien Nobili, Alexandra K Shannon, John C, et al. The prevalence of schistosomiasis in Uganda: A nationally

representative population estimate to inform control programs and water and sanitation interventions. 2019;(Ci):1-21.

27. Sparg SG, Staden JVan, Ja AK, Efficiency of traditionally used South African plants against schistosomiasis. 2001;73:209-214.
28. Kenfack Mérimé Christian, Jatsa Hermine Boukeng, Feussom Nestor Gipwe, Emilienne TiengaNkondo, Ulrich MembeFemoe, Christelle Dongmo Tsague, et al. In vitro Antischistosomal Activity of Clerodendrum umbellatum Poir( Labiateae ) Leaves Aqueous Extract and Derived Fractions against Schistosoma mansoni Adult Worms. 2020;22(December):40-49. doi:10.9734/JAMPS/2020/v22i1030199
29. Rugel Anastasia R, Guzman Meghan A, Taylor Alexander B, Frédérick D Chévalier, Reid S Tarpley, Stanton F McHardy, et al. Drugs and Drug Resistance Why does oxamniquine kill Schistosoma mansoni and not S . haematobium and S . japonicum ? IJP Drugs Drug Resist. 2020;13(April):8-15. doi:10.1016/j.ijpddr.2020.04.001
30. Rousseau JJ, Dickens C. Indigenous plants and schistosomiasis control in south africa :molluscicidal activity of some zulu medicinal plants. Published online 2004.
31. Organisation mondiale de la santé. Utilisation Sur Le Terrain de Molluscicides Dans Les Programmes de Lutte Contre La Schistosomiase.2020
32. Guang Zang and Weizhong Yang. Prevention and Control of Infectious Diseases in BRI Countries, 2021
33. Organisation mondiale de la santé. Lutter Contre Les Maladies TropicalesNégligées Pour Atteindre Les Objectifs de Développement Durable.; 2021.French.
34. Alan Fenwick, David Rollinson and Vaughan Southgate, Schistosomiasis Implementation of Human schistosomiasis control: Challenges and prospects. 2006
35. World Health Organization (WHO). WHO Traditional Medicine Strategy 2014-2023. World Heal Organ. Published online 2013:1-76. doi:2013
36. Yousif Fouad, Wassel Gamila, Boulos Loutty, and Therese Labib, Contribution to in vitro screening of Egyptian plants for schistosomicidal activity Contribution to in vitro screening of Egyptian plants for schistosomicidal activity. 2012;0209. doi:10.3109/13880209.2011.625952
37. Bergquist R, Utzinger J, and Keiser J. Controlling schistosomiasis with praziquantel : How much longer without a viable alternative ? Published online 2017:1-10. doi:10.1186/s40249-017-0286-2
38. Evelyn Asante Kwatia, Lord Gyimah, Abraham, Yeboah Mensah, and Arnold Forkuo Donkor. The Ghanaian Flora as a Potential Source of Anthelmintic and Antischistosomal agents, 2021

39. Nelwan Martin L. Prevention of schistosomiasis, 2021
40. David Rollinsona, Stefanie Knoppb, Sarah Levitza, Russell Stothardd J, Louis-Albert TchuemTchuentée, Amadou Garbab, et al. Time to set the agenda for schistosomias elimination.2013
41. Zanolla D, Perissutti B, Passerini N, et al. A new soluble and bioactive polymorph of praziquantel. 2018;(January):1-10.
42. Nicolas J pierre. Plantesmédicinales du Nord de Madagascar Ethnobotaniqueantakarana et informations scientiifiques.2012. French.
43. Ledoux A, Nicolas J pierre, Gallé J baptiste, Gro S. Quelquesplantesemployées dans le Sud Ouest de Madagascar. 2017;(January 2014).French.
44. Hostettmann K, Marston A, Ndjoko K, Wolfender J luc. The Potential of African Plants as a Source of Drugs. Published online 2001:973-1010.
45. Emmanuella Bema Twumasi, Pearl Ihuoma Akazue, Kwaku Kyeremeh, Theresa Manful Gwira, Jennifer Keiser, Fidelis Cho-Ngwa, et al. Antischistosomal ,antionchocercal and antitypanosomal potentials of some Ghanaian traditional medicines and their constituents. Published online 2020:1-21. doi:10.1371/journal.pntd.0008919
46. Diego Ricardo, Galhardo Duarte, Mahomoodally Mohamed Faouzi, Lobine Devina, Suroowan Shano, and Rengasamy Kana R R. Botanical Products in the Treatment and Control of Schistosomiasis : Recent Studies and Distribution of Active Plant Resources According to A ff ected Regions, 2020.
47. Organisation Ouest Africaine de la Santé. Pharmacopee de l 'Afrique de l 'Ouest, 2020. French.
48. Muya Kalunga, Kalonji Monique, and KahumbaByanga,Survoethnobotanique de quelquesplantesutiliséescontre la schistosomiaseurogénitale à Lubumbashi et environs. 2014;(August). doi:10.1007/s10298-014-0877-z
49. Rolland KipréGueyraud, Offoumou Rostand M'Bai, DieudonnéKigbaforiSilué, AllicoDjaman Joseph, and Guédé Noel Zirihi. Technique d ' extraction des flavonoides par solvants/ Technique for flavonoid extraction by solvents. 2019;d:141-142.
50. Esham R El-Seedi, Shaden AM Khalifa, Azza H Mohamed, Nermeen Yorsi, Chao Zhao, Nabil El-Wakeil, et al. Plant extracts and compounds for combating schistosomiasis.2022
51. Bhatia Manisha and Siddiqui Nasir Ali. Abrus Precatorius (L.): An Evaluation of Traditional Herb. 2014;(January 2013).

52. Fakahany Amany Fakahany EI, Abdelmaabood Amina I, Abd Atef Hamid, Amira T Morsi, Abououf Eman A. In Vitro and in Vivo Activity of Allicin against *Schistosoma Mansoni*. Published online 2020.
53. Sadek Gehan S, Harba Nancy M, and Faheem Mona Faheem, In vitro assessment of *Schistosoma mansonicercaricidal* activities of *Solanum nigrum* and *Callistemon citrinus* leaves extracts and cercarial genetic changes by RAPD-PCR. Published online 2019. doi:10.21608/puj.2019.7468.1029
54. Elizondo-luévano Joel H, Castro-ríos Rocio, Vicente Belen, Pedro Fernandez Soto, JulioLopez Aban, Antonio Muro, et al. In Vitro Antischistosomal Activity of the Argemone Mexican Methanolic Extract and Its Main Component Berberine. 2021;16(1):91-100.
55. Nunes Rosimeire Oliveira, Lúcia Vera Garcia Rehder, Adriana Silva Santos Oliveira, Veronica de Lourde Sierpe Jeraldo, Aricio Xavier Linhares, and Silmara Marque Allgretti, Anthelmintic activity in vitro and in vivo of *Baccharis trimera* (Less) DC against immature and adult worms of *Schistosoma mansoni*. *Exp Parasitol*. 2014;139:63-72. doi:10.1016/j.exppara.2014.02.010
56. Aline Pereira Castro, Ana Carolina Alves de Mattos, Neusa Araújo Pereira, Naira Ferreira Anchieta, Matheus Siqueira Silva, Danielle Ferreira Dias, et al. Potent Schistosomicidal Constituents from *Garcinia brasiliensis*. Potent Schistosomicidal Constituents from *Garcinia brasiliensis*. Published online 2015. doi:10.1055/s-0035-1545927
57. Bawazir Salwa S, Ibtihal H Badawi, and Ehssan A. Hassan, Effects of *Dodonaeaviscosa* on experimental schistosomiasis in mice extract. 2020;1(1):11-19.
58. Mohamed Azza M, Metwally NM, and Mahmoud SS. *Sativa* seeds against *Schistosoma mansoni* different stages. 2005;100(April):205-211.
59. Marwa I Ezzat, Mona M Okba, Sherif H Ahmed, Hossny A El-Banna, Abdelbary Prince, Shanaz O Mohamed, et al. In-depth hepatoprotective mechanistic study of *Phyllanthus niruri*: In vitro and in vivo studies and its chemical characterization. Published online 2020:1-23.
60. Klev Gaitan Sikam, Gervais Mouté Appl, Ntabo Virginia Klen, and Wansi Jean Duplex, Chemistry and pharmacological aspects of Aridanin, a lead compound from *Tetrapleura tetraptera* (Fabaceae). 2022;(September). doi:10.53365/nrfhh/152273
61. Azzam M Ahmed, Aboueldahab MM, Haidar Mohammad, and Taha Hoda. In vitro effects of the stem extracts of the plant *Calotropis procera* on *Schistosoma mansoni* adult worms. 2016;(August 2018).

62. Ahmed H Ngm, Marwa Abou Eldahab, Lotfia M Khalil, Rhama BM Juroud, Ahmad M Azzam, Hoda H Taha, et al. Ameliorative effect of calotropisprocera and/or praziquantel on schistosomamansoni infected mice. 2019;49(3):529-538.
63. Behnaz Asianipour and Mehmet Caian, Determination and Comparison of In Vitro Radical Scavenging Activity of Both Garlic Oil and Aqueous Garlic Extracts and Their In Vivo Antioxidant Effect on Schistosomiasis Disease in Mice. 2018;6(7):820-827.
64. Ibrahim Aly, Hanan Taher, and Faten EL-Feky. Antischistosomal and antioxidant protective role of Carica papaya fruit extracts against Schistosoma mansoni. 2020;16(1):107-117.
65. El-shabasyShrouk MAEl-morsy, Sayed AM El- Tantawy, and Eman A El Shabasy. Antischistosomal effects of Ficus carica leaves extract and/or PZQ on Schistosoma mansoni infected mice. J Parasite Dis. Published online 2021. doi:10.1007/s12639-021-01417-7
- 66 Rizk M, Ibrahim N, and El-Rigal N. Comparative in vivo antioxidant levels in Schistosoma mansoni infected mice treated with Praziquantel or the essential oil of Melaleuca armillaris leaves, 2012
67. Hussein Atef, Rashed Samira, Hayawan Ibrahim El, El-sayed Rabab, and Ali Hemat. Evaluation of the anti-schistosomal effects of Turmeric (Curcuma longa) versus praziquantel in Schistosoma mansoni infected mice 2017;12(4):587-596.
68. Ilona Jean A Almanzor, Dianabelle E Clemente, Raffy Jay C Fornillos, Maricel A Gomez and Facundo Rey M, Ladio. In Vivo Trials of Moringa oleifera Lam. Extracts as Antischistosomal Treatment on Schistosoma japonicum Infected Mice. 2014;56:49-56.
69. Jatsa BH, Ngo Sock TE, Tchuem Tchuente AL, and Kamtchouing P. Evaluation of the in vivo activity of different concentrations of clerodendrum umbellatum powder against schistosomamansoni infection in mice. 2009;6:216-221.
70. Domo GA and Kela SL. The efficacy of Maytenus senegalensis (L) extracts on experimentally infected rats with Schistosoma mansoni. 2009;5(4):157-163.
71. Muema Jackson M, Obonyo Meschack A, Njeru Sospeter N, and Mwatha Joseph K. Mwatha. Antischistosomal Effects of Selected Methanolic Plant Extracts in Swiss Albino Mice Infected with Schistosoma mansoni. 2015;9(1):1-11. doi:10.9734/EJMP/2015/16953
72. Nicolaus O Mbugi, Geoffrey Maina and Musa Chacha. Schistosomacidal Activity of LanneaSchimperii and SearsiaLongipes against Chronicles of Pharmaceutical Science Schistosomacidal Activity of LanneaSchimperii and SearsiaLongipes against Cercariae, Schistosomula and Adult Stage of Schistosoma Mansoni. 2019;(June).

73. Abou M Marwa El Dahab, Shahat Sondos M, Mahmoud Soheir SM, and Mahana Noha A. Experimental Parasitology In vitro effect of curcumin on *Schistosoma* species viability, integument ultrastructure and egg hatchability. *Exp Parasitol.* 2019;199(January):1-8. doi:10.1016/j.exppara.2019.02.010
74. Rayan Rubens da Silva Alves, João Gustavo Mendes Rodrigues, Andrea Teles-Reis, Ranielly Araújo Nogueira, Irla Correia Lima Licá, Maria Gabriela Sampaio Lira, et al. Antiparasitic effects of ethanolic extracts of *Piper arboreum* and *Jatropha gossypifolia* leaves on cercariae and adult worms of *Schistosoma mansoni*. doi:10.1017/S003118202000181X2020
75. Zaki Sheir, Amira A Nasr, Ahmed Massoud, Osama Salama, Gamal A Badra, Hassan El-Shennawy, et al. A safe, effective, herbal antischistosomal therapy derived from Myrrh,
- [ PubMed ] 76. Bagayan M, Zongo D, Savadogo B, Oueda A, Sorgho H, Kabre GB, et al. Prevalence of urinary schistosomiasis in a school environment in a peri-urban neighborhood of the city of Ouagadougou, 2014. French.
77. Narin Preyavichyapugdee, Manao Sangfuang, Saowa Park Chaiyapum, Suwatcharaporn Sriburin, Yupa Pootaeng-on, Phiraphol Chusongsang, et al. Schistosomicidal activity of the crude extract of *Artocarpus lakoocha* Published online 2006:1-15.
78. Ali Hussein Abu Almaaty, Hadeer Abd El-Hak Rashed, Maha Farid Mohammed Soliman, Eman Fayad, Fayez Althobaiti, and Nahla Soliman El-Shenawy. Parasitological and Biochemical Efficacy of the Active Ingredients of *Allium sativum* and *Curcuma longa* in *Schistosoma mansoni* Infected Mice. Published online
79. Jerome Munyangia, Lucile Cornet-Vernetb, Michel Idumboc, Chen Lud, Peter Lutgen, Christian Platform, et al. Effect of *Artemisia annua* and *Artemisia afra* tea infusions on schistosomiasis in a large clinical trial. *Phytomedicine.* 2018;51(October):233-240. doi:10.1016/j.phymed.2018.10.014
80. Loyana Silva Godinho, Lara Soares Alexo de Carvalho, Clarissa Fields Barbosa de Castro, Mirna Meana Dias, Priscilla Faria Pinto, Anthony Eduardo Miller Crotti, et al. Anthelmintic Activity of Crude Extract and Essential Oil of *Tanacetum vulgare* (Asteraceae) against Adult Worms of *Schistosoma mansoni*, 2014
81. Tianshan Wang, Xiaopeng Wu, Yanping Liu, Youping Luo, Xiang Li, Yujie Zhou, et al. Chemical constituents from ethanol extract of *Polyalthiarumphii* branches and their cytotoxicity evaluation. *Rev Bras Pharmacogn.* 2018;28(2):235-238. doi:10.1016/j.bjp.2018.03.004
82. Mariana Araújo Vieira do Carmo, Marina Fidelis, Caroline Andolfato Sanchez, Aline Pereira Castro, Ihosvany Camps, Fabio Antonio Colombo, et al. Camu-camu (*Myrciariadubia*) seeds as a novel source of bioactive compounds with promising

antimalarial and antischistosomicidal properties. *Food Res Int.* 2020;136:109334. doi:10.1016/food.2020.109334

83. Ana C Mengarda, Polyana S Mendonca, Christiane S Morais, Ramon M Cogo, Susana F Mazloum, Maria C Salvadori et al. Antiparasitic activity of pipartine (piperlongumine) in a mouse model of schistosomiasis, 2020:105350. doi:10.1016/j.actatropic.2020.105350
84. Hermine BoukengJatsa, Nestor GipweFeussom, Emilienne TiengaNkondo, Merime Christian Kenfack, Nadege Distele Simo, Joseph Bertin Kadji Fassi, et al. Efficacy of methanolic extract of *Ozoroa pulcherrima* schweinf against schistosomamansoni-induced liver injury in mice *rim e.* published online 2018:4-11. doi:10.1016/j.jtcme.2017.

UNDER PEER REVIEW