

Genetic variability studies of vegetable amaranth (*Amaranthus spp.*) for productivity traits

ABSTRACT

Aims: In the present investigation 60 different genotypes of vegetable amaranth were analysed for genetic variability in productivity traits and noticeable range of variability were reported in herbage yield and quality.

Study design: The layout of experimental plot was prepared according to augmented block design II (Federer, 1956).

Place and Duration of study: The experiment was conducted in Department of Vegetable Science, College of Horticulture, Bengaluru, UHS, Bagalkot, Karnataka during summer *i.e.*, from January 2020 to March, 2022.

Methodology: The experiment consisted of 60 vegetable amaranth genotypes that were collected from KRCCH, Arabhavi, NBPGR New-Delhi, TNAU, Coimbtore and some local collections, the whole plot was divided into 4 blocks, wherein, each block contained 19 beds of 1m x 1m dimension, each representing different genotypes including four check varieties. The genotypes were sown in line at the spacing of 20 cm, accommodating 10 plants per line and hence each bed accommodated 50 plants. All growth and yield parameters were recorded in five randomly selected and tagged plants of each genotype and mean was worked out for further statistical analysis. Variability parameters were worked out as per method given by Burton and Devane (1953), heritability in broad sense and genetic advance as percent of mean was calculated according to the formula given Johnson *et al.* (1995).

Results: The phenotypic coefficient of variation and genotypic coefficient of variation was maximum for number of leaves, stem weight (g), Stem to leaf ratio, herbage yield per plant (g), herbage yield per plot (kg), leaf area per plant (cm²) and leaf size (cm²) minimum in case of plant height and stem thickness as shown in table. High heritability was observed for all the traits, except plant height and stem thickness maximum was recorded for Herbage yield per hectare (99.45 %) and minimum was observed for stem thickness (1.07%). High heritability ($h^2 = 99.45\%$) with high genetic advance (GA=92.47%) as percentage of mean was observed for herbage yield per hectare which indicated that additive gene effects were more important for that trait. High genetic advance as per cent of mean was observed for herbage yield per plant (g) however, lowest for stem thickness (0.27)

Conclusion: Results indicates the preponderance of additive genes and selection will be effective for improvement of these traits having high heritability with genetic advance as percent of mean.

Keywords: Genetic advance Vegetable Amaranth, Genetic advance as percent of mean heritability, GCV, Heritability, PCV

1. INTRODUCTION

Amaranthus [*Amaranthus* spp. (L.)] belongs to the family amaranthaceae and genus *Amaranthus*. The word “Amaranthus” has derived from Greek language which means “one that does not wither” or “never fading” (Anjali *et al.*, 2013). The genus *Amaranthus* consists of a large number of species that differ in morphology, yield and also nutrient status. There are two chromosome groups in amaranthus, n=16 and n=17. The species with n=16 are *Amaranthus hypochondriacus* and *A. caudatus* and the species with n=17 are *Amaranthus tricolor*, *A. spinosus*, *A. viridis*, *A. cruentus* and *A. blitum*. Cultivated amaranthus includes at least eight species grown for multiple purposes viz., *Amaranthus caudatus*L., *A. cruentus*L. and *A. hypochondriacus* L. for grain purpose (Costea *et al.*, 2004); *A. dubious* L., *A. blitum*L., *A. hybridus* and *A. tricolor* L. as leafy vegetable (Omondi *et al.*, 2016). Whereas, *A. retroflexus*L. (redroot pigweed), *A. albus* L. (tumble weed), *A. palmeri*S. Wats. (Palmer amaranth) and *A. spinosus* L. (spiny amaranth) are considered as weed species (Erum *et al.*, 2012). *Amaranthus* being a C₄ plant exhibits higher photosynthetic activity and productivity than C₃ plants (Brown, 1999). *Amaranthus* exhibits broad genetic variability in plant type, number of inflorescences, seed colour, earliness, plant height, seed and green matter yield, resistance to pest and diseases and adaptability to soil, climate, rainfall and day length (Mujica and Jacobsen. 2003) etc., that are worldwide known, along with these there exists a great variability in case of nutrient content of amaranthus leaf. *Amaranthus* is a historically sustained nutritional crop which was disappeared for centuries but is now emerging again and showing great potential for food and nutritional security around the world (Das, 2016). As amaranthus can be used as both vegetable and grain crop and has the ability to be grown in varied climatic conditions, it is considered as a good source of nutritional security for a wide array of population (Das, 2016). In America (South America and Central America) where amaranth is thought to have originated (Sauer, 1950; Sauer, 1967, whereas the amaranth leaves and stems are used as vegetables in South Asian and African countries (Grubben and Denton, 2004; Rastogi and Shukla, 2013).

Further, there is a need to intensify the development of leafy amaranth cultivars with nutrient rich quality. To plan appropriate breeding programme and to evolve high yielding cultivars with higher nutritive value, the plant breeders must possess adequate knowledge on nutrient status of the amaranthus leaves and the variability existing. To develop such programmes there is a need to evaluate the available genotypes, hence the present study has been carried out with the objective of genetic variability for those traits in vegetable amaranth genotypes.

2. METHADODOLOGY

The experiment was conducted in Department of Vegetable Science, College of Horticulture, Bengaluru, UHS, Bagalkot, Karnataka during summer *i.e.*, from January 2020 to March, 2022. The experiment consisted of 60 vegetable amaranth genotypes that were collected from KRCCH, Arabhavi, NBPGR New-Delhi, TNAU, Coimbtore and some local collections are BVA-4, BVA-15, BVA-14, BVA-28, BVA-34, BVA-36, BVA-38, BVA-26, BVA-1, BVA-24, BVA-2, BVA-30, BVA-21, BVA-23, BVA-31, BVA-29, BVA-A-1, BVA-A-4, BVA-A-8, BVA-A-9, BVA-A-6, BVA-A-7, BVA-A-10, BVA-A-3, BVA-A-5, BVA, BVA-39, BVA-40, BVA-41, BVA-42, BVA-43, BVA-44, BVA-45, BVA-46, BVA-18, BVA-19, BVA-22, BVA-53, BVA-52, BVA-5, BVA-7, BVA-10, BVA-11, BVA-12, BVA-13, BVA-14, BVA-15, BVA-16, BVA-17, BVA-49, BVA-32, BVA-33, BVA-28, BVA-27, BVA-25, BVA-20, BVA-A-2, BVA-47, BVA-6, BVA-8 and BVA-35 as follows (Table 1). The layout of experimental plot was prepared according to

augmented block design II (Federer, 1956). The whole plot was divided into 4 blocks, wherein, each block contained 19 beds of 1m x 1m dimension, each representing different genotypes including four check varieties. The genotypes were sown in line at the spacing of 20 cm, accommodating 10 plants per line and hence each bed accommodated 50 plants. All growth and yield parameters were recorded in five randomly selected and tagged plants of each genotype and mean was worked out for further statistical analysis. Variability parameters were worked out as per method given by Burton and Devane (1953), heritability in broad sense and genetic advance as percent of mean was calculated according to the formula given Johnson *et al.* (1995).

3. RESULTS AND DISCUSSION

The evidence from Table 2 depicts the significant difference among the vegetable amaranth genotypes for the different characters at 5% significance. The Table 4 depicts the growth and yield status of the leaf amaranth for plant height (23.10-46.00 cm), the maximum plant height was noticed in BVA-35 (46.0 cm) followed by BVA-33 (45.0 cm) and BVA-A-2 (44.80 cm), whereas, the minimum was recorded in the genotype BVA-53 (23.10 cm). Low level of GCV (4.55 %) and medium level of PCV (13.51 %) were noticed. Low heritability (11.35 %) and low genetic advance as per cent of mean (3.16 %) were also observed, Similar results was recorded by Sagar *et al.* (2018) with Plant height ranged from 29.85 (HUB-15) to 43.70 cm (HUB-22) with an average of 34.88 cm. Moderate estimates of GV (12.91) and PV (13.80), moderate GCV (10.30 %) and PCV (10.65 %) with very high heritability (93.50 %), low GA (7.15) and high GAM (20.522 %) were observed for plant height character and previous studies in amaranthus species for plant height (Buhroyet *al.*, 2017; and Diwan *et al.*, 2017). stem thickness (3.80-7.20) maximum was noticed in Arka Varna followed by Arka Suguna (7.12 mm) and BVA-21 (7.0 mm), whereas, the minimum was observed in the genotype BVA-16 (3.80 mm). Low level of GCV (1.27 %) and medium level of PCV (12.24 %) similar results was recorded by Sagar *et al.* (2018) for stem girth that stem girth ranged from 7.35 (HUB-2) to 16.94 mm (HUB-22) with general mean of 12.52 mm. Low estimates of GV (3.26) and PV (4.42), moderate GCV (14.43%) and PCV (16.80%) with high heritability (73.80%), low GA (3.19) and high GAM (25.54%) were observed for stem girth trait. (Panda *et al.*, 2017; Chattopadhyay *et al.*, 2013; Buhroyet *al.*, 2017; and Diwan *et al.*, 2017), Low heritability (1.07 %) and low genetic advance as per cent of mean (0.27 %) were also resulted, days to horticulture maturity (21.00-35.00 days) the genotype BVA-A-4 took minimum days to harvest (21 days), followed by BVA-53 (22 days) and BVA-A-2 (22 days), the maximum number of days to harvest was observed in the genotype BVA-43 (35 days). Days to harvest of leaves exhibited medium range of GCV (10.60 %) and PCV (11.66 %) with high heritability (74.45 %) and moderate genetic advance as per cent of mean was resulted and same was noticed by Adeniji (2018), Bhanuprathap *et al.* (2018) , number of leaves per plant (7.30-20.10 g) the maximum mean number of leaves was noticed in BVA-10 (20.10) followed by BVA-A-1 (19.10) and BVA-A-4 (16.20), whereas, the minimum was recorded for the genotype BVA-A-9 (7.30) ,Moderate level of GCV (19.83 %) and high level of PCV (23.0 %) was noticed coupled with high heritability (74.38 %) and high genetic advance as per cent of mean (35.28 %) leaf weight per plant (2.20-5.40 g), same was recorded by Sagar *et al.*, (2018) that maximum number of leaves per plant was observed in genotype HUB-19 (21.00) followed by HUB-22 (20.58) and minimum number of leaves per plant was observed in genotype HUB-4 (12.04). Low estimates of GV (5.13) and PV (5.59), moderate GCV (14.90%) and PCV (15.54%) with very high heritability (91.80%), low GA (4.47) and high GAM (29.41%) were observed for number of leaves per plant. stem weight per plant (2.80-15.30 g), The highest stem weight per plant was noticed in BVA-5 (15.30 g) followed by BVA-40 (14.80 g) and BVA-23 (11.20 g), whereas, the minimum was observed in the genotype BVA-41 (2.80 g). High level of GCV (38.81 %) and PCV (39.22 %) was observed along with high heritability (97.93 %) and

high genetic advance as per cent of mean (79.23 %). Stem to leaf ratio (0.70-4.30), The maximum stem to leaf ratio was noticed in BVA-5 (4.30 %) followed by BVA-40 (4.0 %) and BVA-26 (2.80 %), whereas, the minimum was observed in the genotype BVA-A-3 (2.20 %). Higher estimates of GCV (35.24 %) and PCV (36.43 %) was noticed. High heritability (93.56 %) and high genetic advance as per cent of mean (70.32 %) were also identified. dry weight of plant (0.02-0.62 g), A maximum dry weight per plant was noticed in BVA-10 (0.62 kg) followed by BVA-A-8 (0.37 kg) and BVA-6 (0.37 kg), whereas, the minimum was observed in the genotype BVA-22 (0.029 kg). Higher estimates of GCV (44.51 %) and PCV (56.15 %) were indicated along with high heritability (62.86 %) and high genetic advance as per cent of mean (72.81 %) were also observed. The traits viz., stem weight per plant, leaf to stem ratio, fresh weight per plant, dry weight per plant, leaf yield per plant and leaf yield per plot were having the higher estimates of heritability and GAM pronouncing the additive gene action and possibility of improvement of the traits through direct selection. These results were in close agreement with that of Sheela *et al.* (2018), Hassan *et al.* (2013) and Bhanuprathapet *et al.* (2018) in the amaranthus crop, Herbage yield per plant (1.80-28.40 g), The maximum herbage yield per plant was noticed in BVA-10 (28.40 g) followed by BVA-38 (17.60 g) and BVA-A-1 (17.60 g), whereas, the minimum was observed in the genotype BVA-22 (1.80 g). High level of GCV (45.01 %) and PCV (45.13 %) coupled with high heritability (62.86 %) and high genetic advance as per cent of mean (72.81 %) were observed. For leaf yield and component traits the considerable variability was noticed in view of wide range values observed for leaf to stem ratio, fresh weight per plant, dry weight per plant, leaf yield per plant and leaf yield per plot, the same results were recorded by Shahiba *et al.* (2020) in amaranthus. Herbage yield per hectare (2.70-42.60 tons) Foliage yield per plant ranged from 112.99 to 175.22 g with an overall mean of 143.60 g per plant. High estimates of GV (133.64) and PV (352.00), low GCV (8.05%) and moderate PCV (13.06%) with moderate heritability (38.00%), moderate GA (14.67) and GAM (10.21%) for foliage yield per plant similar results was recorded by Sagar *et al.*, (2018) minimum foliage yield per plant was observed in HUB-9 (112.99 g) and maximum was observed in HUB-19 (175.22 g) followed by HUB-18 (164.88 g). Leaf weight per plant ranged from 2.20 to 5.40 g with an average of 3.89 g. The highest leaf weight per plant was noticed in Arka Arunima (5.40 g) followed by BVA-38 (5.0 g) and BVA-A-2 (5.0 g), whereas, the minimum was observed in the genotype BVA-A-3 (2.20 g). Moderate levels of GCV (10.99 %) and PCV (14.13 %) were noticed in combination with high heritability (60.47 %) and medium genetic advance as per cent of mean (17.63 %). Leaf size (20.10-47.50 cm²), the leaf size was noticed to be maximum in BVA-6 (47.50 cm²) followed by BVA-21 (45.9 cm²) and BVA-27 (45.1 cm²), whereas the minimum was observed in the genotype BVA-A-9 (20.10 cm²). Higher estimates of GCV (23.34 %) and PCV (24.86 %) was noticed. High heritability (96.09 %) and high genetic advance as per cent of mean (48.25 %) were also identified. and leaf area per plant (29.33-159.84 cm²), the maximum leaf area per plant was noticed in BVA-A-2 (159.84 cm²) followed by BVA-A-1 (119.57 cm²) and BVA-25 (117.37 cm²) whereas, the minimum was observed in the genotype BVA-A-9 (29.33 cm²). High level of GCV (24.9 %) and PCV (30.72 %) was noticed. High heritability (65.71 %) and high genetic advance as per cent of mean (41.64 %) were also examined.,

The higher estimates of genotypic coefficient of variation (GCV) and phenotypic coefficient of variation (PCV) for the growth and yield parameters of leafy amaranth genotypes are presented in Table 3 and Fig. 1. The magnitude of (PCV and GCV) were found high were noticed for the traits, viz., stem weight per plant, stem to leaf ratio, Herbage yield per plant, dry weight per plant and herbage yield per hectare, leaf area per plant and leaf size which was also confirmed in earlier studies on amaranthus by Varalakshmi and Reddy (1997), Ahammed *et al.* (2012), Chattopadhyay *et al.* (2013), Labiba *et al.* (2018) and Sheela *et al.* (2018). Moderate GCV and PCV were noticed for the traits, leaf weight per plant and days to horticultural maturity as noticed by Adeniji (2018), Banuprathapet *et al.*

(2018) in amaranthus crop. High PCV and moderate GCV were noticed for the traits number of leaves per plant and leaf weight per plant which agreed with interpretations of Hassan *et al.* (2013) and Onuoha *et al.* (2019) in amaranthus.

In the present study, traits with higher magnitude of GCV and PCV values, indicated the presence of high degree of variability in the studied accessions which provide ample scope for improvement through selection. Closeness of genotypic and phenotypic coefficients indicates that these characters are least affected by environment and are stable. However, for traits like plant height and stem girth, the PCV estimates were higher than the GCV estimates which suggested that a considerable portion of variability in these traits could be attributed mainly due to environmental influence.

The traits *viz.*, stem weight per plant, stem to leaf ratio, dry weight per plant, herbage yield per plant, herbage yield per hectare, leaf size and leaf area per plant were having the higher estimates of heritability and GAM as shown in table 3 and fig 2 pronouncing the presence of additive gene action and possibility of improvement of the traits through direct selection. These results are in close agreement with that of Anuja and Mohideen (2007), Ahemmed *et al.* (2012), Hassan *et al.* (2013), Bhanuprathapet *et al.* (2018) and Sheela *et al.* (2018) in the amaranthus crop, whereas, the traits like herbage yield per plant and dry weight per plant were controlled by the non-additive gene action as they were of high heritability and low GAM, hence the direct selection based on this traits would become difficult as they indicated by the results of the studies from Chattopadhyay *et al.* (2013) that pronounced the same in amaranthus crop. The traits *viz.*, stem weight per plant, stem to leaf ratio, dry weight per plant, herbage yield per plant, herbage yield per hectare, leaf size and leaf area per plant showed high GCV, PCV, heritability and GAM pronouncing the additive gene action and possibility of improvement of the traits through direct selection (Anuja, 2010 and Panda *et al.*, 2017).

Table 1. List of amaranthus genotypes used in the study and their source of collection

Sl. No.	Genotype	Source of collection
1.	BVA -1	KRCCH, Arbhavi
2.	BVA -2	KRCCH, Arbhavi
3.	BVA -3	KRCCH, Arbhavi
4.	BVA -4	KRCCH, Arbhavi
5.	BVA -5	KRCCH, Arbhavi
6.	BVA -6	KRCCH, Arbhavi
7.	BVA -7	KRCCH, Arbhavi
8.	BVA -8	KRCCH, Arbhavi
9.	BVA -9	KRCCH, Arbhavi
10.	BVA -10	KRCCH, Arbhavi
11.	BVA -11	KRCCH, Arbhavi
12.	BVA -12	KRCCH, Arbhavi
13.	BVA -13	NBPGR, Selection -1 from IC-469722
14.	BVA -14	NBPGR, Selection- 2 from IC-469722
15.	BVA -15	NBPGR, Selection-3 from IC-469722
16.	BVA -16	NBPGR, Selection-4 from IC-469722
17.	BVA -17	Selection from Gagwad local

Sl. No.	Genotype	Source of collection
18.	BVA -18	NBPGR, Selection from IC-469694
19.	BVA -19	NBPGR, Selection from IC-553743
20.	BVA -20	NBPGR, Selection from IC-469558
21.	BVA -21	NBPGR, Selection from IC-551471
22.	BVA -22	Belgaum local collection
23.	BVA -23	Selection from Arka Varna
24.	BVA -24	TNAU, Coimbatore, TN
25.	BVA -25	NBPGR, Selection from IC-469658
26.	BVA -26	NBPGR, Selection from IC-536714
27.	BVA -27	NBPGR, Selection from IC-551472
28.	BVA -28	Selection-2
29.	BVA -29	TNAU, Coimbatore, TN
30.	BVA -30	Selection from IC-469652
31.	BVA -31	Derivative of IC-553743xVA-16
32.	BVA -32	Bellary local collection
33.	BVA-33	Selection-4 local type
34.	BVA -34	Rajgiri red
35.	BVA -35	F6 derivative of Arka SugunaxIC-469645
36.	BVA -36	Selection-5 from local type
37.	BVA -37	F6 derivative of IC-469645xIC-551461
38.	BVA -38	Collection from Chitradurga local
39.	BVA -39	NBPGR Selection from IC-553744
40.	BVA -40	NBPGR Selection from IC-551463
41.	BVA -41	NBPGR Selection from IC-536714
42.	BVA -42	NBPGR Selection from IC-551473
43.	BVA -43	NBPGR Selection from IC-536698
44.	BVA -44	NBPGR Selection from IC-4696605
45.	BVA -45	Local collection
46.	BVA -46	NBPGR Selection from IC-522214
47.	BVA -47	NBPGR Selection from IC-553749
48.	BVA -48	Selection-3 from local type
49.	BVA -49	F4 derivative of CO-1x1/39
50.	BVA -53	Arka amaranth selection
51.	BVA -A-1	NBPGR Selection from IC-469579
52.	BVA -A-2	Kanakpura local collection
53.	BVA -A-3	NBPGR Selection from IC-469601
54.	BVA -A-4	NBPGR Selection from IC-553731
55.	BVA -A-5	NBPGR Selection from IC-541407
56.	BVA -A-6	NBPGR Selection from IC-551472
57.	BVA -A-7	TNAU, Coimbatore, TN

Sl. No.	Genotype	Source of collection
58.	BVA –A-8	NBPGR Selection from IC-553737
59.	BVA –A-9	NBPGR Selection from IC-469579
60.	BVA –A-10	NBPGR Selection from IC-536713
61.	Check-1	Arka Arunima (IIHR, Bengaluru)
62.	Check-2	Arka Varna (IIHR, Bengaluru)
63.	Check-3	Arka Suguna (IIHR, Bengaluru)
64.	Check-4	Arka Samraksha (IIHR, Bengaluru)
IC Series from NBPGR, New Delhi		
BVA: Bengaluru Vegetable Amaranth		

Table 2. ANOVA for growth and yield parameter in vegetable amaranth genotypes

Sl. No.	Source	df	PH	ST	DHM	NOL	LW	SW	S:L	DW	HY/Pt	HY/PI	HY/ha	LS	LA
1.	Treatment (ignoring Blocks)	63	24.08	0.83	10.71 *	5.69 *	0.49 *	8.50 **	0.46 **	0.01	25.50 **	0.58 **	57.50 **	152.21**	514.68*
2.	Treatment: Check	3	12.20	0.49	6.56	3.79	2.44 **	3.09 **	0.64 **	0.01	23.42 **	0.53 **	52.8 **	144.0**	374.13
3.	Treatment: Test vs. Check	1	179.9 *	22.1 **	76.0**	2.16	5.66 **	66.0 **	1.27 **	0.01	179.90 **	4.06 **	405.5 **	376.60**	2230.96**
4.	Treatment: Test genotypes	59	22.04	0.48	9.81 *	5.85 *	0.30	7.80 **	0.44 *	0.01	23.06 **	0.52 **	51.80 **	132.40**	492.74*
5.	Block (eliminating Treatments)	3	18.3	0.33	0.56	1.80	0.08	0.48	0.04	0.003	0.19	0.004	0.47	5.08	77.95

6.	Error	9	19.5	0.48	2.51	1.50	0.10	0.16	0.03	0.004	0.13	0.002	0.28	152.20**
----	-------	---	------	------	------	------	------	------	------	-------	------	-------	------	----------

Significant * (if $P= 0.05$) Highly significant ** (if $P= 0.01$)

df: Degrees of freedom

LW: Leaf weight (g) HY/Pt: Herbage yield per plant (g)

PH: Plant height (cm)

SW: Stem weight (g)

HY/ha: Herbage yield per hectare (t)

SD: Stem thickness (mm) S:L: Shoot to leaf ratio

LS: Leaf size (cm²)

DHM: Days to horticulture maturity DW: Dry weight (kg) LA: Leaf area per plant (cm²)

NOL: No. of leaves per plant HY/PI: Herbage yield per plot (kg)

Table 3. Estimates of genetic variability characteristics for growth and yield parameters in vegetable amaranth genotypes

Sl. No.	Trait	Range	Mean	PCV (%)	GCV (%)	h^2_{bs} (%)	GA	GAM (%)
A.	Growth parameters							
1.	Plant height (cm)	23.10-46.00	34.76	13.51	4.55	11.35	1.1	3.16
2.	Stem thickness (mm)	3.80-7.20	5.67	12.24	1.27	1.07	0.02	0.27
3.	Days to horticulture maturity	21.00-35.00	26.86	11.66	10.06	74.45	4.81	17.91
B.	Green yield and its component traits							
4.	Number of leaves per plant	7.30-20.10	10.51	23.00	19.83	74.36	3.71	35.28
5.	Leaf weight per plant (g)	2.20-5.40	3.89	14.13	10.99	60.47	0.68	17.63
6.	Stem weight per plant (g)	2.80-15.30	7.12	39.22	38.81	97.93	5.64	79.23
7.	Stem to leaf ratio (%)	0.70-4.30	1.82	36.43	35.24	93.56	1.28	70.32
8.	Dry weight of plant (g)	0.02-0.62	0.20	56.15	44.51	62.86	0.15	72.81
9.	Herbage yield per plant (g)	1.80-28.40	10.64	45.13	45.01	99.43	9.85	92.58
10.	Herbage yield per hectare (tons)	2.70-42.60	15.98	45.07	44.95	99.45	14.78	92.47
11.	Leaf size	20.10-	46.84	24.34	23.86	96.09	10.40	48.25

	(cm ²)	47.50							
12.	Leaf area per plant (cm ²)	29.33-159.84	72.27	30.72	24.90	65.71	30.90	41.64	

GCV: Genotypic coefficient of variation (%), PCV: Phenotypic coefficient of variation (%), GAM: Genetic advance as per cent over mean (%), h^2_{bs} : Heritability (broad sense) (%), GA: Genetic advance

Table 4. Performance of vegetable amaranth genotypes for growth and yield parameters

Sl. No.	Genotype/ Traits	PH	ST	NOL	LW	SW	S:L	DW/PI	HY/Pt	HY/PI	HY/ha	DH M	LS	LA
1.	Arka Arunima	31.00	6.40	9.30	5.40	8.70	1.60	0.20	13.30	2.00	20.10	24.00	67.60	70.92
2.	Arka Varna	30.20	7.20	11.10	4.80	9.20	2.00	0.30	16.80	2.50	25.30	23.30	48.10	86.825
3.	Arka Suguna	33.75	7.12	10.82	4.35	10.52	2.43	0.28	11.23	1.68	16.87	26.25	46.09	93.50
4.	Arka Samraksha	31.70	6.90	10.40	4.80	9.50	2.00	0.20	13.80	2.10	20.70	24.50	53.90	87.65
5.	BVA-32	37.00	5.40	10.20	4.10	3.60	0.90	0.17	10.00	1.50	15.00	26.00	81.20	76.54
6.	BVA-33	45.00	5.40	10.20	4.10	6.40	1.60	0.10	7.60	1.20	11.50	28.00	68.60	69.03
7.	BVA-34	40.00	5.40	10.80	4.10	4.80	1.20	0.29	12.60	1.90	19.00	24.00	51.30	65.84
8.	BVA-35	46.00	5.40	8.60	4.10	9.40	2.30	0.11	13.30	2.00	20.00	30.00	53.20	61.46
9.	BVA-36	36.00	5.80	8.40	3.90	9.70	2.50	0.05	2.60	0.40	4.00	30.00	37.40	38.47
10.	BVA-37	43.00	5.10	10.80	4.00	5.20	1.30	0.24	10.60	1.60	16.00	28.00	44.40	83.03
11.	BVA-38	36.00	5.20	9.60	5.00	10.00	2.00	0.18	17.60	2.70	26.50	26.00	53.80	68.22
12.	BVA-39	34.00	5.00	9.20	4.10	8.20	2.00	0.20	6.90	1.00	10.40	30.00	32.00	54.78
13.	BVA-40	32.50	5.20	10.60	3.70	14.80	4.00	0.12	6.80	1.00	10.30	28.00	45.60	77.59
14.	BVA-41	39.00	5.50	10.40	4.00	2.80	0.70	0.31	16.50	2.50	24.80	28.00	45.30	69.18
15.	BVA-42	36.00	5.30	10.60	4.20	8.40	2.00	0.12	6.70	1.00	10.00	28.00	44.20	77.10
16.	BVA-43	31.00	5.10	11.00	4.00	6.40	1.60	0.31	13.30	2.00	20.00	35.00	55.70	77.64
17.	BVA-44	29.20	4.90	9.80	3.30	5.90	1.80	0.32	11.70	1.80	17.60	30.00	48.00	79.73
18.	BVA-45	36.30	4.50	10.40	4.10	7.40	1.80	0.08	4.50	0.70	6.70	24.00	35.60	80.12
19.	BVA-46	39.10	4.80	14.00	3.00	3.60	1.20	0.28	14.80	2.20	22.30	24.00	44.60	91.03
20.	BVA-18	32.50	6.90	7.90	4.20	5.40	1.30	0.04	4.70	0.70	7.00	30.00	48.10	68.90
21.	BVA-19	33.80	6.40	7.50	3.00	6.00	2.00	0.05	5.70	0.90	8.50	26.00	44.10	38.31
22.	BVA-20	37.00	6.80	11.80	4.50	9.00	2.00	0.09	5.80	0.90	8.80	32.00	57.20	96.43

Sl. No.	Genotype/ Traits	PH	ST	NOL	LW	SW	S:L	DW/PI	HY/Pt	HY/PI	HY/ha	DH M	LS	LA
23	BVA-21	37.00	7.00	8.60	4.20	8.40	2.00	0.05	2.00	0.30	3.00	30.00	60.90	78.88
24	BVA-22	40.50	6.00	7.80	3.90	9.70	2.50	0.03	1.80	0.30	2.70	30.00	40.90	48.81
25	BVA-23	40.00	6.00	11.20	4.10	11.20	2.70	0.15	6.70	1.00	10.00	31.00	58.00	87.92
26	BVA-24	39.50	5.30	11.10	3.90	8.20	2.00	0.07	7.30	1.10	11.00	31.00	31.50	48.95
27	BVA-25	42.00	6.70	14.30	4.10	8.90	2.30	0.10	6.30	0.90	9.40	24.00	63.37	117.37
28	BVA-26	35.00	4.70	10.20	4.10	11.20	2.80	0.27	9.60	1.40	14.30	32.00	37.20	51.31
29	BVA-27	33.00	6.00	10.00	4.10	8.20	2.00	0.20	8.50	1.30	12.80	30.00	67.10	90.26
30	BVA-28	36.00	6.30	8.80	4.10	6.10	1.50	0.24	8.30	1.30	12.50	27.00	46.20	55.97
31	BVA-29	34.00	7.00	9.40	4.10	3.20	0.80	0.34	14.00	2.10	21.00	26.00	46.10	60.01
32	BVA-30	31.00	4.50	9.60	4.10	7.30	1.80	0.27	10.30	1.50	15.50	24.00	33.10	43.53
33	BVA-31	33.00	5.80	9.60	4.30	7.80	1.80	0.05	5.70	0.90	8.50	28.00	52.20	68.29
34	BVA-47	34.20	5.20	9.20	4.10	6.50	1.60	0.29	10.80	1.60	16.10	25.00	55.20	69.64
35	BVA-53	23.10	6.10	9.90	2.60	5.00	1.90	0.26	10.00	1.50	15.00	22.00	50.30	60.49
36	BVA-52	34.00	5.00	9.20	4.10	8.20	2.00	0.20	6.90	1.00	10.40	30.00	32.00	97.66
37	BVA-A-1	33.50	5.90	19.10	3.40	4.00	1.20	0.26	10.30	1.50	15.50	24.00	39.40	97.39
38	BVA-A-2	44.80	6.60	20.10	5.00	9.20	1.80	0.31	13.30	2.00	20.00	22.00	63.50	68.34
39	BVA-A-3	32.00	5.40	9.10	2.20	4.00	1.80	0.32	16.50	2.50	24.80	24.00	33.50	73.54
40	BVA-A-4	28.30	5.40	16.20	3.00	4.50	1.50	0.33	11.70	1.80	17.60	21.00	36.00	53.76
41	BVA-A-5	31.00	6.70	14.10	3.40	5.80	1.60	0.11	13.30	2.00	20.00	27.00	48.50	119.57
42	BVA-A-6	36.00	5.50	8.90	3.40	4.80	1.40	0.24	10.60	1.60	16.00	24.00	43.10	159.84
43	BVA-A-7	26.00	5.60	9.20	3.20	4.20	1.30	0.22	17.60	2.70	26.50	29.00	43.10	45.83
44	BVA-A-8	37.00	4.60	12.60	2.60	4.20	1.60	0.37	17.00	2.60	25.50	23.00	32.50	96.88
45	BVA-A-9	30.00	5.80	7.30	3.40	3.80	1.10	0.34	17.30	2.60	26.00	29.00	18.10	100.90
46	BVA-A-10	41.00	5.70	10.60	3.80	6.40	1.70	0.24	10.60	1.60	16.00	30.00	53.70	57.96
47	BVA-1	32.50	5.40	7.90	3.30	5.20	1.60	0.31	13.30	2.00	20.00	24.00	37.00	53.99
48	BVA-2	31.20	5.00	9.90	3.00	3.40	1.10	0.29	7.70	1.20	11.60	23.00	42.70	67.71
49	BVA-48	37.00	5.00	11.40	4.00	2.90	0.70	0.22	10.40	1.60	15.60	24.00	46.00	29.33
50	BVA-4	30.20	5.80	9.90	4.40	3.50	0.80	0.04	4.70	0.70	7.00	24.00	47.90	66.84
51	BVA-5	35.10	5.10	9.00	3.60	15.30	4.30	0.12	7.80	1.20	11.70	25.00	46.80	64.57
52	BVA-6	34.00	5.60	10.00	4.20	11.00	2.60	0.3	17.00	2.60	25.00	28.00	63.00	96.00

Sl. No.	Genotype/Traits	PH	ST	NOL	LW	SW	S:L	DW/PI	HY/Pt	HY/PI	HY/ha	DHM	LS	LA
2.		00	0	10	0	20	0	7	20	0	80	00	40	3
5 3.	BVA-7	33. 90	5.9 0	11. 10	3.9 0	10. 40	2.7 0	0.3 0	14. 70	2.2 0	22. 10	23. 00	41. 40	66.1 8
5 4.	BVA-8	32. 10	6.1 0	10. 10	3.7 0	7.4 0	2.0 0	0.1 5	9.0 0	1.4 0	13. 50	26. 00	59. 50	82.1 7
5 5.	BVA-9	34. 30	6.2 0	12. 00	3.9 0	7.0 0	1.8 0	0.1 8	12. 80	1.9 0	19. 20	29. 00	38. 70	67.4 2
5 6.	BVA-10	35. 50	6.9 0	10. 20	4.9 0	10. 20	2.1 0	0.6 2	28. 40	4.3 0	42. 60	23. 00	57. 20	84.1 7
5 7.	BVA-11	39. 20	5.0 0	8.6 0	3.7 0	7.4 0	2.0 0	0.1 5	9.6 0	1.4 0	14. 40	24. 00	43. 10	59.4 3
5 8.	BVA-12	40. 10	5.3 0	8.4 0	3.8 0	7.6 0	2.0 0	0.1 6	10. 00	1.5 0	15. 00	30. 00	43. 10	41.7 8
5 9.	BVA-13	29. 20	5.0 0	10. 70	4.3 0	5.5 0	1.3 0	0.1 6	12. 40	1.9 0	18. 60	26. 00	32. 50	59.8 8
6 0.	BVA-14	30. 50	5.7 0	9.8 0	4.3 0	10. 70	2.5 0	0.1 1	11. 40	1.7 0	17. 10	30. 00	38. 70	63.1 7
6 1.	BVA-15	26. 90	6.3 0	9.8 0	3.8 0	4.9 0	1.3 0	0.0 8	4.4 0	0.7 0	6.6 0	32. 00	24. 80	52.6 7
6 2.	BVA-16	32. 60	3.8 0	10. 40	4.0 0	7.6 0	1.9 0	0.2 2	14. 70	2.2 0	22. 10	29. 00	40. 60	67.0 0
6 3.	BVA-17	39. 50	5.2 0	14. 00	3.2 0	5.1 0	1.6 0	0.2 0	14. 00	2.1 0	21. 00	25. 00	48. 90	102. 42
6 4.	BVA-49	30. 50	5.1 0	11. 10	4.0 0	5.5 0	1.4 0	0.0 7	3.8 0	0.6 0	5.7 0	26. 00	54. 80	93.2 0
	Mean	34. 76	5.6 7	10. 51	3.8 9	7.1 2	1.8 2	0.2 0	10. 64	1.6 0	15. 98	26. 86	46. 84	73.3 0
	S.E m ±	6.9 9	1.0 9	1.9 4	0.5 5	0.6 4	0.2 7	0.1 1	0.5 7	0.0 8	0.8 4	2.5 0	3.5 6	2.78
	C.D (5%)	15. 81	2.4 7	4.3 8	1.2 3	1.4 4	0.6 0	0.2 5	1.2 9	0.1 9	1.9 0	5.6 6	8.0 6	46.6 9
	CV (%)	12. 93	11. 77	11. 71	8.6 6	5.3 9	9.0 1	33. 51	3.2 3	3.1 6	3.1 6	5.9 7	4.7 0	17.5 1

PH: Plant height (cm) LW: Leaf weight (g) HY/Pt: Herbage yield per plant (g)
ST: Stem thickness (mm) SW: Stem weight(g) HY/ha: Herbage yield per hectare (t)
DHM: Days to horticulture maturity S: L: Shoot to leaf ratio LS: Leaf size (cm²)

NOL: No. of leaves per plant DW: Dry weight (kg) LA: Leaf area per plant (cm²)
S. E m±: Standard error of mean HY/PI: Herbage yield per plot (kg)
C.D: Critical difference (%) CV: Coefficient of variation (%)




Fig. 1: Genotypic and phenotypic co-efficient of variation (GCV % and PCV %) of growth and yield parameters




Fig. 2. Heritability, genetic advance and genetic advance as percent of mean of growth and yield parameter

4. CONCLUSION

Amaranthus being an underutilized crop is one of the major sources of essential nutrients to the human diet. The parameters like stem to leaf ratio, herbage yield per plot, herbage yield per plant, herbage yield per hectare, stem weight, number of leaves

per plant, dry weight, leaf size, showed the higher levels of PCV and GCV highlighting the respective levels of variability for the traits within genotypes. Further, the traits like plant height, stem to leaf ratio, dry weight of plant, herbage yield per plant, per hectare and per plot, stem weight and number of leaves, exhibited higher levels of heritability coupled with the higher GAM indicating the action of additive genes confirming the positive effect of direct selection based on respective traits, while the traits like days to horticultural maturity and leaf weight were with the higher heritability united with the moderate GAM and the trait leaf moisture exhibited the higher heritability linked with the low levels of GAM revealed the effect of non-additive gene action.

REFERENCES

1. Adeniji OT, 2018, Genetic variation and heritability for foliage yield and yield component traits in edible *Amaranthus cruentus* L. genotypes. *Bangladesh J. Agric. Res.*, 43(3): 513-524.
2. Ahammed AU, Rahman MM and Mian M AK. Genetic variability, heritability and correlation study in stem amaranth (*Amaranthus tricolor*). *Bangladesh Journal of Plant Breeding and Genetics* 2012; 25(2): 25-32.
3. Anjali K, Joshi A, Maloo SR and Sharma R. Assessment of the morphological and molecular diversity in *Amaranthus* spp. *African Journal of Agricultural Research* 2013; 8(19): 2307–2311.
4. Anuja S and Mohideen Mk. Variability, heritability and genetic advance studies in amaranthus (*Amaranthus* spp.). *Asian Journal of Horticulture* 2007; 2(1): 63-66.
5. Anuja, S., 2010, D² analysis in vegetable amaranthus. *Asian J. Bio Sci.*, 6(2): 198-200.
6. Banupratap J, Asati BS, Tripathy B, Bairwa PL and Kumar L. Genetic variability for quantitative characters in vegetable amaranthus (*Amaranthus tricolor* L.). *International Journal of Bioresources and Stress Management* 2018; 9(1):093-097.
7. Burton, G. W. and Devane, R. W., 1953, Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. *Agron. J.*, 45:478-481.
8. Brown RH. Agronomic implications of C₄ photosynthesis. *C₄ plant biology*. 1999; 473 - 507.
9. Buhroy, S., Saraswathi, T. and Ramalingam, J., 2017, Genetic basis of yield and quality variations in vegetable amaranth (*A. tricolor*) to identify the promising genotypes. *Int. J. Curr. Microbiol. App. Sci.*, 6(4): 2104-2111.
10. Chattopadhyay, A., Das, S., Rana, N. P., Seth, T. and Dutta, S., 2013, Estimation of genetic attributes relationships and genetic divergence of vegetable amaranths. *Int. J. Plant Breed.*, 7(2): 111-115.
11. Costea M, Weaver SE and Tardif FJ. The biology of Canadian weeds. *Amaranthus retroflexus* L., *A. powellii* S. Watson and *A. hybridus* L. *Canadian Journal of Plant Science* 2004; 84: 631 - 668.

12. Das S. Amaranths: the crop of great prospect. In *Amaranthus: A promising crop of future*, Springer, Singapore. 2016; 13-48.
13. Diwan, I. S., Shukla, N. and Kurrey, V., 2017, Genetic studies in amaranthus germplasm. *Int. J. Curr. Microbiol. App. Sci.*, 6(8): 2459-2470
14. Erum S, Ambreen F, Naeemullah M, Masood S, Qayyum A and Rabbani MA. Genetic divergence in amaranthus collected from Pakistan *Journal of Animal and Plant Science* 2012;22: 653-658.
15. Grubben GJH and Denton OA. *Plant resources of tropical African Vegetables*, PROTA Foundation, Netherlands 2004; 668.
16. Hasan M, Akther CA and Raihan MS. Genetic variability, correlation and path analysis in stem amaranth (*Amaranthus tricolor* L.) genotypes. *Agriculturists* 2013;11(1): 1-7.
17. Labiba Riyaz Shah, Baseerat Afroza, Azra Lateef, SyesBerjees Zehra, Ummiyah H Masoodi. 2018 Genetic studies in vegetable amaranth (*Amaranthus spp.*) under temperate conditions. *Int. J. Chem. Stud.*, 6(4): 724-726.
18. Mujica A and Jacobsen SE. The genetic resources of Andean grain Amaranths (*Amaranthus caudatus* L., *A. cruentus* L. and *A. hypochondriacus* L.) in America. *Plant Genetic Resources* 2003; 133: 41-44.
19. Johnson, H. W., Robinson, H. F. and Comstock, R.S., 1955, Estimation of genetic and environmental variability in soyabean. *Agron. J.*, 41: 314-318.
20. Omondi EO, Debener T, Linde M, Abukutsa-Onyango M, Dinssa FF and Winkelmann T. Molecular markers for genetic diversity studies in African leafy vegetables. *Advances in Bioscience and Biotechnology* 2016; 7(3): 188.
21. Onuoha, N.O., Antia, A., Steel, J., Antia, R. and Lowen, A.C., 2019, Incomplete influenza A virus genomes occur frequently but are readily complemented during localized viral spread. *Nat. commun.*, 10(1): 1-17.
22. Panda, R. K., Mishra, S. P., Nandi, A., Sarkar, S., Pradhan, K., Das, S., Patnaik, A. and Padhiary, A. K., 2017, Genetic variability and varietal performance in vegetable amaranthus (*Amaranthu ssp.*). *J. Pharmacognosy & Phytochem.*, 6(6): 1250-1256.
23. Rastogi A and Shukla S. Amaranth: A new millennium crop of nutraceuticals values. *Critical Review of Food Science and Nutrition* 2013; 53(2): 109–125.
24. Sauer, J. D., 1950, The grain Amaranths: A survey of their history and classification. *Ann. Mo. Bot. Gard.*, 37(4): 561-632.
25. Sagar, K.P., Hanchinamani, C.N., Imamsaheb, S.J., Nishani, S. and Ramanagouda, S.H., 2018. Genotypic correlation coefficients among growth, yield and quality parameters in *Amaranthus* genotypes (*Amaranthus tricolor* L.). *Int J Curr Microbiol App Sci*, 7(8), pp.2708-2714.
26. Sauer, J. D., 1967, The grain Amaranths and their relatives: a revised taxonomic and geographic survey. *Ann. Mo. Bot. Gard.*, 54(2): 103–137.
27. Sheela, M. N., Revanappa, S., Mansur, Satish, D. and Ajjappalavar, P. S., 2018, Estimates of genetic variability for growth attributes in amaranth (*Amaranthus* L.) species. *Int. J. Curr. Microbiol. App. Sci.*, 7(7): 1495-1500.

28. Shahiba, A. M., Thomas, B. and Chacko, A., 2020, Evaluation of thirty amaranthus genotypes (*Amaranthus tricolor* L.) for different biometric characters. *Int. J. Curr. Microbiol. App. Sci.*, 9(8): 1621-1631
29. Varalakshmi, B. and Reddy, V. V. P., 1997, Variability, heritability and correlation studies in vegetable amaranth. *Indian J. Hortic.*, 54(2): 1

UNDER PEER REVIEW

UNDER PEER REVIEW