

## ***Original Research Article***

### **Performance of Turmeric (*Curcuma longa* L.) and Pigeon pea (*Cajanuscajan* L.) intercropping system under North Eastern Ghat Zone of Odisha, India**

Formatted: Font: 12 pt

Formatted: Font: 12 pt

#### **ABSTRACT**

Turmeric (*Curcuma longa* L.) is an important cash crop grown by tribal farmers of Odisha for their livelihood.

Despite the good agro-climatic condition, the productivity of turmeric in Odisha is much below (2.4 t/ha) as compared to national average (5.1 t/ha). Therefore, the present study was carried out during 2021-22 and 2022-23 season to evaluate the performance of cropping system to find out suitable planting geometry and row proportions. The different treatments taken were viz. T<sub>1</sub>. Sole Turmeric, T<sub>2</sub>- Sole Pigeon pea T<sub>3</sub>. Turmeric + Pigeon pea (3:1) – one row of pigeon pea after three rows of turmeric (Additive) T<sub>4</sub>- Turmeric + Pigeon pea (5:1)- one row of pigeon pea after five rows of turmeric (Additive) T<sub>5</sub>- Turmeric + Pigeon pea (3:1) -one row of pigeon pea after three rows of turmeric (Replacement) T<sub>6</sub>- Turmeric + Pigeon pea (5:1)- one row of pigeon pea after five rows of turmeric (Replacement) T<sub>7</sub>- Turmeric + Pigeon pea (6:2)- Two rows of pigeon pea after six rows of turmeric (Replacement) T<sub>8</sub>- Turmeric + Pigeon pea (10:2)- one row of pigeon pea after ten rows of turmeric (Replacement).- Sole crop of turmeric fresh rhizome (106.04 q/ha) and pigeon pea grain (14.07 q/ha) produced the highest yield when compared with other intercropping systems. Turmeric + Pigeon pea (10:2) geometry gave the highest net return and benefit cost ratio over other intercropping system.

Keywords: Turmeric, pigeon pea, rhizomes, tribal, livelihood.

#### **1.0.1 INTRODUCTION**

Turmeric is an important commercial spice crop grown in India since ancient times and it is named as “Indian saffron”. It is known as the “golden spice” as well as the “spice of life.” It reached China by 700 AD, East Africa by 800 AD and West Africa by 1200. It was introduced to Jamaica in the 18th Century and started becoming popular throughout the world. Turmeric was probably cultivated at first as a dye, and then became valued as a condiment as well as for cosmetic purposes. The Arab traders took turmeric to Europe in 13th century. During his travels in China in 1280, Marco Polo was so impressed by turmeric that he mentioned it as a vegetable that has properties of saffron, but it is not really saffron. Turmeric is mainly known as a spice all over the world. It is used as a spice in curry powder,

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Black

Formatted: List Paragraph, Outline numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0" + Indent at: 0.25"

chicken bouillon, sauces, gravies, dry seasoning, baking mixes, processed cheese pickles, relishes, breadings, soups, beverages and confections (Sasikumar, 2001). Turmeric is now grown in countries like India, China, Pakistan, Bangladesh, Vietnam, Thailand, Philippines, Japan, Korea, Sri Lanka, Nepal, South Pacific Islands, East and West Africa, Malaysia, Caribbean Islands and Central America. The world production of turmeric is 800,000 tonnes in which India holds a share of approximately 75-80% and consumes around 80% of its own production. India is by far the largest producer and exporter of turmeric in the world. Indian turmeric is considered the best in the world market because of its high curcumin content (6.7%). Turmeric occupies about 6% of the total area under spices and condiments in the country. During 2012-2013, the country produced 9,92,900 tonnes of turmeric from an area of 1,95,100 ha (Anonymous, 2013). During the period ~~between~~ from April 2011 to January 2012, India exported 67,000 tonnes of turmeric valued at Rs 6,438 million. From India's total turmeric exports, 65% were exported to UAE, USA, Japan, Sri Lanka, UK, and Malaysia. Andhra Pradesh, Tamil Nadu, Odisha, Karnataka, West Bengal, Gujarat, Meghalaya, Maharashtra and Assam are important states cultivating turmeric. Andhra Pradesh alone occupies 35.0% of the area and 47.0% of the production.

In Odisha, turmeric is an important cash crop grown by tribal families for their livelihood and more than 50% of this crop grower is tribal. Odisha contributes about 21% of India's turmeric cultivation in terms of area and Kandhamal makes up for over 50% of the state's share. Odisha produced 59,361 t from 247,333 ha. Kandhamal district stands first in turmeric area as well as production (28,828 t from 11,088 ha). Koraput is the second largest producing district (7,761 t from 3,168 ha) followed by Nayagarh (5,343 t from 2,473 ha) and Keonjhar (2,937 t from 1,224 ha). Turmeric is a cash crop grown by Kondhatribes of Kandhamal district and Langi Kondha of Gajapati district. The crop is grown in back-yard by the tribals with their indigenous methods of crop production. The crop is mostly produced organically and the farmers get a good return. In view of the potential of turmeric, the state proposed for the Agri Export Zone (AEZ). The zone covers the district of Kandhamal (Ray, 2007). Turmeric is also credited with religious and magical rites in India and certain South-East Asian countries. Traditionally turmeric has been used in India for treatment of a variety of human and veterinary ailments, as a natural dye, as well as preparation of delicious dishes. Though traditional Indian Ayurvedic and Siddha systems of medicine have recognized the medicinal value of turmeric in its crude form since very ancient times, the few decades have witnessed extensive research interests worldwide in the biomedical activity of turmeric and its compounds. Thus

Curcuma is now gaining importance all over the world as a mighty cure to combat a variety of ailments, as the genus carries molecules credited with anti-inflammatory, hypocholesteremic, choleraic, antimicrobial, antirheumatic, antifibrotic, antivenomous, antiviral antidiabetic, antihepatotoxic and anticancerous properties as well as insect repellent activity (Chattopadhyay *et al.*, 2004). Turmeric can be grown in Odisha as an intercrop in coconut and areca nut plantations. Turmeric is the third-largest spice exported from India. In terms of quantity and value, it accounts about 12% and 5% respectively.

Due to the rapidly increasing population of the country, the pressure on agricultural land to get maximum yield per unit is increasing continuously. On the other hand, due to climate change, there is also a significant change in the uncertainty of rainfall. Intercropping farming is considered very useful in dealing with such challenges. In tur growing areas of the country, farmers often plant tur on the bunds of paddy fields. This gives less yield. But if turmeric farming is done with tur, then farmers can earn double.

Intercropping Farming is such a scientific technique, which has been found to be very effective in increasing income of the farmers. In this regard, it is recommended by agricultural experts to do turmeric with tur or ginger with tur or turmeric with drumstick or inter-cultivation of ginger and turmeric with papaya. With this modern technology, farmers can increase their income apart from reducing the risk of farming.

Actually, turmeric can be cultivated easily even in shady environment. Turmeric is a popular medicinal and spice crop. That's why it gets good price in the market. Cultivation of turmeric with tur as an intercropping crop yields income from both the costly crops. The inclusion of tur cultivation in the crop rotation maintains the fertility of the soil, because the roots of pulse crops provide natural fertility to the soil by absorbing nitrogen directly from the atmosphere.

[Here add a paragraph talking about Pigeon pea and some literature about its intercropping with turmeric or other crops. Remember your research was about inter cropping turmeric with pigeon pea](#)

## 2. MATERIALS AND METHODS

One experiment was conducted at ~~RRITS~~, G.Udayagiri, Kandhamal, Odisha to study the performance of cropping

**Comment [p1]:** Write in full and abbreviate in brackets

system to find out suitable row proportions of turmeric and arhar with additive and replacement series. The

~~Experimental Design~~ was Randomized Complete Block Design (RCBD), with three replications. ~~RBD Replication:~~

~~three~~ ~~Treatments: 8~~. The experiment was conducted in medium land site and the ~~The soil is~~ sandy loam in

texture, pH 5.39, low available P<sub>2</sub>O<sub>5</sub>, organic carbon-5.4g/kg, and available K<sub>2</sub>O is 160kg ha<sup>-1</sup>. ~~Observations on~~

~~growth and yield parameters were recorded in each replication and the mean obtained were used for statistical~~

~~analysis (Panse and Sukhatme, 1985). The treatments used were:~~

### ~~Treatments:~~

T<sub>1</sub>. Sole Turmeric,

T<sub>2</sub>- Sole Pigeon pea

T<sub>3</sub>. Turmeric + Pigeon pea ( 3:1) – one row of pigeon pea after three rows of turmeric ( Additive )

T<sub>4</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric ( Additive )

T<sub>5</sub>- Turmeric + Pigeon pea ( 3:1) -one row of pigeon pea after three rows of turmeric (Replacement )

T<sub>6</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric (Replacement )

T<sub>7</sub>- Turmeric + Pigeon pea ( 6:2)- Two rows of pigeon pea after six rows of turmeric (Replacement )

T<sub>8</sub>- Turmeric + Pigeon pea ( 10:2)- one row of pigeon pea after ten rows of turmeric (Replacement )

~~Recommended Ecosystem~~ ~~Rain fed Upland~~ ~~Parameters measured included plant height, number of~~

**Formatted:** Font: Not Bold

~~branches/plant at harvest, number of pods/plant, number of seeds/pods, 100 seed mass, rooting depth, mass of~~

~~mother rhizome, mass of primary rhizome, mass of secondary and tertiary rhizome.~~ ~~The data was subjected to~~

**Formatted:** Font: Not Bold

~~Analysis of Variance (ANOVA) in Statistical Analysis System (SAS) software and the mean were separated by~~

~~Duncan Test.~~

**Comment [p2]:** I assume you subjected your data to ANOVA using a specific software. Please specify the statistical software you used and also specify which Test you used for mean separation, eg Duncan, tukey's, fisher etc

### 3. RESULTS AND DISCUSSION EXPERIMENTAL FINDINGS

The morphological characters of turmeric were found to be significantly influenced by various intercrops. The pigeon-pea as pure crop recorded significantly higher values for plant height (174.9 cm), no of branches/plant at harvest (15.3), no of pods /plant (132.3), no of seeds /pod (4.7), 1000 seed weight-mass (82.11g), mass weight of secondary rhizome /plant (405.1), yield of 14.07 qha<sup>-1</sup> and B:C ratio-1.40. The turmeric as pure crop recorded plant height-78.30cm, Rooting depth-13.11cm, length of leaves—36.31cm, weight of mother rhizome-34.40g, weight of primary rhizome-132.30g, and weight of secondary and tertiary rhizome-75.20g, total weight clump-241.80g, fresh turmeric rhizome-106.04 qha<sup>-1</sup> and B:C ratio -1.37. The turmeric:Pigeonpea(10:2) in T8 having one row of pigeon pea after ten rows of turmeric (Replacement) recorded significantly higher values for plant height, 74.59cm, Rooting depth, 12.46cm, length of leaves—35.46cm, mass weight of mother rhizome 30.60g, mass weight of primary rhizome 127.40g, and mass weight of secondary and tertiary rhizome, 64.20g, total weight clump 221.20g, fresh turmeric rhizome 84.02q ha<sup>-1</sup>, Grain yield of Pigeon pea (qha<sup>-1</sup>) 4.54 and B:C ratio (2.11 ha<sup>-1</sup>), dry turmeric rhizome yield (16.81 qha<sup>-1</sup>), dry turmeric rhizome equivalent yield was 19.91 qha<sup>-1</sup>, and Islam *et al.* (2016) indicated that higher biomass production and consequently more efficient use of land and available resources under intercropping than under sole cropping contributed to the higher turmeric yield. Similar results were obtained by Balashanmugam *et al.* (1988) and Narayanpur and Sulikeri (1996) in turmeric and Kumar *et al.* (2018) in ginger. Paraye *et al.* (2014) reported that ginger equivalent yield and net return was higher in ginger (Raigarh local) + turmeric (Sudershana) in 1:1 row ratio intercropping system.

When the base crop and intercrops were adequately fertilised then there is no yield reduction in the main crop (Nayar and Suja, 2004). Similar results were observed by Kannan *et al.*, (2001), Chattopadhyay *et al.*, (2008) in elephant foot yam intercropped with vegetables. It was interpreted by Singh *et al.*, (2013) that less competition for resources and better scope of intercultural operations at early growth stages were the favourable points, which might have triggered the process of partitioning photosynthates from source to sink resulting in higher yield of main crop. Further, Das and Maharana (1995) explained that elephant foot yam does not compete for light because it is able to tolerate shade. Yield of inter crops - Int. J. Curr. Microbiol. App. Sci (2020) Special Issue 11: 1247-1253-1250 yam sole crop. The higher productivity of the intercrop system compared to the sole crop might have resulted from

complementary and efficient use of growth resources by the component crops as explained by Li *et al.*, (2006).

Willey (1979) also elucidated that mixed cropping was more efficient and productive than sole cropping because of higher combined yield and better energy use efficiency. Ravindranet *et al.*, (2006) also advocated intercropping of tuber crops for better remuneration.

**Table1.**

**Effect of planting geometry on growth and yield attributes of Pigeon pea in turmeric + pigeon pea intercropping system(Pooled over 2021-22 &2022-23)**

Treatments	Plant height (cm) at harvest	No of branches <sup>-1</sup> plant at harvest	No of field pods/plant	No of seeds/for pods	1000- Seed weight (g)
T <sub>1</sub>	-	-	-	-	-
T <sub>2</sub>	174.9	15.3	132.3	4.7	82.11
T <sub>3</sub>	169.4	14.6	130.3	4.6	82.10
T <sub>4</sub>	161.2	14.0	125.7	4.4	81.78
T <sub>5</sub>	166.4	14.5	130.3	4.5	81.98
T <sub>6</sub>	156.1	13.5	122.2	4.3	81.55
T <sub>7</sub>	165.2	14.1	127.8	4.5	81.78
T <sub>8</sub>	151.5	13.1	120.4	4.1	81.34
SEm (±)	5.29	0.46	2.78	0.22	1.65
<u>CD</u> CV(0.05)	10.78	0.95	5.77	0.46	NS

Formatted: Space After: 0 pt, Line spacing: single

T<sub>1</sub>. Sole Turmeric,

T<sub>2</sub>- Sole Pigeon pea

T<sub>3</sub>. Turmeric + Pigeon pea ( 3:1) – one row of pigeon pea after three rows of turmeric ( Additive )

T<sub>4</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric ( Additive )

T<sub>5</sub>- Turmeric + Pigeon pea ( 3:1) -one row of pigeon pea after three rows of turmeric (Replacement )

T<sub>6</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric (Replacement )

T<sub>7</sub>- Turmeric + Pigeon pea ( 6:2)- Two rows of pigeon pea after six rows of turmeric (Replacement )

T<sub>8</sub>- Turmeric + Pigeon pea ( 10:2)- one row of pigeon pea after ten rows of turmeric (Replacement )

**Table2.**

**Effect of planting geometry on growth and yield attributes of Turmeric in turmeric + pigeon pea intercropping system(Pooled over 2021-22 &2022-23)**

Treatments	Plant height (cm)	Rooting depth (cm)	Length of leaves (cm)	Weight of mother rhizome (g)	Weight of primary rhizome (g)	Weight of secondary and tertiary rhizome (g)	Total weight clump (g)
T <sub>1</sub>	78.30	13.11	36.31	34.40	132.30	75.20	241.80
T <sub>2</sub>	-		-	-	-	-	-
T <sub>3</sub>	63.75	8.70	28.45	23.98	107.56	46.76	178.24
T <sub>4</sub>	67.68	11.53	34.56	26.21	126.34	60.67	216.22
T <sub>5</sub>	66.21	9.89	30.18	25.21	119.23	54.45	201.22
T <sub>6</sub>	67.37	10.57	32.10	28.34	121.01	56.28	205.63
T <sub>7</sub>	65.34	9.23	27.42	24.11	117.34	52.75	194.23
T <sub>8</sub>	74.59	12.46	35.46	30.60	127.40	64.20	221.20
SEm (+)	2.15	0.54	1.17	1.24	3.68	2.46	4.55
CD (0.05)	6.30	1.59	3.44	3.64	10.78	7.20	13.34

**Comment [p3]:** Its best to measure leaf area. You can therefore delete this parameter if you do not have leaf area data

T<sub>1</sub>. Sole Turmeric, T<sub>2</sub>- Sole Pigeon pea T<sub>3</sub>. Turmeric + Pigeon pea ( 3:1) – one row of pigeon pea after three rows of turmeric ( Additive )

T<sub>4</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric ( Additive )

T<sub>5</sub>- Turmeric + Pigeon pea ( 3:1) -one row of pigeon pea after three rows of turmeric (Replacement )

T<sub>6</sub>- Turmeric + Pigeon pea ( 5:1)- one row of pigeon pea after five rows of turmeric (Replacement )

T<sub>7</sub>- Turmeric + Pigeon pea ( 6:2)- Two rows of pigeon pea after six rows of turmeric (Replacement )

T<sub>8</sub>- Turmeric + Pigeon pea ( 10:2)- one row of pigeon pea after ten rows of turmeric (Replacement)

**Table3.**

**Yield and economics of Turmeric +Pigeon pea intercropping system (Pooled over 2021-22 &2022-23)**

Treatment	Fresh Turmeric Rhizome (q/ha)	Grain yield of Pigeon pea (q/ha)	Dry Turmeric Rhizome (q/ha)	Turmeric (dry rhizome ) equivalent yield (q/ha)	Gross Return (Turmeric) (Rs/ha)	Gross Return (Pigeon pea) (Rs/ha)	Gross Return (Turmeric &Pigeon pea) (Rs/ha)	Net Return (Rs/ha)	B:C Ratio
T1-Sole turmeric	106.04		22.21		371140	-	371140	40140	1.37

T2- Sole Pigeon pea	-	14.07	-	9.62	-	92862	92862	41863	1.40
T3- T+P(3:1) Additive	42.99	11.13	8.60	16.26	150465	73458	223923	51563	1.51
T4- T+P(5:1)Additive	59.90	7.30	11.83	16.88	209650	48180	257830	57590	1.54
T5T+P(3:1)Replacement	53.22	8.81	10.65	16.66	186270	58146	244416	69736	1.67
T6T+P(5:1)Replacement	61.59	5.97	12.32	16.40	215565	39402	254967	75927	1.87
T7- T+P(6:1)Replacement	52.60	8.28	10.52	16.21	184100	54648	238748	66068	1.71
T8-T+P(10:2) Replacement	84.02	4.54	16.81	19.91	294070	29964	324034	98994	2.11
SEm (±)	6.74	1.08	0.55	0.71	3396	2983	2359	2431	0.04
CV(0.05)	18.04	3.16	1.16	1.50	7199	6323	5001	5154	0.08

**Comment [p4]:** Coefficient of Variation I guess you meant

#### 4.CONCLUSION

~~Details of the Technology Adoption of T~~ Turmeric and pigeon pea (10:2) planting pattern to turmeric and pigeon pea intercropping system was found to be the most stable, productive and remunerative planting geometry for the turmeric and pigeon pea intercropping system.

~~Recommended Ecosystem Rainfed Upland~~

~~Superiority Over Existing Technology~~

**Formatted:** Tab stops: 2.83", Left + Not at 2.88"

~~Productivity~~ Sole crop of turmeric fresh rhizome (106.04 q/ha) and pigeon pea grain (14.07 q/ha) produced the highest yield when compared with other intercropping systems.

**Formatted:** Tab stops: 2.83", Left

~~Profitability~~ Turmeric +Pigeon pea (10:2) geometry gave the highest net return and benefit cost ratio over other intercropping system.

~~Message to Farmers~~ Turmeric and pigeon pea (10:2) planting pattern to turmeric and pigeon pea intercropping system recommended for higher productivity and net return.

~~FUTURE RESEARCH~~ Popularization of the technology by conducting OFT and FLDs through KVK and

**Formatted:** Justified, Line spacing: Double, Tab stops: 2.83", Left

~~Govt. Line Department.~~

#### REFERENCES

Anonymous. National Horticulture Board. Ministry of Agriculture, Govt. of India, Gurgaon; Indian horticultural database. 2013.

Balashanmugam, PV, Shah Ahmed, Chezhiyan N. Studies on relay intercropping in turmeric. South Indian Hort. 1988; 36 : 207–208.

Chattopadhyay I, Biswas K, Bandyopadhyay U, Banerjee RK. Turmeric and curcumin: biological actions and medicinal applications. Curr. Sci. 2004; 87: 44-53.

Das AK., Maharana T. Effect of spacing and intercropping on bunch characters of banana cv. Robusta. Orissa Journal of Horticulture. 1995; 23(1–2): 13 – 17.

Islam, M.R., Molla, M.S.H. and Main, M.A.K. Productivity and profitability of intercropping sesame with turmeric at marginal farmers level of Bangladesh. SAARC J. 2016; Fig. 1

Kannan MP, Jansirani VA, Sathiyamurthy D, Veeraragavathatham Sankar V. Standardization of vegetable intercropping system in Amorphophallus yam. Changing scenario in the production systems of horticultural crops. South Indian Horticulture. 2001; 49: 370–372.

Kumar AK, Reddy MD. Integrated weed management in maize + turmeric intercropping system. Indian J. Weed Sci. 2000; 32 : 59–62.

Li L, Sun JH, Zhang FS, Li XL, Yang SC, Rengel, Z. Wheat/maize or wheat/soybean intercropping I. Yield advantage and interspecific interactions on nutrients. Field Crop Research. 2006; 71: 123–137.

Narayanpur MN, Sulikeri GS. Economics of companion cropping system in turmeric (*Cucurma longa* L.). Indian  
Cocoa, Arec. Spices J. 1996;20: 77–79.

Paraye PM, Mahobia RK, Paikra KK, Singh SP. (2014). Ginger (*Gingiber officinale* Roscur) based intercropping  
system for Chhattisgarh plain zone. Environ. & Ecol 1996; 32(2A): 791-793.

Nayar TVR, Suja G. Production potential of root and tuber crops in multiple cropping systems involving plantation  
crops. Journal of root crops. 2004; 31: 111–114.

Panase VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council of Agricultural Research,  
New Delhi Cr. Ist 1967; pp.100–109 and 152–161.

---

Ray DP. Spices cultivation in eastern and north eastern regions- A perspective, Souvenir &  
Abstracts of National Symposium on Spices and Aromatic Crops held at OUAT, Bhubaneswar,  
2007; (25- 26): 47-74.

Ravindran CS, Nair GM, Nedunchezhiyan M. Intercropping tuber crops is remunerative. Indian  
Horticulture. 2006; 51(6): 25 – 27.

Sasikumar B. Turmeric. In: Peter KV (ed.) Handbook of Herbs and Spices, Cambridge,  
Woodland Publishing. 2001; Vol.1: 297-308.

Singh J, Randhawa GS. Effect of intercropping and mulch on yield and quality of turmeric. Acta Hort. 1988; 188A:  
183–186.

Formatted: Tab stops: 5.2", Left

Willey RW. Intercropping – its importance and research needs. Part I. Competition and yield advantages. Field Crops Abstracts .1979;32(1): 1–10.

UNDER PEER REVIEW