

Original Research Article

Development and Evaluation of Physico-chemical Properties of Moringa-Woodapple-Karonda Mixed Jam

Abstract

The present investigation was conducted to develop mixed jam using moringa leaf powder, wood apple and karonda fruit pulp. This research focuses on the standardization and evaluation of a unique mixed fruit jam comprising wood apple (WK), karonda, and moringa. The study initially standardized the wood apple and karonda mixed jam by blending their pulps in different proportions. Organoleptic scores were used to determine the most favourable blend, with WK₄ (40% wood apple and 60% karonda) emerging as the preferred combination based on appearance, aroma, taste, mouthfeel, and overall acceptability. Subsequently, WK₄ was utilized as the control for the development of moringa-based mix jam. The organoleptic evaluation of the moringa-based jam revealed a decrease in appearance scores with an increase in moringa leaf powder. However, taste and mouthfeel exhibited favourable scores, with WKM₃ (95% WK: 5% Moringa) achieving the highest overall acceptability. The study also explored the physico-chemical properties of the jam, including total soluble solids, titratable acidity, pH, ascorbic acid, total phenol, and sugar content. Notably, WKM₆ (12.5% Moringa) demonstrated the highest gel and rupture strength, indicating a robust texture. The findings contribute to the development of innovative fruit-based products with enhanced nutritional profiles, emphasizing the potential of moringa as a valuable ingredient in mixed fruit jams.

Keywords: Moringa, value-added, textural properties, blend, Organoleptic

1. Introduction

Moringa (*Moringa oleifera*) is the most widely cultivated species of the Moringaceae family. Moringa is usually known as drumstick tree and also by different names in local languages like sahjan, moonga, benzolive tree, kelor, marango, mlonge, mulangay, sajna or Ben oil tree. Moringa is native to the sub-Himalayan tracts of India, Pakistan, Bangladesh and Afghanistan [1]. India is the major producer of moringa crop, with an annual production of 2.2 million tonnes from an area of 38,000 hectares [2]. Among the major states in moringa

production, Andhra Pradesh leads in both area and production followed by Karnataka and Tamil Nadu. Commercial production of the crop has also been adopted in Maharashtra and Gujarat. The plant adapts well to different soil conditions and has an incredible ability to survive in harsh weather and even to drought. It is therefore considered as a fast-growing drought-resistant tree. The rainfed agriculture in arid and drought-prone areas of Bundelkhand could be rejuvenated with the commercial cultivation of moringa.

Moringa stands out as a remarkably versatile plant, offering diverse applications including food, natural medicine, animal feed, natural coagulants, forestry products, fertilizer, alley cropping, and fuel [3,4]. Its extensive utility stems from the fact that nearly every part of the tree holds value, making moringa renowned as one of the most multifunctional plants on the planet. The leaves, fruits, flowers and immature pods are edible and form part of traditional diets in tropical and subtropical nations [5]. The immature green pods called “drumsticks” are widely used as part of the tree. Drumsticks or pods have a slight asparagus taste and are commonly consumed in South India in vegetable preparation or sambar. The flowers of moringa are edible when cooked, and are said to taste like mushrooms and added in vegetable preparation. However, another edible part of the plant i.e. moringa leaves has found less commercial exploitation.

Moringa leaves possess remarkable nutritional and medicinal qualities. The plant has brought massive attention as the ‘natural nutrition of the tropics’ because it has the potential to enhance the consumption of essential nutrients and phytochemicals that promote health. Medicinal properties like anti-tumour, anti-inflammatory, anti-ulcer, anti-atherosclerotic and anti-convulsant activities are important [6,7,8]. Moringa is also utilized for the treatment of home-scale medicine [9]. Moringa leaves are considered to be a causative factor in controlling cholesterol and blood glucose levels and therefore thought to be useful for glucose purification [10]. The leaves contain a high amount of vitamin C, which fights different illnesses including colds and flu. Carotenoids, serving as Vitamin A precursors in the leaves, protect against various diseases such as eye and skin conditions, heart ailments, and diarrhea. Additionally, the leaves contain abundant Calcium for building strong bones and teeth, preventing osteoporosis, and essential Potassium for proper brain and nerve function. Moringa leaves are rich sources of protein and also contain all of the essential amino acids in a good proportion. For individuals lacking protein from animal sources, these leaves could be a valuable resource. The nutritional

content is particularly high in dried leaves, with ten times the vitamin A of carrots, 17 times the calcium of milk, 15 times the potassium of bananas, 25 times the iron of spinach, and nine times the protein of yoghurt, as indicated by [11].

Table 1: Nutritional content of fresh and dried *Moringa oleifera* leaves (per 100gm)

Nutrient	Unit	Fresh leaves	Oven-dried
Moisture	(%)	75.9	6
Energy	(Kcal)	92	271.54
Protein	(g)	6.7	23.78
Carbohydrates	(g)	12.5	28.32
Fat	(g)	1.7	7.014
Fiber	(g)	0.9	11.8
Vitamin C	(mg)	220	56
Calcium	(mg)	440	3467

[12]

To bring this crop into the mainstream of food product categories, it needs to be supplemented with additional value and storability. The moringa leaf-based value-added product diversification can play a vital role in the increased consumption, and utilization of this nutritionally rich crop. These diversified value-added products developed from moringa leaves could help not only to alleviate malnutrition problems but also help to address micronutrient deficiency in the Indian population. Moringa leaves are suitable for consumption in either their fresh or dried form as part of a daily diet, with the advantage of extended storage for dried leaves. Various Moringa leaf products, including Moringa Tea, Moringa Tablets, Moringa Capsules, Moringa Leaf Powder, Moringa Soaps, and Moringa Face products, are commercially utilized as health foods and cosmetic items. Moringa leaves beverage is another consumable product that has potential as health food. So, it is necessary to develop new technologies/products from Moringa leaves that could be hygienically processed and have extended shelf life for further uses. The main aim of the study was to standardize different Moringa leaves-based value-added products for commercial utilization. Products like Ready-to-serve beverages, instant fruit drink powder and jam have potential to be incorporated with moringa leaves. These products need to be developed considering their palatability, appearance and storage life. Therefore, other fruits/medicinal crops in the arid region could be utilized with the moringa

leaves to make it more attractive and increase consumer acceptance. Wood apple and Karonda are the other crops selected for blending with Moringa leaves for the development of different value-added convenience products.

Mix fruit jam is another value-added product having higher commercial value due to its appealing nature. Jam is generally consumed in breakfast food with bread or roti. In India, changes in lifestyle promote value-added products like jam. Although moringa has several nutritional benefits, its utilization in jam is challenging. Therefore, other fruits like wood apple and karonda were explored for the purpose.

Wood apple botanically known as *Feronia lemonia* belonging to the family Rutaceae is one of the hardiest fruits grown in semi-arid and arid regions of India. It is an acidic fruit in unripe condition but gives characteristic flavour when ripe. It contains sweet aromatic pulp, which has 74% moisture, 7.3 g protein, 15.5 g carbohydrates, 170 mg riboflavin, 2 mg vitamin C per 100 g of pulp and minerals, especially 0.17% calcium 0.08% phosphorus and 0.07% iron [13]. The fruit is considered to be a rich source of pectin and therefore found to be most suitable for the preparation of value-added products like jam or jelly. However, the characteristic taste of the fruit needs to be neutralized to make it more acceptable. Mixed fruit jam therefore could be a good alternative for the utilization of wood apples.

Karonda (*Carissa Carandas*) is a native fruit of India and belongs to the family Apocynaceae. This is a rich source of pectin and also minerals, especially iron and calcium. Karonda possesses a higher vitamin C content compared to apples and bananas. The ripe fruit having anti-scorbutic properties is reported to be cooling, acidic and useful in bilious [14]. Fruit is slightly sour and astringent in taste, therefore its cultivation is only confined as a fencing bush and it is not popular as dessert fruit. The fruit has a potential for processing but its commercial utilization was limited by the characteristic acidic taste of the fruit. However, the Fruit can be used for the preparation of mixed jam, jelly, pickle and preserves[15]. Moringa-Wood apple-Karonda mixed jam could thus be one of the nutritive value-added products having a delicious taste with higher consumer acceptability.

2. Materials and Methods

The experiment was conducted at the Post Harvest Technology Department, College of Horticulture, Banda University of Agriculture and Technology, Banda, Uttar Pradesh during the

session 2021-2022 under Complete Randomized Design with six treatments and three replications in each.

Procurement of raw material

Moringa leaves, karonda and wood apple collected from the field of College of Horticulture, BUAT, Banda. Sugar and other ingredients required were procured from the local market of Banda.

Standardization of wood apple and karonda mixed jam

Wood apple and karonda pulp were blended in different proportions for the preparation of jam and the best proportion was selected based on a 9-point hedonic sensory score. The selected proportion was further blended with moringa leaf powder.

Table 2: Standardization of wood apple and karonda mixed jam

Treatments	Fruit pulp (Ratio)		TSS (⁰ Brix)
	Wood apple	: Karonda	
WK ₁	70	: 30	68%
WK ₂	60	: 40	68%
WK ₃	50	: 50	68%
WK ₄	40	: 60	68%
WK ₅	30	: 70	68%

WK: Wood apple and Karonda

Standardisation of Moringa-Wood Apple- Karonda mixed jam


Table 3: Combination of mixed jam from moringa leaf powder, wood apple and karonda


Treatments	Fruit Pulp (%)		TSS (⁰ Brix)
	WK	MLP	
WKM ₁	100%	0% (control)	68%
WKM ₂	97.5%	2.5%	68%
WKM ₃	95%	5%	68%
WKM ₄	92.5%	7.5%	68%
WKM ₅	90%	10%	68%
WKM ₆	88.5%	12.5%	68%

WKM: Wood apple, Karonda and Moringa; **MLP:** Moringa Leaf Powder

WK: Wood apple and Karonda

Flow Chart for Preparation of Moringa-based mix jam:





3. Result and Discussion

Standardization of wood apple and karonda mixed jam

Wood apple and Karonda pulp were blended in different proportions for the preparation of jam and the best proportion was selected based on a 9-point hedonic organoleptic score. A significant difference was recorded among all the other formulations the detail given in table 3.

A gradual increase in appearance scores of all the formulations was recorded with an increase in the proportion of karonda pulp. The similar pattern was observed with the scores of taste and mouth feel. But in the case of aroma scores was decreased from all the formulations were recorded with increasing the proportion of karonda pulp. The highest overall acceptability score was recorded as 8.66 for the formulation WK₄ (40% wood apple and 60% karonda) followed by WK₃ (7.33). The minimum overall acceptability score was recorded in the jam WK₁ which contains 70% wood apple and 30% karonda pulp. WK₄ (40% wood apple and 60% karonda pulp) was adjudged best among all the combinations considering the higher overall acceptability. It was further used for the development of moringa based mix jam. The details given table in 4

Table 4. Organoleptic evaluation of standardized wood apple and karonda mix jam

Parameter Treatment	Appearance	Aroma	Taste	Mouth feel	Overall acceptability
WK ₁	5.66	8.33	5.33	4.66	5.66
WK ₂	6.66	7.66	6.33	5.66	6.33
WK ₃	7.66	6.33	7.33	6.66	7.33
WK ₄	8.33	7.66	8.66	8.33	8.66
WK ₅	8.66	5.66	6.66	6.66	6.00
Mean	7.39	7.12	6.86	6.39	6.79
CD (0.05%)	1.34	1.41	1.04	1.20	1.39

WK: Wood apple and Karonda

Standardisation of Moringa-Wood Apple- Karonda mix jam

Wood apple: karonda pulp blend for mixed jam standardized in table (4) was further blended with moringa leaf powder in five different proportions using WK₄ (40:60) as control.

Organoleptic evaluation of developed moringa based mix jam

Appearance

A gradual decrease in the appearance scores of all the formulations for moringa based mix jam was recorded with an increasing proportion of moringa leaf powder and the observation given in Table No. 4. The maximum score was recorded in WKM₁ (8.30) followed by WKM₂ (7.60), WKM₃ (7.15), WKM₄ (6.31), WKM₅ (5.98) and WKM₆ (5.18). The appearance score was decreased significantly [16] and reported similar result in mixed fruit jam. A similar result was recorded by [17] in strawberry fruit jam. A similar result was reported by [18] in jamun fruit jam and [19] also reported in apple and pear mixed fruit jam.

Aroma

The aroma scores of developed moringa based mix jam have been given in Table 4 and indicate that a significant difference in aroma scores was observed for all the treatments. The maximum score was recorded in WKM₃ (7.55) followed by WKM₄ (6.30), WKM₂ (5.33), WKM₅ (5.26) and WKM₁ (5.15) respectively, while the minimum value was recorded in WKM₆ (4.46). The aroma score was decreased significantly. A similar result was reported by [18] in jamun fruit jam and [19] also reported in apple and pear mixed fruit jam.

Taste

The taste scores of developed moringa based mix jam have been presented in Table 4. Results indicated that the mean score for different jam varied from 5.33 to 8.33. The mean maximum score for taste was recorded as 8.50 for WKM₃ followed by WKM₂ (7.91), WKM₄ (7.56), WKM₁ (7.30), WKM₅ (6.66) and the minimum 5.33 was scored by WKM₆ respectively. The aroma score was decreased significantly. A similar result was recorded by [17] in strawberry fruit jam. A similar result was reported [18] in jamun fruit jam and [19] also reported in apple and pear mixed fruit jam.

Table 5: Organoleptic evaluation of developed moringa based mix jam

Parameter	Appearance	Aroma	Taste	Mouth feel	Texture and body	Overall acceptability
-----------	------------	-------	-------	------------	------------------	-----------------------

Treatment						
WKM ₁ (100%)	8.30	5.15	7.30	6.65	7.06	6.33
WKM ₂ (97WK:5:2.5M)	7.60	5.33	7.91	7.80	7.46	7.00
WKM ₃ (95WK:5M)	7.15	7.55	8.50	8.60	8.61	7.91
WKM ₄ (93WK:7.5M)	6.31	6.30	7.56	7.68	7.03	6.91
WKM ₄ (93WK:7.5M)	5.98	5.26	6.66	6.26	6.15	5.91
WKM ₆ (87.5WK:12.5M)	5.18	4.46	5.33	5.30	5.38	4.58
Mean	6.75	5.67	7.21	7.04	6.94	6.44
CD (0.05%)	0.29	0.37	0.65	0.66	0.42	0.72

WKM: Wood apple, Karonda and Moringa

Mouth feel

The data presented in Table5 shows the effect of moringa leaf powder on all moringa based mix jam. The maximum mouth feel mean score was recorded as 8.60 for WKM₃ followed by WKM₂ (7.80), WKM₄ (7.68), WKM₁ (6.65), WKM₅ (6.26) and the minimum (5.30) was scored by WKM₆ respectively. The mouth feels score differed significantly in most of the different combinations of moringa based fruit jam. A similar result was reported by [18] in jamun fruit jam and [19] also reported in apple and pear mixed fruit jam.

Texture and body

The texture and body mean score of the developed moringa based mix jam was recorded and mentioned in Table5 The maximum score was recorded in WKM₃ (8.61) followed by WKM₂ (7.46), WKM₁ (7.06), WKM₄ (7.03), WKM₅ (6.15) and minimum score was recorded in WKM₆ respectively. The texture and body score was differ significantly. A similar result was reported [18] in jamun fruit jam and [19] also reported a similar result in apple and pear mixed fruit jam.

Overall acceptability

The Overall acceptability mean score of different combinations of moringa based mix jam was found to range from 4.58 to 7.91 in Table 4. Among the products studied, WKM₃ displayed the highest score for overall acceptability, closely followed by WKM₃, WKM₂, WKM₄, WKM₁, WKM₅ and WKM₆. In most of the jam combinations, overall acceptability score differed significantly. A similar result was recorded by [17] in strawberry fruit jam and [20] reported in pineapple and mango mixed fruit jam. [18] reported in jamun fruit jam and [19] also reported similar results in apple and pear mixed fruit jam.

Physico-chemical properties of moringa based mix jam

Total soluble solid (⁰Brix)

The total soluble solids of the products are the index of sweetness. TSS of different combinations of the mix jam as determined by the refractometer was found to be in the range of 68.10 to 70.53 ⁰Brix given in Table 5. Among the previously studied products, WKM₁ displayed maximum TSS content (70.53 ⁰Brix), thoroughly followed by WKM₆ (70.06 ⁰Brix), WKM₅ (68.33 ⁰Brix), WKM₂ (68.26 ⁰Brix), WKM₃ (68.1 ⁰Brix) and WKM₄ (67.50 ⁰Brix). In most of the treatment combinations, TSS content differed significantly. A similar result was reported by [21] in fruit jam [16] reported a similar result in mixed fruit jam and [19] also reported a similar result in apple and pear mixed fruit jam.

Titrateable acidity (%)

The examination of titrateable acidity was undertaken to ascertain physico-chemical alterations during the preparation process. The titrateable acidity of samples WKM₁, WKM₂, WKM₃, WKM₄, WKM₅ and WKM₆ were 1.10, 1.10, 1.16, 1.18, 1.07 and 1.12% respectively in Table 5. Maximum and minimum acidity were found to be in WKM₄ and WKM₅ (1.07% and 1.18%). In most of the treatment combinations acidity content differed non-significantly. A similar result was reported by [21] in strawberry jam [20] reported a similar result in pineapple and mango mixed fruit jam and [19] also reported a similar result in apple and pear mixed fruit jam.

pH

The pH measurement was conducted to determine if a consistently low pH was maintained, as it could serve as an effective preservation method and contribute to the textured gel structure of the jam. Maximum pH value was in WKM₆ (3.35) followed by WKM₅, WKM₄,

WKM₃, WKM₂ and WKM₁, (3.34, 3.21, 3.11, 3.05, 3.04) respectively in Table 5. pH value differed significantly in most of the different combinations of moringa based mix jam. Similar result were reported by [22] in Mulberry and Roselle Mixed Fruits Jam and [23] reported similar result in fruit jam. [24] also reported similar result in moringa powder and apple mixed jam.

Table 6: TSS, titratable acidity and pH content of moringa based mix jam

Treatment	TSS (^oBrix)	Titratable Acidity (%)	pH
WKM ₁	70.53	1.10	3.04
WKM ₂	68.26	1.10	3.05
WKM ₃	68.10	1.16	3.11
WKM ₄	67.50	1.18	3.21
WKM ₅	68.33	1.07	3.34
WKM ₆	70.06	1.12	3.35
Mean	68.80	1.12	3.18
C.D.(0.05%)	1.31	N/A	0.07

WKM: Wood apple, Karonda and Moringa

Ascorbic acid (mg/100g)

The ascorbic acid content of different combination of moringa based mix jam was found to be range from 7.59 to 11.14 mg/100g. Among the products studied, WKM₁ displayed highest ascorbic acid content (11.14mg/100g), closely followed by WKM₂ (9.24 mg/100g), WKM₃ (8.71 mg/100g), WKM₄ (8.30 mg/100g), WKM₅ (7.88 mg/100g) and WKM₆ (7.59 mg/100g) respectively in Table 5. In most of the treatment combinations, ascorbic acid content differed significantly. A similar result was recorded by [20] in pineapple and mango mixed fruit jam and [25] reported a similar result in mixed fruit jam.

Total phenol (mg/100g)

The total phenol content of different combination of moringa based mix jam was found to be range from 98.6 to 201.54 mg/100g given in Table 5. In most of the treatment combinations, total phenol content differed significantly. Among the products studied, WKM₆ displayed highest total phenol content (201.54 mg/100g), closely followed by WKM₅ (167.92 mg/100g),

WKM₄ (153.71mg/100g), WKM₃ (118.75 mg/100g), WKM₂ (110.45 mg/100g) and WKM₁ (98.60 mg/100g) respectively. WKM₁ showed the lowest value of total phenol content 98.60 mg/100g.

Table 7: Ascorbic acid and Total phenol content of moringa based mix jam

Treatment	Ascorbic Acid (mg/100g)	Total phenol (mg/100g)
WKM ₁	11.14	98.60
WKM ₂	9.24	110.45
WKM ₃	8.71	118.75
WKM ₄	8.30	153.71
WKM ₅	7.88	167.92
WKM ₆	7.59	201.54
Mean	8.81	141.83
C.D.	0.74	8.55

WKM: Wood apple, Karonda and Moringa

Reducing sugars (%)

The reducing sugars of developed products were the index of sweetness. Reducing sugar of different blends of moringa based mix jam was found to be ranged from of 34.74 to 43.27 percent given in Table 5. Amongst the developed products studied, WKM₆ displayed the lowest reducing sugar content (34.74%), closely followed by WKM₅ (35.42%), WKM₄ (37.29%), WKM₃ (39.47%) and WKM₂ (41.80%) respectively in Table 6. WKM₁ showed the highest value of reducing sugars content 43.27 per cent. In most of the treatment combinations, total sugar content different significantly. A similar result was reported by [18] in jamun fruit jam and [25] also reported similar result in black-plum fruit jam.

Non-reducing sugars(%)

Non-reducing sugars of different combinations of moringa based mix jam were found to be ranged from 18.62 to 20.02 percent given in Table 6. Among the products studied, WKM₆ displayed the lowest non-reducing sugar content (18.62%), closely followed by WKM₃

(19.04%), WKM₂ (19.07%), WKM₄ (19.41%) and WKM₅ (19.88%) respectively. WKM₁ showed the highest value of non-reducing sugar content 20.02 percent. In most of the treatment combinations total sugar content is different non-significantly. A similar result was reported by [18] in jamun fruit jam and [25,26] also reported a similar result in black-plum fruit jam.

Total sugars (%)

The total sugars of developed products were the index of sweetness. Total sugars of different blends of moringa based mix jam were found to be a range of 63.51 to 67.36 per cent given in Table8. Amongst the products studied, WKM₄ existing lowest total sugar content (63.40%), closely followed by WKM₃ (63.51%), WKM₂ (64.52%), WKM₅ (65.64%) and WKM₁ (67.29%) respectively. WKM₆ presented the maximum value of total sugar content of 67.36 percent. In most of the treatment combinations total sugar content different significantly. A similar result was also reported by [23] in fruit jam [16] reported similar result in mixed fruit jam and [19,26,27] also reported similar result in apple and pear mixed fruit jam.

Table 8: Reducing, non-reducing and total sugar content of moringa based mix jam

Treatment	Reducing sugars (%)	Non-reducing sugars (%)	Total sugars (%)
WKM ₁	43.27	20.02	67.29
WKM ₂	41.80	19.07	64.52
WKM ₃	39.47	19.04	63.51
WKM ₄	37.29	19.41	63.40
WKM ₅	35.42	19.88	65.64
WKM ₆	34.74	18.62	67.36
Mean	38.66	19.34	65.29
CD(0.05%)	1.55	N/A	1.78

WKM: Wood apple, Karonda and Moringa

Textural properties of developed moringa based mix jam

Analysis of jam texture was performed, using a TA-XT-plus texturometer (Stable Micro Systems, at Banda University of Agriculture and Technology, Banda). The following conditions were applied: Return to Start (Set Dist), Test Mode: Compression, Pre-Test Speed: 3.00 mm/sec,

Test Speed: 2.00 mm/sec, Post-Test Speed: 10.00 mm/sec, T.A. Variable No: 5: 0.0 g, Target Mode: Distance, Distance: 10.000 mm, Strain: 10.0 %, Trigger Type: Auto (Force), Trigger Force: 10.0 g, Probe: SMS P/36R; P/36R, Batch: TEST, Points per second: 500. Prior to the analyses, the samples were acclimated at room temperature. Each sample underwent analysis in ten repetitions. Jam texture was established using the following texture parameters: gel strength (force at a point in the initial stage of penetration, where little deformation has occurred), rupture strength (the rupture point of the gel), brittleness (mm) and adhesiveness (area under the negative region of the curve). The results were calculated using Texture Exponent software (Stable Micro Systems, Banda University of Agriculture and Technology, Banda).

Pectin is the main factor analyzing jam consistency and its content and type have an effect on gel strength (g), jam rupture strength (g) and brittleness (mm), which were measured, along with adhesiveness (g.sec.).

Gel strength (g)

Right after production, the gel strength values for the examined jam ranged from 127.09 to 561.85g (see Table 6). Among the all jam studied, WKM₂ displayed the highest value of gel strength (561.85g), closely followed by WKM₄ (429.18g), WKM₃ (342.81g), WKM₁ (145.89g) and WKM₅ (134.63g). WKM₆ showed the lowest value of gel strength 127.09g. In most of the treatment combinations gel strength differed significantly.

Rupture strength (g)

After preparing mixed jam the values of rupture strength were recorded within the range of 435.86 to 2021.52g (Table 6). Among all jam studied WKM₂ displayed the highest value of rupture strength (2021.52g), closely followed by WKM₄ (1032.79g), WKM₁ (969.92g), WKM₃ (888.61g) and WKM₅ (635.78g). WKM₆ Showed the lowest value of rupture strength 435.86g. in most of the treatment combinations rupture strength value differed significantly.

Brittleness (mm)

In the present study brittleness value was recorded in different combinations of mixed fruit jam result showed that the brittleness value was 10 mm for all jam except WKM₃ 9.65 mm. Brittleness differed significantly in most of the different combinations of moringa leaves-based jam. The detail given in Table7.


Adhesiveness (g.sec)

Adhesiveness of different combinations of mixed jam as determined by the texturometer was found to be range of -1130.72 to -369.22g.sec in table9. Among the all jam studied WKM₂ displayed lowest adhesiveness (-1130.72g.sec), closely followed by WKM₄ (-531.16g.sec), WKM₁ (-499.25g.sec), WKM₅ (-414.97g.sec), WKM₃ (-410.03g.sec) and WKM₆ showed highest value of adhesiveness. In most treatment combinations adhesiveness differed significantly. A similar result was reported by [27] in strawberry jam.

Table 9: Textural properties of developed moringa based mix jam

Parameters Treatments	Gel strength (g)	Rupture strength (g)	Brittleness (mm)	Adhesiveness (g.sec)
WKM ₁ (Control)	145.89	969.92	10	-499.25
WKM ₂ (2.5%)	561.85	2021.52	10	-1130.72
WKM ₃ (5%)	342.81	888.61	9.65	-410.03
WKM ₄ (7.5%)	429.18	1032.79	10	-531.16
WKM ₅ (10%)	134.63	635.78	10	-414.97
WKM ₆ (12.5%)	127.09	435.86	10	-369.22
Mean	290.24	997.41	9.94	-550.89
S.D.	167.09	501.46	0.31	264.02
C.V.	57.57	50.28	1.31	-47.93

WKM: Wood apple, Karonda and Moringa;



Wednesday, 29 September, 2021
Page 1 of 2

Fig. 1: Textural properties of developed moringa based mixed jam

Reference:

1. Fahey JW. *Moringa oleifera*: a review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. *Trees for life Journal*. 2005;1(5):1-15.
2. Balakumbahan R, Sathiyamurthy VA and Janavi GJ. Moringa Leaf–A Super Food. *Biotica Research Today*. 2020;2(6):438-440.
3. Anjula P, Pradheep K, Rita G, Nayar ER and Bhandari DC. Drumstick tree (*Moringa oleifera* Lam.): A multipurpose potential species in India. *Genetic Resources and Crop Evolution*. 2011;58(3):453-460.
4. Pandey A and Negi PS. Traditional uses, phytochemistry and pharmacological properties of *Neolamarckiacadamba*: A review. *Journal of Ethnopharmacology*. 2016; 181:118-135.
5. Siddhuraju P and Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agricultural and Food Chemistry*. 2003;51(8):2144-2155.
6. Chumark P, Khunawat P, Sanvarinda Y, Phornchirasilp S, Morales NP, Phivthong-Ngam L, Ratanachamnong P, Srisawat S, Klai-upsorn SP. The in-vitro and ex-vivo antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *Journal of Ethnopharmacology*. 2008;116(3):439-446.

7. Danmalam HU, Abubakar Z, Katsayal UA. Pharmacognostic Studies on the Leaves of *M. stenopetala*, Niger. *Journal of Natural Produce and Medicine*. 2001;5:4-9.
8. Dahiru D, Onubiyi JA, Umaru HA. Phytochemical screening and antiulcerogenic effect of *Moringa oleifera* aqueous leaf extract. *African Journal of Traditional, Complementary and Alternative Medicines*. 2006;3(3):70-75.
9. Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves: An overview. *International Journal of Molecular Sciences*. 2015;16(6):12791-12835.
10. Ramachandran C, Peter KV, Gopalakrishnan PK. Drumstick (*Moringa oleifera*): a multipurpose Indian vegetable. *Economic Botany*. 1980;52(1):276-283.
11. Mahmood KT, Mugal T, Haq IU. *Moringa oleifera*: a natural gift-A review. *Journal of Pharmaceutical Sciences and Research*. 2010;2(11):775-778.
12. Sahay S, Yadav U, Srinivasamurthy S. Potential of *Moringa oleifera* as a functional food ingredient: A review. *Magnesium*. 2017;8(9):4-90.
13. Thakur N, Chugh V, Dwivedi SV. Wood apple: An underutilized miracle fruit of India. *The Pharma Innovation Journal*. 2020;9(10):198-202.
14. Bajpai R, Yadav M, Mure S, Kushwah RS. Browning analysis of different karonda processed products during storage. *Plant Archives*. 2015;15(1):339-342.
15. Banik BC, Dey AN, Pradhan S, Thapa N, Bhowmick N. Seed germination of karonda (*Carissa carandas* L.). In II International Symposium on Wild Relatives of Subtropical and Temperate Fruit and Nut Crops, 2014;1074:23-26.
16. Khan S, Amin R, Ali S, Zeb TF, Haider MS, Tarar OM. Comparative analysis of the nutritional components of different plants of *Moringa oleifera* from Karachi Pakistan. *Pakistan Journal of Science*. 2020;72(4):271-275.
17. Khan U, Ullah J, Saeed B, Ali F. Effect of potassium sorbate and sodium benzoate on the quality and shelf-life of strawberry jam during storage. *Journal of Agricultural and Biological Sciences*. 2014;9(12):454-458.
18. Shahnawaz M, Shiekh SA. Analysis of viscosity of jamun fruit juice, squash and jam at different compositions to ensure the suitability of processing applications. *International Journal of Plant Physiology and Biochemistry*. 2011;3(5):89-94.

19. Durrani Y, Zeb A, Ayub M, Ullah J. Development of diet jam from apple grown in swat (NWFP). *Sarhad Journal of Agriculture*.2008;24(3):461-467.
20. Oo K,Than S. Study on physico-chemical properties and shelf-life of mixed pineapple and mango jam under ambient storage. *International Journal Advance Research Publication*. 2019;3(8):4-8.
21. Khan RU, Afridi SR, Ilyas M, SohailM, Abid H. Development of strawberry jam and its quality evaluation during storage. *Pakistan Journal of Biochemistry and Molecular Biology*.2012;45(1):23-25.
22. Wongchalat R,Chatthongpisut R. Nutritional Value and Anthocyanins of Mulberrry and Roselle Mixed Fruits Jam. In *Applied Mechanics and Materials*. Trans Tech Publications Ltd.2017;855:65-69.
23. Chawafambira A, Sedibe MM, Mporfu A,Achilonu M. Probiotic potential, iron and zinc bioaccessibility, and sensory quality of Uapacakirkiana fruit jam fermented with *Lactobacillus rhamnosusyoba*. *International Journal of Food Science*. 2020;6:3-11.
24. Rajput H, Prasad SGM, Srivastava P, Singh N,Morya S. Development of fresh *Moringa oleifera* leaf jam and its physico-chemical properties. *Development*. 2017;2(6): 234-238.
25. Selvamuthukumaran M, Khanum F. Processing seabuckthorn fruit for antioxidant rich jam development and shelf stability assessment. *International journal of traditional knowledge*.2014;13:110-114.
26. Pandey A, Negi PS. Traditional uses, phytochemistry and pharmacological properties of *Neolamarckiacadamba*: A review. *Journal of Ethnopharmacology*. 2016;181:118-135.
27. Kopjar M, Piližota V,Tiban, N,Šubarić D, Babić J,Ačkar Đ,Sajdl M. Strawberry jams: Influence of different pectin's on color and textural properties. *Czech Journal of Food Sciences*.2007;27:20–28.