

INFLUENCE OF ABIOTIC FACTORS ON THE POPULATION DYNAMICS AND SPECIES COMPOSITION OF STEM BORER COMPLEX IN RICE ECOSYSTEM

ABSTRACT

A field experiment was carried out at the Murjhad Research Farm, College of Agriculture, Balaghat (M.P.) during *Kharif* 2022 and *Kharif* 2023 to ascertain the population dynamics of stem borer complex, their relative abundance and correlation with abiotic factors in order to propose ecologically and financially feasible measures. Results pertaining to the dynamics of species composition of stem borers of rice reveal that all the three species of stem borer of rice *viz.*, yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), white stem borer, *Scirpophaga innotata* (Walker) were prevalent during the crop season. The pooled data of the total population of stem borer complex observed during the 30th SMW was 8.0 larvae/ 20 hills attaining two peaks, the first peak during the 36th SMW (77.0 larvae/ 20 hills) and the second peak during the 42nd SMW (96.0 larvae/ 20 hills) and thereafter, the population started decreasing and reached 68.5 larvae/ 20 hills during the 43rd SMW and it disappears at the time of harvest. The per cent dead heart incidence was first observed during the 30th SMW (2.19% dead hearts) with its peak during the 36th SMW (16.47% dead hearts) and thereafter, it started decreasing and reached 6.06 per cent dead hearts during the 38th SMW and further, it disappears. The per cent white ears incidence was first observed during the 37th SMW (1.51% white ears) with its peak during the 42nd SMW (23.42% white ears) and thereafter, it started decreasing and reached 12.82 per cent white ears during the 43rd SMW and further, it disappears. The pooled data revealed that a total of 676 larvae stem borer complex were observed throughout the crop growth. Among them, 492 larvae of *Scirpophaga incertulas* were observed comprising 72.78 per cent, *Sesamia inferens* population was 124.5 larvae comprising 18.42 per cent and *Scirpophaga innotata* population was 59.5 larvae comprising 8.80 per cent among the stem borer complex. The pooled data revealed that population of *Scirpophaga incertulas* was found significantly positive correlated ($r= 0.495$) with sunshine hours. The per cent dead heart was found significantly positive correlated with minimum temperature and morning relative humidity ($r= 0.471$ and 0.606 , respectively) and significantly negative correlated with evening relative humidity ($r= -0.625$). The per cent white ear was found significantly positive correlated with evening relative humidity and sunshine hours ($r= 0.663$ and 0.492 , respectively) and significantly negative correlated with morning relative humidity and wind velocity ($r= -0.741$ and -0.484 , respectively).

Keywords: seasonal incidence, stem borer complex, rice, *Scirpophaga incertulas* (Walker), *Sesamia inferens* (Walker), *Scirpophaga innotata* (Walker), abundance, population dynamics, abiotic factors

1. INTRODUCTION

Rice (*Oryza sativa* L.) belongs to the family Gramineae/Poaceae and is an important cereal crop in the world being one of the most important cereal crops worldwide feeding more than 50% of the human population [1]. World rice production in 2021 was 502.98 million metric tons in an area of 165.25 million hectares. China and India are considered the main producers of paddy rice worldwide. In 2021, India's paddy rice production amounted to over 195 million metric tons in an area of about 45 million hectares after China [2]. In Madhya Pradesh, paddy is grown in an area of about 21.17 mha with a production of 44.14 million tons and productivity of 2085 kg/ha [3], while in the Balaghat region, it is grown on about 3.10 lakh ha area with a production of 10.25 million mt and the productivity of 3305 kg/ha [4]. Considering the area, production and productivity of Balaghat district is referred to be the "Paddy Bowl" of Madhya Pradesh.

The productivity of rice crop is influenced by several biotic and abiotic factors. The rice crop is subjected to considerable damage by nearly 300 species of insect pests, among them 23 species are serious pests of rice [5]. Yield loss due to insect pests of rice have been estimated to be 25% [6]. About 40% damage is caused by stem borer alone [7]. The young larva of yellow rice stem borer initially enters into the leaf sheath and feeds for two to three days, after which the larva enters the basal region usually 5 to 10 cm above water level and feeds inside the stem which causes central shoot to die *i.e.*, dead heart at vegetative stage and white ear head is formed by boring at the peduncle node during the heading stage of the crop [8]. The yellow stem borer, *Scirpophaga incertulas* (Walker) is monophagous while, pink stem borer, *Sesamia inferens* (Walker) and white stem borer, *Scirpophaga innotata* (Walker) are polyphagous pests of the Gramineae. The prevalence and population fluctuations of paddy stem borers are influenced by a number of abiotic variables [9]. The changing climatic

scenario with modern cultivation practices in rice crop has made yellow stem borer the most predominant species followed by pink stem borer and white stem borer in many rice growing regions of India, that causes dead heart, white ear and results in yield reduction [10,11,12]. Similarly, some other researcher *viz.*, [13,14,15,16,17,18,19,20,21] have also reported population dynamics of paddy stem borer. In view of the above facts, the present study was conducted to examine the population dynamics of stem borer complex, their relative abundance and correlation with abiotic factors in rice ecosystem.

2. MATERIALS AND METHODS

A field study was carried out at the Murjhad farm of the College of Agriculture, Balaghat (M.P.) during the *Kharif* seasons of 2022 and 2023 in order to investigate the population dynamics of the rice stem borer complex, which includes the yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), and white stem borer, *Scirpophaga innotata* (Walker) in the Balaghat region. For **this purpose**, the nursery was raised with a well-established susceptible variety JRB-1 in the second fortnight of June. For transplanting, the main field was prepared by ploughing once by **tractor drawn disc plough** and second ploughing was done by mould board plough. In the second fortnight of July, the rice was transplanted with 15cm (plant to plant) x 15cm (row to row) spacing in 15m x 20m (300 sq.m.) plot area. Before transplanting the seedlings into main plots, the recommended doses of fertilizers (N:P:K: 100:60:40) were applied. All the **recommended agronomic** practices were conducted but there was no application of any pesticides during **the crop** season.

The observations on incidence of stem borer complex of rice crop were recorded at weekly intervals starting from the first appearance of symptoms of stem borer complex up to the harvesting of the crop. A total of 5 spots (5m x 6m each) were fixed for the seasonal incidence of stem borer complex in 300 m² plot area. Observations were taken from 20 randomly selected hills in each fixed spot. Infested tillers (dead hearts and white ear heads) were carefully uprooted and brought to the laboratory for their splitting and identification of larval species. The larvae were kept under observation for the confirmation of the species of stem borers of rice. The species were identified based on larval characters mentioned in table 1 that were described by [22]. After the confirmation of stem borers species, the number of larvae were counted and computed. Per cent dead heart and white ears were also computed by using the following formula:

$$\text{Dead heart (\%)} = \frac{\text{No. of dead hearts per hill}}{\text{Total no. of tillers per hill}} \times 100$$

$$\text{White ear head (\%)} = \frac{\text{white ear head per hill}}{\text{Total no. of tillers per hill}} \times 100$$

Further, the relative abundance of the stem borer species based on larval incidence during the study period was assessed by the following formula:

$$\text{Relative abundance (\%)} = \frac{\text{Total no. of individuals of each species}}{\text{Total no. of individuals of all species}} \times 100$$

For the correlation and regression studies of stem borer complex with weather parameters, observations on meteorological parameters *viz.*, maximum and minimum temperature, morning and evening relative humidity, sunshine hours, wind velocity, evaporation and rainfall were also recorded from Krishi Vigyan Kendra, Badgaon, Balaghat, M.P. and Water Resources Department, Rajegaon, Balaghat, M.P.

3. RESULTS AND DISCUSSION

The pooled data presented in table 1 revealed that **all the** three stem borer species *viz.*, yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), and white stem borer, *Scirpophaga innotata* (Walker) were prevalent during *Kharif* 2022 and *Kharif* 2023. [11] supported the present results and revealed that species of stem borer of rice *viz.* yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), and white stem borer, *Scirpophaga innotata* (Walker) were prevalent during the crop season.

The incidence of *Scirpophaga incertulas* was first observed during the 30th SMW (8.0 larvae/ 20 hills). Further, it gradually increased and attained two peaks, first peak during the 36th SMW (48.0 larvae/ 20 hills) and second peak during the 42nd SMW (70.5 larvae/ 20 hills) and thereafter, the population started decreasing and reached 53.0 larvae/ 20 hills during 43rd SMW and it disappears during harvest. The incidence of *Sesamia inferens* was first observed during the 31st SMW (2.0 larvae/ 20 hills). Further, it gradually increased and attained two peaks, first peak during the 36th SMW (19.0 larvae/ 20 hills) and second peak during the 42nd SMW (16.0 larvae/ 20 hills) and thereafter, the population started decreasing and reached 12.5 larvae/ 20 hills during the 43rd SMW and it disappears during harvest. The incidence of *Scirpophaga innotata* was first observed during the 31st SMW (1.0 larvae/ 20 hills). Further, it gradually, increased and attained two peaks, first peak during the 36th SMW

(10.0 larvae/ 20 hills) and second peak during the 42nd SMW (9.5 larvae/ 20 hills) and thereafter, the population started decreasing and reached 3.0 larvae/ 20 hills during the 43rd SMW and it disappears during harvest. Considering as a whole, the total population of stem borer complex observed during the 30th SMW was 8.0 larvae/20 hills. Further, it gradually increased and attained two peaks, first peak during the 36th SMW (77.0 larvae/20 hills) and the second peak during the 42nd SMW (96.0 larvae/ 20 hills) and thereafter, the population started decreasing and reached 68.5 larvae/20 hills during 43rd SMW and it disappears during harvest. The present results are supported by the findings of [23] who reported that the catching of *Scirpophaga incertulas* moth was commenced as early as 32nd standard week (2nd week of August) with its peak during 37th standard week. Similarly, [24] also recorded maximum population of *S. incertulas* Walker in the month of September. Additionally, [25] studied on the stem borer complex revealed that only two stem borer species viz., yellow stem borer (*Scirpophaga incertulas* Walker) and pink stem borer (*Sesamia inferens* Walker) were prevalent in different rice ecosystems (upland, irrigated and shallow water) during the year 2007. The current results are in consistent with those of [26] who recorded activity of yellow stem borer started from 30th standard week and continued upto 41st standard week, meanwhile it reached peak twice in 34th and 37th standard week. Thereafter, population of yellow stem borer declined and finally no population found. Also, [27] reported that the population of *Scirpophaga incertulas* began to show up from 26th standard week and proceeded till 43rd standard week the highest population was recorded during 34th to 38th standard week.

The per cent dead heart incidence was first observed during the 30st SMW (2.19% dead hearts). Further, it gradually increased and attained its peak during the 36th SMW (16.47% dead hearts) and thereafter, the per cent dead heart incidence started decreasing and reached 6.06 per cent dead hearts during the 38th SMW and further, it disappears. The per cent white ears incidence was first observed during the 37th SMW (1.51% white ears). Further, it gradually increased and attained its peak during the 42nd SMW (23.42% white ears) and thereafter the per cent white ears incidence started decreasing and reached 12.82 per cent white ears during 43rd SMW and further it disappears. The present results are in consistent with those of [23] who reported incidence of dead heart started at 34th standard week (4th week of August) and reached the peak at 38th standard week (3rd week of September). Subsequently, [28] also reported that per cent incidence of DH and WEH was highest between third to fourth week of September and first fortnight of November, respectively. [29] also supported the present results and showed that first trace of dead heart (DH) was noted from 31st SMW. Incidence of DH gradually increases attaining the maximum at about 39th SMW. After 39th SMW no incidence of white head (WH) was noted. Numerically DH was abundant at about 39th SMW. The maximum incidence WH was counted at about 44th SMW.

The data on the relative abundance of stem borer complex is presented in table 2. The pooled data revealed that a total of 676 larvae stem borer complex were observed throughout the crop growth. Of them, 492 larvae of *Scirpophaga incertulas* were observed comprising 72.78%, *Sesamia inferens* population was 124.5 larvae comprising 18.42% and *Scirpophaga innotata* population was 59.5 larvae comprising 8.80% among the stem borer complex. The findings of the present study are supported by [25] who reported that among the stem borer species, the composition of yellow stem borer was higher at late tillering and panicle initiation stages, but declined at maturity stage. However, the composition of pink stem borer was slightly increased at panicle initiation stage and dominated at the maturity stage. Similarly, [11] found that yellow stem borer was dominant over other species of stem borer and recorded 87.00% to 93.00% at 30 DAT during two *kharif* seasons, respectively. The population of yellow stem borer was 82.50% to 89.00% at 50 DAT and 85.00% to 91.50% at 90 DAT. Additionally, [12] revealed that larval abundance of both yellow stem borer and pink stem borer was maximum in 41st standard meteorological week (61 and 51 larvae in 100 plants showing white ear, respectively) and the per cent share of yellow stem borer and pink stem borer was 54.46% and 45.53% and the occurrence of both the species was minimum during 33rd and 34th standard meteorological weeks (0 and 1, 2 and 1, respectively) with the per cent share of 0.00 and 100%, 66.66 and 33.33%, respectively.

Correlation studies of stem borer complex with weather parameters are presented in table 3. The data revealed that population of *Scirpophaga incertulas* was found significantly positive correlated ($r = 0.495$) with sunshine hours. The larval population of *Sesamia inferens* and *Scirpophaga innotata* was not found significant with any weather parameter. The per cent dead heart was found significantly positive correlated with minimum temperature and morning relative humidity ($r = 0.471$ and 0.606 , respectively) and significantly negative correlated with evening relative humidity ($r = -0.625$). The per cent white ear was found significantly positive correlated with evening relative humidity and sunshine hours ($r = 0.663$ and 0.492 , respectively) and significantly negative correlated with morning relative humidity and wind velocity ($r = -0.741$ and -0.484 , respectively). The results in the present correlation study are in line with the findings of [24] who reported that the value of multiple correlations for the insect was 0.949. Meteorological factors were also responsible for the

dynamics of the populations of insects. Also, [30] inferred that the occurrence of *Scirpophaga incertulas* exhibits a critical positive relationship with morning RH (0.11), Aver. Humidity (0.542) and evening Humidity (0.296), whereas, a huge negative connection with Tmin(0.650) and Tmax (-0.699) against Dead heart incidence though Relative Humidity morning($r=0.047$), Aver. Humidity (0.945), Relative Humidity Evening (0.296), Min. Temperature (-0.256) and Maximum temperature (-0.648) against White Ear Headrate showed positive relationship. Also, the present results are supported by [29] who observed that Tmin ($r=0.561$), Relative Humidity Morning (0.561), Relative Humidity Evening ($r=-0.564$), Sunshine ($r=0.463$) and Evaporation (-0.591) showed significant positive correlation against Dead Heart and also positive relationship between White Ear Head(WEH)and abiotic factors such as Tmin (0.589), Relative Humidity Morning (0.618), Relative Humidity Evening (-0.569), Sunshine (0.362) and Evaporation (-0.798) was recorded. [31] also reported that YSB had non-significant positive correlation with sunshine hours while, other parameters exhibited non-significant negative correlation.

4. CONCLUSION

The pooled data revealed that three stem borer species viz., yellow stem borer, *Scirpophaga incertulas* (Walker), pink stem borer, *Sesamia inferens* (Walker), and white stem borer, *Scirpophaga innotata* (Walker) were prevalent during Kharif 2022 and Kharif 2023. The total population of stem borer complex observed during the 30th SMW was 8.0 larvae/ 20 hills. Further, it gradually increased and attained two peaks, first peak during the 36th SMW (77.0 larvae/ 20 hills) and the second peak during the 42nd SMW (96.0 larvae/ 20 hills) and thereafter, the population started decreasing and reached 68.5 larvae/ 20 hills during 43rd SMW and it disappears during harvest. The per cent dead heart incidence was first observed during the 31st SMW (2.19% dead hearts). Further, it gradually increased and attained its peak during the 36th SMW (16.47 % dead hearts) and thereafter the per cent dead heart incidence started decreasing and reached 6.06% dead hearts during the 38th SMW and further, it disappears. The per cent white ears incidence was first observed during the 37th SMW (1.51% white ears). Further, it gradually increased and attained its peak during the 42nd SMW (23.42% white ears) and thereafter, the per cent white ears incidence started decreasing and reached 12.82 per cent white ears during the 43rd SMW and further, it disappears. The pooled data revealed that a total of 676 larvae stem borer complex were observed throughout the crop growth. Of them, 492 larvae of *Scirpophaga incertulas* were observed comprising 72.78 per cent, *Sesamia inferens* population was 124.5 larvae comprising 18.42 per cent and *Scirpophaga innotata* population was 59.5 larvae comprising 8.80 per cent among the stem borer complex. The data revealed that population of *Scirpophaga incertulas* was found significantly positive correlated ($r=0.495$) with sunshine hours. The larval population of *Sesamia inferens* and *Scirpophaga innotata* was not found significant with any weather parameter. The per cent dead heart was found significantly positive correlated with minimum temperature and morning relative humidity ($r=0.471$ and 0.606 , respectively) and significantly negative correlated with evening relative humidity ($r=-0.625$). The per cent white ear was found significantly positive correlated with evening relative humidity and sunshine hours ($r=0.663$ and 0.492 , respectively) and significantly negative correlated with morning relative humidity and wind velocity ($r=-0.741$ and -0.484 , respectively).

Table 1 The species were identified based on larval characters described by [21]

Species	Head	Body	Prothoracic shield	Crochets
<i>Scirpophaga incertulas</i> (Yellow stem borer)	Yellowish brown	Creamy yellow 20-25 mm 1 st abdominal segment white	Yellowish brown	Biordinal, sometimes almost uniordinal,
<i>Scirpophaga inferences</i> (Pink stem borer)		Creamy yellow 20-25 mm	Yellowish brown, anterior margin tinged with dark colour	arranged in an ellipse
<i>Scirpophaga innotata</i> (White stem borer)	Black to blackish brown	Dull white tinged with pink grey with longitudinal stripes 17-22mm	Black to blackish brown	Almost triordinal arranged in a circle

Table 3 Relative abundance of stem borer complex in rice ecosystem during *Kharif 2022* and *Kharif 2023*

S. No.	Stem borer Species	Total Larval Population			Relative Abundance (%)		
		2022	2023	Pooled	2022	2023	Pooled
1.	<i>Scirpophaga incertulas</i>	534	450	492	72.75	72.82	72.78
2.	<i>Sesamia inferens</i>	134	115	124.5	18.26	18.61	18.42
3.	<i>Scirpophaga innotata</i>	66	53	59.5	08.99	8.58	8.80
	Total	734	618	676	100	100	100

Table 4 Correlation coefficients (r) of stem borer complex with weather parameters in rice ecosystem during *Kharif 2022* and *Kharif 2023*.

Stem borer complex		Correlation coefficients (r)							
		Maximum Temperature (°C)	Minimum Temperature (°C)	Morning RH (%)	Evening RH (%)	Rainfall (mm)	Evaporation (mm)	Wind Velocity (kmph)	Sunshine Hours (hr)
<i>Scirpophaga incertulas</i>	2022	0.179 ^{NS}	-0.052 ^{NS}	-0.372 ^{NS}	0.327 ^{NS}	-0.046 ^{NS}	0.416 ^{NS}	-0.168 ^{NS}	0.500*
	2023	0.186 ^{NS}	-0.024 ^{NS}	-0.359 ^{NS}	0.314 ^{NS}	0.000 ^{NS}	0.415 ^{NS}	-0.137 ^{NS}	0.485*
	Pooled	0.183 ^{NS}	-0.039 ^{NS}	-0.367 ^{NS}	0.322 ^{NS}	-0.025 ^{NS}	0.417 ^{NS}	-0.154 ^{NS}	0.495*
<i>Sesamia inferens</i>	2022	0.292 ^{NS}	0.054 ^{NS}	-0.127 ^{NS}	0.025 ^{NS}	-0.247 ^{NS}	0.273 ^{NS}	-0.192 ^{NS}	0.421 ^{NS}
	2023	0.252 ^{NS}	0.027 ^{NS}	-0.076 ^{NS}	-0.011 ^{NS}	-0.199 ^{NS}	0.261 ^{NS}	-0.116 ^{NS}	0.399 ^{NS}
	Pooled	0.264 ^{NS}	0.175 ^{NS}	0.005 ^{NS}	-0.080 ^{NS}	-0.036 ^{NS}	0.265 ^{NS}	-0.120 ^{NS}	0.267 ^{NS}
<i>Scirpophaga innotata</i>	2022	0.263 ^{NS}	0.144 ^{NS}	-0.015 ^{NS}	-0.059 ^{NS}	-0.031 ^{NS}	0.259 ^{NS}	-0.152 ^{NS}	0.276 ^{NS}
	2023	0.260 ^{NS}	0.207 ^{NS}	0.027 ^{NS}	-0.101 ^{NS}	-0.041 ^{NS}	0.266 ^{NS}	-0.083 ^{NS}	0.252 ^{NS}
	Pooled	0.275 ^{NS}	0.042 ^{NS}	-0.104 ^{NS}	0.008 ^{NS}	-0.227 ^{NS}	0.269 ^{NS}	-0.159 ^{NS}	0.414 ^{NS}
Dead Hearts (%)	2022	0.090 ^{NS}	0.468*	0.605**	-0.620**	0.121 ^{NS}	0.094 ^{NS}	0.259 ^{NS}	-0.125 ^{NS}
	2023	0.103 ^{NS}	0.468*	0.595**	-0.623**	0.103 ^{NS}	0.161 ^{NS}	0.270 ^{NS}	-0.104 ^{NS}
	Pooled	0.095 ^{NS}	0.471*	0.606**	-0.625**	0.116 ^{NS}	0.117 ^{NS}	0.264 ^{NS}	-0.119 ^{NS}
White Ears (%)	2022	0.237 ^{NS}	-0.237 ^{NS}	-0.694**	0.615**	-0.213 ^{NS}	0.058 ^{NS}	-0.463 ^{NS}	0.400 ^{NS}
	2023	0.209 ^{NS}	-0.479 ^{NS}	-0.757**	0.687**	-0.241 ^{NS}	0.355 ^{NS}	-0.481*	0.593**
	Pooled	0.232 ^{NS}	-0.345 ^{NS}	-0.741**	0.663**	-0.231 ^{NS}	0.184 ^{NS}	-0.484*	0.492*

*Significant at 5% level of significance; **Significant at 1% level of significance; NS= Non-significant

References

1. Fukagawa NK, Ziska LH. Rice: Importance for global nutrition. *Journal of Nutritional Science and Vitaminology*. 2019; 65(Supplement):S2-S3.
2. FAO 2021. Food and Agriculture Organization of the United Nations. <http://www.fao.org/faostat/en/#data/QCL>.
3. Anonymous; 2021a. State/Season-wise Area, Production and Productivity of Rice in India (2020-2021). DES, MoA&FW, GoI. <https://www.indiastatagri.com/table/agriculture/state-season-wise-area-production-productivity-ric/1423615> (Accessed on 26-05-2023)
4. Anonymous. 2021b; Agricultural Statistics. Farmer Welfare and Agriculture Development Department, M.P. https://mpkrishi.mp.gov.in/Englishsite_New/pdfs/201920_n.pdf.
5. Pasalu IC, Katti G. Advances in ecofriendly approaches in rice IPM. *Journal of Rice Research*. 2006;1(1):83-90.
6. Dhaliwal GS, Jindal V, Dhawan AK. Insect pest problems and crop losses: changing trends. *Indian Journal of Ecology*. 2010;37(1):1-7.
7. Kumar SD. Rice Production in India: Analysis of Trend, Constraints and Technologies. In: *Climate Change and Future Rice Production in India*. India Studies in Business and Economics. Springer, Singapore. 2019; https://doi.org/10.1007/978-981-13-8363-2_2
8. Gupta SP, Singh RA, Singh AK. Field efficacy of granular insecticides and single compound sprays against pests in rice. *Indian Journal of Entomology*. 2006;68(2):150-151.
9. Gagan J, Lakhi R, Ram S. Population dynamics of paddy stem borers in relation to biotic and abiotic factors *Annals of Biology*. 2009;25(1):47-51.
10. Sampath KM, Chitra S, Mohan M, Padmavathi, Subaharan K, Katti G. Emergence pattern reproductive biology and courtship behaviour of rice pink stem borer, *Sesamia inferens* (Walker) (Noctuidae; Lepidoptera). *Agrotechnology*. 2014;2(4):60.
11. Kumar A, Misra AK, Satyanarayana P, Choudhary SK. Dynamics of Species Composition of Stem Borers in Rice Crop. *The Ecoscan*. 2016;10(1&2):223-226.
12. Pallavi D, Sharanabasappa, Megaladevi P. Relative abundance of yellow stem borer and pink stem borer on paddy. *Journal of Entomology and Zoology Studies*. 2018;6(4):668-67.
13. Sharma AK, Kumar N, Naveen, Tare S, Nayak S, Seervi S. Phototactic Response and Taxonomic Distribution of Predaceous Species of Paddy Ecosystem. *Biological Forum – An International Journal*. 2023;15(3):91-94.
14. Mishra YK, Sharma AK, Sharma K. Efficacy of combination insecticides against rice stem borer *Scirpophaga incertulas* Wlk. *Indian Journal of Entomology*. Online published. 2021; Ref. No. e21012.
15. Mishra YK, Sharma AK, Bhowmick AK, Saxena AK, Kurmi A. Seasonal Incidence of Insect Pest Species of Paddy Collected through Light Trap. *International Journal of Current Microbiology and Applied Sciences*. 2019;8(4):381-393.
16. Sharma AK, Mandloi R, Bisen UK, Thakur AS. Relative analysis on phototactic insect pests and predatory species of paddy ecosystem. *Journal of Zoology and Entomology Studies*. 2019;7(1):1547-1551.
17. Meena SK, Sharma AK, Aarwe R. Seasonal Incidence and Population Dynamics of Major Insect Pest Species of Paddy Collected in Light Trap in Relation to weather Parameters. *International Journal of Microbiology and Applied Sciences*. 2018;7(8):1705-1715.
18. Singh S, Sharma AK. Population dynamics of major insect pests of rice. *Indian Journal of Entomology*. 2018;80(4):1700-1702.
19. Sharma AK, Mandloi R, Bisen UK. Effect of abiotic factors on seasonal fluctuation of major phototactic insect pests of rice. *Journal of Zoology and Entomology Studies*. 2017;8(6):160-165.
20. Sharma AK, Muchhala Y, Pachori R. Impact of Abiotic Factors on Population Dynamics of Major Predatory and Parasitic Fauna of paddy. *The Ecoscan*. 2015;9(1&2):597-600.
21. Sharma AK, Bisen S, Bisen UK. Comparative Analysis on Activity of Major Predatory and Insect Pest Species of Paddy in two Distinct (Forming-Ecological) Locations through Light Trap. *The Ecoscan*. 2015;9(1&2):81-84.
22. Kok LT, Varghese C. The four major lepidopterous rice stem borer in Malaya. *Malaya Agriculture Journal*. 1996;45(3):275-288.
23. Mandal P, Roy K, Saha G. Weather based prediction model of *Scirpophaga incertulas* (Walk.). *Annual Plant Protection Science*. 2011;19(1):20-24.
24. Sharma MK, Atsedewoin A, Fanta S. Forewarning models of the insects of paddy crop. *International Journal of Biodiversity and Conservation*. 2011;3(8):367-375.

25. MishraMK, SharmaRC, SinghRB, SinghRP. Monitoring of yellow stem borer, *Scirpophaga incertulas* Walker in rice through light and pheromone traps. *Agriculture & Biological Research*. 2012;28(2):135-139.
26. KumarA, MisraAK, SatyanarayanaP, KumarJ. Population dynamics and management of yellow stem borer (*Scirpophaga incertulas* Walker) with insect sex-pheromone trap. *International Journal of Plant Protection*. 2015;8(1):157–161.
27. SomashekaraH, Javaregowda. Population build-up of paddy yellow stem borer (*Scirpophaga incertulas* Walker) in relation to different weather parameters. *Karnataka Journal of Agricultural Sciences*. 2015;28(2):282-283.
28. ChatterjeeS, DanaI, GangopadhyayC, MondalP. Monitoring of yellow stem borer, *Scirpophaga incertulas* (Walker) using pheromone trap and light trap along with determination of field incidence in kharif rice. *Journal of Crop and Weed*. 2017;13(3):156-159.
29. Mondal IH, Chakraborty K. Observation on the impact of environmental parameters on rice yellow stem borer, *Scirpophaga incertulas* (Walker) and its natural enemies at Murshidabad, West Bengal, India. *Journal of Entomology and Zoology Studies*. 2017;5(6):1656-1663.
30. Roshan DR, Raju SVS, Singh KN. Effect of environmental factors on population dynamics of *Scirpophaga incertulas* Walker and its management with novel insecticides in rice ecosystem. *Journal of experimental Zoology*. 2016;19(1):327-332.
31. Hatwar NK, Jalgaonkar VN, Wade PS, Naik KV, Thantharate SH, Kinjale RS. Seasonal incidence of yellow stem borer, *Scirpophaga incertulas* Walker infesting rice and its correlation with weather parameters. *Journal of Entomology and Zoology Studies*. 2021;9(1):263-266.