

Exploring the Potential of Nanotechnology in Fisheries and Aquaculture: Opportunities and Implications

Abstract

Aquaculture has emerged as one of the fastest-growing global food industries in recent years, playing a vital role in meeting the increasing demand for animal protein. However, several challenges, such as disease prevalence, chemical contamination, environmental degradation and inefficient feed utilization, have significantly impeded the sector's ability to contribute effectively to global food security. To address these challenges, considerable efforts have been made to leverage advancements in science and technology within the field of aquaculture. Nanotechnology has emerged as a promising and effective tool to tackle these challenges across various aspects of aquaculture, including fish nutrition, biotechnology, genetics, reproduction, pathology and environmental sustainability. The utilization of emerging nano-materials, such as nano-elements enriched feed, has notably enhanced fish growth and has been applied in aquatic systems to mitigate pollutants, thereby reducing treatment costs. Furthermore, the integration of genetically manipulated techniques with nano-biotechnology has led to significant advancements in fish genetics research. Innovative applications of nanotechnology, such as nano-sensors, DNA nano-vaccines and drug delivery systems, have revolutionized fish health, reproduction and immune system management. Additionally, nanotechnology is increasingly being employed in the fish processing industry to ensure sterile packaging, precise flavoring and high-quality products. Moreover, the utilization of bio-nano-engineering techniques for optimal utilization of fishery wastes and the implementation of green nanoparticles signify a promising era in post-harvesting practices. This comprehensive review aims to delve into nanotechnology's diverse and promising applications in aquaculture, exploring its potential and the challenges it presents.

Keywords:

1. Introduction

Aquaculture stands out as the most rapidly expanding sector within agriculture, boasting considerable growth prospects. It serves as a vital source of easily digestible protein, healthy fat,

and a diverse array of essential micronutrients for nourishing worldwide populations[1]. Moreover, aquaculture holds significant potential for rural employment, livelihood improvement and offers substantial opportunities for generating income through high export earnings, thereby boosting national GDP. The production of farmed finfish and shellfishes has experienced rapid growth, it is because of adoption of advanced and intensive aquaculture techniques[2]. As aquaculture farming systems have expanded, there has been a reduced dependence on wild seedstock and fish supplies[3]. The intensive practice of aquaculture results in pond eutrophication, characterized by the accumulation of an excessive nutrient load[4]. The escalating impacts of global climate change, such as rising temperatures and increased levels of methane and CO₂, pose a significant threat to aquaculture[1].

Nanotechnology has arisen as a hopeful solution, introducing fresh avenues for exploration. It encompasses the development and utilization of materials at nanoscale dimensions (1-100 nm) with exceptional properties, offering novel possibilities for application[5]as illustrated in Fig. 1. Nanotechnology applications have permeated every aspect of human life, yielding numerous value-added products [6].Becoming a crucial interdisciplinary field, nanotechnology seamlessly integrates principles from physics, chemistry and biology[7]. Furthermore, nanotechnology garners global public interest due to its rapid evolution and widespread applications across various fields including engineering, electronics, agriculture, medicine, food industry and environmental sectors[8]. Nano-sized materials possess unique physicochemical properties that enable them to withstand high pressure and temperature, facilitating their involvement in numerous applications[9]. The revolution of nanotechnology has expanded to encompass fish farming, contributing to their growth and addressing associated challenges[10,11]. Nanotechnology applications in fisheries comprise both direct and indirect approaches.

Direct applications of nanotechnology primarily focus on enhancing various aspects of fish, including growth, reproduction and health. Nano-materials play a crucial role in efficiently delivering nutrients, trace elements and vitamins as feed supplements, such as selenium (Se) [12–14], iron (Fe) [15,16], zinc (Zn) [17–19]and vitamin-C[20,21].

Feeding fish with nanoform of these ingredients enhances their absorption and allows them to pass through the intestinal wall, thereby improving growth performance, reproduction, and innate immunity in fish [22,23]. Utilizing various nano-materials for controlling fish diseases proves to

be more effective than traditional antibiotics and chemicals, which often lead to adverse side effects like the development of resistant bacterial strains, water pollution and the accumulation of unwanted chemical residues. Nano-materials exhibit antiviral properties [24], act as antibacterial agents [25–28], demonstrate antifungal effects [29,30] and serve as anti-parasitic agents [31,32]. Additionally, chitosan and poly lactic-glycolic acid (PLGA) nanoparticles are significant in facilitating drug and hormone delivery, as well as vaccination [33–35]. Certainly, nanovaccination offers numerous advantages compared to conventional methods, ensuring sustained release and enhancing stability, bioavailability and residence time [36–38]. Moreover, a notable application of nanotechnology in fish farming is its utilization for rapid and efficient diagnosis of various fish pathogens [39,40].

In today's era, nanotechnology has flourished into a multi-billion-dollar industry, experiencing rapid growth exemplified by the presence of over a thousand products incorporating nanomaterials in the market. Over the past decade, the international market has witnessed the introduction of more than 300 nanofood products [41]. The economic influence of nanotechnology sectors was estimated to exceed \$3 trillion by 2020, with approximately 6 million individuals employed within these industries [8]. Currently, there exists an abundance of published literature offering a thorough examination of the various uses of nanotechnology within the field of aquaculture (Fig. 2.) [42].




Fig. 1. Applications of nanotechnology in aquaculture

2. Enrichment in feed

In natural habitat, fish source their food from phytoplankton, zooplankton and smaller organisms. However, in aquaculture settings, it becomes necessary to add complementary additives in fish feed to significantly increases the growth and production. Fish feed typically necessitates a balance of protein (32%), carbohydrates (20–35%), fats (4–6%), fiber (<4%), and dietary energy (8.5–9.5%) and these nutritional requirements may vary depending on the species and developmental stage offish[43]. Inclusion of minute concentrations of organic and/or inorganic substances into fish feed has been a common practice to enhance growth and immunity. The use of feed supplements stands out as one of the crucial applications of nanotechnology in fish farming. Nano-scale materials possess the advantage of being readily absorbed in small doses, traversing the gastrointestinal tract and small intestine to enter into the bloodstream. Once distributed throughout vital organs, these nano-materials function more effectively compared to their bulk counterparts [22]. Introducing nano-metals such as selenium (Se), zinc (Zn) and iron (Fe) as feed supplements can offer several benefits for the survival,

growth, and overall health of fish[23]. Additionally, organic polymers such as chitosan play a significant role in the delivery of micronutrients [10].

Nutritional supplements and feed formulation are vital in commercial aquaculture for fish growth. Nanoparticles can modify feed consumption by enhancing flavor, color, or attractants. They also improve the bioavailability of water-insoluble vitamins and carotenoids when processed with nanoparticles for dietary supplements [44,45]. Several market-available products utilize advanced micro/nano-encapsulation systems to enhance the delivery and bioavailability of nutrients in aquaculture. For instance, UbiSol-Aqua™ Delivery System Technology, created by Zymes LLC (USA) and NovaSOL from AQUANOVA (Germany). UbiSol-Aqua™ efficiently solubilizes various bioactive compounds such as coenzyme Q10, vitamins A and B, squalene and fish oil (EPA/DHA). AQUANOVA's 'solubilisate' liquid carrier solutions enable the transportation of active raw materials in highly compact capsules[46]. Nano/microcarriers play a crucial role in protecting, encapsulating, stabilizing and delivering valuable bioactive compounds essential for fish growth. Numerous Indian startups have ventured into manufacturing nanotechnology products for aquaculture nutrition. Notably, Filo Life Sciences, an Indian origin nanotech company, produces a range of important nutritional and nutraceutical products for aquaculture including ColloidAG Aqua (Nano silver), Fabgrow Aqua (Nano PUFA), Nanomin Aqua (Nano trace minerals), NannoCAL Aqua (Nano calcium), NanoPHOS Aqua (Nano phosphorus)[47].




Fig. 2. Visual depiction showcasing the implementation of nanotechnology within the realm of fishery.


3. Water filtration and remediation

Water treatment is crucial for sustainable aquaculture. Global water contamination, exacerbated by urban, industrial and agricultural waste discharge, as well as the misuse of antibiotics and synthetic compounds in fisheries, poses a significant health hazard for consumption[42].

Nano-enabled technologies are utilized for water purification, employing nanomaterials like activated carbon or alumina with additives such as zeolite and iron-containing compounds to foster aerobic and anaerobic biofilm for ammonia, nitrites and nitrate removal in aquaculture. Additionally, ultrafine nanoscale iron powder is effective in cleaning less toxic carbon compounds like trichloroethane, carbon tetrachloride, dioxins and polychlorinated biphenyls, advancing nano-aquaculture practices [48].

Researchers created magnetic konjac glucomannan (KGM) aerogels for arsenite water decontamination. The system exhibited a pH-dependent capacity and green step characteristics[49]. Recently, graphene oxide (GO) and graphene nanosheets (GNs) have garnered significant global attention for their effectiveness in removing various contaminants from water[50,51]. Hybrid GO-TiO₂ materials have been particularly focused on for environmental and energy applications, including adsorption of heavy metal ions and organic dyes from wastewater[52,53]. TiO₂ is a promising candidate for wastewater treatment due to its non-toxic nature, chemical and biological stability, low cost and efficient photocatalytic properties. Numerous studies have explored its photocatalytic activity for inactivating pathogenic organisms like bacteria, viruses and algae[52,54,55]. The scientific community's attention to water treatment and contaminant removal is appreciated. Elevated concentrations of these contaminants pose potential threats to human health, as they can accumulate in aquatic animal tissues, notably fish, which are often consumed to combat cardiovascular diseases (CVDs) and cancer[56]. Due to their habitat and feeding habits, fish are highly vulnerable and extensively exposed to the harmful effects of pollutants in aquatic environments, with no means of escaping their impacts[57,58]. Reports indicate increased accumulation of heavy metals (Hg, Cd, and Pb) in marine animal tissues, attributed to both natural processes (e.g., volcanic activity) and human

activities [59]. Likewise, research has shown that F-1 toxicity leads to enzyme dysfunction, gastric issues and immune system impairment in experimental fish[60], as well as habitat degradation and destruction of the freshwater snail *Physella acuta*[61].Utilization of a 3D RGO hydrogel assessed for removing Hg and F⁻¹ from water. The aerogel showed high adsorption capacities of 185 mg/g for Hg⁺² and 31.3 mg/g for F⁻¹, indicating potential for environmental pollution



management[62].Additionally, Figure 2 demonstrates the suggested mechanism by which various nanoparticle-based photocatalytic adsorbents and hydrogel biofilms function effectively in the practical purification of water, showcasing examples of fluoride (F⁻), nitrate (NO₃⁻), and coliforms (*E. Coli*) removal from contaminated water[42].

Fig. 3. The operational mechanism of nanoparticle-loaded adsorbents and hydrogel films for eliminating F⁻, NO₃⁻, and coliforms (*E. coli*) from polluted water.

4. Fish harvesting

Fishing lures are often painted to attract fish with light, but they usually only reflect light in one direction. To improve effectiveness, lures are colored and coated with a polyimide film, increasing fish-catching probability by two to three times compared to uncoated lures[48].

5. Biofouling control

Nanotechnology can enhance aquaculture and shrimp culture productivity by improving disease control, feed formulation, and biofouling management. By incorporating metal oxide nanoparticles like ZnO, CuO and SiO₂ into coatings or paints, biofouling from unwanted bacteria, invertebrates (such as mussels and barnacles) and algae (like seaweeds and diatoms) can be monitored and controlled effectively. This creates antifouling surfaces and boosts antifouling control efficiency [63,64]. The antifouling technology can be applied in fishing and aquaculture networks, aquaculture tank antibacterial substances and new marine product packaging materials.

6. Removal of heavy metals

Ligand-based nano coating enables efficient and cost-effective heavy metal removal by high absorption capacity and renewability. Crystal clear technology purifies water by bonding multiple metal layers to one substrate[65]. Nanomaterials, like metal oxide nanoparticles, are commonly used for efficient heavy metal removal from water due to their high reactivity and large surface area. Ongoing research aims to develop new synthesis methods and practical applications, such as composite materials or granular oxides, to enhance heavy metal removal effectiveness and understand underlying mechanisms such as XAS and NMM[66,67]. Comprehensive research investigating the impact of humic acid and fulvic acid on heavy metal removal using a range of nanomaterials from aqueous solutions. The study focused particularly on iron-based, carbon-based and photocatalytic nanomaterials [68]. The study also examined interaction mechanisms and environmental implications of humic acid and fulvic acid. Chitosan nanoparticles are utilized as adsorbents for heavy metal removal. Recent research focuses on removing heavy metals from clays like kaolinite, bentonite and montmorillonite using chitosan nanoparticles, with studies on nano chitosan-clay composites for metal ion removal recently reported [69,70]. Chitosan-magnetite nanocomposites were also suggested for removing heavy metals from aqueous solutions[71,72].A schematic illustration is provided to highlight the process of zinc oxide nanoparticle-mediated degradation of significant pollutants (Fig. 4.)[47].




Fig. 4. Nanoremediation by zinc oxide nanoparticles.

7. Fish reproduction

Producing monosex tilapia (*Oreochromis sp.*) fish farms aim to prevent uncontrolled spawning and overcrowding. This is achieved through the application of steroid hormones, which can have unintended effects on fish consumers[73]. Nanotechnology has played a role as an effective drug carrier in delivering fadrozole (an estrogen synthesis inhibitor) to *O. niloticus*. Feeding *O. niloticus* with fadrozole overloaded on PLGA nanoparticles at 50–500 ppm concentrations for one month resulted in 100% male fish at both 350 and 500 ppm levels[74]. Nanotechnology plays a crucial role in improving reproduction in various fish species. Specifically, nano-chitosan has been utilized to deliver and release reproductive hormones[75]. A composite of chitosan-nano gold extended the duration of reproductive hormone presence in salmon blood and enhanced egg fertilization rates[35]. Nanoparticles of eurycomanone, extracted from *Eurycoma longifolia*, were combined with chitosan nanoparticles and injected into *Clarias magur* fish. Within 7 days, this resulted in increased gonado-somatic index (GSI), Ca and Se concentrations, reproductive capacity and endocrine hormone gene expression levels [33]

8. Nanotechnology for fish quality testing

Quality testing is an essential post-harvesting technology and nanotechnology can be applied for quality improvement of fishes.

8.1. Fish freshness testing

A quantum dot-based nanosensor was created to address fishery product freshness concerns. Tin oxide quantum dots (SnO₂ QDs) were synthesized, dispersed in a colloidal solution and deposited onto indium–tin-oxide (ITO) glass[47]. This facilitated the development of a fish freshness biosensor by immobilizing xanthine oxidase. The sensor displayed heightened sensitivity, faster response time and an extensive linear range in electrochemical output [76].

8.2 Determination of formalin in fish samples

Formalin poses a significant threat to fish food safety in modern times. A market survey conducted in Bangladesh revealed formalin contamination in 50% of total fish samples, with 70% of rohu (*L. rohita*) affected. This preservative is commonly used to maintain freshness during transportation [77]. A report from the Somdet market in Kalasin Province, Thailand, highlighted the varying levels of formalin contamination across different fish products. White shrimp, mackerel, Shishamo fish, squid and other samples were found to have higher contamination levels[78]. A formaldehyde nano-biosensor was designed using formaldehyde dehydrogenase enzyme and nanomaterials like carbon nanotubes and chitosan. This sensor ensures precise detection of the health hazard, with quick response times (≥ 5 s), high sensitivity (1-10 ppm) and reproducibility [79]. A reproducible formaldehyde biosensor was created by coating an ionic liquid, nanoscale gold, and chitosan onto a glassy carbon electrode. This biosensor detects formalin in the tissues of *Lutjanus malabaricus* and *Thunnus tonggol*[80].

Fish packaging

Nanotechnology enhances fish packaging by delaying spoilage and extending shelf life. Nanomaterials in biodegradable packaging act as antimicrobial agents, remove oxygen, inhibit degradation and stabilize products[81–83]. Nano-composites derived from natural biopolymers (proteins, lipids, polysaccharides) are favored as healthy packaging alternatives to toxic petrochemical plastics[84].


Delivery of vaccines

Vaccines are essential in aquaculture for defending against pathogens and safeguarding host animal from different infections. Oral or injectable methods are the most reliable ways of

vaccination in fisheries. However, traditional injection practices with oil/water formulations often lead to adverse effects and occasional fish mortality [85]. To address these issues, the scientific community has proposed nano-delivery systems as safer and more effective alternatives for vaccine delivery in fish. Various encapsulation techniques, including alginate particles, have been developed and tested for oral vaccine delivery to aquatic animals[86]. Alginate, derived from brown algae or bacterial polysaccharides, is a copolymer of β -D-mannuronic acid (M) and α -L-guluronic acid (G). Its mechanical stability and mucoadhesive properties make it ideal for oral administration, enabling direct contact with epithelial cell walls[87,88]. For application in fish, alginate particles are generally produced by emulsification[89]that is one of the fastest methods for NP preparation and is readily scalable[90]and to a lesser extent by other methodologies such as the orifice-ionic gelation and the spray method. Reports from different researchers presented alginate as an antigen adjuvant [91,92], survival and weight promoter of fish[93,94]. Additionally, alginate administration has demonstrated enhanced immune-stimulant responses in carp (*Cyprinus carpio* L.) and brown-marbled grouper (*Epinephelus fuscoguttatus*)[95–97]. It has also been shown to bolster defense against *V. anguillarum* in turbot (*Scophthalmus maximus* L.)[98]and against iridovirus and *Streptococcus sp.* in orange-spotted grouper (*Epinephelus coioides*) and brown-marbled grouper[95,97]. Chitosan (CS), derived from crustaceans and insects, is the second most abundant biopolymer. Its unique biological properties: bioadhesive, biodegradable, biocompatible and non-toxic—make CS-based formulations widely used in drug delivery, bio-nanosensors, edible coatings and various medical fields such as dentistry and surgery[99,100]. Furthermore, the oral administration of nano-chitosan encapsulated DNA or overloaded with DNA or recombinant DNA vaccine has been shown to positively enhance the immune response against certain fish pathogens, such as the European sea bass (*Dicentrarchus labrax*) against nodavirus[101], Asian sea bass (*Lates calcarifer*) against *V. anguillarum*[37], black seabream (*Acanthopagrus schlegelii*) against *V. parahemolyticus*[102]and Shrimp (*P. monodon*) against the white spot syndrome virus[63]. Similarly, nano-PLGA was employed in oral DNA vaccines for Japanese flounder (*Paralichthys olivaceus*) against lymphocytic disease virus[103]and for rainbow trout (*Oncorhynchus mykiss*) against infectious hematopoietic necrosis virus[104]. The use of nanotechnology in fish vaccination is crucial for reducing reliance on toxic and

carcinogenic chemical adjuvants. Oral or immersion vaccinations are preferred over injections to minimize stress[38].

Nanoparticles offer promising potential for enhancing siRNA delivery systems in molecular therapeutics. An assessment examined PAMAM nanoparticle-mediated delivery of siRNA to zebrafish cardiac wounds, quantitatively analyzing nanoparticle distribution in cardiac cells and



monitoring their role in cellular uptake[105]. A schematic diagram of nanoparticle-mediated siRNA delivery in the fish system is presented as Fig. 5.

Fig. 5. Nanoparticle mediated Si RNA delivery in fish system

Nanotechnology devices for aquatic environment management

Nanotechnology application in seawater shrimp aquaculture demonstrated that nanodevices reduced water exchange rates, improved water quality, increased shrimp survival rates and ultimately boosted yield [106,107]. Among various nanodevices, nanonet treatment stood out as the most effective, with a 100% increase in fish survival rate and significant reductions in water nitrite and nitrate levels, with nitrite dropping to just a quarter of the control group. Nanotechnology also raised water pH and substantially enhanced water efficiency [108]. In China, Nano-863 is a popular high-tech agricultural product widely utilized in livestock, crop and aquaculture sectors. It incorporates high-temperature sintered nanomaterials with excellent light-absorbing properties into a ceramic carrier[106].

Nanotechnology as a new tool in fish diseases

Nanotechnology operates at the nanometer scale, with one nanometer (nm) equaling one billionth of a meter, according to the International System of Units (SI)[84,109]. New materials, with distinct properties derived from their small size, form, surface area, conductivity, or surface chemistry, find extensive applications in textiles, electronics, engineering and medicine [110,111]. The term "nanomaterial" derives from "nano," originating from the Greek word for "dwarf." It typically refers to materials ranging in size from 1 to 100 nm[112]. Nanotechnology offers solutions to prevent and monitor diseases in aquaculture, enhancing its benefits. This includes developing antibacterial surfaces, using nanosensors for pathogen detection and delivering veterinary products through fish foods[113]. Nano-trace elements exhibit usage levels up to 100% higher than conventional inorganic trace elements, primarily due to their direct penetration into the animal body [11,42,113].

Aquaculture scientists are developing a sensitive immunodiagnosics protocol using nanoscale gold attached to alkaline phosphatase (ALP) conjugated antibodies for early detection of the white spot syndrome virus (WSSV) in shrimp[114]. Nanosensors provide effective solutions for identifying pathogens, including important aquaculture viruses like aquabirna, salmonid alpha and betanoda[115]. Graphene oxide (GO) enables rapid and real-time detection of white spot syndrome virus (WSSV) in shrimp samples through electrochemical immunosensors, with an exceptionally low detection limit of 1.36×10^{-3} copies μL^{-1} , offering an alternative to PCR-based techniques[116].

Drug delivery for health management

Disease outbreaks pose significant obstacles to the sustainability and development of aquaculture[11,48,117]. Nanotechnology plays a crucial role in providing novel perspectives for disease diagnosis and health management in aquaculture [64]. Solid core drug delivery systems, involving coating solid nanoparticles with a fatty acid shell, are effective for protecting labile or thermo-sensitive drugs[118]. Porous nanomaterials serve as effective pharmaceutical delivery matrices. Mesoporous silica particles, for example, can be utilized for the controlled release of drugs[119].

Oral nano-delivery systems associated with nanoparticles offer several advantages, such as enhanced control over drug release[120], direct targeting of specific tissues[121], improved

bioavailability of pharmaceuticals with low absorption rates[122], stabilization of drugs through prolonged residence time in the gut[123]and increased absorption capability due to higher dispersion rates at the molecular level[124].

Delivery of nutrients

While nutraceuticals significantly enhance growth and immunological parameters in fish, their incorporation entails higher costs compared to minimal dietary requirements. Thus, careful management is essential to avoid wastage and maximize their utilization[125]. Nanotechnology enhances the delivery of dietary supplements and nutraceuticals in fisheries, improving nutrient bioavailability and efficacy by enhancing solubility and protecting against harsh gut conditions. For instance, adding 1 mg of nano-Selenium (Se) per kg of diet significantly boosts growth and antioxidant defense in common carp (*Cyprinus carpio*) compared to controls[12]. Supplementation of selenium (Se), zinc (Zn) and manganese (Mn) nanoparticles in early weaning diets enhanced stress resistance and bone mineralization in gilthead seabream (*Sparus aurata*)[126]. Solid lipid nanoparticles (SLNPs) containing 6-COUM demonstrated superior uptake compared to competitor 6-coumarin-loaded pectin microparticles (MPs) in gilthead seabream (*Sparus aurata* L.) cell types, including SAF-1 cells and primary head-kidney (HK) cultures. This suggests SLNPs as promising nanocarriers for delivering biologically active substances in fish [127]. In rainbow trout, the addition of iron nanoparticles (NPs) and *Lactobacillus casei* as a probiotic to the diet led to notable enhancements in growth parameters [128]. Similarly, supplementation of the diet with 16 mg kg⁻¹ of MnO nanoparticles (NPs) significantly boosted both growth and the antioxidant defense system in freshwater prawn (*Macrobrachium rosenbergii*) [129]. Copper nanoparticles at 20 mg kg⁻¹ notably enhanced growth, biochemical constituents, digestive enzymes, antioxidants, metabolic enzymes and immune response in freshwater prawn post-larvae (*Macrobrachium rosenbergii*) [130]and red sea bream, (*Pagrus major*) [131]. The hepatoprotective and antioxidant effects of *Azolla microphylla*-based gold nanoparticles (GNPs) against acetaminophen (APAP)-induced toxicity in freshwater common carp fish (*Cyprinus carpio* L.) has been investigated [132].GNPs notably improved metabolic enzyme levels, hepatotoxic markers, oxidative stress markers, tissue enzymes, hepatic ion levels, and abnormal liver histology. Thus, *Azolla microphylla* phytochemically synthesized GNPs were recommended as effective protectors against

acetaminophen-induced hepatic damage in freshwater common carp fish [132]. The effects of different levels (0.5%, 1% and 1.5% of the diet) of Aloe vera nanoparticles on the growth, survival rate and body composition of Siberian sturgeon examined. They found that supplementing the diet with 1% Aloe vera nanoparticles significantly improved the growth indices of Siberian sturgeon compared to controls[133].

Toxicity

Nano-materials' heightened reactivity and unique properties generate increased production of reactive oxygen species, leading to cellular dysfunction as they easily penetrate cell membranes and interact with intracellular organelles[134]. In general, when comparing the physiological effects of metals and nanometals, it is found that nanometals cause more toxic effects[135]. Various aquatic organisms exhibit diverse toxic responses to both nanometals and bulk forms, with metal nanoparticles demonstrating notably higher toxicity compared to other nanoparticle types. Exposure of Medaka fish (*Oryziaslatipes*) to nano-Se (3.2 mg/l) for 2 days resulted in 100% mortality, whereas sodium selenite at 2.0 mg/l caused only 10% mortality and 80% at 8.0 mg/l. Nano-Se accumulated six times more in fish liver compared to selenite[136]. Certain nano-materials employed in water remediation, such as nZVI, have been documented to induce toxicity in microorganisms, crustaceans, fish larvae, and other aquatic and soil-dwelling organisms[137]. Zebrafish exposed to both titanium dioxide nanoparticles (nano-TiO₂) and cypermethrin experienced increased accumulation of cypermethrin, leading to neurotoxic effects and reduced fish larvae locomotion[138]. Additionally, nano-TiO₂ enhanced uptake of an organophosphate compound in zebrafish tissues, decreasing reproductive hormones and inhibiting reproduction [139].

Conclusion

The emergence of nanotechnology presents a dual-edged sword, showcasing both its immense potential and inherent risks. The unique properties of nanoparticles give rise to both positive and negative effects. In the realm of fish culture, nanotechnology holds promise for revolutionizing various aspects, including enhancing growth performance, productivity, disease control and diagnosis, water purification, pollutant remediation and extending shelf life through improved packaging techniques. Despite its potential benefits, the application of nanotechnology in fish

culture poses challenges, primarily concerning the potential toxicity of nanoparticles and their adverse effects on fish and non-targeted organisms, as well as their environmental impact. However, the concept of green nanotechnology offers eco-friendly alternatives to toxic metal nanoparticles. To ensure the safe and responsible application of nanoparticles in fish culture, it is imperative to address existing gaps in knowledge regarding the environmental fate of nanoparticles and their potential accumulation in the food chain. Comprehensive assessments are necessary to guarantee the safety of nanoparticle usage for fish, environmental sustainability and human consumption.

References

- [1] (Fisheries and Aquaculture Department) FAO, The future of food and agriculture trends and challenges, Food Agric. Organ. U. N. Rome (2017).
- [2] FAO, The state of world fisheries and aquaculture 2022. Towards blue transformation, Food and Agriculture Organization of the United Nations Rome, Italy, 2022.
- [3] R.L. Naylor, R.J. Goldberg, J.H. Primavera, N. Kautsky, M.C. Beveridge, J. Clay, C. Folke, J. Lubchenco, H. Mooney, M. Troell, Effect of aquaculture on world fish supplies, *Nature* 405 (2000) 1017–1024.
- [4] L.S. Herbeck, D. Unger, Y. Wu, T.C. Jennerjahn, Effluent, nutrient and organic matter export from shrimp and fish ponds causing eutrophication in coastal and back-reef waters of NE Hainan, tropical China, *Cont. Shelf Res.* 57 (2013) 92–104.
<https://www.sciencedirect.com/science/article/pii/S0278434312001288> (accessed February 25, 2024).
- [5] F. Matteucci, R. Giannantonio, F. Calabi, A. Agostiano, G. Gigli, M. Rossi, Deployment and exploitation of nanotechnology nanomaterials and nanomedicine, in: *AIP Conf. Proc.*, AIP Publishing, 2018.
- [6] S.T. Stern, M.N. Martinez, D.M. Stevens, When is it important to measure unbound drug in evaluating nanomedicine pharmacokinetics?, *Drug Metab. Dispos.* 44 (2016) 1934–1939.
- [7] K. Subramani, A. Elhissi, U. Subbiah, W. Ahmed, Introduction to nanotechnology, in: *Nanobiomaterials Clin. Dent.*, Elsevier, 2019: pp. 3–18.
- [8] X. He, H. Deng, H. Hwang, The current application of nanotechnology in food and agriculture, *J. Food Drug Anal.* 27 (2019) 1–21.
- [9] V. Aruoja, H.-C. Dubourguier, K. Kasemets, A. Kahru, Toxicity of nanoparticles of CuO, ZnO and TiO₂ to microalgae *Pseudokirchneriella subcapitata*, *Sci. Total Environ.* 407 (2009) 1461–1468.

- [10] K. Khosravi-Katuli, E. Prato, G. Lofrano, M. Guida, G. Vale, G. Libralato, Effects of nanoparticles in species of aquaculture interest, *Environ. Sci. Pollut. Res.* 24 (2017) 17326–17346.
- [11] A.I.S. Luis, E.V.R. Campos, J.L. de Oliveira, L.F. Fraceto, Trends in aquaculture sciences: from now to use of nanotechnology for disease control, *Rev. Aquac.* 11 (2019) 119–132.
- [12] S. Ashouri, S. Keyvanshokoo, A.P. Salati, S.A. Johari, H. Pasha-Zanoosi, Effects of different levels of dietary selenium nanoparticles on growth performance, muscle composition, blood biochemical profiles and antioxidant status of common carp (*Cyprinus carpio*), *Aquaculture* 446 (2015) 25–29.
- [13] K.U. Khan, A. Zuberi, S. Nazir, I. Ullah, Z. Jamil, H. Sarwar, Synergistic effects of dietary nano selenium and vitamin C on growth, feeding, and physiological parameters of mahseer fish (*Tor putitora*), *Aquac. Rep.* 5 (2017) 70–75.
- [14] I.F. Xia, J.S. Cheung, M. Wu, K.-S. Wong, H.-K. Kong, X.-T. Zheng, K.-H. Wong, K.W. Kwok, Dietary chitosan-selenium nanoparticle (CTS-SeNP) enhance immunity and disease resistance in zebrafish, *Fish Shellfish Immunol.* 87 (2019) 449–459.
- [15] P. Akbary, A. Jahanbakhshi, Nano and macro iron oxide (Fe_2O_3) as feed additives: effects on growth, biochemical, activity of hepatic enzymes, liver histopathology and appetite-related gene transcript in goldfish (*Carassius auratus*), *Aquaculture* 510 (2019) 191–197.
- [16] V. Srinivasan, P.S. Bhavan, G. Rajkumar, T. Satgurunathan, T. Muralisankar, Effects of dietary iron oxide nanoparticles on the growth performance, biochemical constituents and physiological stress responses of the giant freshwater prawn *Macrobrachium rosenbergii* post-larvae, *Int J Fish Aquat Stud* 4 (2016) 170–182.
- [17] A. Awad, A.W. Zagloul, S.A. Ahmed, S.R. Khalil, Transcriptomic profile change, immunological response and disease resistance of *Oreochromis niloticus* fed with conventional and Nano-Zinc oxide dietary supplements, *Fish Shellfish Immunol.* 93 (2019) 336–343.
- [18] H. Faiz, A. Zuberi, S. Nazir, M. Rauf, N. Younus, Zinc oxide, zinc sulfate and zinc oxide nanoparticles as source of dietary zinc: comparative effects on growth and hematological indices of juvenile grass carp (*Ctenopharyngodon idella*), *Int. J. Agric. Biol.* 17 (2015).
- [19] M. Tawfik, M. Moustafa, I.M.K. Abumourad, E. El-Meliegy, M. Refai, Evaluation of nano zinc oxide feed additive on tilapia growth and immunity, in: 15th Int. Conf. Environ. Sci. Technol. Rhodes Greece, 2017: pp. 1–9.
- [20] A. Alishahi, A. Mirvaghefi, M.R. Tehrani, H. Farahmand, S. Koshio, F.A. Dorkoosh, M.Z. Elsabee, Chitosan nanoparticle to carry vitamin C through the gastrointestinal tract and induce the non-specific immunity system of rainbow trout (*Oncorhynchus mykiss*), *Carbohydr. Polym.* 86 (2011) 142–146.
- [21] E. Jiménez-Fernández, A. Ruyra, N. Roher, E. Zuasti, C. Infante, C. Fernández-Díaz, Nanoparticles as a novel delivery system for vitamin C administration in aquaculture, *Aquaculture* 432 (2014) 426–433.

- [22] A. Bhattacharyya, S.J. Reddy, M.M. Hasan, M.M. Adeyemi, R.R. Marye, H. Naika, Nanotechnology-a unique future technology in aquaculture for the food security, *Int. J. Bioassays* 4 (2015) 4115–4126.
- [23] U.O. Chris, N.B. Singh, A. Agarwal, Nanoparticles as feed supplement on Growth behaviour of Cultured Catfish (*Clarias gariepinus*) fingerlings, *Mater. Today Proc.* 5 (2018) 9076–9081.
- [24] A.R. Ochoa-Meza, A.R. Álvarez-Sánchez, C.R. Romo-Quiñonez, A. Barraza, F.J. Magallón-Barajas, A. Chávez-Sánchez, J.C. García-Ramos, Y. Toledano-Magaña, N. Bogdanchikova, A. Pestryakov, Silver nanoparticles enhance survival of white spot syndrome virus infected *Penaeus vannamei* shrimps by activation of its immunological system, *Fish Shellfish Immunol.* 84 (2019) 1083–1089.
- [25] C. Nikapitiya, S.H.S. Dananjaya, B.C.J. De Silva, G.-J. Heo, C. Oh, M. De Zoysa, J. Lee, Chitosan nanoparticles: A positive immune response modulator as display in zebrafish larvae against *Aeromonas hydrophila* infection, *Fish Shellfish Immunol.* 76 (2018) 240–246.
- [26] M. Raissy, M. Ansari, In vitro antimicrobial effect of silver nanoparticles on *Lactococcus garvieae* and *Streptococcus iniae*, *Afr J Microbiol Res* 5 (2011) 4442–4445.
- [27] P. Swain, S.K. Nayak, A. Sasmal, T. Behera, S.K. Barik, S.K. Swain, S.S. Mishra, A.K. Sen, J.K. Das, P. Jayasankar, Antimicrobial activity of metal based nanoparticles against microbes associated with diseases in aquaculture, *World J. Microbiol. Biotechnol.* 30 (2014) 2491–2502.
- [28] M. Tello-Olea, S. Rosales-Mendoza, A.I. Campa-Córdova, G. Palestino, A. Luna-González, M. Reyes-Becerril, E. Velazquez, L. Hernandez-Adame, C. Angulo, Gold nanoparticles (AuNP) exert immunostimulatory and protective effects in shrimp (*Litopenaeus vannamei*) against *Vibrio parahaemolyticus*, *Fish Shellfish Immunol.* 84 (2019) 756–767.
- [29] P. Kalatehjari, M. Yousefian, M.A. Khalilzadeh, Assessment of antifungal effects of copper nanoparticles on the growth of the fungus *Saprolegnia* sp. on white fish (*Rutilus frisii kutum*) eggs, *Egypt. J. Aquat. Res.* 41 (2015) 303–306.
- [30] M.I. Shaalan, M.M. El-Mahdy, S. Theiner, M. El-Matbouli, M. Saleh, In vitro assessment of the antimicrobial activity of silver and zinc oxide nanoparticles against fish pathogens, *Acta Vet. Scand.* 59 (2017) 1–11.
- [31] K.M. Barakat, H.S. El-Sayed, Y.M. Gohar, Protective effect of squilla chitosan–silver nanoparticles for *Dicentrarchus labrax* larvae infected with *Vibrio anguillarum*, *Int. Aquat. Res.* 8 (2016) 179–189.
- [32] M. Saleh, A.-A. Abdel-Baki, M.A. Dkhil, M. El-Matbouli, S. Al-Quraishy, Antiprotozoal effects of metal nanoparticles against *Ichthyophthirius multifiliis*, *Parasitology* 144 (2017) 1802–1810.
- [33] I.A. Bhat, I. Ahmad, I.N. Mir, R.A.H. Bhat, P. Gireesh-Babu, M. Goswami, J.K. Sundaray, R. Sharma, Chitosan-eurycomanone nanoformulation acts on steroidogenesis pathway genes to increase the reproduction rate in fish, *J. Steroid Biochem. Mol. Biol.* 185 (2019) 237–247.

- [34] F. Fenaroli, D. Westmoreland, J. Benjaminsen, T. Kolstad, F.M. Skjeldal, A.H. Meijer, M. van der Vaart, L. Ulanova, N. Roos, B. Nystrom, Nanoparticles as drug delivery system against tuberculosis in zebrafish embryos: direct visualization and treatment, *ACS Nano* 8 (2014) 7014–7026.
- [35] M.A. Rather, R. Sharma, S. Gupta, S. Ferosekhan, V.L. Ramya, S.B. Jadhao, Chitosan-nanoconjugated hormone nanoparticles for sustained surge of gonadotropins and enhanced reproductive output in female fish, *PloS One* 8 (2013) e57094.
- [36] S. Kitiyodom, T. Yata, J. Yostawornkul, S. Kaewmalun, N. Nittayasut, K. Suktham, S. Surassmo, K. Namdee, C. Rodkhum, N. Pirarat, Enhanced efficacy of immersion vaccination in tilapia against columnaris disease by chitosan-coated “pathogen-like” mucoadhesive nanovaccines, *Fish Shellfish Immunol.* 95 (2019) 213–219.
- [37] S.R. Kumar, V.I. Ahmed, V. Parameswaran, R. Sudhakaran, V.S. Babu, A.S. Hameed, Potential use of chitosan nanoparticles for oral delivery of DNA vaccine in Asian sea bass (*Lates calcarifer*) to protect from *Vibrio* (*Listonella*) *anguillarum*, *Fish Shellfish Immunol.* 25 (2008) 47–56.
- [38] A. Rivas-Aravena, Y. Fuentes, J. Cartagena, T. Brito, V. Poggio, J. La Torre, H. Mendoza, F. Gonzalez-Nilo, A.M. Sandino, E. Spencer, Development of a nanoparticle-based oral vaccine for Atlantic salmon against ISAV using an alphavirus replicon as adjuvant, *Fish Shellfish Immunol.* 45 (2015) 157–166.
- [39] A. Elsheshtawy, N. Yehia, M. Elkemary, H. Soliman, Direct detection of unamplified *Aeromonas hydrophila* DNA in clinical fish samples using gold nanoparticle probe-based assay, *Aquaculture* 500 (2019) 451–457.
- [40] M. Saleh, H. Soliman, O. Haenen, M. El-Matbouli, Antibody-coated gold nanoparticles immunoassay for direct detection of *Aeromonas salmonicida* in fish tissues, *J. Fish Dis.* 34 (2011) 845–852.
- [41] J. Ramsden, *Applied nanotechnology: the conversion of research results to products*, William Andrew, 2018.
- [42] B.R. Shah, J. Mraz, *Advances in nanotechnology for sustainable aquaculture and fisheries*, *Rev. Aquac.* 12 (2020) 925–942.
- [43] M.H. Li, E.H. Robinson, Use of cottonseed meal in aquatic animal diets: a review, *North Am. J. Aquac.* 68 (2006) 14–22.
- [44] S. Singha, K. Das, N. Jha, Nano-systems for micro-nutrient delivery in aquaculture: a critical analysis, *Ann Aquac Res* 4 (2017) 1046.
- [45] J. Jampilek, J. Kos, K. Kralova, Potential of nanomaterial applications in dietary supplements and foods for special medical purposes, *Nanomaterials* 9 (2019) 296.

- [46] M. Aklakur, M. Asharf Rather, N. Kumar, Nanodelivery: an emerging avenue for nutraceuticals and drug delivery, *Crit. Rev. Food Sci. Nutr.* 56 (2016) 2352–2361.
- [47] B. Sarkar, A. Mahanty, S.K. Gupta, A.R. Choudhury, A. Daware, S. Bhattacharjee, Nanotechnology: A next-generation tool for sustainable aquaculture, *Aquaculture* 546 (2022) 737330.
- [48] M.A. Rather, R. Sharma, M. Aklakur, S. Ahmad, N. Kumar, M. Khan, V.L. Ramya, Nanotechnology: a novel tool for aquaculture and fisheries development. A prospective mini-review, *Fish. Aquac. J.* 16 (2011) 3.
- [49] S. Ye, W. Jin, Q. Huang, Y. Hu, B.R. Shah, Y. Li, B. Li, Development of Mag-FMBO in clay-reinforced KGM aerogels for arsenite removal, *Int. J. Biol. Macromol.* 87 (2016) 77–84.
- [50] E. Motamedi, M.T. Atouei, M.Z. Kassaee, Comparison of nitrate removal from water via graphene oxide coated Fe, Ni and Co nanoparticles, *Mater. Res. Bull.* 54 (2014) 34–40.
- [51] L. Liu, Z. Cui, Q. Ma, W. Cui, X. Zhang, One-step synthesis of magnetic iron–aluminum oxide/graphene oxide nanoparticles as a selective adsorbent for fluoride removal from aqueous solution, *RSC Adv.* 6 (2016) 10783–10791.
- [52] C. Hu, Y. Lan, J. Qu, X. Hu, A. Wang, Ag/AgBr/TiO₂ visible light photocatalyst for destruction of azo dyes and bacteria, *J. Phys. Chem. B* 110 (2006) 4066–4072.
- [53] R. Atchudan, T.N.J.I. Edison, S. Perumal, D. Karthikeyan, Y.R. Lee, Effective photocatalytic degradation of anthropogenic dyes using graphene oxide grafting titanium dioxide nanoparticles under UV-light irradiation, *J. Photochem. Photobiol. Chem.* 333 (2017) 92–104.
- [54] J. Park, E. Kettleson, W.-J. An, Y.J. Tang, P. Biswas, Inactivation of *E. coli* in water using photocatalytic, nanostructured films synthesized by aerosol routes, *Catalysts* 3 (2013) 247–260.
- [55] K. Ouyang, K. Dai, S.L. Walker, Q. Huang, X. Yin, P. Cai, Efficient photocatalytic disinfection of *Escherichia coli* O157: H7 using C70-TiO₂ hybrid under visible light irradiation, *Sci. Rep.* 6 (2016) 25702.
- [56] I. Sioen, S. De Henauw, F. Verdonck, N. Van Thuyne, J. Van Camp, Development of a nutrient database and distributions for use in a probabilistic risk–benefit analysis of human seafood consumption, *J. Food Compos. Anal.* 20 (2007) 662–670.
- [57] S. Mahboob, H.A. Al-Balawi, F. Al-Misned, S. Al-Quraishy, Z. Ahmad, Tissue metal distribution and risk assessment for important fish species from Saudi Arabia, *Bull. Environ. Contam. Toxicol.* 92 (2014) 61–66.
- [58] Y.S. Saleh, M.-A.S. Marie, Assessment of metal contamination in water, sediment, and tissues of *Arius thalassinus* fish from the Red Sea coast of Yemen and the potential human risk assessment, *Environ. Sci. Pollut. Res.* 22 (2015) 5481–5490.

- [59] G. Dugo, L. La Pera, A. Bruzzese, T.M. Pellicanò, V.L. Turco, Concentration of Cd (II), Cu (II), Pb (II), Se (IV) and Zn (II) in cultured sea bass (*Dicentrarchus labrax*) tissues from Tyrrhenian Sea and Sicilian Sea by derivative stripping potentiometry, *Food Control* 17 (2006) 146–152.
- [60] P. Manna, M. Sinha, P.C. Sil, A 43 kD protein isolated from the herb *Cajanus indicus* L attenuates sodium fluoride-induced hepatic and renal disorders in vivo, *BMB Rep.* 40 (2007) 382–395.
- [61] J.A. Camargo, Á. Alonso, Ecotoxicological assessment of the impact of fluoride (F⁻) and turbidity on the freshwater snail *Physella acuta* in a polluted river receiving an industrial effluent, *Environ. Sci. Pollut. Res.* 24 (2017) 15667–15677.
- [62] S. Wu, L. Kong, J. Liu, Removal of mercury and fluoride from aqueous solutions by three-dimensional reduced-graphene oxide aerogel, *Res. Chem. Intermed.* 42 (2016) 4513–4530.
- [63] S. Rajeshkumar, C. Venkatesan, M. Sarathi, V. Sarathbabu, J. Thomas, K.A. Basha, A.S. Hameed, Oral delivery of DNA construct using chitosan nanoparticles to protect the shrimp from white spot syndrome virus (WSSV), *Fish Shellfish Immunol.* 26 (2009) 429–437.
- [64] R.D. Handy, G. Cornelis, T. Fernandes, O. Tsyusko, A. Decho, T. Sabo-Attwood, C. Metcalfe, J.A. Steevens, S.J. Klaine, A.A. Koelmans, Ecotoxicity test methods for engineered nanomaterials: practical experiences and recommendations from the bench, *Environ. Toxicol. Chem.* 31 (2012) 15–31.
- [65] L. Farnen, Commercialization of nanotechnology for removal of heavy metals in drinking water, in: *Nanotechnol. Appl. Clean Water*, Elsevier, 2009: pp. 115–130.
- [66] M. Hua, S. Zhang, B. Pan, W. Zhang, L. Lv, Q. Zhang, Heavy metal removal from water/wastewater by nanosized metal oxides: a review, *J. Hazard. Mater.* 211 (2012) 317–331.
- [67] R. Kumar, J. Chawla, Removal of cadmium ion from water/wastewater by nano-metal oxides: a review, *Water Qual. Expo. Health* 5 (2014) 215–226.
- [68] W.-W. Tang, G.-M. Zeng, J.-L. Gong, J. Liang, P. Xu, C. Zhang, B.-B. Huang, Impact of humic/fulvic acid on the removal of heavy metals from aqueous solutions using nanomaterials: a review, *Sci. Total Environ.* 468 (2014) 1014–1027.
- [69] C.M. Futralan, C.-C. Kan, M.L. Dalida, C. Pascua, M.-W. Wan, Fixed-bed column studies on the removal of copper using chitosan immobilized on bentonite, *Carbohydr. Polym.* 83 (2011) 697–704.
- [70] S. Pandey, S.B. Mishra, Organic–inorganic hybrid of chitosan/organoclay bionanocomposites for hexavalent chromium uptake, *J. Colloid Interface Sci.* 361 (2011) 509–520.
- [71] M. Namdeo, S.K. Bajpai, Chitosan–magnetite nanocomposites (CMNs) as magnetic carrier particles for removal of Fe (III) from aqueous solutions, *Colloids Surf. Physicochem. Eng. Asp.* 320 (2008) 161–168.

- [72] Z. Fang, Y. Zhao, R.D. Warner, S.K. Johnson, Active and intelligent packaging in meat industry, *Trends Food Sci. Technol.* 61 (2017) 60–71.
- [73] Z.A. El-Greisy, A.E. El-Gamal, Monosex production of tilapia, *Oreochromis niloticus* using different doses of 17 α -methyltestosterone with respect to the degree of sex stability after one year of treatment, *Egypt. J. Aquat. Res.* 38 (2012) 59–66.
- [74] H.D. Joshi, V.K. Tiwari, S. Gupta, R. Sharma, W.S. Lakra, U. Sahoo, Application of nanotechnology for the production of masculinized Tilapia, *Oreochromis niloticus* (Linnaeus, 1758), *Aquaculture* 511 (2019) 734206.
- [75] W.T. Abbas, Advantages and prospective challenges of nanotechnology applications in fish cultures: a comparative review, *Environ. Sci. Pollut. Res.* 28 (2021) 7669–7690.
- [76] K.K. Reza, M.K. Singh, S.K. Yadav, J. Singh, V.V. Agrawal, B.D. Malhotra, Quantum dots based platform for application to fish freshness biosensor, *Sens. Actuators B Chem.* 177 (2013) 627–633.
- [77] R. Uddin, M.I. Wahid, T. Jesmeen, N.H. Huda, K.B. Sutradhar, Detection of formalin in fish samples collected from Dhaka City, Bangladesh, (2011).
- [78] T. Suwanaruang, Formalin contaminated in seafood and frozen meat at Somdet Market, Kalasin Province, *J. Environ. Prot.* 9 (2018) 1286–1293.
- [79] N. Marzuki, F.A. Bakar, A.B. Salleh, L.Y. Heng, N.A. Yusof, S. Siddiquee, Electrochemical biosensor immobilization of formaldehyde dehydrogenase with nafion for determination of formaldehyde from Indian mackerel (*Rastrelliger kanagurta*) fish, *Curr. Anal. Chem.* 8 (2012) 534–542.
- [80] B. Noor Aini, S. Siddiquee, K. Ampon, Development of formaldehyde biosensor for determination of formalin in fish samples; malabar red snapper (*Lutjanus malabaricus*) and longtail tuna (*Thunnus tonggol*), *Biosensors* 6 (2016) 32.
- [81] B. Kuswandi, Environmental friendly food nano-packaging, *Environ. Chem. Lett.* 15 (2017) 205–221.
- [82] J.-W. Rhim, H.-M. Park, C.-S. Ha, Bio-nanocomposites for food packaging applications, *Prog. Polym. Sci.* 38 (2013) 1629–1652.
- [83] H.S. Jiang, L. Yin, N.N. Ren, L. Xian, S. Zhao, W. Li, B. Gontero, The effect of chronic silver nanoparticles on aquatic system in microcosms, *Environ. Pollut.* 223 (2017) 395–402.
- [84] E. Can, V. Kizak, M. Kayim, S.S. Can, B. Kutlu, M. Ates, M. Kocabas, N. Demirtas, Nanotechnological applications in aquaculture-seafood industries and adverse effects of nanoparticles on environment, *J. Mater. Sci. Eng.* 5 (2011).
- [85] J. Ji, D. Torrealba, À. Ruyra, N. Roher, Nanodelivery systems as new tools for immunostimulant or vaccine administration: targeting the fish immune system, *Biology* 4 (2015) 664–696.

- [86] P.H.M. Joosten, E. Tiemersma, A. Threels, C. Caumartin-Dhieux, J. Rombout, Oral vaccination of fish against *Vibrio anguillarum* using alginate microparticles, *Fish Shellfish Immunol.* 7 (1997) 471–485.
- [87] W.R. Gombotz, S.F. Wee, Protein release from alginate matrices, *Adv. Drug Deliv. Rev.* 64 (2012) 194–205.
- [88] A. Sosnik, Alginate particles as platform for drug delivery by the oral route: state-of-the-art, *Int. Sch. Res. Not.* 2014 (2014).
- [89] C.A.G. Leal, G.A. Carvalho-Castro, P.S.C. Sacchetin, C.O. Lopes, A.M. Moraes, H.C.P. Figueiredo, Oral and parenteral vaccines against *Flavobacterium columnare*: evaluation of humoral immune response by ELISA and in vivo efficiency in Nile tilapia (*Oreochromis niloticus*), *Aquac. Int.* 18 (2010) 657–666.
- [90] C.P. Reis, R.J. Neufeld, F. Veiga, A.J. Ribeiro, Preparation of drug-loaded polymeric nanoparticles, in: *Nanomedicine Cancer*, Jenny Stanford Publishing, 2017: pp. 171–214.
- [91] M. Tafaghodi, T.S.G. SAJADI, M. Payan, Alginate microsphere as a delivery system and adjuvant for autoclaved *Leishmania major* and Quillaja saponin: preparation and characterization, (2007).
- [92] O. Borges, M. Silva, A. de Sousa, G. Borchard, H.E. Junginger, A. Cordeiro-da-Silva, Alginate coated chitosan nanoparticles are an effective subcutaneous adjuvant for hepatitis B surface antigen, *Int. Immunopharmacol.* 8 (2008) 1773–1780.
- [93] K. Fujiki, H. Matsuyama, T. Yano, Protective effect of sodium alginates against bacterial infection in common carp, *Cyprinus carpio* L., *J. Fish Dis.* 17 (1994) 349–355.
- [94] S.-T. Chiu, R.-T. Tsai, J.-P. Hsu, C.-H. Liu, W. Cheng, Dietary sodium alginate administration to enhance the non-specific immune responses, and disease resistance of the juvenile grouper *Epinephelus fuscoguttatus*, *Aquaculture* 277 (2008) 66–72.
- [95] A.-C. Cheng, Y.-Y. Chen, J.-C. Chen, Dietary administration of sodium alginate and κ-carrageenan enhances the innate immune response of brown-marbled grouper *Epinephelus fuscoguttatus* and its resistance against *Vibrio alginolyticus*, *Vet. Immunol. Immunopathol.* 121 (2008) 206–215.
- [96] H.B. Huttenhuis, A.S. Ribeiro, T.J. Bowden, C. Van Bavel, A.J. Taverne-Thiele, J.H. Rombout, The effect of oral immuno-stimulation in juvenile carp (*Cyprinus carpio* L.), *Fish Shellfish Immunol.* 21 (2006) 261–271.
- [97] S.-P. Yeh, C.-A. Chang, C.-Y. Chang, C.-H. Liu, W. Cheng, Dietary sodium alginate administration affects fingerling growth and resistance to *Streptococcus* sp. and iridovirus, and juvenile non-specific immune responses of the orange-spotted grouper, *Epinephelus coioides*, *Fish Shellfish Immunol.* 25 (2008) 19–27.

- [98] J. Skjermo, Ø. Bergh, High-M alginate immunostimulation of Atlantic halibut (*Hippoglossus hippoglossus* L.) larvae using *Artemia* for delivery, increases resistance against vibriosis, *Aquaculture* 238 (2004) 107–113.
- [99] P.K. Dutta, J. Dutta, V.S. Tripathi, Chitin and chitosan: Chemistry, properties and applications, (2004).
- [100] I.A. Sogias, A.C. Williams, V.V. Khutoryanskiy, Why is chitosan mucoadhesive?, *Biomacromolecules* 9 (2008) 1837–1842.
- [101] Y. Valero, E. Awad, F. Buonocore, M. Arizcun, M.Á. Esteban, J. Meseguer, E. Chaves-Pozo, A. Cuesta, An oral chitosan DNA vaccine against nodavirus improves transcription of cell-mediated cytotoxicity and interferon genes in the European sea bass juveniles gut and survival upon infection, *Dev. Comp. Immunol.* 65 (2016) 64–72.
- [102] L. Li, S.-L. Lin, L. Deng, Z.-G. Liu, Potential use of chitosan nanoparticles for oral delivery of DNA vaccine in black seabream *Acanthopagrus schlegelii* Bleeker to protect from *Vibrio parahaemolyticus*, *J. Fish Dis.* 36 (2013) 987–995.
- [103] J. Tian, X. Sun, X. Chen, J. Yu, L. Qu, L. Wang, The formulation and immunisation of oral poly (DL-lactide-co-glycolide) microcapsules containing a plasmid vaccine against lymphocystis disease virus in Japanese flounder (*Paralichthys olivaceus*), *Int. Immunopharmacol.* 8 (2008) 900–908.
- [104] M. Adomako, S. St-Hilaire, Y. Zheng, J. Eley, R.D. Marcum, W. Sealey, B.C. Donahower, S. LaPatra, P.P. Sheridan, Oral DNA vaccination of rainbow trout, *Oncorhynchus mykiss* (Walbaum), against infectious haematopoietic necrosis virus using PLGA [Poly (D, L-Lactic-Co-Glycolic Acid)] nanoparticles, *J. Fish Dis.* 35 (2012) 203–214.
- [105] H. Wang, W. Tang, R. Zhang, S. Ding, Analysis of enzyme activity, antibacterial activity, antiparasitic activity and physico-chemical stability of skin mucus derived from *Amphiprion clarkii*, *Fish Shellfish Immunol.* 86 (2019) 653–661.
- [106] A.A. Tayel, B.E. Elsaied, A.M. Diab, Nanotechnology for Aquaculture, *Nanosci. Sustain. Agric.* (2019) 479–544.
- [107] W. Jun-qiang, Summary report on experiment of strong nanometer 863 biological assistant growth unit in sea shrimp farming, *Mod. Fish. Inf.-Chang.* 18 (2003) 12–15.
- [108] A. Liu, Y. Cao, M. Dai, Z.W. Liao, Nanomaterial application in carp aquaculture experiment, *Fish Mod.* 2 (2008) 24–27.
- [109] J.F. Mongillo, *Nanotechnology* 101 Greenwood Press, Westport CT (2007) 4–13.
- [110] M. Abhilash, Potential applications of Nanoparticles, *Int. J. Pharma Bio Sci.* 1 (2010) 1–12.

- [111] A. Hernando Grande, Nanotecnología y nanopartículas magnéticas: La física actual en la lucha contra la enfermedad, *Rev. Real Acad. Cienc. Exactas Físicas Nat.* 101 (2007) 321–327.
- [112] M. Rai, A. Ingle, Role of nanotechnology in agriculture with special reference to management of insect pests, *Appl. Microbiol. Biotechnol.* 94 (2012) 287–293.
- [113] M. Muruganandam, S.R. Chipps, P.R. Ojasvi, On the advanced technologies to enhance fisheries production and management, *Acta Sci. Agric.* 3 (2019) 216–222.
- [114] C. Thiruppathiraja, S. Kumar, V. Murugan, P. Adaikkappan, K. Sankaran, M. Alagar, An enhanced immuno-dot blot assay for the detection of white spot syndrome virus in shrimp using antibody conjugated gold nanoparticles probe, *Aquaculture* 318 (2011) 262–267.
- [115] M. Crane, A. Hyatt, Viruses of fish: an overview of significant pathogens, *Viruses* 3 (2011) 2025–2046.
- [116] A. Natarajan, K.S. Devi, S. Raja, A. Senthil Kumar, An elegant analysis of white spot syndrome virus using a graphene oxide/methylene blue based electrochemical immunosensor platform, *Sci. Rep.* 7 (2017) 46169.
- [117] X. Yang, S. Zhang, J. Liu, Q. Gao, S. Dong, C. Zhou, Deep learning for smart fish farming: applications, opportunities and challenges, *Rev. Aquac.* 13 (2021) 66–90.
- [118] J. Mitchell, V. Trivedi, Pharmaceutical nanomaterials: the preparation of solid core drug delivery systems (SCDDS), *J. Pharm. Pharmacol.* 62 (2010) 1457–1458.
- [119] M. Strømme, U. Brohede, R. Atluri, A.E. Garcia-Bennett, Mesoporous silica-based nanomaterials for drug delivery: evaluation of structural properties associated with release rate, *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.* 1 (2009) 140–148.
- [120] J.H. Eldridge, C.J. Hammond, J.A. Meulbroek, J.K. Staas, R.M. Gilley, T.R. Tice, Controlled vaccine release in the gut-associated lymphoid tissues. I. Orally administered biodegradable microspheres target the Peyer's patches, *J. Controlled Release* 11 (1990) 205–214.
- [121] P. Jani, G.W. Halbert, J. Langridge, A.T. Florence, Nanoparticle uptake by the rat gastrointestinal mucosa: quantitation and particle size dependency, *J. Pharm. Pharmacol.* 42 (1990) 821–826.
- [122] A.T. Florence, A.M. Hillery, N. Hussain, P.U. Jani, Nanoparticles as carriers for oral peptide absorption: studies on particle uptake and fate, *J. Controlled Release* 36 (1995) 39–46.
- [123] S.E. Peters, C.H. Brain, Benefits of a soy lecithin based nanotechnology for the animal and human food industry, in: ACS Publications, 2009.
- [124] V.J. Mohanraj, Y. Chen, Nanoparticles-a review, *Trop. J. Pharm. Res.* 5 (2006) 561–573.
- [125] FOE, Friends of the Earth International: Negotiating a North-South Identity, (2008).

- [126] M.S. Izquierdo, W. Ghrab, J. Roo, K. Hamre, C.M. Hernández-Cruz, G. Bernardini, G. Terova, R. Saleh, Organic, inorganic and nanoparticles of Se, Zn and Mn in early weaning diets for gilthead seabream (*Sparus aurata*; Linnaeus, 1758), *Aquac. Res.* 48 (2017) 2852–2867.
- [127] A. Trapani, D. Mandracchia, C. Di Franco, H. Cordero, P. Morcillo, R. Comparelli, A. Cuesta, M.A. Esteban, In vitro characterization of 6-Coumarin loaded solid lipid nanoparticles and their uptake by immunocompetent fish cells, *Colloids Surf. B Biointerfaces* 127 (2015) 79–88.
- [128] N. Mohammadi, A. Tukmechi, The effects of iron nanoparticles in combination with *Lactobacillus casei* on growth parameters and probiotic counts in rainbow trout (*Oncorhynchus mykiss*) intestine., *J. Vet. Res.* 70 (2015).
- [129] A. Asaikkutti, P.S. Bhavan, K. Vimala, M. Karthik, P. Cheruparambath, Dietary supplementation of green synthesized manganese-oxide nanoparticles and its effect on growth performance, muscle composition and digestive enzyme activities of the giant freshwater prawn *Macrobrachium rosenbergii*, *J. Trace Elem. Med. Biol.* 35 (2016) 7–17.
- [130] T. Muralisankar, P.S. Bhavan, S. Radhakrishnan, C. Seenivasan, V. Srinivasan, The effect of copper nanoparticles supplementation on freshwater prawn *Macrobrachium rosenbergii* post larvae, *J. Trace Elem. Med. Biol.* 34 (2016) 39–49.
- [131] M.F. El Basuini, A.M. El-Hais, M.A.O. Dawood, A.-S. Abou-Zeid, S.Z. El-Damrawy, M.-S. Khalafalla, S. Koshio, M. Ishikawa, S. Dossou, Effects of dietary copper nanoparticles and vitamin C supplementations on growth performance, immune response and stress resistance of red sea bream, *Pagrus major*, *Aquac. Nutr.* 23 (2017) 1329–1340.
- [132] S. Kunjiappan, C. Bhattacharjee, R. Chowdhury, Hepatoprotective and antioxidant effects of *Azolla microphylla* based gold nanoparticles against acetaminophen induced toxicity in a fresh water common carp fish (*Cyprinus carpio* L.), (2015).
- [133] M. Sharif Rohani, M. Haghghi, S. Bazari Moghaddam, Study on nanoparticles of Aloe vera extract on growth performance, survival rate and body composition in Siberian sturgeon (*Acipenser baerii*), Iran. *J. Fish. Sci.* 16 (2017) 457–468.
- [134] V. Kumar, N. Sharma, S.S. Maitra, In vitro and in vivo toxicity assessment of nanoparticles, *Int. Nano Lett.* 7 (2017) 243–256.
- [135] B.J. Shaw, R.D. Handy, Physiological effects of nanoparticles on fish: a comparison of nanometals versus metal ions, *Environ. Int.* 37 (2011) 1083–1097.
- [136] H. Li, J. Zhang, T. Wang, W. Luo, Q. Zhou, G. Jiang, Elemental selenium particles at nano-size (Nano-Se) are more toxic to Medaka (*Oryzias latipes*) as a consequence of hyper-accumulation of selenium: a comparison with sodium selenite, *Aquat. Toxicol.* 89 (2008) 251–256.

[137] M. Stefaniuk, P. Oleszczuk, Y.S. Ok, Review on nano zerovalent iron (nZVI): from synthesis to environmental applications, *Chem. Eng. J.* 287 (2016) 618–632.

[138] M. Li, Q. Wu, Q. Wang, D. Xiang, G. Zhu, Effect of titanium dioxide nanoparticles on the bioavailability and neurotoxicity of cypermethrin in zebrafish larvae, *Aquat. Toxicol.* 199 (2018) 212–219.

[139] X. Ren, X. Zhao, X. Duan, Z. Fang, Enhanced bio-concentration of tris (1, 3-dichloro-2-propyl) phosphate in the presence of nano-TiO₂ can lead to adverse reproductive outcomes in zebrafish, *Environ. Pollut.* 233 (2018) 612–622.

UNDER PEER REVIEW