

# **Impact of NPK on the Ecophysiological attributes of *Thespesia populnea***

## **ABSTRACT**

A field experiment was conducted to investigate the effect of different levels of NPK on the performance of *Thespesia populnea*. A total of 17 treatments consisted of four levels of Nitrogen (50, 75, 100 and 125 g N plant<sup>-1</sup>), two levels each of Phosphorus (50 and 75 g P plant<sup>-1</sup>), and Potassium (25 and 50 g K plant<sup>-1</sup>) were tried in RBD replicated three times, to standardize the fertilizer schedule for the tree crop taken for investigation. The experimental soil was red sandy loam, non-calcareous, neutral in reaction, low in available nitrogen, phosphorus and medium in available potassium with low organic carbon content. Ecophysiological parameters viz., photosynthetic rate (A), transpiration rate (E), stomatal conductance (g<sub>s</sub>) and Intercellular CO<sub>2</sub> concentration (C<sub>int</sub>). Growth attributes were influenced by N, P and K at different levels. Among the various treatments application of 100 : 75 : 50 g NPK plant<sup>-1</sup> (T<sub>12</sub>) significantly influenced the growth attributes followed by 100 : 75 : 25 g NPK plant<sup>-1</sup> (T<sub>11</sub>).

## **Introduction**

Forests are not just trees, but part of an ecosystem that interlinks life, economics and societies. Forest provides a multiplicity of environmental services. As per Indian State of Forest Report (ISFR) 2013, the forest cover of the country is 6,97,898 sq.km. (69.79 million ha) which is 21.23 per cent of the geographical area of the country while the tree cover of the country is estimated to be 91,266 sq.km. (9.13 million ha) which is 2.78 per cent of the geographical area. The total forest and tree cover of the country as per 2013 assessment is 7,89,164 sq.km. (78.92 million ha) which is 24.01 per cent of the geographical area of the country. There is an increase of 5,871 sq. km. in the forest and tree cover of the country in comparison to 2011

assessment. The total growing stock of India's forest and trees outside forest is estimated as 5,658.046 million m<sup>3</sup> which comprises 4,173.362 million m<sup>3</sup> inside the forests and 1,484.68 million m<sup>3</sup> outside the forests. India is home to 17 per cent of world human population and 18 per cent of livestock population which causes severe pressure to the Indian forests. Nearly 40 per cent of domestic fuel needs of people and 30 per cent of fodder needs for cattle population in the country are met from forests. The demand and supply gap of timber, fuelwood and fodder is widening whereas the raw material requirement of wood based industries is on the raise. The National Forest Policy (1988) suggested that the wood based industries will have to make their own arrangements for the supply of raw materials. Besides this, most of the Government policies and acts stress mostly on the conservation of existing natural vegetation. Hence, the importance of industrial plantations has gained momentum across the country to meet out the industrial raw material needs. In order to fulfill the domestic and industrial demands of wood, the only viable option is to increase the area under agroforestry and industrial plantations. The role of indigenous species and fast growing species in this regard need not be overemphasized as selection of suitable tree species plays a major role in the success of agroforestry. *Thespesia populneais* one such indigenous tree species which has multiple uses.

*Thespesia populneais* a fast growing evergreen tree belonging to the family Malvaceae which is commonly known as Tulip tree or Portia tree and it is called as Puvarasu in Tamil. *Thespesia populneais* a small to medium-sized evergreen tree of the coastal tracts of India and Myanmar which grows upto 20 m with a dense crown. It is widely planted as a roadside tree in tropical regions and recognized by its large yellow flowers with purple centres. The flowers open and close on the same day, and the yellow flowers turn dark red, purple or pink as the day progresses. Bark is dark

brown, rough, deeply cracked, fissured and slash yellow coloured. It prefers neutral soils with pH 6.0-7.4 and thrives well on sandy coastal soils, but also grows on volcanic soils, soils derived from limestone and rocky headlands. It does not do well on upland, acidic clays. It is very tolerant to salt spray, steady coastal winds and highly resistant to dry wood termites. It tolerates heavier soils, soil salinity and occasional inundation, but does not grow on permanently inundated soils. In order to obtain robust and healthy plantations, proper fertilizer management is essential under field conditions. Lack of studies in this aspect necessitated the present investigation to study the Impact of NPK on the Ecophysiological attributes of the species

## **Review of Literature**

### **Ecophysiological parameters**

#### **1. Photosynthetic rate**

Estimates of the canopy photosynthetic capacity of forests provide information about their role in global carbon cycles, involving biomass production (Bassow and Bazzaz, 1997). Comparison of 51 clones of *Casuarina equisetifolia* in terms of growth and eco physiological characters revealed that five fast growing clones recorded high photosynthetic rate and thus showed a higher productivity (Balasubramanian and Gurumurthi, 2001). Chen *et al.* (2001) stated that the photosynthetic characters of *Phyllostachys heterocyclus* cv. *pubescens* provenances recorded increase in Net Photosynthetic Rate (NPR) with increasing latitude and all provenances had two peaks of NPR in one year.

Rao (2005) studied the effect of shade on net photosynthetic rate of five important tree species viz., *Albizia lebbek*, *Dalbergia sissoo*, *Leucaena leucocephala*, *Shorea robusta* and *Tectona grandis*. Net photosynthetic rate (P,

$\mu \text{ mol m}^{-2} \text{ s}^{-1}$ ) decreased with increasing shade in all the species during all the three seasons. The photosynthetic rate of *Dalbergia sissoo* was reported to be maximum as compared to *Acacia auriculiformis* and *Albizia lebbek* (Srivastava and Ram, 2009).

Saraswathi and Paliwal (2008) investigated the net photosynthesis rate ( $P_n$ ) in six month old seedlings of *Albizia lebbek* and *Cassia siamea* under different levels of drought stress. *Cassia siamea* had a transpiration rate of  $2.56 \text{ m mol m}^{-2} \text{ s}^{-1}$  and increased photosynthesis rate was observed in *A. lebbek* of  $4.35 \mu \text{ mol m}^{-2} \text{ s}^{-1}$  at 9.00 hrs.

Highest net photosynthetic rate of  $6.61$  and  $6.76 \mu \text{ mol m}^{-2} \text{ s}^{-1}$  was recorded in the treatment which received 25 per cent more fertilizer than the soil test recommendation value of 93.75: 225: 100 of NPK  $\text{kg ha}^{-1}$  in *Casuarina* species (Baranidharan *et al.*, 2013).

Photosynthesis, photorespiration and  $\text{CO}_2$  compensation point comparison of two year old seedlings of *Thuja orientalis* from six seed sources showed significant variation in net photosynthetic rate among different sources (Wu and Ma, 1988).

Highly significant variation was reported by Kundu and Tigerstedt (1999) among ten provenances of *Azadirachta indica* in net photosynthesis rate ranged from 8.14 to  $15.13 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ . They concluded that whole plant phytomass production in neem seedlings was associated with photosynthesis and stomatal characteristics during the early stages of growth.

## **2. Transpiration rate**

In an afforested plantation, *Gmelina arborea* @ 21 MAP recorded significantly higher transpiration rate of  $4.81 \text{ m mol m}^{-2} \text{ s}^{-1}$  followed by *Dalbergia*

*sissoo* ( $4.32 \text{ m mol m}^{-2} \text{ s}^{-1}$ ), *Tectona grandis* ( $4.01 \text{ m mol m}^{-2} \text{ s}^{-1}$ ) and *Bambusa vulgaris* var. *vulgaris* ( $2.47 \text{ m mol m}^{-2} \text{ s}^{-1}$ ). The lowest transpiration rate of  $1.67 \text{ m mol m}^{-2} \text{ s}^{-1}$  was registered by *Swietenia macrophylla* (Narendra Babu, 2012). Transpiration rate from several Australian semiarid zone species was studied in the field by Kreeb (1966) using the torsion balance technique and he concluded that among the different species studied, the maximum transpiration rate was observed in *Eucalyptus camaldulensis* ( $5.0$  and  $7.9 \text{ m mol m}^{-2} \text{ s}^{-1}$ ).

Dolley and Grieve (1966) studied *E. marginata* forest over a period of four months to evaluate the transpiration rate which ranged from  $0.34$  to  $1.10 \text{ m mol m}^{-2} \text{ s}^{-1}$  respectively per day. It was found that transpiration rate had a positive correlation with air temperature. A field study done by Ujala Singh (2005) studied the transpiration rates of different *Eucalyptus* species and concluded that *E. urophylla* and *E. grandis* had the highest transpiration rate with  $180$  and  $165 \text{ g day}^{-1} \text{ plant}^{-1}$  respectively. On the other hand, *E. ochrophloia* recorded the lowest value of  $65 \text{ g day}^{-1} \text{ plant}^{-1}$ .

Highly significant differences in transpiration rate and stomatal conductance of leaves were reported and strongly correlated among seed sources of *Ginkgo biloba* (Chiang *et al.*, 1996). Suman Chandra (2003) observed leaf to air temperature difference ( $dT$ ) of  $0.72 \text{ m mol m}^{-2} \text{ s}^{-1}$  and transpiration rate ( $Tr$ ) of  $2.64 \text{ m mol m}^{-2} \text{ s}^{-1}$  in *Aesculus indica* and lowest in *Toona ciliata* of  $0.41 \text{ m mol m}^{-2} \text{ s}^{-1}$  and  $1.54 \text{ m mol m}^{-2} \text{ s}^{-1}$  respectively. He concluded a positive correlation between leaf to air temperature difference ( $dT$ ) and rate of transpiration ( $Tr$ ).

Baranidharan *et al.* (2013) observed that a fertilizer dose of 25 per cent more than the soil test recommendation value of  $93.75: 225: 100$  of NPK  $\text{kg ha}^{-1}$  recorded

the highest transpiration rate value of 2.20 and 2.38  $\text{m mol m}^{-2} \text{s}^{-1}$  and was significantly superior when compared with other nutrient levels in *Casuarina* species.

### 3. Stomatal conductance

Stomata are the points of exit for the water vapour from leaves. Under conditions of adequate supply of water, stomata opening in the morning with increasing radiant flux and maintain a relatively constant aperture until radiant flux declines in the late afternoon. So, there was a correlation exists between stomatal opening and radiant flux (Gikloot *et al.*, 2012).

Stomatal conductance did not show any variation in three provenances of rosewood studied and appeared to be more influenced by the photon flux density during the morning. However, in afternoon, the stomatal behavior appeared to be controlled by specific humidity deficit (Rosa *et al.*, 1998). Ladjalet *et al.* (2007) studied greenhouse grown Cedar seedlings subjected to two different watering regimes and observed that the dry set showed lower net photosynthesis and stomatal conductance than the plants in the well watered set.

De Costa *et al.* (2000) investigated stomatal conductance in nine tree species *viz.*, *Alstonia macrophylla*, *Filiciumdecipiens*, *Macaranga peltata*, *Tectona grandis*, *Acronychia pedunculata*, *Semicarpusnigro-viridis*, *Swietenia macrophylla*, *Mesua ferrea* and *Terminalia catappa* growing under different levels of natural shade, i.e. open, medium shade and full shade. According to them, the total leaf stomatal conductance varied significantly with tree species and shade levels and the highest stomatal conductance was observed in *S. nigroviridis* and *T. catappa*, with 92.03 and 78.18  $\text{m mol m}^{-1} \text{s}^{-1}$  respectively. The rest of the species had recorded lower stomatal conductance values which ranged from 34 to 44  $\text{m mol m}^{-2} \text{s}^{-1}$ .

The stomatal conductance and transpiration were studied for tropical rain forest species and desert species in biosphere. The results showed that the stomatal conductance and transpiration of rain forest species decreased from  $127.4 \pm 65.6 \text{ m mol m}^{-2} \text{ s}^{-1}$  and  $2.04 \pm 0.61 \text{ m mol m}^{-2} \text{ s}^{-1}$  to  $61.3 \pm 30.5 \text{ m mol m}^{-2} \text{ s}^{-1}$  and  $1.54 \pm 0.65 \text{ m mol m}^{-2} \text{ s}^{-1}$  respectively (Jiang Gaominget *al.*, 1997).

Okuto and Ouma (2010) reported variation in stomatal conductance among provenances of *Syzygiumcumini* seedlings from three districts of Kenya *viz.*, Vihiga, Kisumu and Siaya. Among them, stomatal conductance was similar for Kisumu and Siaya but higher than Vihiga.

#### **4. Inter cellular CO<sub>2</sub> concentration (CINT)**

Esser (2004) correlated Inter cellular CO<sub>2</sub> concentration with the photosynthetic efficiency of three species.

Suresh kumar (2011) and Narendra Babu (2012) have also confirmed that Inter cellular CO<sub>2</sub> concentration in *Tectona grandis* and *Gmelina arborea* was lower than that of *Dalbergia sisoo* and *Bambusa vulgaris* var. *vulgaris* during 9 MAP respectively.

#### **Materials and Methods**

The present study was conducted in order to standardize the fertilizer requirement for *Thespesia populnea* (L.) Selex Correa. With the view of fulfilling the objectives envisaged, experiments were carried out at Forest College and Research Institute, Mettupalayam, Tamil Nadu during 2014-2015.

The materials used and the methods followed during the course of investigation are furnished in this chapter.

## MATERIALS

### Species under study- *Thespesia populnea*

#### i) Species description

Species	:	<i>Thespesia populnea</i>
Family	:	Malvaceae
Tamil name	:	Poovarasu
Common name	:	Indian Tulip Tree, Portia tree, Pacific Rosewood
Trade name	:	Seychelles Rose wood

#### ii) Botanical features

It is a small to medium sized, fast-growing, evergreen tree with a dense spreading crown attaining a height of 20 m with a clear bole height of 0.6 - 2 m, grey to brown bark, fissured, often knobby and fibrous. Leaves are broad 8-15 cm x 6.10 cm, heart-shaped like pipal or poplars, entire, petiole 2.5-10 cm long, stipules subulate, deciduous. Flowers axillary, solitary, bisexual, bell-shaped, 5-7.5 cm across, pedicels 5 -7.5 cm long. Petals 5, yellow with deep maroon in the centre, fading to purplish-pink before falling. Fruit a globose capsule, 2.5 - 4 cm in diameter, flattened covered with disk like persistent calyx at the base, black when ripe, containing 1-3 egg-shaped seeds about 1 cm long.

*Thespesia populnea* anchors sandy coastal soils and protect them from erosion .The tree is used to stabilize bunds for ponds in prawn production (Hanum and van der Maesen, 1997). The wood may produce a yellow dye used to dye wool in East and South-East Asia, and the leaves are used to make a black dye (Clark and Thaman, 1993). Large tulip trees are made into small canoes in ancient times in

Hawaii (Abbott, 1992). The wood is also used for boat building in South Asia and elsewhere in the Pacific.

### **Location**

The experiment was conducted in J block, of Forest College and Research Institute, Mettupalayam (Coimbatore district). The geographical location is 11°19' N latitude and 77°56' E longitude at an altitude of 320 m above mean sea level

### **Climate and Weather**

The mean annual rainfall of the experimental site is 913.9 mm. The maximum temperature ranges from 30.5 to 36.6°C and the minimum temperature ranges from 17.1 to 23.5°C. The mean maximum and minimum temperature are 34.5°C and 20.63°C respectively.

### **Assessment of eco-physiological behavior**

Ecophysiological characters were assessed by using a Portable Photosynthesis System (PPS, model LCpro + Photosynthesis System CO<sub>2</sub> gas analyzer, UK) to assess the eco-physiological behaviour of selected tree species. The measurements were made on fully matured leaves (5-6 leaves from the bud). The ecophysiological parameters viz., transpiration rate, stomatal conductance, Intercellular CO<sub>2</sub> Concentration (CINT) and photosynthetic rate were measured on a sunny day between 10.00 AM and 11.00 AM as per the procedure described below.

### **Photosynthetic rate (A)**

The photosynthetic rate of trees was measured using the Portable Photosynthesis System (PPS, model LCpro + Photosynthesis System CO<sub>2</sub> gas analyzer, UK). The PPS measures the uptake of CO<sub>2</sub> and estimates the photosynthetic productivity using Infra-Red Gas analyzer (IRGA) and expressed in  $\mu\text{ mol m}^{-2} \text{ s}^{-1}$ .

### **Transpiration rate (E)**

The transpiration rate was measured using Portable Photosynthesis System (PPS, model LCpro<sup>+</sup> Photosynthesis System CO<sub>2</sub> gas analyzer, UK) and expressed as  $\text{m mol m}^{-2} \text{ s}^{-1}$ .

### **Stomatal conductance (gs)**

Stomatal conductance was measured using the Portable Photosynthesis System (PPS, model LCpro + Photosynthesis System CO<sub>2</sub> gas analyzer, UK) and expressed in  $\text{m mol m}^{-2} \text{ s}^{-1}$ .

### **Intercellular CO<sub>2</sub> Concentration (CINT)**

Intercellular CO<sub>2</sub> concentration measures the photosynthetic efficiency of plant. It was measured using Portable Photosynthesis System (PPS, model LCpro + Photosynthesis System CO<sub>2</sub> gas analyzer, UK) and expressed in ppm.

## **Results and Discussion**

### **Ecophysiological attributes**

Portable Photosynthesis System (LC Pro<sup>+</sup>) was used to observe changes in ecophysiological behaviour of *Thespesia populnea* and four parameters viz., photosynthetic rate (A), transpiration rate (E), stomatal conductance (g<sub>s</sub>) and intercellular CO<sub>2</sub> (CINT) were recorded and tabulated in Table 1.

**Table 1. Influence of NPK on the ecophysiological behaviour of *Thespesia populnea* at 6 MAP**

Treatments	Photosynthetic rate ( $\mu \text{ mol m}^{-2} \text{ s}^{-1}$ )	Transpiration rate ( $\text{m mol m}^{-2} \text{ s}^{-1}$ )	Stomatal conductance ( $\text{m mol m}^{-2} \text{ s}^{-1}$ )	Inter Cellular CO <sub>2</sub> Concentration (ppm)
T <sub>1</sub> -50:50:25 g NPK plant <sup>-1</sup>	9.94	3.82	0.11	175
T <sub>2</sub> -50:50:50 g NPK plant <sup>-1</sup>	10.67	3.84	0.12	178
T <sub>3</sub> -50:75:25 g NPK plant <sup>-1</sup>	11.62	4.42	0.12	188
T <sub>4</sub> -50:75:50 g NPK plant <sup>-1</sup>	11.86	4.83	0.13	190
T <sub>5</sub> -75:50:25 g NPK plant <sup>-1</sup>	12.27	5.06	0.14	198
T <sub>6</sub> -75:50:50 g NPK plant <sup>-1</sup>	12.63	5.35	0.14	205
T <sub>7</sub> -75:75:25 g NPK plant <sup>-1</sup>	12.93	5.53	0.15	208
T <sub>8</sub> -75:75:50 g NPK plant <sup>-1</sup>	12.95	5.61	0.15	214
T <sub>9</sub> -100:50:25 g NPK plant <sup>-1</sup>	13.14	6.15	0.16	218
T <sub>10</sub> -100:50:50 g NPK plant <sup>-1</sup>	13.21	6.22	0.16	225
T <sub>11</sub> -100:75:25 g NPK plant <sup>-1</sup>	14.13	6.96	0.21	256
T <sub>12</sub> -100:75:50 g NPK plant <sup>-1</sup>	15.19	7.26	0.24	263
T <sub>13</sub> -125:50:25 g NPK plant <sup>-1</sup>	13.54	6.41	0.17	232
T <sub>14</sub> -125:50:50 g NPK plant <sup>-1</sup>	13.68	6.50	0.18	245
T <sub>15</sub> -125:75:25 g NPK plant <sup>-1</sup>	13.69	6.56	0.19	250
T <sub>16</sub> -125:75:50g NPK plant <sup>-1</sup>	15.10	7.04	0.22	259
T <sub>17</sub> -Absolute control	9.45	3.62	0.10	167
<b>SEd</b>	<b>1.78</b>	<b>0.20</b>	<b>0.01</b>	<b>1.32</b>
<b>CD(P=0.05)</b>	<b>3.61</b>	<b>0.42</b>	<b>0.03</b>	<b>2.68</b>

### Photosynthetic rate (A)

Photosynthetic rate which is a measure of productive potential of a tree was measured and the results are presented in the Table 1. During 6 MAP, the application of 100:75:50 g NPK plant<sup>-1</sup> (T<sub>12</sub>) recorded the highest photosynthetic rate of 15.19  $\mu \text{ mol m}^{-2} \text{ s}^{-1}$  followed by the application of 125:75:50 g NPK plant<sup>-1</sup> (T<sub>16</sub>) which

recorded  $15.10 \mu \text{ mol m}^{-2} \text{ s}^{-1}$  whereas the treatment  $T_{17}$  which did not receive any fertilizers registered the lowest photosynthetic rate of  $9.45 \mu \text{ mol m}^{-2} \text{ s}^{-1}$ .

### **Transpiration rate (E)**

The transpiration rate exhibited significant variation among the different treatments imposed (Table 1). At 6 MAP, the highest transpiration rate of  $7.26 \text{ m mol m}^{-2} \text{ s}^{-1}$  was observed due to application of  $100:75:50 \text{ g NPK plant}^{-1}$  ( $T_{12}$ ) followed by the application of  $125:75:50 \text{ g NPK plant}^{-1}$  ( $T_{16}$ ) and  $100:75:25 \text{ g NPK plant}^{-1}$  ( $T_{11}$ ) with the values of  $7.04$  and  $6.96 \text{ m mol m}^{-2} \text{ s}^{-1}$  respectively. The lowest transpiration rate of  $3.62 \text{ m mol m}^{-2} \text{ s}^{-1}$  was registered where no fertilizer were imposed.

### **Stomatal conductance (gs)**

Stomatal conductance recorded significant variation among the various treatments and the results are presented in Table 1. During 6 MAP, the highest stomatal conductance of  $0.24 \text{ m mol m}^{-2} \text{ s}^{-1}$  was reported due to application of  $100:75:50 \text{ g NPK plant}^{-1}$  ( $T_{12}$ ) followed by the application of  $125:75:50 \text{ g NPK plant}^{-1}$  ( $T_{16}$ ) and  $100:75:25 \text{ g NPK plant}^{-1}$  ( $T_{11}$ ) with the values of  $0.22$  and  $0.21 \text{ m mol m}^{-2} \text{ s}^{-1}$  respectively. The treatment  $T_{17}$  recorded the lowest stomatal conductance of  $0.10 \text{ m mol m}^{-2} \text{ s}^{-1}$  where no nutrients were applied.

### **Inter Cellular CO<sub>2</sub> Concentration (CINT)**

Among the different treatments, the maximum inter cellular CO<sub>2</sub> concentration (CINT) of  $263 \text{ ppm}$  was observed in  $T_{12}$  followed by  $T_{16}$  and  $T_{11}$  with the values of  $259$  and  $256 \text{ ppm}$  whereas the treatment  $T_{17}$  which did not receive any fertilizers registered lowest inter cellular CO<sub>2</sub> concentration (CINT) of  $167 \text{ ppm}$ .

### **Ecophysiological attributes**

The ecophysiological behaviour of plants plays a vital role in the growth of plants and are important factors for changing environmental conditions (Camposeo *et al.*, 2011). The influence of different levels of nutrients on the parameters *viz.*, photosynthetic rate (A), transpiration rate (E), stomatal conductance ( $g_s$ ) and Inter-cellular  $CO_2$  concentration (C<sub>int</sub>) are discussed.

### **Photosynthetic rate (A)**

At 6 MAP, significant difference in photosynthetic rate was observed due to the influence of various treatments imposed. Photosynthetic rate was highest in T<sub>12</sub> (100: 75: 50 g NPK plant<sup>-1</sup>) which recorded 15.19  $\mu\text{ mol m}^{-2} \text{ s}^{-1}$  followed by T<sub>16</sub> (125:75:50 g NPK plant<sup>-1</sup>) which recorded 15.10  $\mu\text{ mol m}^{-2} \text{ s}^{-1}$  and were *on par*.

Similar findings were also observed by Baranidharan *et al.* (2013) who observed the highest net photosynthetic rate of 6.61 and 6.76  $\mu\text{ mol. m}^{-2} \text{ s}^{-1}$  when *Casuarina* species was applied with 25 per cent more of fertilizer over the soil test recommendation value of 93.75: 225: 100 of NPK kg ha<sup>-1</sup>.

Mohan Raj (2008) observed the highest photosynthetic rate of 8.21  $\text{m mol. m}^{-2} \text{ s}^{-1}$  in Eucalyptus at 18 MAP in heavy fertilizer dose application (150:200:100 g NPK kg<sup>-1</sup> of soil).

Increased fertilizer application increased the photosynthetic rate in trees (Kao and Tsai, 1999) who stated that an increase in nitrogen availability significantly increased the photosynthetic rate in *Kandeliacandel* grown under saline conditions. Sitka spruce (*Picea sitchensis*) seedlings supplied with high nitrogen dose recorded higher photosynthetic rate than seedlings supplied with low nitrogen dose (Murray *et al.*, 2000). The current finding also state that increased application of N, P and K enhanced the photosynthetic rate in *Thespesia populnea*.

### **Transpiration rate (E)**

Significant difference in transpiration rate was observed at 6MAP due to the influence of various treatments imposed. The transpiration rate was highest in T<sub>12</sub> (100: 75: 50 g NPK plant<sup>-1</sup>) which recorded 7.26 m mol. m<sup>-2</sup> s<sup>-1</sup>. This result is in tune with Mohan Raj (2008) who recorded the highest transpiration rate in Eucalyptus (2.63 m mol. m<sup>-2</sup>s<sup>-1</sup>) at 18 MAP in heavy fertilizer dose application.

Baranidharan *et al.* (2013) observed that 25 per cent more of fertilizer application over the soil test recommendation value of 93.75: 225: 100 of kg NPK ha<sup>-1</sup> in *Casuarina* species recorded the highest transpiration rate value of 2.20 and 2.38 m mol. m<sup>-2</sup> s<sup>-1</sup> and was significantly superior to all the other nutrient levels recorded.

In *Gmelina arborea*, significantly higher transpiration rate of 4.81 m mol. m<sup>-2</sup> s<sup>-1</sup> followed by *Dalbergia sissoo* (4.32 m mol. m<sup>-2</sup> s<sup>-1</sup>), *Tectona grandis* (4.01 m mol m<sup>-2</sup> s<sup>-1</sup>) and *Bambusa vulgaris* var. *vulgaris* (2.47 m mol m<sup>-2</sup> s<sup>-1</sup>). The lowest transpiration rate of 1.67 m mol m<sup>-2</sup> s<sup>-1</sup> was registered by *Swietenia macrophylla* at 21 MAP (Narendra Babu, 2012). Transpiration rate from several Australian semiarid zone species was studied in the field by Kreeb (1966) using the torsion balance technique and he concluded that among the different species studied, the maximum transpiration rate was observed in *Eucalyptus camaldulensis* (5.0 and 7.9 m mol m<sup>-2</sup> s<sup>-1</sup>).

### **Stomatal conductance rate (g<sub>s</sub>)**

Stomatal conductance rate exhibited significant difference at 6 MAP due to the influence of various treatments applied. The highest stomatal conductance rate was found to be associated with the application of 100: 75: 50 g NPK plant<sup>-1</sup> (T<sub>12</sub>) which recorded the value of 0.24 m mol. m<sup>-2</sup> s<sup>-1</sup> followed by 100:75:25 g NPK plant<sup>-1</sup>

<sup>1</sup>(T<sub>11</sub>) which recorded 0.21 m mol. m<sup>-2</sup> s<sup>-1</sup> and which were *on par* with each other. This is in line with findings of De Costa *et al.*(2000) who reported that the stomatal conductance in nine tree species *viz.*, *Alstonia macrophylla*, *Filiciumdecipiens*, *Macaranga peltata*, *Tectona grandis*, *Acronychia pedunculata*, *Semicarpusnigroviridis*, *Swietenia macrophylla*, *Mesua ferrea* and *Terminalia catappa*growing under different levels of natural shade, i.e. open, medium shade and full shade. According to them, the total leaf stomatal conductance (g<sub>s</sub>) varied significantly with the tree species and shade levels and the highest (g<sub>s</sub>) was observed in *S. nigroviridis* and *T. catappa*, with values of 92.03 and 78.18 m mol. m<sup>-1</sup> s<sup>-1</sup> respectively.

Stomata are the points of exit for the water vapour from leaves. Under conditions of adequate water supply, stomata open in the morning with increasing radiant flux and maintain a relatively constant aperture until radiant flux declines in the late afternoon. So, there was a correlation present among the stomatal opening and radiant flux (Giklooet *al.*, 2012). They also reported that the leaf stomatal conductance during January was consistently higher than during September (0.08 to 0.14 vs 0.02 to 0.08 mol m<sup>-2</sup> s<sup>-1</sup>).

Ladjalet *al.* (2007) studied greenhouse grown Cedar seedlings subjected to two different watering regimes and observed that dry set showed lower net photosynthesis and stomatal conductance than the plants in the well watered set. Measurements of diameter, growth, CO<sub>2</sub> uptake and stomatal conductance at the age of 4-5 years showed that Victorian provenances of *Eucalyptus camaldulensis* was better adapted to semi-arid conditions.

### **Inter cellular CO<sub>2</sub> concentration (C<sub>INT</sub>)**

The inter cellular CO<sub>2</sub> concentration exhibited significant difference at 6 MAP due to the influence of various treatments applied. The highest inter cellular CO<sub>2</sub> concentration was found to be associated with the application of 100 : 75 : 50 g NPK plant<sup>-1</sup> (T<sub>12</sub>) they recorded the value of 263 ppm followed by 125:75:50 g NPK plant<sup>-1</sup>(T<sub>16</sub>) which recorded 259 ppm and which were *on par*.

Inter cellular CO<sub>2</sub> is also an indicator of plant productivity in which it is highly related to plant photosynthetic rate. Esser (2004) correlated Inter cellular CO<sub>2</sub> concentration with the photosynthetic efficiency of three species. Similar result was observed by Suresh Kumar (2011) and Narendra Babu (2012) have also confirmed that Inter cellular CO<sub>2</sub> concentration in *Tectona grandis* and *Gmelina arborea* was lower than that of *Dalbergia sisoo* and *Bambusa vulgaris* var. *vulgaris* during 9 MAP respectively.

### **Summary and Conclusion**

Nutrient management is one of the prime factors which plays a vital role in the growth, development and successful completion of life cycle in crop plants. Judicious integration of organic and inorganic fertilizers has been proved to be a viable and economically feasible technology for enhancing the initial growth and development of plants.

Growth components *viz.*, plant height, collar diameter, volume index and biomass yield were recorded. Growth analysis was done by estimating LAI and ecophysiological parameters *viz.*, photosynthetic rate (A), transpiration rate (E), stomatal conductance (g<sub>s</sub>) and Inter cellular CO<sub>2</sub> concentration (C<sub>int</sub>) were also recorded. The growth attributes were influenced by N, P and K at different levels.

Among N, P and K levels, application of 100: 75: 50 g NPK plant<sup>-1</sup> (T<sub>12</sub>) significantly influenced the growth attributes followed by 100: 75: 25 g NPK plant<sup>-1</sup> (T<sub>11</sub>).

## Reference

- Abbott, I.A. 1992. Laau Hawaii: Traditional Hawaiian uses of plants. Bishop Museum Press, Honolulu.
- Adalarasan, R. 2002. Integrated nutrient management studies on Casuarina (*Casuarina equisetifolia*) seedlings. **M.Sc. Thesis**, Tamil Nadu Agricultural University, Coimbatore.
- Adalarasan, R. 2008. Carbon sequestration potential of teak (*Tectona grandis*) as influenced by fertilization and irrigation. **Ph.D. Thesis**, Tamil Nadu Agricultural University, Coimbatore, India.
- Balasubramanian, A. and K. Gurumurthi. 2001. Divergence studies in *Casuarina equisetifolia* for grouping of productive clones. *Casuarina improvement and utilization*. **IFGTB**, **57**: 62.
- Baranidharan, K., M. Vijayabhama and K.K. Suresh. 2013. Intensive silvicultural packages for Casuarina species. Lap Lambert, Academic Publishing, Germany. pp. 292.
- Bassow, S.L. and F.A. Bazzaz. 1997. Intra- and inter-specific variation in canopy photosynthesis in a mixed deciduous forest. **Oecologia**, **109**: 507-515.
- Chiang, C., W. YaNan, C. Kuo Chen, H. Kuo Chung, C.H. Chiang, Y.N. Wang, K.C. Chang and K.C. Hwang. 1996. A study of the germination of seed, cutting and characteristics of seedling in *Ginkgo biloba* L. **Quart. J. Experimental Forest of National Taiwan University**, **10**: 29-36.
- Clark, W.C. and R.R. Thaman (eds.) 1993. Agroforestry in the Pacific Islands: Systems for sustainability. United Nations University Press, New York.
- De Costa, Abeysinghe and A.G. Chandrapala. 2000. Relationship between stomatal conductance and leaf water potential in selected forest tree species growing under different levels of natural shade in the mid-country wet zone. **Journal of Natural Science Foundation Sri Lanka**, **28(1)**: 63-78.
- Dinesh Kumar and N. Ahmed. 2010. Response of nitrogen and potassium fertigation to 'Waris' almond (*Prunus dulcis*) under north western Himalayan region of India. **Indian J. Agric. Sci.**, **80**: 512-516.
- Dolley, D. and B.J. Grieve. 1966. Measurement of sap flow in a eucalyptus by thermo-electric methods. **Aust. For. Res.**, **2**: 3-27.
- Esser Gerd. 2004. Stomatal response of some trees and shrubs from various tropical environments to variable CO<sub>2</sub> concentrations. **Tropical Ecology**, **45(1)**: 113-121.
- Gikloo, Samandari and Behnam Elhami. 2012. Physiological and morphological responses of two almond cultivars to drought stress and cyclone. **International Research Journal of Applied and Basic Sciences**, **3(5)**: 1000-1004.
- Hanum, I.F. and L.J.G. van der Maesen (Eds.). 1997. Plant Resources of South-East Asia No. II. Auxiliary Plants. Backhuys Publishers, Leiden, The Netherlands.
- Jiang Gaoming, Lin Guang Hui and Bruno D.V. Marino. 1997. Changes in stomatal conductance, transpiration and water use efficiency of ten species experienced

- in high CO<sub>2</sub> concentrations in biosphere. **Acta Botanica Sinica**, **39(6)**: 546-553.
- Kao, W.Y. and H.C. Tasi. 1999. The photosynthesis and chlorophyll a fluorescence in seedlings of *Kandeliacandel* (L.) Druce., grown under different nitrogen and NaCl control. **Photosynthetica**, **37(3)**: 405-412.
- Kreeb, K. 1966. Transpiration messenger an engine australiaschen Immergrünen in Bereich des feuchten *Eucalyptus walde*s und der *Mulgabuschsavanne*. **Oecol. Plant**, **1**: 235-244.
- Kundu, S.K. and P.M.A. Tigerstedt. 1999. Variation in net photosynthesis, stomatal characteristics leaf area and whole plant phytomass production among ten provenances of neem (*Azadirachta indica*). **Tree Physiol.**, **19**: 47-52.
- Ladjal Mehdi, Nathalie Deloche, Roland Hucand Michel Ducrey. 2007. Effects of soil and air drought on growth, plant water status and leaf gas exchange in three Mediterranean cedar species: *Cedrus atlantica*, *C. brevifolia* and *C. libani*. **Trees - Structure and Function**, **21(2)**: 201-215.
- Murray, M.B., R.I. Smith, A. Friend and P.G. Jarvis. 2000. Effect of elevated CO<sub>2</sub> and varying nutrient application rates on physiology and biomass accumulation of Sitka spruce (*Picea sitchensis*). **Tree Physiology**, **20(7)**: 421-434.
- Narendra Babu, S. 2012. Quantifying carbon sequestration potential in fast growing trees through afforestation. **M.Sc. Thesis**, Tamil Nadu Agricultural University, Coimbatore.
- Okuto, J.L. and G. Ouma. 2010. Provenance variation of photosynthesis of seedlings of java plum (*S. cuminii*) in Western Kenya. **Agriculture and Biology Journal of North America**, **(6)**: 1260-1263.
- Rao, P.B. 2005. Effect of shade on seedling growth of five important tree species in Tarai region of Uttaranchal. **Bulletin of the National Institute of Ecology**, **15**: 161-170.
- Saraswathi, S.G. and K. Paliwal. 2008. Diurnal and seasonal trends in photosynthetic performance of *Dalbergia sissoo* Roxb. and *Hardwickiabinata* Roxb. from a semi-arid ecosystem. **Photosynthetica**, **46**: 248-254.
- Srivastava, N.K. and L.C. Ram. 2009. Bio-restoration of coal mine spoil with fly ash and biological amendments. Published in edited book on Sustainable Rehabilitation of Degraded Ecosystems (Chaubey, O.P., Vijay Bahadur and P.K. Shukla (Eds.). Aavishkar Publishers, Distributors, Jaipur, Raj. 302 003 India. pp. 77-99.
- Ujala Singh. 2005. Comparative study of transpiration rates by different *Eucalyptus* species under various soil moisture and climatic conditions. **Indian Journal of Forestry**, **28(2)**: 112-115.