

Pre-sowing treatments on seeds of forest tree species for overcome the germination problems

Abstract

Production of the quality planting material has get more emphasis under sub-mission on agroforestry (SMAF) with aiming to increase tree cover outside the forest area. Establishment of the tree species depends on the quality of the planting material, available soil, water for irrigation and the adopted protection measures. Most of the tree species get problem with germination of seed due to external and internal factors and causes seed dormancy. Different kind of pre-sowing treatments were tested and applied for the different kind of tree species by researchers of the forestry. Scarification (mechanical, acid), water soaking (hot/cold), the application of chemicals and plant growth regulators, or alternate wetting and drying prior to sowing effectively break seed dormancy and improve seed germination for producing required quality planting material. Auxins (IAA, IBA, 2-4D, 4-CPA), Gibberellic acid (GA3), Cytokinins (Kinetin, Zeatin, Benzyl adenine), Ethylene (Ethereal) and Abscisic acid (Dormins, Phaseic Acid) are plant growth regulators also used in different concentrations for improving tree seed germination.

Keywords: Pre-sowing treatment, seed dormancy, plant growth regulators, growth hormones, stratification, and scarification.

Introduction

Plantation of the forest tree species has emerged as a crucial initiative in forestry sector to meet the increasing demand for timber, non-timber forest produce (NTFP), fodder, fuel wood etc. both from natural forests and plantation forestry. Establishment of plantations recognizing the current need for sustainable alternatives for depleting natural forests, and gained significance for industrial and domestic wood production on a large as well as small scale. The vital decision in initiating a plantation project is the selection of the suitable tree species. The availability of high-quality planting material stands as a significant determinant influencing the choice of species for large-scale afforestation. Therefore, the nursery practices of the suitable tree species is the key for successful establishment of plantations. A critical concern in this context is the insufficient awareness regarding suitable pre-sowing seed treatments for various tree species. This knowledge gap poses a significant obstacle to the mass production and conservation efforts of many species. Addressing this concern necessitates focused research into innovative seed technologies, considering the growing importance of this domain in the global crop protection markets, as emphasized by Sharma *et*

al. (2015). Thus, a profound understanding of seed science, encompassing aspects such as seed collection, storage, viability, and germination, becomes imperative for enhancing the success of restoration endeavours.

Seed germination constitutes the process through which the embryo evolves into a seedling under favourable conditions. It involves a series of physiological, biochemical, and morphological changes that collectively lead to germination. This intricate process can be delineated into three major phases: Phase I, characterized by rapid water imbibition; Phase II, dominated by the reactivation of metabolism, cell elongation, and chitting; and Phase III, marked by rapid cell division coinciding with radicle growth. Phase II, in particular, is pivotal, as it involves the reactivation of crucial physiological and biochemical processes such as the hydrolysis of food reserves, macromolecular biosynthesis, respiration, reorganization of subcellular structures, and cell elongation, all of which play a fundamental role in initiating germination (Bonsager *et al.*, 2010).

Seed germination can be classified into two types based on the outcome of the cotyledons:

1. **Epigeal Germination:** This process involves the cotyledon being pushed out of the soil due to the rapid growth and elongation of the hypocotyl, as seen in species like *Acacia*, *Adenantha*, *Albizia*, *Cassia*, *Dipteryx*, *Diphysa*, *Erythrina*, *Gliricidia*, *Haematoxylum*, *Hymenaea*, *Hymenolobium*, *Parkia*, *Parkinsonia*, *Pterocarpus*, *Samanea saman*, *Sclerolobium*, *Tamarindus*, *Casuarina*, *Annona*, *Cordia alliodora*, *Myrcia*, *Capparis*, *Cedrela*, *Melia*, *Zizyphus*, *Anacardium excelsum*, *Jacaranda*, *Ulmus*, *Ilex* and *Elaeocarpus*.
2. **Hypogeal Germination:** In this type, cotyledons remain below the soil due to the rapid elongation of the epicotyl. This predominantly occurs in monocotyledonous seeds, E.g. *Calophyllum*, *Swartzia*, *Syzygium*, *Swietenia*, *Prunus*, *Sapindus*, *Terminalia catappa* etc.

Seed germination and quality are influenced by both internal and external factors. Genetic makeup, seed viability and dormancy are internal factors inherited from parent plants, influencing germination potential and subsequent growth. Seed source variations among species can significantly affect seed quality and growth (Kumar *et al.*, 2022; Kumar *et al.*, 2021). External factors consider the growing environmental conditions, including temperature, moisture, light, and soil properties, also influence germination and seedling establishment (Penfield *et al.*, 2005; Srivastava, 2002). Additionally, external interventions such as seed priming, scarification, and chemical treatments can modify seed coat properties, break dormancy, or enhance seedling growth. The interaction between internal genetic factors and external environmental conditions is paramount in determining the success of seed germination and subsequent seedling establishment. This review specifically focuses on pre-sowing treatments for seeds of forest tree species to address germination challenges.

The Mechanism of Seed Germination

Germination, a complex biological process, initiates with a crucial step known as imbibition, where the seed absorbs water. Following imbibition, activated enzymes commence the breakdown of starch into sugars, providing nourishment to the embryo. The initial sign of germination is the swelling of the radicle, marking the onset of growth. The duration for seedlings to emerge varies based on seed size. Species with large seeds possess substantial food reserves, enabling them to germinate at greater depths and extend their epicotyl to the soil surface. Conversely, seedlings of small-seeded species emerge more rapidly, typically closer to the soil surface.

The germination process can be categorized into epigeous and hypogeous types. In epigeous germination, the hypocotyl elongates, and the cotyledons rise above ground. On the other hand, hypogeous germination involves the elongation of the epicotyl, with the cotyledon(s) remaining belowground. Epigeous species, like beans and onions, witness the cotyledons turning green and engaging in photosynthesis until they eventually detach. In contrast, hypogeous species, such as peas and corn, exhibit cotyledons that remain belowground throughout the germination process. This intricate interplay of imbibition, enzymatic activity, and seedling emergence underscores the diverse strategies employed by plant species during germination, influenced by factors like seed size and environmental conditions.

Factors Affecting Seed Germination

Seed germination can be influenced by a combination of internal and external factors (Figure 1). Genetic characteristics, seed viability and dormancy are internal factors inherited from parent plants, impacting germination potential and subsequent growth. External factors encompass moisture, temperature, oxygen, light, and soil conditions. Conversely, internal factors influencing the germination process include the embryo maturity of the seed, seed viability, and seed dormancy. Understanding and manipulating these factors are crucial for optimizing the germination process in tree seedlings.




Figure 1: Factors influencing seed germination and quality

Source: adopted from El-Ramady *et al.* (2023)

Seed Dormancy

Seed dormancy, the physiological state preventing a viable seed from germinating even under favourable conditions (Umarani and Vannangamudi, 2005; Willan 1985; Baskin and Baskin, 2004), exhibits considerable variation in storage duration among species. While some seeds can be stored for several months, others rapidly lose viability. Comprising an outer protective coat and an inner embryo developing into the plant, seeds initiate germination when water permeates the seed coat, resulting in swelling and subsequent plant growth.

Types of Dormancies

1. *Physiological Dormancy*: Predominantly observed in species like Fagus, Pinus, and eucalyptus, physiological dormancy is initiated by either the embryo or surrounding endosperm tissues. Freshly matured seeds with non-deep physiological dormancy either cannot germinate at any temperature or only germinate over a narrow temperature range.
2. *Morphological Dormancy*: Seeds with underdeveloped embryos, requiring time for growth and germination (Umarani and Vannangamudi, 2005; Finch and Leubner 2006).
3. *Physical Dormancy*: Seed coats impermeable to water due to a layer of thick-walled, cutinized palisade-like macroscleroid cells. Various treatments, such as mechanical and

chemical scarification or soaking in boiling water, are employed to induce germination in water-impermeable seed coats. For water-permeable seed coats, cold stratification, treatment with gibberellic acid (GA₃), or sodium hypochlorite, scarification, or heat treatment can be applied (Baskin and Baskin, 2020).

Various methods have been employed to induce germination in seeds with water-impermeable seed coats, including mechanical scarification, chemical scarification, and soaking in boiling or hot water. Conversely, for seeds with water-permeable coats, a range of treatments can be applied, such as cold stratification, treatment with gibberellic acid (GA₃), or sodium hypochlorite, along with scarification or heat treatment (Baskin and Baskin, 2020). Some seeds naturally overcome dormancy over time, while others require specific pre-treatments. Seed dormancy poses a significant hurdle to germination, necessitating pre-sowing treatments. Techniques like scarification (mechanical, acid), water soaking (hot/cold), the application of chemicals and plant growth regulators, or alternate wetting and drying prior to sowing effectively break seed dormancy.

1. Breaking Seed-Coat Dormancy: Preferred procedures to overcome seed-coat dormancy include puncturing or scarifying the seed coat through methods like piercing, nicking, chipping, or filing with a knife, needle, or sandpaper.

2. Breaking Embryo Dormancy: Several recommended treatments exist to overcome embryo dormancy:

a) Pre-chilling (Cold Stratification): Seeds are placed in containers on a moistened germination substrate and kept at 3°-5°C in a refrigerator for 7-14 days. Seeds derived from cold-climate species, particularly perennials accustomed to prolonged snow cover and harsh winter conditions; necessitate significantly lower temperatures over extended periods for effective stratification. An illustrative case involves a perennial herb adapted to the extended cold winters of mountainous regions, demonstrating enhanced germination rates. Specifically, exposure to a low temperature of 32 °F (0 °C) for a duration of 7-months resulted in a notably higher germination rate, reaching 60% (Cuenca-Lombrana *et al.*, 2018).

b) Treatment with Hormones: Dormancy can be broken by directly applying the plant hormones: abscisic acids (ABA), gibberellin (GA₃), ethylene, and jasmonic acid (JAS). Gibberellin (GA₃) is the most effective in inducing germination (Shu *et al.*, 2016). To assess germination, germination test paper is dampened with a 0.05% gibberellic acid (GA₃) solution, prepared by dissolving 500 mg of GA₃ in 1 litre of water. The mechanism involves softening the endosperm, promoting embryo growth, and facilitating seedling emergence by breaking through the seed coat.

c) Chemical Scarification: Seed impermeability, governed by the seed coat, can be addressed through chemical scarification. The impermeability of seed coats to water poses a challenge

for the germination of seeds from tree species in dry and semi-arid regions (Kheloufi *et al.*, 2018). Dormancy can be overcome by soaking seeds in chemicals such as acidic or basic solutions, organic solvents, alcohol, disinfectants, or boiling water. Common acids employed for scarifying seeds include phosphoric acid (25 to 75% concentration depending on the seed type), potassium nitrate, nitric acid, and sulphuric acid (96%).

For potassium nitrate treatment, a 0.2% solution is prepared by dissolving 2 g of potassium nitrate (KNO₃) in 1 litre of water. This solution is used to moisten the germination paper at the initiation of the test. This specific treatment aids in seed scarification, facilitating water absorption and breaking dormancy barriers for improved germination outcomes.

d)Light Treatment: Exposing seeds to strong cool-white and infrared light mimics the natural environment during spring and summer. The light source, close to the seeds, should be on for at least 8 hours daily in a moist environment under low temperature. After light treatment, seeds should be incubated in the dark until germination. Positive results have been observed with both cool-white and red light, while germination rates are lower in blue and green light (Penfield *et al.*, 2005; Liand Khan, 2005).

The role of light varies across different seed sizes. Generally, medium and large-sized seeds exhibit a neutral photoblastic nature, allowing them to germinate in both light and darkness. The germination process for these seeds is primarily governed by factors such as soil moisture and temperature, with light having minimal influence. Conversely, some seeds, termed negatively photoblastic, necessitate darkness for successful germination. Notably, sizable seeds, approximately 1 inch (2.5 cm) in length, fall into this category, demonstrating a preference for germination in the absence of light. For this species, exposure to light impedes germination, attributed to the presence of phytochrome A in the seeds. Phytochrome A plays a crucial role by diminishing the production of germination-stimulating enzymes upon exposure to light (Goggin *et al.*, 2011).

On the other end of the spectrum, tiny seeds, measuring less than ¼ inch (6 mm), present distinct light-dependent germination requirements. These little seeds typically mandate a minimum of 8 hours of light exposure to initiate germination. This need arises from their limited food reserves, requiring efficient utilization for soil penetration and light access. Notably, certain vegetable and herb seeds fall into this category, relying on light to break seed dormancy. The mechanism involves photoreceptors, specifically phytochrome B, which responds to light stimuli. Phytochrome B acts by suppressing the production of abscisic acid, a key dormancy-maintaining factor, and simultaneously promoting the synthesis of gibberellins hormone (Leite and Takaki, 2001; Socolowski and Takak, 2004; Attriet *al.*, 2015), thereby facilitating germination in these small-sized seeds.

e) **Hot or Cold Water:** This is an economical and farmer-friendly practice was identified for enhancing germination. This treatment, applied to most medium-sized dry seeds, improves germination when seeds absorb adequate water and soften the seed coat. Heat treatment is moderately effective in breaking dormancy. The study suggests 1-30 minutes of acid scarification and one-hour (55-77°C) heat treatment for optimizing germination based on species (Koutouan-Kontchoiet *al.*, 2020). Seeds require soaking in water for 24-48 hours, serving to soften hard seeds and leach out chemical inhibitors (Umarani and Vanangamudi, 2005).

f) **Stratification method (warm and/or cold)** - Stratification is the process of breaking down the hard outer coating of a seed to promote germination. This process involves exposing the seed to a moist environment, which softens the outer shell, making it easier for the seed to sprout. There are two types of stratification: cold stratification and warm stratification (D'Esteet *al.*, 2019).

g) **Scarification**- Scarification is the process of breaking or weakening the hard outer coating of a seed to promote germination. This process involves physically damaging the seed coat by scratching, nicking, or rubbing it with sandpaper or a file. Scarification is commonly used for seeds that have a hard outer coating that prevents water and air from reaching the embryo (D'Esteet *al.*, 2019).

Example of Seed Pre-Treatment for Improved Germination

Soaking in Cold Water: Soaking seeds in cold water for 24 to 48 hours prior to sowing has been explored as an effective pre-treatment method. According to Hossain *et al.* (2005), *Terminalia chebula* seeds exhibited the highest germination percentage (66.7%) with a 48-hour cold-water treatment, further increasing to 73% when de-pulped before the cold-water treatment. A study by Viswanath *et al.*, 2021, on *Terminalia paniculata*, also supported the positive impact of soaking seeds in water for 24 hours. Soaking *Jatropha* seeds in water overnight proved effective for enhancing germination (Sunil *et al.*, 2016).

Soaking in Hot Water: For seeds with hard coatings, soaking in boiling hot water has proven to be an advantageous method. Commonly applied to seeds with robust shells like *Albizia*, *Acacia*, *Cassia*, *Poplars* and *Leucaena leucocephala*, this treatment involves soaking seeds in hot water for 2-48 hours, depending on the species. Compared to traditional methods like sulphuric acid, hot water treatment is considered a better, safer, and more cost-effective alternative. Notably, *Subabul* seeds soaked in hot water for 12 hours exhibited a significantly increased germination percentage (42.22%) (Hamad and Anwer 2021). Similarly, *Sapindus mukorssi* seeds soaked in hot water for 10 seconds achieved the highest germination rate (72%). For *Acacia mearnsii*, immersion in hot water at 80°C, either for 5 minutes or

simply at 80°C, was found to be the most optimum treatment, resulting in total germination rates of 83 % (São José 2019) and for *Acacia auriculiformis* (Azad *et al.*, 2011), respectively.

Alternate Wetting and Drying: This method proves effective for seeds with thicker seed coats (Dwivedi, 1993). Gupta and Pathanath (1975) reported that teak seeds from certain localities require no treatment, while others got benefit from alternate wetting and drying procedure. Similarly, Pamei *et al.* (2017) found that *Tectonagrandis* seeds subjected to alternate wetting (12 hrs) and drying (12 hrs) for 8 days exhibited the highest germination percentage (43.3%), along with optimal seedling height and root length. For *Buchananialanzan* seeds, the highest germination of 86.7% was achieved with alternate wetting (24 hrs) and drying (12 hrs) (Thounaojam and Dhaduk 2020).

Knuckling and Sand Paper Scarification (Mechanical): Mechanical treatments play a crucial role in breaking the impermeable and hard seed coats of species such as *Acacia catechu*, *A. nilotica*, *Albizia spp.*, *Cassia fistula* etc. Research indicates that knuckling the distal ends of *Gliricidiasepium* seeds enhances germination without causing damage to the embryo (Alamgir and Hossain, 2005; Chichaghareet *al.*, 2020). The scarification of seeds from *Melia azedarach* and *Acacia auriculiformis* with sandpaper resulted in impressive germination rates of 80% and 78%, respectively (Azad *et al.*, 2010 & 2011). *Tamarindus indica* recorded a germination percentage of 82.3% following scarification (Azad *et al.*, 2013), while *Parkiabiglobosa* demonstrated a remarkable 91.7% germination rate (Okunlolaet *al.*, 2011). Additionally, *Schleicheraoleosa* seeds, when rubbed with sandpaper at the distal end, exhibited a 42% germination rate (Tanjinaet *al.*, 2014). Nail clipping one side of *Albizia saman* seeds at the distal end resulted in the highest germination rate of 50% (Alamgir and Hossain, 2005). Manual cutting of seeds also proved effective, with *Prosopisjuliflora* and *Dalbergiasissoo* achieving 100% and 90.4% germination, respectively (Asif *et al.*, 2020).

Nitrogenous Compounds: The impact of nitrate compounds on seed germination, in conjunction with other factors like temperature or light, is well-established (Eremrena and Mensah, 2016). Potassium nitrate (KNO₃) stands out as a widely used chemical to release seed dormancy and stimulate germination. Recommended solutions of 0.1–0.2% KNO₃ have proven effective in optimizing germination for species exhibiting low or shallow dormancy patterns (Gashi *et al.*, 2012). KNO₃ is also a frequently applied priming chemical, known to enhance seed germination by improving both uniformity and speed of germination (Tapfumaneyiet *al.*, 2023; Thongtipet *al.*, 2022). Moreover, the application of KNO₃ has shown utility in mitigating the adverse effects of salt on seed germination, seedling growth, mitotic activity, and chromosomal aberrations (Çavuşoğlu *et al.*, 2017) and also effective in improving emergence and seedling growth under drought (Ali *et al.*, 2020).

Acid Scarification: Soaking seeds in concentrated sulphuric acid is a common method for softening the seed coat. Sulphuric acid is believed to disrupt the seed coat and expose the lumens of the macrosclereids cells, facilitating water imbibition (Nikoleave, 1977). *Semicarpusanacardium* seeds treated with 98% concentrated H₂SO₄ for 1 minute recorded the highest germination percentage at 74.67%. Olatunji *et al.* (2013) reported that *Acacia auriculiformis* seeds treated with concentrated H₂SO₄ for 5-10 minutes exhibited the highest germination percentage of 92-96%. Oxidation of phenolic compounds through hypochlorite or hydrogen peroxide, or decreased polyphenol oxidase activity, can enhance seed germination when seed coats are impermeable (Gendreau and Corbineau, 2009).

A study on breaking the dormancy of *Prunus yedoensis* seeds also found the best germination rate (40%) when the seeds were chemically treated, washed, and stratified for 9 weeks (Kim, 2019).

Seed Priming Treatments: Seed priming, a preparatory treatment before sowing, which involves hydrating seeds sufficiently to initiate metabolic processes before germination without allowing radicle emergence (Rashid *et al.*, 2004; Rehman *et al.*, 2011). Natural priming has proven beneficial for on germination of tropical trees such as *Albizia saman*, *Cedrela odorata*, and *Swietenia macrophylla*, while hydropriming exhibited positive effects on *Enterolobium cyclocarpum* seeds (Peraza *et al.*, 2018). Inoculation of *Acacia senegal* (L.) Willd. seeds with rhizobium also exhibited positive effects on the germination process (Singh *et al.*, 2011).

Hormonal Regulation in Seed Germination

Plant growth regulators play a pivotal role in seed germination, with hormones such as GA, ABA, auxin, and ethylene (table 1) demonstrating significant influence (Han and Yang, 2015). Ethylene might be crucial in regulating dormancy through interaction with GA metabolism. Seed treatments with GA solutions have been found to enhance germination parameters, improving seed vigour and uniform germination (Maharana *et al.*, 2018). Additionally, applying plant growth regulators like GA₃, IAA, and IBA, along with cold stratification, has been effective in breaking dormancy and enhancing germination in various species (Dhiman *et al.*, 2015; Iralu *et al.*, 2019).

Plant Growth Regulators:

a. Abscisic Acid (ABA): ABA inhibits the germination of non-dormant seeds and is a crucial factor inducing dormancy, accumulating in maturing seeds. ABA biosynthesis genes directly influence seed germination and respond to abiotic stresses, with varying ABA levels at different developmental stages regulating processes like seed maturation and germination

(Vishal and Kumar, 2018; Martínez-Andújar *et al.*, 2011). ABA is synthesized at radical, blocks embryo growth, and maintains dormancy in the seed (Yang *et al.*, 2022).

b. Gibberellin (GA3): GA stimulates seed germination in many species, particularly overcoming dormancies with chilling and light requirements. The effectiveness of GA3 treatments depends on species, concentrations used, and the timing of application (Cornea-Cipciganet *al.*, 2020). Gibberellins are multifunctional phytohormones influencing germination, flowering, sex determination, and seed and fruit development (Cornea-Cipciganet *al.*, 2020). GA3 has shown significant effects on seed germination and seedling growth in various species, showcasing its potential in overcoming seed dormancy (Hemalatha and Chaudhari, 2021; Dilip *et al.*, 2017). For example, pre-soaking Anola seeds in a GA3 solution at 200 ppm resulted in the highest germination at 88 percentages (Chiranjeevi *et al.*, 2017). Maximum germination percentage (34.66%) was recorded in the sandalwood seeds treated with 300ppm GA3 (Hemalatha and Chaudhari 2021). Maharana *et al.* (2018) studied seed treatments of *Gmelinaarborea*Roxbwith GA solution and recorded enhanced germination percent and other seed germination parameters and thus improved seed vigour and uniform germination. The application of GA3 at 80 ppm for a duration of 12 hours significantly enhances both seed germination and seedling growth in Kagzi Lime, offering a viable recommendation for improved growth and yield (Dilip *et al.*, 2017). Furthermore, *Jatropha curcas* seeds exhibited a notably high germination percentage when soaked in GA3 solutions, particularly at 300 ppm for 8 hours and 400 ppm for 10 hours (Pawar *et al.*, 2010).

c. Auxin: Auxin, like IAA, IBA, NAA, and 2, 4-D, plays a role in the later stages of embryo expansion during seed germination. These auxins are commonly used in seed soaking to enhance germination and break dormancy, with NAA and IBA reported to stimulate germination (Attriet *al.*, 215; Dhiman *et al.*, 2015).

Table 1: Various Plant Growth Regulators (PGR) and their classes along with functions

PGR	Classes	Functions	Site of production
Auxin	IAA (Indole-3-acetic acid), NAA (1-Naphthaleneacetic acid), IBA (Indole-3-butyric acid), 2-4D (2,4-Dichlorophenoxyacetic acid), 4-CPA (4-Chlorophenoxyacetic acid)	(a) Involved in Apical dominance (b) stimulates Cell division and enlargement (c) Shoot and root growth (d) Plant growth movement (e) Parthenocarpy (f) Abscission (g) root induction	Meristem of apical buds, embryo of seed, young expanding leaves
Gibberellin	Gibberellic acid (GA3)	(a) Prevent genetical dwarfism (b) Regulation in bolting	Immature seeds

		and flowering (c) Production of parthenocarpic fruit (d) Germination (e) Increase flower and fruit size	
Cytokinins	Kinetin, Zeatin, Benzyl adenine	(a) Cell and organ enlargement (b) Seed germination (c) Development of bud and shoot growth (d) Flower induction (e) delay senescence	Root apex, endosperm of seeds, young fruits
Ethylene	Ethanal	(a) Ripening of fruit (b) Seedling growth and emergence (c) Abscission of leaf.	Ripe fruits, flowers and leaves and nodes of stem
Abscissic acid	Dormins, Phaseic Acid	(a) Abscission (b) Maintaining Dormancy (c) Inhibit seed germination and development (d) stimulate stomatal closure	In the roots of the plant as well as the terminal buds at the top of the plant

Mechanistic Insights into Growth Hormone Influence on Seedling Germination:

Plant growth regulators are the chemical substances, which govern all the factors of development and growth within plants (Srivastava, 2002). Seed coats impermeable to oxygen pose a challenge to germination, but hormonal treatments have proven effective in overcoming this barrier. A crucial balance between Abscissic Acid (ABA) and Gibberellic Acid (GA) regulates dormancy, with ABA synthesized at the radical to block embryo growth and maintain dormancy (Chenet *al.*, 2020; Sabaghet *al.*, 2021). Gibberellins, notably gibberellic acid (GA₃), are pivotal hormones widely employed to stimulate seed germination by breaking dormancy. GA₃, a common form utilized in seed treatments, initiates germination by promoting enzyme production and cell elongation in the embryo. It operates by overcoming seed dormancy and stimulating various enzyme production, which is helpful playing a crucial role in seed germination.

Gibberellins reduce abscissic acid levels, a hormone associated with seed dormancy. Further, to overcoming seed dormancy, signalling the seed that the conditions are conducive for germination. Simultaneously, they enhance enzyme production required for germination, promoting cell elongation and division, crucial for embryo growth and seed coat penetration. Gibberellins induce the synthesis of enzymes like amylases and proteases as stimulating enzyme. Amylases break down stored starch into sugars, providing energy for the growing embryo, while proteases assist in breaking down proteins into essential amino acids

for seedling development. The promotive effect of GA₃ on seed germination might be due to its participation in the activity of alpha-amylase which catalyses the starch conversion into simple carbohydrates and chemical energy is liberated which is used in the activation of embryo (Chenet *et al.*, 2020; Sabaghet *et al.*, 2021).

Cytokinins are primarily found in growing tissues like roots, embryos, and fruits, where cell division occurs. They regulate various functions including root development, shoot meristem formation and maintenance, organ development, seed germination, fruit development, senescence delay, and stress response (Sharma *et al.*, 2022). As a class of phytohormones, cytokinins participate in regulating plant growth, physiological activities, and yield, playing a crucial role in responding to abiotic stresses like drought, salt, and temperature variations (Li *et al.*, 2021; Prasad, 2022). Cytokinins delay leaf senescence, promote mitosis, and stimulate meristem differentiation in shoots and roots. They often work in conjunction with auxin or other hormones, such as gibberellins, to regulate plant development. For instance, they contribute to balancing apical dominance by inhibiting lateral buds while promoting bushier growth. Cytokinins, along with auxins, are key phytohormones that regulate plant growth and development from the cellular to the whole-plant level (Schaller *et al.*, 2015). Their diverse functions and interactions are best described as 'complementary' (Schaller *et al.*, 2015; Großkinsky and Petrášek, 2019).

IAA (Indole-3-acetic acid) is a ubiquitous endogenous auxin in plants, vital for root growth and seedling establishment post-germination. Auxin specifically stimulates the development of the radicle (embryonic root) and the primary root system. It plays a central role in almost every facet of root development, controlling transitions between cell division, growth, and differentiation, and establishing the root apical meristem. (Zhang *et al.*, 2022; Roychoudhry and Kepinski, 2022).

The role of ethylene in regulating seed germination and seedling growth varies depending on its interaction with other internal and external signals. Ethylene plays a crucial role in breaking seed dormancy and regulating the germination process. It contributes significantly to promoting seed germination effectively in certain species. It counteracts the effects of ABA by regulating ABA metabolism and signalling pathways. Ethylene's role in dormancy release is significant in numerous species, with effective concentrations for germination of dormant seeds ranging from 0.1 to 200 $\mu\text{L L}^{-1}$ (Corbineau *et al.*, 2014). In a complex signalling network, ethylene interacts with ABA and GAs, both of which are essential regulators of germination and dormancy. Further, ethylene promotes seed germination through antagonistic interactions with ABA and synergistic actions with GAs (Ahmed *et al.*, 2020).

Why we need Seed Pre-treatments?

Efficient nursery management involves strategic pre-treatment of seeds, particularly those with a longer germination period exceeding one week. This proactive approach not only conserves time but also minimizes resource allocation. Conducting germination tests is essential to determine the optimal seed orientation for sowing, especially when dealing with species prone to twisted shoots or roots upon emergence, such as Mahogany and Mango. Emphasizing the pre-treatment of seeds with prolonged germination periods contributes to accelerated production timelines in the nursery. This not only reduces nursery costs but also facilitates optimal planting schedules for farmers, aligning with the most favourable planting times. Some important tree species and their pre-sowing treatments listed in Table 2.

Additionally, a prudent nursery practice involves germination rate testing for seeds stored for more than one month. Sowing 100 seeds in a shaded germination bed, followed by regular watering, allows for an accurate assessment of germination rates. The outcome guides the determination of the number of seeds required per container to ensure a single plant per pot. For instance, if only 25 out of 100 seeds germinate, adjusting the sowing strategy to include four seeds per container aligns with effective resource utilization (Alamgir and Hossain, 2005; Azad *et al.*, 2010 & 2011; Tanjina *et al.*, 2014; Statwick, 2016).

Controlling germination

Creating optimal conditions for germination is paramount to successful seed development. The key factor in this process is maintaining consistent humidity around the seed. To achieve this, various coverings, such as sieved soil, sand, rice hulls, or pine needles, can be utilized to keep the seeds moist. It is crucial, however, not to add supplementary fertilizer to the substrate, as this could elevate the risk of diseases like damping-off (Sarvade *et al.*, 2014a & 2014b; Lamichhane *et al.*, 2017; Raphael *et al.*, 2023).

Germinating seedlings typically draw all the necessary nutrients from the cotyledons or, in the case of palms, from the first leaf developed inside the seed. Consequently, a substrate devoid of fertilizer, like sand, is generally deemed an ideal germinating medium. Small seedlings are particularly prone to damping-off, a condition caused by fungi present on the seed surface or in the substrate (Sarvade *et al.*, 2014a & 2014b; Lamichhane *et al.*, 2017; Raphael *et al.*, 2023).

To counteract damping-off and other potential contaminants, seed sterilization becomes imperative. One effective method involves immersing the seed in a 10% solution of bleach (1 tablespoon of bleach plus 9 tablespoons of water) for 30 minutes. Alternatively, hydrogen peroxide serves as an efficient seed sterilant with advantages such as lower toxicity than bleach. It softens the seed coat, facilitating easier entry of water and oxygen, potentially enhancing germination. Seeds can be soaked directly in the antiseptic for up to four

hours(Lamichhane *et al.*, 2017).However, it is essential to exercise caution during sterilization, as both bleach and hydrogen peroxide may require experimentation to determine the most effective solution strength and soaking time. Additionally, it is important to recognize that sterilizing the substrate may eliminate beneficial fungi, bacteria, and insects. Some of these microorganisms are crucial for breaking down organic matter or aiding plants in nutrient absorption. Striking the right balance in sterilization practices is crucial to fostering a healthy germination environment (Ochieno, 2022).

Conclusion

Seed treatments offer significant potential in overcoming seed dormancy, improving germination rates and uniformity, and reducing emergence time. The field of seed technologies is rapidly advancing within global crop protection markets (Sharma *et al.*, 2015).While numerous studies have assessed the efficacy of various methods in breaking dormancy, there is a crucial need for further research. Specifically, investigations should focus on exploring combinations of treatments and leveraging plant growth regulators for enhanced dormancy breaking outcomes. The diverse array of plant species necessitates pre-treatments to address specific dormancy types effectively. The implementation of various pre-sowing treatments not only influences the rate of seed germination but also allows for a substantial reduction in the germination period, transforming it from months to a matter of weeks. The profound understanding of dormancy and its manipulation holds significant practical implications, particularly in the mass production and propagation of high-quality seedling stocks for both indigenous and exotic trees.Ensuring the availability of planting material for afforestation and reforestation, particularly for endangered and valuable forest species, necessitates pre-sowing treatments for dormant seeds. Additionally, to meet the demand for high-quality planting material of commercially valuable species, seed pre-treatment is crucial to breaking dormancy and expediting germination to take up large-scale plantation projects. Furthermore, there is a need for research to standardize treatments aimed at breaking dormancy in responding to abiotic stresses like drought, salt, and temperature variations.

Table. 2List of tree species, seed viability and their pre-treatment

Common Name	Scientific Name	Seed pre-treatment	Time period	References
Harada	<i>Terminalia chebula</i> Retz.	Cold water	48 hrs	Hossain <i>et al.</i> , 2005
Kindal	<i>Terminalia paniculata</i> Roth	Normal Water	24 hrs	Viswanath <i>et al.</i> , 2021
Subabul	<i>Leucaena leucocephala</i> (Lam.) de Wit	hot water	12 hours	Hamad and Anwer, 2021
Ritha	<i>Sapindus mukorossi</i> Gaertn.	hot water		Attriet <i>et al.</i> , 2015
black wattle, late black wattle or green wattle	<i>Acacia mearnsii</i> De Wild.	hot water (80°C)	5 min	São José <i>et al.</i> , 2019
Australian Babul	<i>Acacia auriculiformis</i> A. Cunn. ex Benth.	hot water (80°C); 98% conc. H ₂ SO ₄	5-10 min	Azad <i>et al.</i> , 2011; Bisht and Ahlawat, 1999; Olatunji <i>et al.</i> , 2012
Mangium	<i>A. mangium</i> Willd.	hot water (80°C) for 2 min followed by cold water for 24 hrs	2 min	Bisht and Ahlawat, 1999
Bakain	<i>Melia azedarach</i> L.	hot water (80°C)	10 min	Azad <i>et al.</i> , 2010
Pyinkado	<i>Xylocarpus zeyheri</i> Roxb. Taub.	hot water (80 °C) immersion	10 min	Azad <i>et al.</i> , 2006b
White siris	<i>Albizia procera</i> (Roxb.) Benth.	Hot water 80 °C)	10 min	Ali <i>et al.</i> , 1997
Woman's tongue tree	<i>Albizia lebeck</i> (L.) Benth.	hot water 80 °C) for 10 minute	10 min	Azad <i>et al.</i> , 2006a
Gum arabic tree	<i>Acacia nilotica</i> (L.) P.J.H. Hurter & Mabb.	hot water 80 °C); sulphuric acid treatment	10 min 45 min	Nasr <i>et al.</i> , 2013
Western myall	<i>Acacia papyrocarpa</i> Benth.	hot water 100°C	5 min	Pound <i>et al.</i> , 2014
Australian blackwood	<i>Acacia melanoxylon</i> R.Br.	hot water 100°C	60 second	Burrows <i>et al.</i> , 2009
Teak	<i>Tectona grandis</i> L.f.	Alternate	(12-	Gupta <i>et al.</i> ,

		wetting and drying	12hrs)	1975; Bisht and Ahlawat, 1999; Pamei <i>et al.</i> , 2017
Chironji	<i>Buchananialanzan</i> Spreng.	Alternate wetting and drying	(24-12 hrs)	Thounaojam and Dhaduk 2020
Tamarind tree	<i>Tamarindus indica</i> L.	Scarified with sand paper		Azad <i>et al.</i> , 2013
Kusum tree	<i>Schleichera oleosa</i> Lour.	Scarified with sand paper		Tanjina <i>et al.</i> , 2014
Cow tamarind, monkey pod	<i>Albizia saman</i> (Jacq.) Merr.	Scarified with Nail clipping (one side of the seeds)		Alamgir and Hossain 2005
Vilaitikeekar, Mesquite	<i>Prosopis juliflora</i> (Sw.) DC.	manual cutting		Asif <i>et al.</i> , 2020
Shisham; North Indian rosewood	<i>Dalbergia sissoo</i> Roxb.	manual cutting		Asif <i>et al.</i> , 2020
Bombay blackwood; Indian rosewood	<i>Dalbergia latifolia</i> Roxb.	tap water	12 hours	Kumar and Chavan, 2018
Khair	<i>Acacia catechu</i> (L. f.) Willd.	hot water (80°C)	10 min	Das, 2014
Golden Shower Tree; Amaltas	<i>Cassia fistula</i> L.	Conc. H ₂ SO ₄	2 Min	Humtsoet <i>et al.</i> , 2018
Marking nut	<i>Semecarpus anacardium</i> Linn.	98% conc. H ₂ SO ₄	1 min	Panda and Hazra, 2009
Anola; Indian gooseberry	<i>Phyllanthus emblica</i> L.	GA ₃ 200ppm		Chiranjeevi <i>et al.</i> , 2017
Physic Nut, Barbados Nut	<i>Jatropha curcas</i> L.	GA ₃ 50 ppm or CaCl ₂ , Shell cracking	12 hours	Sunil <i>et al.</i> , 2016; Marcello <i>et al.</i> , 2015
West Indian lime, bartender's lime, Omani lime, or Mexican lime	<i>Citrus aurantifolia</i> (Christm.) Swingle	GA ₃ 80 ppm	12 hours	Dilip <i>et al.</i> , 2017
Gamhar; khamer	<i>Gmelina arborea</i> Roxb.	GA ₃		Maharana <i>et al.</i> , 2018
Sandalwood; Indian Chandan	<i>Santalum album</i> L.	GA ₃ @ 300ppm	12 hrs	Hemalatha and Chaudhari, 2021
Chaulmoogra; Hydnocarpus Oil Tree	<i>Hydnocarpus pentandra</i> (Buch.-Ham.) Oken	GA ₃ @ 350 ppm	overnight	Bhat <i>et al.</i> , 2020
Tamanu, oil-nut,	<i>Calophyllum inophyllum</i>	Seeds	-	Gunaga <i>et al.</i> ,

mastwood, beach calophyllum	L.	without pericarp; mechanical scarification		2011
Ber, Indian Jujube	<i>Zizyphus mauritiana</i> Lam.	GA ₃ @ 400 ppm	24 hr	Kumar <i>et al.</i> , 2020
Herb				
Chickpea milkvetch	<i>Astragalus cicer</i> L.	physical scarification	-	Statwick, 2016
Chili pepper; Cayenne Pepper Plant	<i>Capsicum annum</i> L.	Sulphuric acid	10 min	Evans and Blazich, 1999
Brazil macca	<i>Mimosa bimucronata</i> (DC.) Kuntze	hot water at 80 °C with post-treatment immersion in water at room temperature for 24 hours	1 min	Giasson <i>et al.</i> , 2019
Aswagandha	<i>Withania somnifera</i> (L.) Dunal	GA ₃ 100ppm	48 hrs	Madhumita and Ghosh, 2004
Rose root; Arctic root	<i>Rhodiola rosea</i> L.	GA ₃ 100ppm	12 or 18 hrs	Aiello and Fusani, 2008
Eastern gamagrass	<i>Tripsacum dactyloides</i> (L.) L.	GA ₃ 1000 ppm	24 hrs	Rogis <i>et al.</i> , 2004
Indian Noni, vomit fruit, awl tree	<i>Morinda citrifolia</i> L.	GA ₃ at 800 ppm	24 hours	Chandra and Sagar, 2013

References

- Ahammed GJ, Gantait S, Mitra M, Yang Y and Li X. 2020. Role of ethylene crosstalk in seed germination and early seedling development: A review. *Plant Phys. and Bioch.* 151: 124–131. <https://doi.org/10.1016/j.plaphy.2020.03.016>
- Aiello N and Fusani P. 2008. Effect of pre-chilling and gibberellic acid on seed germination of roseroot. *SementiElette.* 50:33-35.
- Alamgir M and Hossain MK. 2005. Effect of pre-sowing treatments on *Albizia procera* (Roxb.) Benth seeds and initial development of seedlings in the nursery. *J. of For. and Env.* 3: 53-60.
- Ali M, Akhter S and Kamaluddin M. 1997. Study on the bearing of water treatment on seed germination and seedling growth study of *Albizia procera*. *Ind. For.* 123(8): 764-768.

- Ali MM, Javed T, Mauro RP, Shabbir R, Afzal I and Yousef A. F. 2020. Effect of seed priming with potassium nitrate on the performance of tomato. *Agriculture*. 10(11): 1–10.
- Asif JM, Ali A, Mazhar MZ, Tanvir A, Zia B, Anmbreen I and Mahr MS. 2020. Effect of different pre-treatments on seed germination of *Prosopis juliflora* and *Dalbergia sissoo*: a step towards mutation breeding. *J. of For. Sci.* 66(2): 80-87.
- Attri V, Pant KS, Dhima R, Lal C and Sarvade S. 2015. Effect of seed size and pre-sowing treatments on growth parameters and biomass of *Sapindus mukorossi* (Gaertn.) seedlings under nursery condition. *Env. Ecol*, 33: 46-49.
- Azad MS, Musa ZA and Matin A. 2010. Effect of pre-sowing treatments on seed germination of *Melia azedarach*. *J. of For. Res.* 21(2): 193-196.
- Azad S, Manik MR, Hasan S and Matin A. 2011. Effect of different pre-sowing treatments on seed germination percentage and growth performance of *Acacia auriculiformis*. *J. of For. Res.* 22(2): 183-188.
- Azad MS, Islam MW, Matin MA and Bari MA. 2006a. Effect of pre-sowing treatment on seed germination of *Albizia lebbek* (L.) Benth. *South Asian J. of Agri.* 1 (2): 32-34.
- Azad MS, Matin MA, Islam MW and Musa MZA. 2006b. Effect of pre-sowing treatment on seed germination of *Lohakath* (*Xylocarpus parlati* Craib and Hutch.). *Khulna Univ. Stud.* 7(2): 33-36.
- Azad MS, Nahar Nand Matin MA. 2013. Effects of variation in seed sources and pre-sowing treatments on seed germination of *Tamarindus indica*: A multi-purpose tree species in Bangladesh. *For. Sci. Pract.* 15: 121-129.
- Baskin CC and Baskin JM. 2020. Breaking seed dormancy during dry storage: a useful tool or major problem for successful restoration via direct seeding? *Plants*. 9(5):636.
- Baskin JM and Baskin CC. 2004. A classification system for seed dormancy. *Seed Sci. Res.* 14, 1–16.
- Bhat V, Rajasekharan PE, Shetty GR, Nadukeri S and Ganapati M. 2020. Effect of pre sowing seed treatments on germination and seedling growth parameters of *Hydnocarpus pentandra* (Buch. Ham.) Oken. *Electr. J. of Pl. Breeding*. 11(1): 331-334.
- Bisht NS and Ahlawat SP. 1999. State Technology. SFRI, Information Bulletin No 7, Department of Environment & Forests, Government of Arunachal Pradesh. 30p.
- Bønsager BC, Shahpiri A, Finnie C and Svensson B. 2010. Proteomic and activity profiles of ascorbate–glutathione cycle enzymes in germinating barley embryo. *Phytochemistry*. 71(14-15): 1650-1656.
- Burrows GE, Virgona JM and Heady RD. 2009. Effect of boiling water, seed coat structure and provenance on the germination of *Acacia melanoxylon* seeds. *Aust. J. Botany*. 57(2):139-147.

- ÇavuşoğluK, Cadil Sand ÇavuşoğluD. 2017. Role of potassium nitrate (KNO₃) in alleviation of detrimental effects of salt stress on some physiological and cytogenetical parameters in *Allium cepa* L. *Cytologia*. 82(3): 279-286.
- Chandra KJ and Sagar DDG. 2013. A review on the propagation methods of a miracle fruit *Morinda citrifolia* L. *Ind. J. of Pl. Sci.* 2(2): 78-84.
- ChenK, LiGuo- Jun, BressanRA, Song CP, ZhuJK and Zhao Y. 2020. Abscisic acid dynamics, signaling, and functions in plants^{FA}. *J. of Integ. Pl. Biol.* 62(1): 25-54.
- Chichaghare AR, Dey S, Prasad S and Singh Y. 2020. Study of germination behaviour and growth response of *Gliricidia sepium* towards various pre-seed treatments in Vindhyan region, India. *J. of Pharm. and Phytoch.* 9(5): 673-676.
- Chiranjeevi MR, Muralidhara BM, Sneha MK, Shivan and Hongal. 2017. Effect of growth regulators and biofertilizers on germination and seedling growth of Aonla (*Emblica officinalis* Gaertn). *Int. J. Curr. Microbiol. App. Sci.* 6(12): 1320-1326.
- Corbineau F, Xia Q, Bailly C, and El-Maarouf-Bouteau H. 2014. Ethylene, a key factor in the regulation of seed dormancy. *Front. in Pl. Sci.* 5: 539. <https://doi.org/10.3389/fpls.2014.00539>
- Cornea-Cipcigan M, Pamfil D, Sisea CR and Mărgăoan R. 2020. Gibberellic acid can improve seed germination and ornamental quality of selected *Cyclamen species* Grown under short and long days. *Agronomy*. 10(4):516. <https://doi.org/10.3390/agronomy10040516>
- Cuena-Lombraña A, Porceddu M, Dettori CA and Bacchetta G. 2018. Discovering the type of seed dormancy and temperature requirements for seed germination of *Gentiana lutea* L. subsp. *lutea* (Gentianaceae). *J. of Pl. Ecol.* 11(2): 308-316.
- D'Este VR, Axelsson J, Yndgaard F and Øivind SS. 2019. Stratification, scarification and gibberellic acid treatments of garden angelica (*Angelica arcangelica*) seeds. *Seed Sci. and Tech.* 47(3): 243-247.
- Das N. 2014. The Effect of seed sources variation and pre-sowing treatments on the seed germination of *Acacia catechu* and *Elaeocarpus floribundus* Species in Bangladesh. *Int. J. of For. Res.* 2014:1-8. Article ID 984194.
- Dhiman R, Gupta NK, Attri V and Sarvade S. 2015. Effect of soaking conditions with growth regulators and nutrient solutions on germination behaviour of Ratela: a fuelwood species of Shiwalik Hills. *Envir. and Ecol.* 33(1): 41-45.
- Dilip WS, Singh D, Moharana D, Rout S and Patra SS. 2017. Influence of gibberellic acid (GA₃) on seed germination and seedling growth of Kagzi Lime. *J. of Sci. Agri.* 1: 62-69.
- Dwivedi AP. 1993. *A Textbook of Silviculture*. International Book Distributors. Dehradun. 505p.

- El-Ramady H, Abdalla N, Sári D, Ferroudj A, Muthu A, Prokisch J, Fawzy ZF, Brevik EC and Solberg SØ. 2023. Nano-farming: promising solutions for the future of the global agricultural industry. *Agronomy*. 13(6):1600. <https://doi.org/10.3390/agronomy13061600>
- Eremrena PO and Mensah SIJ. 2016. Effect of plant growth regulators and nitrogenous compounds on seed germination of pepper (*Capsicum frutescens* L). *Appl. Sci. Environ. Manage.* 20(2):242-250.
- Evans EX and Blazich PA. 1914. *Overcoming Seed Dormancy: Trees and Shrubs*. North Carolina Cooperative Extension Service. 1/99 HIL-8704. 3p.
- Finch-Savage WE and Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *New Phytol.* 171(3): 51-523.
- Flores J, González-Salvatierra Cand Jurado E. 2016. Effect of light on seed germination and seedling shape of succulent species from Mexico. *J. of Pl. Ecol.* 9(2): 174-179. <https://doi.org/10.1093/jpe/rtv046>
- Gashi B, Kasamedin A, Valbona M and Efigjeni K. 2012. Effect of gibberellic acid and potassium nitrate on seed germination of the resurrection plants *Ramonda serbica* and *Ramonda thalictroides*. *Afri. J. of Biot.* 11(20): 4537-4542.
- Gendreau E and Corbinea F. 2009. Physiological aspects of seed dormancy in woody ornamental plants. *Prop. of Ornam. Plants.* 9(3): 151-159.
- Giasson C, Baretta CRDM, Sobral LS and Baldissera R. 2019. Dormancy breaking, germination, and production of *Mimosa bimucronata* (DC.) Kuntze seedlings. *Cerne*. 25(1): 68-75.
- Goggin DE, Powles SB, Toorop PE and Steadman KJ. 2011. Dark-mediated dormancy release in stratified *Lolium rigidum* seeds is associated with higher activities of cell wall-modifying enzymes and an apparent increase in gibberellin sensitivity. *J. of Plant Phys.* 168(6): 527-533.
- Großkinsky DK and Petrášek J. 2019. Auxins and cytokinins – the dynamic duo of growth-regulating phytohormones heading for new shores. *New Phytol.* 221(3): 1187-1190. <https://doi.org/10.1111/nph.15556>
- Gunaga N, Doddabasava and Vasudeva R. 2011. Enhancement of seed germination through proper pre-sowing treatment in *Calophyllum inophyllum*, an important forest resource of the Western Ghats. *Karnataka J. Agric. Sci.* 24(3):413-414.
- Gupta BN and Pathanath PG. 1975. Factors affecting germination of teak seeds of eighteen Indian origins. *Indian For.* 101:584-587.
- Hamad SO and Anwer LS. 2021. Effect of pre-treatments and sowing depths on germination and early growth of *Leucaena leucocephala* seeds. *Zanco J. of Pure and Appl. Sci.* 33(s1): 53-61.

- Hemalatha M and Chaudhari SB. 2021. Effect of pre sowing treatments on seedgermination and its parameters in sandalwood(*Santalum album* L.). *J. of Pharmac. and Phytoch.* 10(1): 92-95.
- Hong TD and Ellis RH. 1996. A protocol to determine seed storage behaviour. *In: JMM Engels, Toll J (eds). IPGRI Technical Bulletin No. 1. International Plant Genetic Resources Institute, Rome, Italy.*62p.
- Hong TD, Linington S and Ellis RH. 1996. Seed storage behaviour: a compendium. *Handbooks for gene banks: No. 4. International Plant Genetic Resources Institute, Rome, Italy.*104p.
- Hossain MA, Arefin MK, Khan BM and Rahman MA. 2005. Effects of seed treatments on germination and seedling growth attributes of Horitaki (*Terminaliachebula* Retz.)in the nursery. *Res. J. of Agri. and Biolo. Sci.* 1(2):135-141.
- HumtsoeCT, KhareN, Rout S and Debbarma R. 2018. Influence of different pre-treatments methods on seedgermination and seedling growth performance of goldenshower tree (*Cassia fistula* L.). *J. of Plant Devel. Sci.* 10(11): 617-621.
- Iralu V, Barbhuyan H and Upadhaya K. 2019. Ecology of seed germination in threatened trees: a review. *Ener. Ecol. and Envir.* 4: 189-210.
- Kheloufi A, Mansouri L, Aziz N, Sahnoune M, Boukemiche SandAbabsa B. 2018. Breaking seed coat dormancy of six tree species. *Reforesta.* (5): 4-14.
- Kim D. 2019. Practical methods for rapid seed germination from seed coat-imposed dormancy of *Prunusyedoensis*. *Sci. Hortic.* 243: 451-456. doi: 10.1016/j.scienta.2018.08.039
- Koutouan-KontchoiMN, PhartyalSS, Rosbakh S,KouassiEK and PoschlodP. 2020. Seed dormancy and dormancy-breaking conditions of 12 West African woody species with high reforestation potential in the forest-savanna ecotone of Côte d'Ivoire. *Seed Sci. and Techn.* 48(1): 101-116.
- Kumar A, Gurjar PKS, KashyapA, MandloiV andPartetiA. 2020. Response of pre-sowing seed treatments on growth of Ber (*Zizyphus mauritiana* L.). *Int. J. of Chem. Studies.* 8(2): 2368-2371.
- Kumar U and Chavan RL. 2018. Effect of pre-sowing treatments on germination parameters of *Dalbergialatifolia*Roxb.*J. Farm Sci.* 31(1): 99-101.
- Kumar R, Kumar A, Banyal R, Kumar M, Singh A, Yadav RK, Dobhal Sand Sharma S. 2022. Seed and seedling diversity delimitation and differentiation of Indian populations of *Meliadubia* cav. *Saudi J. of Biol. Sciences.* 29(1): 489-498.<https://doi.org/10.1016/j.sjbs.2021.09.004>

- Kumar R, Mehta H, Kumar A, Bhardwaj AK, Kaushal R, Dobhal S, Gupta AK, Banyal R, Kumar M, Kumar Sand Verma K. 2021. Seed source variation affects the growth, biomass, carbon stock, and climate resilience potential: A case study of *Celtis australis* in Indian Himalayas. *Global Eco. and Cons.* 26: e01469.
<https://doi.org/10.1016/j.gecco.2021.e01469>
- Lamichhane JR, Dürr C, Schwanck AA, Robin MH, Sarthou JP, Cellier V, Messéan A and Aubertot JN. 2017. Integrated management of damping-off diseases. A review. *Agron. Sustain. Dev.* 37:10. DOI 10.1007/s13593-017-0417-y
- Leite Idalina T de A and Takaki M. 2001. Phytochrome and temperature control of seed germination in *Muntingia calabura* L. (Elaeocarpaceae). *Brazilian Arch. of Biol. and Techn- An International Journal.* 44(3): 297-302.
- Li SM, Zheng HX, Zhang XS and Sui N. 2021. Cytokinins as central regulators during plant growth and stress response. *Plant Cell Reports.* 40(2): 271–282.
<https://doi.org/10.1007/s00299-020-02612-1>
- Li W, Liu X, Khan MA and Yamaguchi S. 2005. The effect of plant growth regulators, nitric oxide, nitrate, nitrite and light on the germination of dimorphic seeds of *Suaeda salsa* under saline conditions. *J. of Plant Research.* 118: 207-214.
- Madhumita M and Ghosh P. 2004. Effect of chemicals and soaking duration on seed germination of ashwagandha (*Withania somnifera*). *J. of Inter –academicia.* 8 (2):302-304.
- Maharana R, Dobriyal MJ, Behera LK, Gunaga RP and Thakur NS. 2018. Effect of pre seed treatment and growing media on germination parameters of *Gmelina arborea* Roxb. *Indian J. of Ecol.* 45(3): 623-626.
- Marcello L, Nicla C, Luca G and Maurizio M. 2015. Effects of pre-sowing treatments on *Jatropha curcas* seed germination and seedling growth. *African J. of Agri. Research.* 10(26): 2553-2561.
- Martínez-Andújar C, Ordiz MI, Huang Z, Nonogaki M, Beachy RN and Nonogaki H. 2011. Induction of 9-cis-epoxycarotenoid dioxygenase in *Arabidopsis thaliana* seeds enhances seed dormancy. *Proc. Natl. Acad. Sci. U.S.A.* 108: 17225-17229. doi: 10.1073/pnas.1112151108
- Milberg P, Andersson Land Thompson KL. 2000. Large-seeded species are less dependent on light for germination than small-seeded ones. *Seed Sci. Research.* 10(1): 99-104.
<https://doi.org/10.1017/s0960258500000118>
- Nasr SMH, Savadkoobi SK and Ahmadi E. 2013. Effect of different seed treatments on dormancy breaking and germination in three species in arid and semi-arid lands. *For. Sci. Pract.*, 15(2): 130–136.

- Nikolaeva MG. 1977. Factors controlling the seed dormancy pattern. *In*: Khan A.A. (ed). The Physiology and Biochemistry of Seed Dormancy and Germination. Elsevier N.Y., USA: 51–74.
- Ochieno DMW. 2022. Soil sterilization eliminates beneficial microbes that provide natural pest suppression ecosystem services against *Radopholus similis* and *Fusarium Oxysporum* V5w2 in the Endosphere and Rhizosphere of tissue culture banana plants. *Front. Sustain. Food Syst.* 6:688194. doi: 10.3389/fsufs.2022.688194
- Okunlola AI, Adebayo RA and Orimogunje AD. 2011. Methods of breaking seed dormancy on germination and early seedling growth of African locust bean (*Parkia biglobosa*) (Jacq.) Benth. *Journal of Hort. and For.* 3(1): 1-6.
- Olatunji D, Maku JO and Odumefun OP. 2013. The effect of pre-treatments on the germination and early seedlings growth of *Acacia auriculiformis* Cunn. Ex. Benth. *African J. of Plant Sci.* 7(8): 325-330.
- Pamei K, Larkin A and Kumar H. 2017. Effect of different treatments on the germination parameters and seedling quality index of *Tectona grandis* (Teak) under nursery condition. *Int. J. of Chem. Studies.* 5(5): 2418-2424.
- Panda BM and Hazra S. 2009. Seedling culture of *Semecarpus anacardium* L. *Seed Sci. and Biotech.* 3(2): 54-59.
- Pawar VB, Gore RV, Patil VK and Narsude PB. 2010. Effect of gibberellic acid on seed germination and growth of *Jatropha curcas* L. *The Asian J. of Horti.* 5(2): 311-313.
- Penfield S, Josse EM, Kannangara R, Gilday AD, Halliday KJ and Graham IA. 2005. Cold and light control seed germination through the bHLH transcription factor Spatula. *Current Biol.* 15(22): 1998-2006.
- Peraza-Villarreal H, Sánchez-Coronado ME, Lindig-Cisneros R, Tinoco-Ojanguren C, Vela-zquez-Rosas N, Cámara-Cabrales L and Orozco-Segovia A. 2018. Seed priming effects on germination and seedling establishment of useful tropical trees for ecological restoration. *Tropical Cons. Science.* 2018;11. doi:10.1177/1940082918817886
- Pound LM, Ainsley PJ and Facelli JM. 2014. Dormancy-breaking and germination requirements for seeds of *Acacia papyrocarpa*, *Acacia oswaldii* and *Senna artemisioides* ssp. *coriacea*, three Australian arid-zone Fabaceae species. *Australian J. of Botany.* 62:546–557.
- Prasad R. 2022. Cytokinin and its key role to enrich the plant nutrients and growth under adverse conditions-an update. *Front. in Genetics.* 13: 883924.
<https://doi.org/10.3389/fgene.2022.883924>

- RaphaelMR, SaidiB, PhilemonMF, Mtui H andKudraA. 2023. Pre-sowing treatments to improve seed germinationand seedling growth of *Commiphoraswynnertonii*(Burr.) and *Synadeniumglaucescens* (Pax.). *Journal of Med. Plants Res.* 17(7): 225-241.
- Rashid A, Harris D, Hollington P and Ali S.2004. On-farm seed priming reduces yieldlosses of mungbean (*Vignaradiata*)associated with mungbean yellow mosaicvirus in the North West Frontier Provinceof Pakistan. *Crop Prot.* 23:1119-1124.
- Reddy MC, Bhargavi C, Rajendra MP, Indu Kand Babu H. 2022. A review on seed pre-sowing treatments in forestry applications. *Journal of Plant Deve. Sci.* 14(3): 243-251.
- Rehman HU, Maqsood S, Basra A andFarooq M. 2011. Field appraisal of seedpriming to improve the growth, yield andquality of direct seeded rice. *Turk. J. Agric.*35: 357-65.
- Rogis C, Gibson LR, Knapp AD and Horton R. 2004. Enhancing germination of astern gamagrass seed with stratification and gibberellic acid. *Crop Sci.* 44 (2): 549-552.
- Roychoudhry Sand Kepinski S. 2022. Auxin in root development. *Cold Spring Harb.Perspe. in Biol.* 14(4): a039933. <https://doi.org/10.1101/cshperspect.a039933>
- Sabagh AE, Mbarki S, Hossain A, Iqbal MA, Islam MS, Raza A, Llanes A, Reginato M, Rahman MA, Mahboob W, Singhal RK, Kumari A, Rajendran K, Wasaya A, Javed T, Shabbir R, Rahim J, Barutçular C, Habib Ur Rahman M, Raza MA, Ratnasekera D, Konuskan Ö l, Hossain MA, Meena VS, Ahmed S, Ahmad Z, Mubeen M, Singh K, Skalicky M, Brestic M, Sytar O, Karademir E, Karademir C, Erman M and Farooq M. 2021. Potential role of plant growth regulators in administering crucial processes against abiotic stresses. *Front. Agron.* 3:648694. doi: 10.3389/fagro.2021.648694
- São José JFB, Volpiano CG, Vargas LK, Hernandes MAS, Lisboa BB, SchlindweinG and Sampaio JAT. 2019. Influence of hotwater on breaking dormancy, incubation temperatureand rhizobial inoculation on germination of *Acacia mearnsii* seeds.*Australian For.* 82(3): 157-161.
- Sarvade S, Mishra HS, Kaushal R, Chaturvedi S and Tewari S. 2014b. Wheat (*Triticum aestivum* L.) yield and soil properties as influenced by different agri-silviculture systems ofTeraiRegion,Northern India. *Int. J. of Bio-resource and Stress Manag.* 5(3): 350–355.
- Sarvade S, Mishra HS, Kaushal R, Chaturvedi S, Tewari S and Jadhav TA. 2014a. Performance of wheat (*Triticum aestivum* L.) crop under different spacings of trees and fertility levels. *African J. of Agri. Res.* 9(9): 866-873.
- Sawada Y, Aoki M, Nakaminami K, Mitsuhashi W, Tatematsu K, Kushiro T, Koshiba T, Kamiya Y, Inoue Y, Nambara Eand Toyomasu T. 2008. Phytochrome and gibberellin-mediated regulation of abscisic acid metabolism during germination of photoblastic lettuce seeds. *Plant Phys.*146(3): 1386-1396. <https://doi.org/10.1104/pp.107.115162>

- Schaller GE, Bishopp A and Kieber JJ. 2015. The yin-yang of hormones: cytokinin and auxin interactions in plant development. *Plant Cell*. 27:44-63.
- Sharma A, Singh P, Kumar S, Kashyap PL, Srivastava AK, Chakdar H, Nageena Singh R, Kaushik R, Saxena AK and Sharma AK. 2015. Deciphering diversity of salt-tolerant bacilli from saline soils of eastern Indo-gangetic plains of India. *Geomicro. J.* 32(2):170-180.
- Sharma KK, Singh US, Sharma P, Kumar A and Sharma L. 2015. Seed treatments for sustainable agriculture-A review. *Journal of App. and Natu. Sci.* 7(1):521-539. <https://doi.org/10.31018/jans.v7i1.641>
- Sharma S, Kaur Pand Gaikwad K. 2022. Role of cytokinins in seed development in pulses and oilseed crops: Current status and future perspective. *Frontiers in Genetics*. 13: 940660. <https://doi.org/10.3389/fgene.2022.940660>
- Shu K, Liu XD, Xie Q and He ZH. 2016. Two faces of one seed: hormonal regulation of dormancy and germination. *Molec.Plant*. 9(1): 34-45.
- Singh SK, Pancholy A, Jindal SK and Pathak R. 2011. Effect of plant growth promoting rhizobia on seed germination and seedling traits in *Acacia Senegal*. *Ann. For. Res.* 54(2): 161-169.
- Socolowski F and Takak M. 2004. Germination of *Jacaranda mimosifolia* (D. Don - Bignoniaceae) seeds: effects of light, temperature and water stress. *Brazilian Archives of Biology and Technology- An Intern. Journal*. 47(5): 785-792.
- Srivastava LM. 2002. Plant growth and development. Hormones and environment. Academic Press. 140p. ISBN 0-12-660570-X
- Statwick 2016. Germination pre-treatments to break hard-seed dormancy in *Astragalus cicer* L. (Fabaceae). *PeerJ* 4:e2621; DOI 10.7717/peerj.2621
- Sunil N, Keshavulu K, Kumar V, Sudhir E, Yashoda and Chakrabarty SK. 2016. Germination enhancement studies in *Jatropha curcas* L. by chemical and physical methods. *Int. J. of Agri. and For.* 6(3): 132-136.
- Tanjina HGN, Hossain M, Kalimuddin BM and Alam NIS. 2014. Effect of pre sowing treatments on germination and initial growth of seedlings of kusum (*Schleichera oleosa*). *Int. J. of Usuf. Mnst*, 15(1): 3-9.
- Tapfumaneyi L, Dube P, Mavengahama S and Ngezimana W. 2023. Effect of gibberellic acid and potassium nitrate seed treatments on the emergence and seedling vigour of Amaranth and *Cleome gynandra*. *Agrosystems, Geosci. Envir.* 6: e20359. <https://doi.org/10.1002/agg2.20359>

- Thongtip A, Mosaleeyanon K, Korinsak S, Toojinda T, Darwell CT, Chutimanukul P and Chutimanukul P. 2022. Promotion of seed germination & early plant growth by KNO₃ and light spectra in *Ocimum tenuiflorum* using plant factory. *Scientific Reports*. 12: 6995.
- Thounaojam AS and Dhaduk HL. 2020. Enhancement of seed germination in Chironji (*Buchanania lanzan* Spreng) through physical and chemical treatments. *Journal of Pharmac. and Phytoc.* 9(1): 1354-1359.
- Umarani R and Vanangamudi K. 2005. Pre-storage treatments to improve viability in *Casuarina equisetifolia* seeds. *Madras Agric. J.* 92(7-9): 484-491.
- Vishal B and Kumar PP. 2018. Regulation of seed germination and abiotic stresses by gibberellins and abscisic acid. *Front. Plant Sci.* 9:838. doi: 10.3389/fpls.2018.00838
- Viswanath SC, Pillai PKC, Sinny Francis and Hrideek TK. 2021. Seed germination behaviour of *Terminalia paniculata* Roth (Combretaceae), an economically important endemic tree to peninsular India. *Current Botany*. 12: 110-114.
- Willan RL. 1985. *A Guide to Forest Seed Handling*. FAO Forestry Paper 20/2. Rome. 379p.
- Yang D, Zhao F, Zhu D, Chen X, Kong X, Wu Y, Chen M, Du J, Qu LJ and Wu Z. 2022. Progressive chromatin silencing of ABA biosynthesis genes permits seed germination in *Arabidopsis*. *Plant Cell*. 34(8):2871-2891. doi: 10.1093/plcell/koac134. PMID: 35522002; PMCID: PMC9338806.
- Yang F, Fan Y, Wu X, Cheng Y, Liu Q, Feng L, Chen J, Wang Z, Wang X, Yong T, Liu W, Liu J, Du J, Shu K and Yang W. 2018. Auxin-to-gibberellin ratio as a signal for light intensity and quality in regulating soybean growth and matter partitioning. *Front. Plant Sci.* 9: 1-13.
- Zhang Q, Gong M, Xu, X, Li H and Deng W. 2022. Roles of auxin in the growth, development, and stress tolerance of horticultural plants. *Cells*. 11(17). <https://doi.org/10.3390/cells11172761>