

Original Research Article

An Assessment of Hepatoprotective Activity of *Aloe barbadensis* Rat Model with Safety Profile Analysis

Abstract

The art or practice of employing herbs and herbal medications to preserve health and prevent, treat, Aloe lipid profiles of the *Aloe barbadensis* extract. Groups 5 and 6 showed statistically significant results ($P < 0.05$) in the renal function test. Groups 4 and 5 exhibited significant levels of creatinine. When considering the SGPT and SGOT, it was seen that the SGOT levels in groups 5 and 6 were statistically significant ($P < 0.05$). On the other hand, the SGOT levels were found to be significant in groups 6 and 7. In the instance of the lipid profile test, the outcome was found to be statistically significant ($P < 0.05$) in groups 4, 5, and 6 for the total cholesterol level. In the case of triglyceride levels, however, there is no result that can be considered statistically significant. In the case of groups 5 and 6, the HDL level exhibited statistically significant ($P < 0.05$) outcomes, but the LDL level showed statistically significant results in the case of group 6.

Keywords: *Aloebadensis*; SGPT; Herbal medicine; Triglyceride, phytopharmacology; cholesterol.

Introduction

Of all the human organs, the largest glandular organ, the liver, is responsible for the most functions. The whole blood supply of a person passes through the liver many times a day. Human metabolism relies heavily on the liver. [1]. The most common kind of liver disease, known as hepatotoxicity, is caused by medicines and ranks high among animal and human causes of mortality and disability. [2]. Liver cells are especially vulnerable to damage from things like drug and alcohol misuse, toxic chemicals, viral or parasite infections, and elevated levels of reactive oxygen species (ROS) (OH, H₂O₂, O₂). [2]. People take ascorbic acid and the tripeptide L-glutathione (L-cysteine, glycine, and L-glutamate) orally as a dietary supplement

Comment [Ma1]: Correct sentence formation

Comment [Ma2]: Put in subscript

Comment [Ma3]: Put in subscript

Comment [Ma4]: Put in subscript

because they can neutralize free radicals. Many people value them for their anti-oxidant, cleaning, and immune-boosting properties. [3]. As an alternative, they might cause allergic reactions, including dermatitis, and gastrointestinal problems like bloating, diarrhea, stomach pains, and difficulty breathing owing to airway constriction. Chronic liver disease (CLD) affects 1.5 billion people globally; in the United States, the incidence has increased by 31% in the last several years among adults aged 45 to 64 [4]. There may be therapeutic applications for some chemical components obtained from medicinal plants, according to experts in the field. Thus, scientists are continuously seeking out novel plant-based medicines, such as herbal remedies, to treat a broad variety of diseases [5]. Herbalism focuses on the practical use of therapeutic herbs, as opposed to phytotherapy, which is centered on scientific research. The vast variety of compounds found in plants, many of which have therapeutic properties, has made them an important part of human medicine for thousands of years [6]. Medicinal plants have a wide range of pharmacological and therapeutic effects due to the abundance of chemical components they contain. Some examples of these components include tannins, glycosides, alkaloids, saponins, polysaccharides, essential oils, terpenoids, resins, and plant lipids [7–9]. Plant genetic modification can achieve the targeted therapeutic impact by adjusting the amounts of chemical components. Improving the manufacture of secondary metabolites, such as alkaloids, is one use of reverse genetics [10].

Comment [Ma5]: Add years

One member of the Liliaceae family is the evergreen succulent Aloe vera (*A. barbadensis*). "Young lady" is the literal translation of Aloe vera's Sanskrit name, Kumari. Directing the flow of menstruation is a common use of aloe. It balances the monthly cycle-controlling Apana Vata, according to Ayurveda. Many more areas, including Mexico, nations around the Pacific Rim, South America, Central America, and the Caribbean. Its many medicinal and economic uses have led to its global cultivation. On a global scale, several species are farmed for significant economic gain. Indian states such as Rajasthan, Haryana, Punjab, Andhra Pradesh, Gujarat, Maharashtra, and Tamil Nadu are home to this species [11]. Polyphenol, glycoside, alkaloids, terpenoids, triterpenoids, phlobatannins, saponin, flavonoids, steroids, carbohydrate, and anthroquinone are all present [12, 13]. It has a wide range of pharmacological effects. Research has shown that this molecule has a variety of beneficial effects, including antibacterial, antidiabetic, cytotoxic, cardioprotective, bone protective, anti-inflammatory, skin protective, hepatoprotective, and neuroprotective properties [14, 15].

Comment [Ma6]: Check the spelling

Our current research aims to assess *Aloe barbadensis*'s hepatoprotective effects.

MATERIALS AND METHODS

Plant Collection and Extract Preparation

Aloe barbadensis were collected from local market of Dhaka. The material was authenticated by the University of Dhaka's Department of Pharmacy. *Aloe barbadensis* was air-dried and severely crushed. The powder was then extracted for 15 days in 50% ethanol. The extract was filtered at three-day intervals. The extracted material was dried in a rotary evaporator at a low temperature and pressure. Finally, the crude residue was subjected to the required pharmacological testing.

Drugs and Chemicals

Carbon tetrachloride (CCl₄), a well-known hepatotoxicity causing chemical, was purchased from the Sigma firm in the United States. The typical anti-oxidant medication silymarin was purchased as Livasil 140 mg from Incepta Pharmaceuticals Ltd.

Comment [Ma7]: Put in subscript

Experimental Animal Procurement, Nursing, and Grouping

A total of 100 male rats weighing between 120 and 150 grams were obtained from Jahangirnagar University in Savar, Dhaka. Each of them was housed in a climate-controlled environment (temperature 25±3°C, relative humidity 55±5%, and a 12-h light/dark cycle) at the University of Dhaka's Institute of Nutrition & Food Science (INFS). They were given a conventional food and were permitted to drink clean water. All of the animals were maintained in this habitat for at least one week prior to the research for adaptation. All experimental methods followed the recommendations of the Institutional Animals Ethics Committee (IEAC).

Animal Model Sample Size Detection

There were 100 rats in all, and they were randomly divided into ten groups of ten. In every study, the rats were randomly allocated to one of the groups. To make sure the study was genuine, we utilized 10 rats per group. Conversely, we watched the rat closely each day while it was mating. The study's control groups were both positive and negative.

Comment [Ma8]: Provide the reference

Dose Selection and Route of Administration for Respective Study

Carbon tetrachloride (CCl₄) is a common chemical agent used in laboratories to study a range of liver diseases, both acute and chronic. Trichloromethyl free radical (CCl₃·), a CYP2E1 isozyme-produced CCl₄ metabolite, reacts with cellular lipids and proteins to form trichloromethylperoxy radical, which attacks lipids on the endoplasmic reticulum membrane faster than the trichloromethyl free radical, causing lipid peroxidation and lobular necrosis. A single oral treatment of CCl₄ mixed with olive oil as a vehicle in a 1:1 ratio (3 ml/kg of rat body weight) produced hepatic damage in all animal groups except the usual control group. *Aloe barbadensis* extracts were administered to animals with hepatic injury as a post-treatment. The extract was administered orally in various quantities.

Comment [Ma9]: Check spelling

Comment [Ma10]: Check spelling

Comment [Ma11]:

Comment [Ma12]: Put subscript

Comment [Ma13]: Put subscript

Evaluation of Hepato-Protective Activity

For this experiment, 100 rats were randomly picked and equally divided into fourteen groups

Comment [Ma14]: Check once 10 or 14

Table 1: Application of treatment efficacy

Group Number	Group Specification	Treatment species	Dose treatment species (mg/kg)	Abbreviation of Groups
1	Negative Control	Physiological saline	10 ml/kg	N
2	CCl ₄ Control	N/A	N/A	A
3	CCl ₄ + S ₁₀	Silymarin	10	S ₁₀
4	CCl ₄ + AB ₂₀₀	<i>Aloe barbadensis</i>	200	AB ₂₀₀
5	CCl ₄ + AB ₄₀₀	<i>Aloe barbadensis</i>	400	AB ₄₀₀
6	CCl ₄ +AB ₆₀₀	<i>Aloe barbadensis</i>	600	AB ₆₀₀
7	S ₁₀	Silymarin	10	S ₁₀
8	AB ₂₀₀	<i>Aloe barbadensis</i>	200	AB ₂₀₀
9	AB ₄₀₀	<i>Aloe barbadensis</i>	400	AB ₄₀₀
10	AB ₆₀₀	<i>Aloe barbadensis</i>	600	AB ₆₀₀

Biological Sample Collection.

We punctured a rat's tail to obtain blood samples for measuring blood glucose levels. On the other hand, blood was drawn from the animal as soon as its heart was punctured and transferred to a microcentrifuge tube after the killing. The samples were centrifuged at 5,000 rpm for 5 minutes to create the supernatant fluid. Biochemical testing subsequently required the transfer of this fluid to an additional microcentrifuge tube. We carefully took the kidney and liver from the animal after sacrifice and cleaned them in ice-cold saline to assess their function.

Estimation of Biochemical Parameters

The blood glucose level was measured using a glucometer. Aside from the Humaluzer 3000, lipid profile, kidney, and liver function tests were performed. In addition, the gluconeogenic and glycolytic enzyme activity of kidney and liver samples was examined

Statistical analysis:

All of our findings (raw data) in terms of numerical parameters were recorded and analyzed on a broadsheet using the MS Excel application. The gathered data were subjected to descriptive statistics, with the findings reported as mean SD. To evaluate statistical significance, we used the SPSS 16 software's "One-way Anova test" to interpret inter-group heterogeneity in terms of several biological factors. The occurrences are considered statistically significant since the 'p' value was less than 0.05 ($p < 0.05$).

Comment [Ma15]: Put capital letters

Results and discussion:

Table 2. Lipid Profile Function Test result

Group no.	Group status	Kidney Function test		Liver Function test		Lipid Profile Function Test			
		Creatinine (mg/dl)	Urea	SGOT (u/l)	SGPT (u/l)	Cholesterol (mg/dl)	Triglyceride (mg/dl)	LDL (mg/dl)	HDL (mg/dl)

Comment [Ma16]: Check spelling

1	Negative Control	0.5±0.04	26.52±3.41	35.24±0.82	35.24±0.82	96.32±4.19	45.35±2.63	38.25±3.91	69.26±4.33
2	CCl ₄ Control	2.8±0.07	94.59±5.52	104.59±3.91	98.24±5.27	152.49±6.22	106.79±5.58	89.57±3.20	43.46±2.39
3	CCl ₄ + S ₁₀	1.4±0.06	42.63±4.53	72.72±3.29	70.29±4.19	109.42±5.82	58.67±4.23	60.54±4.11	57.49±3.24
4	CCl ₄ + AB ₂₀₀	2.3±0.05*	91.69±4.50	100.42±2.68	96.59±4.80	141.39±6.49*	102.31±3.17	89.19±4.53	46.48±1.26
5	CCl ₄ + AB ₄₀₀	1.9±0.05*	87.23±4.39*	96.58±4.79*	92.51±3.24*	137.25±3.26*	96.56±4.91	87.39±3.18	50.24±2.49*
6	CCl ₄ +A B ₆₀₀	1.6±0.07	84.25±5.26*	91.79±2.29*	88.57±2.08*	132.55±4.50*	90.23±2.87	84.61±3.22*	53.19±3.16*
7	S ₁₀	0.7±0.01	27.30±4.19	44.39±2.45	36.28±1.83	97.20±5.40	42.66±1.92	39.91±4.02	69.17±3.19
8	AB ₂₀₀	0.6±0.02	28.24±3.14	42.43±3.39	38.91±2.53	93.28±4.50	44.05±0.87	42.45±2.39	71.26±2.18
9	AB ₄₀₀	0.09±0.07	30.26±0.89	45.39±2.93	34.91±3.29	100.59±3.26	46.29±1.56	35.63±1.92	72.40±3.19
10	AB ₆₀₀	0.08±0.03	26.82±1.57	44.50±1.26	34.56±2.21	96.59±2.49	44.18±2.36	40.49±2.37	68.46±2.20

Note: Each value represents the mean ± SEM. (n=5). One- way ANOVA followed by Dunnett's t test. ***P<0.001, **P<0.01, *P<0.05 compared with control.

Both end-stage renal disease (ESRD) and severe chronic kidney disease (CKD) are medical conditions that may lead to cognitive impairment. The link between chronic kidney disease (CKD) and cognitive impairment is a significant problem that the public health community must address [16]. The prevalence of chronic kidney disease (CKD) in the United States went from 10% in the years 1988–1994, to 13% in the years 1999–2004, and it is possible that this trend may continue in the years to come. There was a statistically significant difference in the levels of creatinine between groups 4 and 5, whereas the findings of the kidney function test were statistically significant between groups 5 and 6. There were many studies that came up with the same results [17, 18]. There was a statistically significant relationship between the SGPT and SGOT levels; however, only the SGOT levels were significant for groups 6 and 7, in comparison to the SGPT levels. The findings of several studies [19–21] were consistent with one another. According to the findings of the lipid panel test, groups 4, 5, and 6 exhibited values that were statistically significant for total cholesterol levels. On the other hand, when it comes to triglyceride levels, there is no statistically significant influence of this kind. Researchers found statistically significant levels of HDL in cases 5 and 6, while case 6 showed statistically significant levels of LDL. In a number of studies [22–24], researchers came to the same results.

Comment [Ma17]: Cases or groups?

Conclusion:

The results of this experiment revealed that the ethanolic extract of *Aloe barbadensis* had hepatoprotective effects. The hepatoprotective effects of the ethanolic extract of *Aloe barbadensis* were determined over the course of the experiment. By lowering the buildup of lipids and liver problems, this extract helps to reduce the negative effects that a diet high in fat may have on the body. In order to determine which component of the entire extract really provides the anti-hyperlipidemic action through a screening approach, more research is necessary.

References:

1. Bhawna S, Kumar SU. Hepatoprotective activity of some indigenous plants. *Int J Pharm Tech Res.* 2009 Oct;4:1330-4.

2. Nadeem M.P.C, Dandiya P.C, K.V., Pasha M., Imran D., Balani K, Vohora S.B., Hepatoprotective activity of Solanum nigrum fruits. *Fitoterapia.*, 1997, 68, 245:51
3. FM SS, Juliana AB, Bornila M, Puja B, Nur-Neasha D, Rafat T. An Assessment of Hepato-Protective Activity of Psidium guajava Fruit Extract against Hepatic Injured Rodent Model. *Asian Journal of Medical Principles and Clinical Practice.* 2023 Oct 7;6(2):240-5.
4. Islam M, Rupak AH, Nasrin N, Chowdhury MM, Sen P, Foysal AU, Uddin MJ, Ferdous J, Tahsin MR, Aktar F, Kabir S. An Evaluation of Potential Hepato-Protective Properties of HylocereusUndatus Fruit in Experimental Rat Model. *Biomedical Journal of Scientific & Technical Research.* 2022;43(2):34405-16.
5. Baroi JA, Hossian MR, Chowdhury MM, Dolon NN, Maliha F, Rupak MA, Lima NN, Ullah MR, Tahsin R. An assessment of anti-hyperlipidemic potentialities of ethanolic extract of hemidesmusindicus in high fat induced rat model. *Asian Journal of Food Research and Nutrition.* 2023 Jul 9;2(4):323-30.
6. Mim II, Peya FY, Chowdhury MM, Khan TR, Mandal SK, Maliha F, Alam M, Rahman T, Tashin R. An evaluation of anti-diabetic activity of ethanolic extract of asparagus racemosus in alloxan induced rat model. *International Journal of Advances in Nephrology Research.* 2023 Aug 2;6(1):60-8.
7. Baroi JA, Hossian MR, Chowdhury MM, Dolon NN, Maliha F, Rupak MA, Lima NN, Ullah MR, Tahsin R. An assessment of anti-hyperlipidemic potentialities of ethanolic extract of hemidesmusindicus in high fat induced rat model. *Asian Journal of Food Research and Nutrition.* 2023;2(4):323-30.
8. Yang L, Stöckigt J. Trends for diverse production strategies of plant medicinal alkaloids. *Natural product reports* 2010;27(10):1469-1479.
9. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of Pharmacognosy and Phytochemistry.* 2013;1(6):168-182. Rupak MA, Chowdhury MM, Shurov

10. Rupak MA, Chowdhury MM, Shurovi FS, Ferdous J, Tahsin MR, Sarif S, Hasan MM, Chowdhury JA, Kabir S, Chowdhury AA, Aktar F. An Evaluation of Analgesic and Anti-Inflammatory Activity of Ethanolic Extract of *Cynodon Dactylon* on Stressed Rodent Model. *Biomedical Journal of Scientific & Technical Research*. 2022;42(3):33550-7
11. Heinemann C, Hylekama V, Van Johan ET, Janssen DB, Busscher HJ, Van Der Mei HC, Reid G (2000). Purification and characterization of a surface-binding protein from *Lactobacillus fermentum* RC-14 that inhibits adhesion of *Enterococcus faecalis* 1131. *FEMS Microbiol Lett*, 90: 177–180.
12. Mahadevi M, Vinotha P, Uma V. Anti-hyperglycemic activity of Aloe vera ethanolic extract against mouse 3T3-L1 adipocyte cell line. *International Journal of Applied Chemical and Biological Sciences*. 2020 Nov 7;1(3):58-66.
13. Bista R, Ghimire A, Subedi S. Phytochemicals and antioxidant activities of Aloe Vera (*Aloe barbadensis*). *J Nut Sci Heal Diet*. 2020 Jun;1(1):25-36.
14. Sánchez M, González-Burgos E, Iglesias I, Gómez-Serranillos MP. Pharmacological update properties of Aloe vera and its major active constituents. *Molecules*. 2020 Mar 13;25(6):1324.
15. Liu C, Cui Y, Pi F, Cheng Y, Guo Y, Qian H. Extraction, purification, structural characteristics, biological activities and pharmacological applications of acemannan, a polysaccharide from aloe vera: A review. *Molecules*. 2019 Apr 19;24(8):1554.
16. Elias MF, Elias PK, Seliger SL, Narsipur SS, Dore GA, Robbins MA. Chronic kidney disease, creatinine and cognitive functioning. *Nephrology Dialysis Transplantation*. 2009 Aug 1;24(8):2446-52.
17. Yanardağ, R., Bolkent, Ş., Özsoy- Sağan, Ö. and Karabulut- Bulan, Ö., 2002. The effects of chard (*Beta vulgaris* L. var. *ciela*) extract on the kidney tissue, serum urea and creatinine levels of diabetic rats. *Phytotherapy Research: An International Journal Devoted to*

Pharmacological and Toxicological Evaluation of Natural Product Derivatives, 16(8), pp.758-761.

18. Shimmi SC, Jahan N, Sultana N. Effect of Ashwagandha (*Withaniasomnifera*) root extract against gentamicin induced changes of serum urea and creatinine levels in rats. *Journal of Bangladesh Society of Physiologist*. 2011;6(2):84-9.
19. Yuneldi RF, Saraswati TR, Yuniwanti EY. Profile of SGPT and SGOT on male rats (*Rattus norvegicus*) hyperglycemic after giving Insulin leaf extract (*Tithoniadiversifolia*). *Biosaintifika: Journal of Biology & Biology Education*. 2018 Dec 19;10(3):519-25.
20. Mahdi C, Pratama CA, Pratiwi H. Preventive study garlic extract water (*allium sativum*) toward SGPT, SGOT, and the description of liver histopathology on rat (*Rattus norvegicus*), which were exposed by rhodamine B. *InIOP Conference Series: Materials Science and Engineering 2019 Jun 1* (Vol. 546, No. 6, p. 062015). IOP Publishing.
21. Hossain MS, Ahmed M, Islam A. Hypolipidemic and hepatoprotective effects of different fractions of methanolic extract of *Momordicacharantia* (Linn.) in alloxan induced diabetic rats. *International Journal of Pharmaceutical Sciences and Research*. 2011 Mar 1;2(3):601.
22. Yousofvand N, Soltany A. Effects of hydroalcoholic extract of dill (*Anethum graveolens*) on the serum levels of blood lipids cholesterol, triglycerides, LDL and HDL in male NMRI mice. *J PharmaceutChemBiol Sci*. 2015 May;3:114-21.
23. Nofianti T, Nurmayasari S, Priatna M, Ruswanto R, Nurfatwa M. The effect of the ethanolic extract of Asam Jawa leaf (*Tamarindus Indica* L.) in total cholesterol, triglyceride, LDL and HDL concentration on
24. Mirzaie H, Johari H, Najafian M, Kargar H. Effect of ethanol extract of root turnip (*Brassica rapa*) on changes in blood factors HDL, LDL, triglycerides and total cholesterol in hypercholesterolemic rabbits. *Advances in Environmental Biology*. 2012 Sep 1:2796-812.

UNDER PEER REVIEW

