

Floristic diversity and carbon stock of woody stands in some sacred forests in the West Cameroon region

Abstract

The vegetation of West Cameroon has been almost entirely destroyed, and the only remaining forests are mostly so-called "sacred" forests, which play a role in regulating the climate. The aim of this study was to assess the wood diversity and carbon sequestered by sacred forests in the Ndé division. The inventories were carried out on 34 plots of 900 m² each. To do this, all woody individuals with a diameter ≥ 10 cm at 1.30 m above ground level were counted. Floristic diversity was assessed using diversity indices. Allometric equations were used to estimate biomass in order to deduce the carbon stored by these forests. At the end of the surveys, 77 species belonging to **57 genera** and 32 families were recorded. *Dracaena arborea* and *Cola* sp. were the most common species (64.71% and 94.12% respectively). The density of individuals varied from 299 stems/ha to 341 stems/ha in the forests studied. Basal area varied from 42.19 m²/ha to 53.28 m²/ha. Shannon index values are low in both forests (around 2 ± 0.38 bit), indicating low specific diversity at the sites studied. Carbon values ranged from 286.89 tC/ha to 215.67 tC/ha in these sacred forests. *Cola* sp. and *Dracaena arborea* recorded the highest carbon values. Of the species recorded, **05** are vulnerable. This study reveals the importance of sacred forests for the conservation of biodiversity and their capacity to store carbon in order to contribute to the mitigation of green house gases.

Key words: Sacred forests, Diversity, Carbon stock, West Cameroon, Green house gases

No of genera ????

Checked
by
POWO

Please checked the number

UNDER PUBLICATION

Introduction

Please put Ref.

Forest ecosystems cover 31% of the planet's land area, equivalent to around 4 billion hectares (Fao, 2011). The largest fraction is in the tropical zone (1.8 billion hectares). Tropical forests are major reservoirs of biological diversity (Thatchou *et al.*, 2015). They play a major role in regulating the climate and conserving biodiversity throughout the world. They provide a wide range of ecosystem services to humans, such as firewood, timber, medicines and fruit. However, these tropical forests are under human pressure not only for agricultural needs, but also for industrial timber exploitation, to the point where they are experiencing annual deforestation rates of 0.17% (Thatchou *et al.*, 2015). This leads to massive losses in biodiversity, reduced ecosystem services and greenhouse gas emissions (Molto, 2012). In addition, some of these forests are called "sacred" and seem to be less influenced by human activities; they also contribute to the management of natural resources and the conservation of biodiversity (Wild and Mcleod, 2012). These sacred forests are found in South and South-East Asia, Latin America and Africa (Fode, 2019; Garcia *et al.*, 2006). They are the site of traditional and religious rituals. The sacred forests of sub-Saharan African countries are characterized by very high biological diversity due to their position in the intertropical zone (Hounto *et al.*, 2016). They provide non-timber forest products for food, medicinal and ritual needs, as well as regulating services (Tiokeng *et al.*, 2019; Ngougni, 2017). The natural forests of the western highlands have been almost totally destroyed in favor of agriculture. Most of the existing relic forests are sacred forests. These play a significant role in the management and conservation of biological diversity. They have long been protected by prohibitions handed down from generation to generation (Tiokeng, 2007; Kitoko *et al.*, 2020).

Delete

In Cameroon, several studies have been carried out in the sacred forests of the western highlands. For example, the work of Noumi and Tiam (2016) in the Oku sacred forest showed that *Piptadeniastrum africanum* is the most dominant species; that of Ngougni (2017) reveals that the biodiversity of the sacred forests of Bafou, Baleveng and Bamendou is dominated by species such as *Albizia* sp. and *Aleurites moluccana*. Frederic *et al.* (2021) studied the woody diversity and carbon storage rate of the sacred forests of Kouoghap in Batoufam; their study showed that the value of carbon stored by this forest is 6.40 tC. ha⁻¹ and the most abundant species are *Austranella congolensis*, *Dacryodes buttnerii*, *Vitex grandifolia* and *Polyscias fulva*. Very few studies have been carried out in the Ndé department. This is the case of the work by Mounmeme (2021), which revealed the presence of the endemic species *Dacryodes igaganga* in the sacred forests of the village of Bazou. However, eucalyptus forestry, bush fires, agrarian pressure and non-compliance with prohibitions have been identified as the main threats to certain sacred sites. All this previous work is still fragmentary and does not give a more complete picture of the vegetation of the sacred forests of the Western Highlands. Furthermore, no study has yet been carried out on the biomass and carbon sequestered by the sacred forests of Bangangté and Balengou.

Checked by POWO

Please put Ref.

Please checked

It is with this in mind that this study was initiated in the sacred forests of Bangangté and Balengou in order to contribute to knowledge of the floristic diversity and carbon stock sequestered by the sacred forests of these two localities. More specifically, the aim is to assess the diversity and structure of the woody stands in these sacred forests and to estimate the quantities of carbon sequestered by these sacred forests.

Study area

The Western Highlands of Cameroon cover an area of 13,890 km² and reach an altitude of over 2,669 m (Kuete and Dikoume, 2000). The sacred forests selected are located in the Ndé department. The Bangangté group includes the sacred forests of Mandja (5° 9'49.27 "N, 10°32'34.41 "E, altitude 1271 m) and Banekane (5° 7'23.62 "N, 10°34'15. 85 "E, Alt: 1273 m), while the Balengou area has the sacred forests of Ndepla (5° 6'4.10 "N, 10°25'9.21 "E, Alt 1298 m) and Ndiop (5° 4'48.00 "N, 10°24'32.02 "E, Alt: 1379m) (Figure 1). The climate in this department is monsoonal and sheltered in the locality of Bazou, while in the localities of Bangangté it is sub-equatorial and not very humid. Annual rainfall fluctuates around 1,600 mm in Bazou and between 1,400 and 2,500 mm in Bangangté (PCD Bangangté, 2015; PCD Bazou, 2013).




Figure 1: Location of study sites in the department of Ndé (Esther, 2020)

Vegetation sampling and floristic inventory

Put []

Thirty-four [30 m x 30 m (900 m²)] plots were established in the sacred forests located between 1264 m and 1327 m altitude. The plots were positioned so as to cover all the different habitats present. In order to better assess the variation of species in the study sites, the plots were paired by grouping. Thus, 17 plots each were sampled in the Bangangté group (1264 m) and the Balengou group (1327 m) respectively. Only trees and shrubs with a diameter at breast height (1.30 m) greater than or equal to 10 cm were considered.

Any stem bifurcated before 1.30 m above the ground was treated as a secondary stem and also counted. For species with buttresses, diameters were taken at 0.30 m above the buttresses. Some species were identified on site using well-known vegetative characters and

field guides such as **Letouzey's Manuel de botanique** forestière d'Afrique tropicale (1982). Unidentified species were collected and brought back to the Yaoundé Herbarium for identification by comparison with the herbarium samples.

Please put Ref.

Data analysis

The data were analyzed for parameters such as species richness, Shannon-Weaver index (H'), Pielou equitability index, Simpson index, basal area, density, importance value index (IVI), Sorensen index and biomass.

Please put Ref.

- Species richness (S) is the total number of species present in the population considered in a given ecosystem (**Ramade, 2003**). It is mainly expressed by the total number of species observed per unit area.

- Shannon-Weaver index (H'): This quantifies specific diversity, taking into account the proportions of each species (**Shannon, 1948**). The higher the value of this index, the greater the diversity. A value greater than 3.5 bits indicates high floristic diversity (Kent & Cooker, 1992; Ngougni, 2017). In a community of S species, the Shannon-Weaver index (H') is calculated according to the following formula:

$$H' = -\sum_{i=1}^S \frac{N_i}{N} \log_2 \left(\frac{N_i}{N} \right)$$

Please put ref.

In this formula, H' is the Shannon diversity index,

N_i: the number of individuals of a species i with i ranging from 1 to S (total number of species),

N: the total number of individuals of all species combined in the environment,

S: the species richness of the plot or biotope under consideration

Log: the logarithm to base 2.

H' = 0 if all the individuals in the stand are represented by one and the same species, or if each species in a stand is represented by a single individual; H' will be at its maximum when all the individuals are equally distributed over all the species.

-The Pielou Equitability Index (EQ): This is the ratio between the observed diversity and the maximum possible diversity, given the total number of species N. It is used to compare the diversity of two quadrats with different diversity indices (Smith & Smith, 2001).

$$EQ = \frac{H'}{\log_2 N}$$

With H': Shannon's diversity index, log₂ the logarithm to base 2

- Simpson's diversity index (1-D)

This gives the probability that two individuals selected at random from a population belong to the same species. The formula used to calculate it is :

$$D = \sum_{i=1}^S \frac{N_i(N_i-1)}{N(N-1)}$$

It is a diversity index that varies between 0 and 1. It tends towards a value of 0 to indicate minimum diversity, and a value of 1 to indicate maximum diversity (Schlaepfer, 2002).

Please put Ref.

stand structure was determined by parameters such as density, basal area and the distribution of individuals by diameter classes.

The **density** of individuals is the number of stems counted per hectare.

The **basal area** corresponds to the sum of the cross-sections at 1.30 m from the ground of the individuals in the stand. It is calculated per hectare and expressed in m²/ha. It is calculated using the following formula: $S = \pi D^2 / 4$, where S is the basal area of plot i, D: the diameter of the individual measured at 1.30 m from the ground and $\pi = 3.1416$.

Importance value (IVI) of Curtis and McIntosh (1950) which corresponds to the sum of the relative density (Dr), relative dominance (Gr) and relative frequency (Fr) of each species. It is calculated by the formula: $IVI = Dr + Gr + Fr$. It is used to characterise plant communities. The IVI provides information on the position that each taxonomic group occupies in relation to all the species within a plant community.

Please checked

Do / Gr

Please put Ref.

The conservation status of each species has been determined using the International Union for Conservation of Nature (IUCN) Red List. This is an essential indicator of the health of the world's biodiversity; it is a tool for informing and catalysing action to conserve biodiversity. The diversity indices were calculated using past 4.03 software.

The allometric equation of Djomo et al (2011) was used to estimate above-ground biomass: $AGB = \exp(-2,436 + 0,139 \times (\ln D)^2 + 0,737 \times \ln(D^2 H) + 0,279 \times \ln(\rho))$

With: AGB = Above-ground biomass; ρ = Specific density; DBH = Diameter at Breast Height.

Below-ground biomass was calculated using the allometric equation of Cairns et al (1997), which is a function of above-ground biomass:

$$BGB = \exp(-1,0587 + 0,8836 \times \ln(ABG))^{36 \times \ln(ABG)} \times \ln(ABG)$$

ABG: Above-ground biomass BGB: Below-ground biomass

The total biomass was evaluated by adding the above-ground biomass to the below-ground biomass: $TB = AGB + BGB$

The carbon stock is obtained from the recommendations of the IPCC guidelines (IPCC, 2006) by multiplying the above-ground biomass obtained by the default value of 0.47.

Results

Please check the no of genera

Floristic composition and species diversity of the sacred forests

The inventories enabled 980 individuals to be counted in 77 species, 56 genera and 32 families. The number of individuals varied from 522 to 458 (490 ± 45.3) per zone. Shannon diversity indices ranged from 2.03 ± 0.26 bits in the Balengou sacred forest to 2 ± 0.38 bits in the Bangangté sacred forest (Table 3). The Pielou equitability index is 0.88 ± 0.07 in Bangangté and 0.85 ± 0.05 in Balengou.) Basal area was $53.28 \text{ m}^2 \cdot \text{ha}^{-1}$ in the Balengou forest and $42.19 \text{ m}^2 \cdot \text{ha}^{-1}$ in the Bangangté forest.

Table 1: number of individuals, species, genera, land areas and diversity indices recorded at the study sites.

	Ni	Ne	Ng	D(tiges/ha)	ST (m ² /ha)	H'(bit)	Eq	D'
Bangangté	458	55	38	299	42,19	2±0,38	0,88±0,07	0,82±0,08
Balengou	522	48	41	341	53,28	2,03±0,26	0,85±0,05	0,82±0,06

Ni: number of individuals, Ne: number of species, Ng: number of genera, H': Shannon index, D': Simpson index, ST: basal area

Family importance values

The most important families according to family importance value (FIV) in the Bangangté sacred forest are Asparagaceae (FIV = 42.88%), Burseraceae (31.98) and Euphorbiaceae (31.88%), whereas in Balengou, Malvaceae (FIV = 66.12%), Burseraceae (34.50%) and Olacaceae (30.99%) (Table 2).

Table 2: Most important families (IVI ≥ 8%) in the two forests

	Families	Dr	Fr	Do	IVI
Bangangté	Asparagaceae	12,2 3	7,27	23,38	42,88
	Burseraceae	6,77	7,27	17,93	31,98
	Euphorbiaceae	12,8 8	12,73	6,27	31,88
	Moraceae	9,39	8,48	6,89	24,77
	Malvaceae	8,08	9,09	6,57	23,74
	Anacardiaceae	6,77	6,67	8,74	22,18
	Fabaceae	6,11	5,45	5,32	16,88
	Ericaceae	4,15	2,42	5,26	11,83
	Arécaceae	4,15	3,64	2,74	10,53
Balengou	Araliaceae	3,28	4,85	1,88	10,00
	Malvaceae	19,3 5	8,51	38,26	66,12
	Burseraceae	11,3 0	6,91	16,29	34,50
	Olacaceae	13,6 0	7,98	9,41	30,99
	Moraceae	9,00	13,30	6,17	28,47

Fabaceae	4,41	7,98	5,91	18,30
Bignoniaceae	6,70	7,45	3,09	17,24
Magnoliaceae	7,66	5,32	3,30	16,28
Euphorbiaceae	5,17	7,98	1,97	15,12
Calophyllaceae	3,83	4,26	2,23	10,31
Araliaceae	3,45	4,79	1,40	9,64

Dr: relative density; Fr: relative frequency; Do: relative dominance; IVI: species importance value.

The most common families are Euphorbiaceae (12.73%) in Bangangté and Moraceae (13.30%) in Balengou, while the most dominant are Asparagaceae (23.38) in Bangangté and Malvaceae (38.26) in Balengou.

Importance Value Index

Analysis of table 3 shows that relative density, relative frequency, relative dominance and Importance Value Index vary from one species to another.

Table 3: Importance value (IVI $\geq 8\%$) of the ten most important species in the two forests.

<i>Species</i>	Dr	Fr	Do	IVI
<i>Dracaena arborea</i> C. Koch	12,01	6,67	23,35	42,03
<i>Canarium schweinfurthii</i> Engl.	5,68	5,45	17,51	28,64
<i>Bridelia ferruginea</i> Benth.	6,11	4,24	4,87	15,23
<i>Agauria salicifolia</i> (Lam.) Oliv	4,15	2,42	5,26	11,83
<i>Ficus</i> sp.	4,37	3,03	4,20	11,60
Bangangté <i>Elaeis guineensis</i> Jacq.	4,15	3,64	2,74	10,53
<i>Polyscias fulva</i> (Hiern) Harms	3,28	4,85	1,88	10,00
<i>Acacia albida</i> Delile	4,59	3,03	1,98	9,60
<i>Macaranga spinosa</i> Müll.Aeg.	3,71	4,85	0,81	9,37
<i>Vitex doniana</i> Sweet	2,84	2,42	2,77	8,03
Balengou <i>Cola</i> spp.	19,35	38,26	8,51	66,12
<i>Strombosia glaucescens</i> Engl.	8,24	6,42	4,79	19,45

Checked by
POWO

Bangangté

Balengou

<i>Canarium schweinfurthii</i> Engl.	6,32	7,48	4,79	18,59
<i>Magnolia macrophylla</i> Michx.	7,66	3,30	5,32	16,28
<i>Aucoumea klaineana</i> Pierre	4,98	8,81	2,13	15,92
<i>Markhamia lutea</i> (Benth.) K.Schum.	6,13	2,66	6,38	15,18
<i>Albizia zygia</i> (D.C.) .J. F.Macbr.	3,07	4,65	4,79	12,51
<i>Strombosiopsis tetrandra</i> Engl.	5,36	2,98	3,19	11,54
<i>Mammea africana</i> Sabine	3,83	2,23	4,26	10,31
<i>Polyscias fulva</i> (Hiern) Harms	3,45	1,40	4,79	9,64

Dr: relative density, Fr: relative frequency, Do: relative dominance, IVI: species importance value.

Among the most important species are *Dracaena arborea* (42.03%), *Canarium schweinfurthii* (28.64%) and *Bridelia ferruginea* (15.23%) in the Bangangté sacred forest, while in the Balengou sacred forest we find *Cola* spp. (66.12%), *Strombosia glaucescens* (19.45%) and *Canarium schweinfurthii* (18.59%). Generally speaking, the species with the highest relative importance values also have significant relative abundance and relative dominance values. The most frequent species are *Dracaena arborea*, *Canarium schweinfurthii* and *Polyscias fulva* in the Bangangté sacred forest and *Cola* spp, *Aucoumea klaineana* and *Canarium schweinfurthii* in Balengou.

In
Italics

Checked
by
POWO

Diameter classes




Figure 2: Distribution of individuals (Ni) by diameter class in the two localities

Figure 2 shows the distribution of individuals by diameter class. The distribution is bell-shaped, with a peak in the 20 to 50 cm diameter class and a reduction in individuals with diameters greater than 50 cm.

Floristic similarity between sites

Comparison of the stands gives a Sorensen coefficient value of less than 50% (table 4). This result shows that, from a floristic point of view, the stands compared do not belong to the same plant community (K being less than 50) (Sorensen = 47.79%).

Table 4: Floristic similarity between the two localities

	Bangangté	Balengou
Bangangté	1	47,79
Balengou	47,79	1

Biomass and carbon stock estimates across the study area

The highest biomass values were observed in the Balengou sacred forest, with a high carbon stock of 286 TC/ha. The lowest value was obtained in the Bangangté sacred forest (Table 5).

Table 5: Total biomass and carbon stock recorded in each forest.

	ABG (tMS/ha)	BGB (tMS/ha)	BT (tMS/ha)	ST (t c/ha)
Bangangté	348,84	110,03	458,87	215,67
Balengou	463,70	146,70	610,40	286,89

ABG: above-ground biomass, BGB: below-ground biomass, BT: total biomass, ST: carbon stock.

Certain species showed the highest carbon values. these include: *Cola* spp. (123,1 tC/ha), *Aucoumea klaineana* (27,61 tC/ha), *Albizia zygia* (15,94 tC/ha), *Strombosia glaucescens* (15,016 tC/ha) à Balengou; puis de *Dracaena arborea* (56,12 tC/ha), *Canarium scheinfurthii* (45,75 tC/ha), *Agauria salicifolia* (13,03 tC/ha), *Erbroma oblongum* (11,95 tC/ha) à Bangangté. Globally, the species showing the highest carbon values differ from one forest to another.

Special-status species recorded in the forests studied

The general list of species recorded reveals the presence of 5 species of special status (Table 6). This table shows that 3 species are vulnerable. These are species such as *Khaya ivorensis*, *Lophira alata* and *Eribroma oblongum* and near-threatened (NT) species such as *Magnolia macrophylla* and *Milicia excelsa*. The rest of the species do not yet appear to have been reviewed by the International Union for Conservation of Nature.

Checked by POWO

Checked

Table 6: Species with special status

Famille

Espèces

UICN

Magnoliaceae	<i>Magnolia macrophylla</i> Michx.	NT
Malvaceae	<i>Eribroma oblongum</i> Mast.	VU
Méliaceae	<i>Khaya ivorensis</i> A. Chev.	VU
Ochnaceae	<i>Lophira alata</i> Banks ex C.F.Gaertn	VU
Urticaceae	<i>Milicia excelsa</i> (Welw.) Beth. & Hook.f.	NT

Checked by POWO

VU = Vulnerable; NT = Near Threatened

Discussion

Species richness

The results of the inventories carried out within the stands of the Bangangté and Balengou sacred forests revealed an almost equal number of species, ranging from 48 species in Balengou to 55 species in Bangangté; this reflects the low species richness of the sites. These values could be explained by the presence of these two forests in the same geographical area, with very little variation in altitude (1264 m and 1327 m respectively in Bangangté and Balengou). This species richness is higher than that obtained by Tiokeng *et al.* (2020), which varied between 21 and 26 species in certain sacred forests of the West Cameroon Highlands, and Noumi and Tagne (2016) in the sacred forests of Mount Oku (31 species). However, these species richnesses are lower than those found in the Benin sacred forest (89 species) by Hounto *et al.* (2016).

The most diverse botanical families within the two forest patches in terms of species are Moraceae (6 species), Euphorbiaceae (5 species) and Malvaceae (4 species). The high diversity within these families would indicate good regeneration of species from these different families in the study area. Similar results were recorded by Ngougni (2017) and Tiokeng (2008) in the sacred forests of Menoua and Bamboutos. The Burseraceae are also among the most important families; their presence in these sacred forests is a sign that they are the result of dense Atlantic forests (Gonmadje *et al.*, 2011; Tiokeng *et al.*, 2019). There is also a relative abundance of the *Cola* genus in Balengou. The trees in Balengou are dominated by the *Cola* sp. **genus**, which in ancient times would have been a species with high financial returns for the village.

Delete

Individual density

In Italics

Delete / Please put Ref.

The densities obtained at the different study sites were 341 stems/ha at Balengou and 299 stems/ha at Bangangté. The difference between the two could be explained by the low anthropogenic influence noted in the sacred forest of Balengou compared with that of Bangangté. These values are lower than those of Frédéric *et al.* (2018) in the Batoufam sacred forest (429 stems/ha); then of Tiokeng *et al.*, 2019 in the Mbing Mekoup sacred forest (926 individuals/ha), in the Bamendou sacred forest (878 individuals/ha) by Tiokeng *et al.* (2020) and of Ali *et al.* (2014) in the sacred forests of south-east Benin (103 individuals/ha).

Basal area

Please give Ref.

The basal area values were 53.28m²/ha and 42.19m²/ha respectively in the sacred forests of Balengou and Bangangté. The difference in basal area between the two sites suggests that the size of the diameter of the individuals and even the densities varied from one sacred forest to another. These values are close to those obtained by Mounmeni (2021) in the

sacred forests of Bazou (30.35 m²/ha); then by Tiokeng *et al.* (2020) in the sacred forest of Mbing Mekoup and Bamendou (45 m²/ha and 39 respectively), Djomo (2015) found in the east of Cameroon in the unexploited area of the Yokadouma communal forest (35.30 m²/ha) and Tajeukem *et al.* (2014) in the Gribe community forest in south-east Cameroon (49.70 m²/ha). However, these values are lower than those of Noumi and Tiam (2016) in the sacred forest of Mont Oku (205.02 m² /ha) and **Noumi (2012)** in the sacred forest of Kouoghap (90.36 m² /ha).

Please put Ref.

Diversity

The mean values of the Shannon index obtained in the Bangangté (2±0.38 bits) and Balengou (2.03±0.26 bits) sacred forests are low. This would indicate low specific diversity in the sacred forests studied. According to Kent and Coker (1992), rich communities have Shannon diversity values greater than or equal to 3.5 bits. This may be due to the low specific variation regularly observed in high-altitude zones such as the case of the present study, but also to the surface area of these forests, which is sometimes very small. These results are close to those obtained by Frederic *et al.* (2018) in the Batoufam sacred forest (2.98 bits) and by Ngougni (2017) in the Bafou sacred forest (2.53 bits).

Please put Ref.

The Piélou equitability index is 0.88 and 0.85 at Bangangté and Balengou respectively. These values are said to be optimal according to **Odum (1976)** because they are between 0.6 and 0.8. They reflect the stability of these plant communities. These results are comparable to those of Tiokeng *et al.* (2015) in the forests of the Lebialem Highlands (West Cameroon) (0.52 and 0.96) and those of Kengne *et al.* (2018) in the logged Kompia and Nkolenyeng Community Forests in East Cameroon. The Simpson diversity index values recorded in these sacred forests are 0.82 in Bangangté and 0.82 in Balengou. These results are average and reflect the average diversity of these sites. The results of this study are similar to those recorded by Mounmemi (2021) in the sacred forests of Bazou (0.94).

Vegetation structure

The diametric structures showed a high density of individuals in the 20-50 cm diameter class, which would indicate unstable regeneration over time and relative disturbance of the stand. This type of vegetation distribution was also observed by Ngounni (2017) in the sacred forests of Bafou. However, work carried out by other researchers in the sacred forests of Cameroon and Benin has instead shown an inverted J-shaped structure, where the number of individuals decreases as the diameter class increases (Tiokeng *et al.*, 2020; Frederic *et al.*, 2018; Noumi and Tagne, 2016; **Agbani *et al.*, 2018**).

Carbon stock

Delete

The carbon stocks obtained are higher in the Balengou sacred forest (286.89 tC/ha) than in the Bangangté sacred forest (215.67 t C/ha). These differences are thought to be linked to the abundance of large-diameter trees in Balengou compared with Bangangté. The carbon values recorded in this study are lower than those of Ngougni (2017), which are respectively 207.46 t C/ha in the Bafou sacred forest and 168.58 t C/ha in the Baleveng sacred forest. However, they are higher than those found by Tsalefack *et al.* (2016) in the Baleng forest reserve (West Cameroon) (120 t C/ha), by Frédéric *et al.* (2018) in the sacred forests of Batoufam (128 t C/ha) and by Ngomeni *et al.* (2021) in robusta coffee (*Coffea canephora* var. *robusta*) agroforests in Ayos (67.84±45.41 t C/ ha) and Nkongsamba (41.94±15.50 t C/ ha). In

general, the differences observed in these values could be due to the methodological approach adopted by each author, but also to the structure of each stand studied.

Conclusion

This study, carried out in the sacred forests of Balengou and Banganté, shows a low species richness in both sites. The most predominant species and families are sometimes different from one sacred forest to another. The distribution of individuals by diameter class is characteristic of a stand disturbed by human activity. The carbon stocks sequestered by these forests vary from one forest to another. These quantities also differ from one species to another. According to the International Union for Conservation of Nature (IUCN) red list of species, there are 5 species in a vulnerable situation in these sacred forests, which calls for even greater attention to be paid to them. This work shows that the sacred forests of West Cameroon, despite their relatively small surface area, are of great importance in the fight against climate change and in maintaining phytodiversity. Management strategies for these relic forests should be strengthened in order to optimize their stewardship.

References:

- Adingra O.M.M. et Kassi J.N'D. Dynamique De La Végétation De Bamo Et Stocks De Carbone Dans La Mosaïque De Végétation. European Scientific Journal June 2016 edition vol.12, No.18 ISSN: 1857 – 7881
- Anjah G. M., Nguetsop V. F., Tsombou F. M. & Njoya M. T. M. 2015. Effect of field capacity of sacred forest soils on regeneration of *Aframomum melegueta* on Western Highlands in Cameroon. *Journal of Horticulture and Forestry*. Vol. 7(5), pp. 141-148.
- Cairns M. A., Brown S., Helmer E. H. & Bamgardner G. A., 1997. Root biomass allocation in the world's upland forests, *Oecologia*. 3, 11p
- Djomo A.N., Knohl A. & Gravenhorst G., 2011. Estimations of total ecosystem carbon pools distribution and carbon biomass current annual increment of a moist tropical forest. *For. Ecol. Manage.*, 261(8), 1448-1459
- Djomo, A.N. (2015). A Structure Analysis for Ecological Management of Moist Tropical Forests. *International Journal of Forestry Research*, Article ID 161645. 12 p. <http://dx.doi.org/10.1155/2015/161645>.
- Esther L. 2020, Access to drinking water and sanitation in the Nde department (Western region - Cameroon). Doctoral thesis, University of Yaoundé I (Cameroon), <https://tel.archives-ouvertes.fr/tel-02937356>
- FAO. 2011. State of the world's forests. FAO, Rome, Italy. 164 p.
- Fode S. S. 2019 The sacred forests of Guinea: integration of ecology for the conservation of a national heritage. [Master's thesis, Paul Sabatier University - Toulouse III]. French.ffNNT: 2018(U30272ff . ff tel-02339014ff
- Frederic C., Tchatchouang L., Djomo C. C., Tajeukem V. C., Djibrilla P. and Happi J.Y 2018. Studies on Diversity, Structure and Carbon Stocks from Three Pools in the Kouoghap Sacred Forest, Hedgerows and Eucalyptus Plantations in the Batoufam Locality , West Cameroon. *Applied Ecology and Environmental Sciences* Vol. 6 6(4): 160-169,

- Garcia C., Pascal J-P., Cheppudira G. Kushalappa, (2006). The sacred forests of Kodagu in India: ecology and religion. *Wood and forests of the tropics*, No. 288(2)
- Intergovernmental Panel on Climate Change (IPCC), 2006. Guide for the national greenhouse gas inventory; agriculture, forestry and other land use. Institute for Global Environmental Strategies, Japan, 4: 446-452
- Hounto, G., Tente, B., Yabi, F., and Yabi, I. 2016. Diversity and ethnobotanical knowledge of plant species in the sacred forest of Badjamè and related areas in southwest Benin. *Scientific and Technical Review Forest and Environment of the Congo Basin-RIFFEAC*, 7, 28–36.
- Kamga Y.B., Nguetsop V.F., Momo S.M.C. and Riera B. 2018. Floristic Diversity of Woody Plants and Structure of *Garcinia Kola* Heckel Formations in the Central and Eastern Regions, Cameroon. *European Scientific Journal* 14(21): 451-484.
- Kengne O., Zapfack L., Garcia C., Noiha N. et Nkongmeneck B.-A. 2018. Diversité Floristique Et Structurale De Deux Forêts Communautaires Sous Exploitation Au Cameroun: Cas De Kompia Et Nkolenyeng. *European Scientific Journal* édition Vol.14, No.24
- Kent, M., & Coker, P. (1992). *Vegetation Description and Analysis: A Practical Approach* (pp. 167-169). New York: John Wiley and Sons.
- Kimpolo L., Ndzai F.S., & Koubouana F. (2021). Floristic Wealth and Carbon Storage of the Agricultural Zone of the Kissila Forest in the Congolese Mayombe. *European Scientific Journal*, ESJ, 17(40), 1.
- Kitoko R. N., Nzo B. J. K. N'Y. and Mukwela R. N. 2020 Floristic inventory of the sacred forest of Wuya in the province of Kongo-central in the Democratic Republic of Congo. *Africa Science* 16(1) (2020) 218 – 225
- Kuete M. & Dikoumé A. (2000). *Space, power and conflicts in the Western Highlands of Cameroon*. Space and Society, CERHT. Yaoundé University Press. 149 pp.
- Letouzey R. 1968. *Phytogeographic study of Cameroon*. *Encycl.Biol.* Ed. P. Lechevalier
- Mounmemi K, 2021, Floristic and ethnobotanical inventory mission report in the sacred forests of Bazou (western region of Cameroon), [https://sgp.undp.org/all documents/country-documents/1659-rapport -of-mission-of%E2%80%99floristic and-ethnobotanical-inventory-in-the-sacred-forests-of-bazou-\(region-of-1%E2%80%99west-cameroon\)/file.html](https://sgp.undp.org/all documents/country-documents/1659-rapport -of-mission-of%E2%80%99floristic and-ethnobotanical-inventory-in-the-sacred-forests-of-bazou-(region-of-1%E2%80%99west-cameroon)/file.html)
- Nana Richard, 2009. Perceptions and approaches for better conservation of sacred places in Bamiléké country: case of the commune of Bangangté [Thesis presented with a view to obtaining the professional master's degree, in environmental impact study at CRESA FORET-BOIS in Yaoundé-Cameroon].
- Ngomeni A. F., Nomo Bidzanga E. L., Avana M. L., Tchamba M. N. Et Chimi C. D. 2021. Carbon sequestration potential of agroforests based on robusta coffee (*Coffea canephora* var. robusta) in the production basins of Cameroon. *Int. J. Biol. Chem. Sci.* 15(6): 2652-2664, 2021

- Ngougni L., 2017 Floristic diversity, structure, biomass and uses of some sacred forests in the highlands of western Cameroon: implications in the conservation of biodiversity, Master's thesis University of Dschang 89 p
- Noumi E. & Tagne T. G. A., 2016. Floristic Inventory of Woody Species of the Oku Sacred Forest in the North-West Cameroon, Theoretical and Philosophical Approach. *International Journal of Current Research in Biosciences and Plant Biology*, Vol. 3:66-91
- Nzetchou N. 2022, Ethnobotanical study and identification of Non-Wood Forest Products of plant origin from the locality of Tonga (West Cameroon), Master's thesis University of Dschang 75p
- Piélou, E.C., 1966. Species diversity and Pattern diversity in study of ecological succession. *J. Theor. Biol.* 10, 370-383.
- PNDP, CAFER, 2015. Bangangté municipal development plan. 319 p
- PNDP, SAF Foundation, 2013. Bazou municipal development plan. 334 pp.
- Sorensen T., 1948. A method of establishing groups of equal amplitude in plant sociology based on similarity of species content. *K. Danske Vidensk. Selsk.* Vol. 5: 1-34
- Smith T. M and Smith R. L., 2001. Ecology and Field Biology. Benjamin Cummings, Pearson education, an imprint of Addison Wesley Longman, Inc. United States of America. 6th ed. 771 p
- Tajeukem, V.C., Fongzossie, F.E., Kemeuze, V.A. and Nkongmeneck, B.-A. (2014). Vegetation structure and species composition at the northern periphery of the Boumba-Bek National Park, Southeastern Cameroon. *African Study Monographs* 49 : 13- 46.
- Tchiengue B., 2004. Ecological and floristic study of the vegetation of a massif of the Cameroon line: Mont Koupe, 3rd cycle Doctoral Thesis. 238 pp.
- Thatchou B. Sonwa D. J., Ifo S. and Tiani A. M.. 2015 Deforestation and forest degradation in the Congo Basin Status report, current causes and perspectives. (CIFOR), Bogor, Indonesia C.
- Tiokeng B, Lilie Ngougni M, Nguetsop V.F., Solefack M, Zapfack L. 2020. Sacred Forests in the Western Highlands of Cameroon: Interest in the Conservation of Biodiversity. *European Scientific Journal*, ESJ, 16 (36), 1.
- Tiokeng B, Zapfack L, Nguetsop VF, Saha Z, Nchongboh CG, Douanla RN; (2019) Sacred Forests in the Western Highlands of Cameroon: Ethnobotanical role and indigenous biodiversity conservation. *Advanced Research Journal of Multidisciplinary Discoveries*. 35(11) p. 54-59
- Tiokeng B., 2008. Diversity, structure, use and local mode of conservation of some sacred forests in the Western Highlands. Master Thesis University of Dschang 120 p
- Tiokeng B., Mapongmetsem P.M., Nguetsop V.F. and Tacham W N. 2015. Floristic biodiversity and natural regeneration in the Lebialem Highlands (West Cameroon). *International Journal of Biological and Chemical Sciences* 9(1): 56-68.

Tsalefack, M., Fogaing, R., Youta H., J., Kana, C..E., Ngouanet, C. 2016. The forests of Western Cameroon between conservation and agroforestry in the perspective of REDD+. International Conference: Spatial observation for sustainable management of forests and land in Central and West Africa, GEOFORAFRI, 26-28, Abidjan, Ivory Coast, 22p

Wild, R., and McLeod. 2012. Sacred Natural Sites: Guidelines for Protected Area Managers. Gland, Switzerland, xii(IUCN), 108pp.

UNDER PEER REVIEW