

ORIGINAL RESEARCH ARTICLE

**CANOPY MANAGEMENT OF THE GREAT BANYAN TREE IN ACHARYA
JAGADISH CHANDRA BOSE INDIAN BOTANIC GARDEN, HOWRAH**

ABSTRACT

The present communication deals with the ~~prop~~-prop root training of Great Banyan Tree (GBT) of Acharya Jagadish Chandra Bose Indian Botanic Garden (AJCBIBG), Botanical Survey of India, Howrah. The GBT spans an area of 4.86 acres and features over 4400 aerial prop roots, covering a canopy circumference of 543 meters. Notably, the tree has survived three major cyclones, including the latest one, Amphan, on May 20, 2020, which significantly impacted a major portion of the tree. To address the damage caused by all cyclones and facilitate the expansion of the GBT, the AJCBIBG ~~authority~~-Authority is continuously working to train the prop roots for improvement of canopy management, using indigenous techniques developed by the staff over a decade for this age-old tree.

Keywords: AJCBIBG, Conservation, Miniature Forest, Prop root training, Walking tree, West Bengal, [India](#)

1. INTRODUCTION

The Great Banyan Tree (GBT) is a living legend, an 'emblem' of Botanical Survey of India, is situated at Division No. 23 of Acharya Jagadish Chandra Bose Indian Botanic Garden (AJCBIBG), Howrah [1, 2, 3] and it is botanically known as *Ficus benghalensis* L. (Moraceae) native to Indian subcontinent and also introduced into many sub-tropical areas due to its multiple uses [4]. Every year, it attracts lakhs of visitors, including national and international delegates ~~in~~ to the AJCBIBG. The tree is also known as "Walking Tree", "Miniature Forest", and "Immortal Tree" and the garden is also considered as one of the best landscaped gardens of the world [95].

The area occupied by the tree is about 19667 sq. m. (4.86 acres) ~~at present~~. The present crown of the tree has a circumference of 543 m and highest branch rises to 26 m. Presently, it has 4412 aerial roots reaching down to the ground as prop roots (Plate 1). In 1925, the main trunk was removed due to a fungal attack; at that time, it attained a girth of 16.5 m at 1.7 m height from the ground [3]. It is also worth mentioning that the temperature under the canopy area ~~is~~ always ~~varies (low)~~ by 2 to 5⁰C ~~degrees Celsius than the temperature from~~ of the outer area.

Formatted: Font color: Red

Formatted: Superscript

There is no clear history about the origin of this plant in the garden but scrutiny of literature [1, 5, 6, 7] reveals that the tree existed on a date palm tree (*Phoenix dactylifera* L.) in 1782 before the foundation of the garden in 1787 by Colonel Robert Kyd. By this way, it is





Plate 1: a-b. A view of Great banyan tree; c. Satellite image of GBT; d & f. Inside view showing prop roots; e. Canopy view from inside.

assumed that the tree is more than 250 years old. The tree also survived and its clonal colonies remained healthy after the three cyclones of 1864, 1867 and 2020. The GBT is standing tall ~~and~~ as senior most citizen ~~of the garden~~ thereby enhancing the beauty and serenity of ~~the~~ garden. With this significance of GBT and to increase massive dimension, canopy circumference and increased number of prop roots, continuous training is warranted to prop roots to give mechanical strength of the tree in desirable area. ~~Hence, in~~ The present communication ~~deals with a~~ methodology ~~that~~ has been standardized for prop root training of GBT.

Formatted: Indent: First line: 0"

Formatted: Font color: Red

2. Importance of Prop Roots Training

Prop roots, synonymously known as columnar roots in *Ficus* spp., are a type of adventitious roots that grow above the ground from the stem of a plant. They reach down and anchor the plant firmly into the soil. These roots play a vital role in stabilizing and providing mechanical support to the tall or heavy aerial branches and the canopy of the plants [89]. These prop roots help to keep the tree balanced and prevent it from toppling over, especially in strong winds or heavy rain. So, prop roots are trained as it will help to divert the different prop roots at desired positions strengthening of mechanical support to the plant. If it is not properly trained, the prop roots will come at haphazard positions and also get ~~damages~~ ~~damaged~~ naturally or even physically by the outsiders ~~people~~. Prop roots can also help in gaining better canopy coverage area through nutrient absorption. Since they are in direct contact with the soil, they can take up water and minerals for the plant, in some cases, they even form additional root systems, making the plant ~~even~~ more efficient in ~~gathering~~ nutrients ~~absorption~~.

As it has fungus attack history, new prop roots will get physical damage and ~~their~~ tips get easily dried in ~~hot~~ summer, and they become very prone to environmental hazards. ~~So, it~~ is, ~~therefore~~, very essential to protect the prop roots in a right way ~~through by~~ caging the roots with bamboos. ~~So, in~~ a nutshell, prop roots are like nature's way of giving ~~the~~ plants a helping hand to stand tall and strong while also ensuring ~~they have~~ a constant supply of essential ~~nutrients and water, but~~ training ~~those of~~ prop roots is like giving ~~the~~ right direction to the natural growth of the tree.

3. MATERIALS AND METHODS

Materials required for prop root training: Bamboo shoots [Shelf-life durability, hardening and hollow core structure bamboo (*Dendrocalamus* spp.) of 3-5 ft. of length shoots were used to give mechanical support]; leaf moulds (Natural leaf moulds collected as organic litter from the GBT were used as ~~the~~ manure along with soil and sand inside the hollow bamboo structure); coconut rope; pruning knife; coir; soil; water; funnel etc.

Techniques:

The step-wise standardized methodology is discussed below for ~~the~~ establishment of prop roots (Plates 2 & 3).

- First bamboo shoots were cut vertically into desired length (3-5 ft.). Then it was also cut longitudinally into 2 to 4 halves and the internodes were removed to make the outer and inner surface smooth for better handling. A groove was also cut at the edge on larger half of the shoot.
- Now coconut rope was tied at two to three places on the bamboo ~~shoots~~ to hold the halves and to make a tube-like structure.
- A manure mixture containing leaf moulds, soil and sand in specified ratio was prepared. The proportion of each material was indigenously ~~decided-prepared~~ by the experienced staff of AJCBIBG. Mention main constituents and proportion used
- Roots were checked from ~~the~~ hanging branches of the tree for their readiness for training. whether they are ready for insertion or not. After checking, the healthy roots were selected depending upon its nature, growth, size, etc.
- Secondary and tertiary branching of the prop roots were removed by cutting through a knife to insert into the hollow bamboo tube. This step is necessary It is performed for easy penetration, direction as well as considering the robust growth of one root over the multiple ones. [It is noted that, if multiple branches of the prop roots were selected then the tip of the root become prone to infection and gets easily dried, so to prevent this, only one healthy primary branched prop root is selected].
- One side of the bamboo tube was blocked with banyan leaves; so that it will hold the manure mixture and provide better growth medium and drainage. It is done keeping in view that if water is logged then growth of the root can be hampered and manure mixture become soggy ~~so it which may will~~ deprive the root from getting ~~the~~ necessary oxygen and dilute the nutrients in the manure mixture.

Formatted: Indent: Left: 0", First line: 0"

Formatted: Font color: Red

Formatted: Highlight

- ~~New~~ Half of the bamboo hollow tube was filled with the manure mixture. It is finally ready for insertion of the desired prop root.
- A selected healthy prop root was inserted within the hollow bamboo tube. Then on top of it, manure mixture is poured and it was shaken properly to make it tight inside without any gap.

UNDER PEER REVIEW



Plate 2: a. Selected bamboos shoots; b. Cutting of selected bamboo shoots into desired length; c. Two halves of bamboo joined together to make a hollow tube-like structure; d. Preparation of manure mixture; e. Removal of secondary and tertiary branches of prop root with knife; f. One side of tube blocked with banyan leaves.




Plate 3: a. Filling up of bamboo tube with manure mixture; b. Insertion of selected prop root inside bamboo tube; c. Dripping of water through the bamboo tube; d. Tying-up of prop root with the grooved part of the tube; e. Prop roots emerges from the lower side of the hanging tube; f. A view of a branch showing trained prop roots.

- Water was poured ~~inside it~~ from the top of the tube until it dripped down from the other side of the tube. It is done to check whether drainage is working properly or not. If it is not dripping, then the tube was shaken until adjusted. Prop root is now set for growth inside the tube.
- The groove that was cut on the upper side of the bamboo shoot was now tied up with the inserted prop root so that it could not move. ~~After that~~ Then the whole bamboo tube was tied at two to three places for better holding and ~~also to ensure~~ that the whole structure remains intact and withstands ~~several~~ environmental hazards. ~~Now,~~ The whole bamboo along with its inserted prop root is attached with the main branch through a coconut rope in a hanging condition.
- Once ~~it was observed that~~ the tip of the growing root comes from other side of the tube, if it ~~doesn't~~ does not touch the ground, then the whole process was repeated depending upon the distance of the branch from the top to the ground, the procedure may be repeated ~~repetition can be~~ twice, thrice and so on.
- Before touching the soil, the bamboo pole was slanted for providing better mechanical strength and to increase the ~~more~~ canopy area.


4. RESULTS AND DISCUSSIONS

The canopy circumference is directly ~~depending~~ dependent on the proper establishment of prop roots of GBT. Hence, ~~the~~ continuous attention has to be paid to GBT to increase number of prop roots. ~~While reviewing literature,~~ it has been observed that canopy circumference of this tree increases throughout its age. Chakraverty et al. [78] reported that the canopy circumference was 240 m in 1850, which increased to 377 m in 1900, in 1986 it reached 420 m. The latest recorded data for canopy circumference of the tree is 543 m in October, 2023. The detailed changes in canopy circumference are presented below (~~Plate-~~ Fig. 1).



PlateFigure41: Changes in Canopy Circumference (m) during 1850-2023

The changes in number of prop root shows variation from 1850 to 2023. In 1850, it was only 89 prop roots, which increased to 464 prop roots in 1900 and 1825 prop roots in 1986 [7] and as per recent data in October, 2023, the prop root count was 4412. The detailed changes in number of prop roots are presented below ([Plate 5Fig. 2](#)).



PlateFigure52: Changes in number of Prop roots from 1850 to 2023

Throughout its life, the tree shows great variation in its highest branch due to ageing and different natural calamities. In 1850, it was 20.5 m, which increased up to 30.5 m in 1968 (Highest till date) and 24.5 m in 1986 [7]. Recent survey shows the highest branch of the tree from ground is 26 m in October, 2023. The changes in height of the tallest branch are presented below ([Plate 6Fig. 3](#)).

Formatted: Justified




Plate 6 Figure 3: Changes in height(m) of the tallest branch during 1850-2023

5. CONCLUSION

The Great Banyan Tree (GBT) at AJCBIBG stands as a botanical marvel, recognized globally for its monumental size and significance. With its sprawling canopy covering 19667 sq. m. and boasting 4412 aerial prop roots, the GBT symbolizes resilience and longevity. Despite facing various natural challenges and even losing its main trunk to a fungal attack in 1925, the tree has continued to thrive, standing tall as a testament to nature's enduring strength. ~~C~~The continuous efforts of the garden staff to manually train the prop roots ~~have~~ ~~has~~played a crucial role in maintaining and enhancing the tree's structural integrity. By employing indigenous techniques, such as the innovative method standardized for prop root training, detailed in this communication, the garden authorities have successfully improved canopy management and provided essential mechanical support ~~for to~~the GBT. Moreover, the meticulous documentation of the tree's growth over the years reflects its remarkable resilience and adaptability.

From its humble beginnings as a sapling on a Date Palm tree in the late 18th century to its current stature as a towering presence in the garden, the GBT continues to captivate visitors from around the world. ~~Overall, t~~The success of ~~the~~ prop root training methodology and the continued growth and expansion of the GBT underscore its significance as a botanical treasure and a living testament to the power of nature's resilience and human ingenuity in conservation efforts.

Formatted: Superscript

REFERENCES

1. Chowdhury, H. J. and Pandey, D. S. 2007. Plants of Indian Botanic Garden. Botanical Survey of India, Kolkata.
2. Sarkar, G. 2012. I am the Identity and I am the Emblem (The Great Banyan Tree speaks). ENVIS 17 (2), 2-3.
3. Anonymous, 2023. Acharya Jagadish Chandra Bose Indian Botanic Garden. EIACP, Botanical Survey of India, Kolkata.
4. Patrut, R.T., Garg, A., Patrut, A., Woodborne, S., Rakosy, L. and Ratiu, I.A. 2023. Radiocarbon analysis of the Indian banyan (*Ficus benghalensis* L.) at Narora. Current Science 124(10), 1175-1180. doi: 10.18520/cs/v124/i10/1175-1180.
5. Barman, R. D., Saha, T., Swamy, J. and Singh, D. 2023. Ex-situ Conservation of Palms with Special Reference to Endemic and IUCN Red List Species in Acharya Jagadish Chandra Bose Indian Botanic Garden, Howrah, India. Journal of Experimental Agriculture International, 45(12), 223–234. <https://doi.org/10.9734/jeai/2023/v45i122283>
6. Hooker, J.D. 1841. Himalayan Journals. Ward, Lock, Bowden & Co. London.
67. Benthall, A.P. 1946. The Trees of Calcutta and its Neighbourhood. Thacker Spink & Co Ltd. London.
78. Chakraverty, R. K. and Mukhopadhyay, D. P. 1987. The Great Banyan Tree. Nelumbo 29(1-4), 59-70.
89. Ramakrishna, T.M., Poornima, S. and Vimala, R. 2015. A Report on Different Types of Aerial Roots. World Journal of Pharmacy and Pharmaceutical Sciences 4 (06), 794-797.
9. ~~Barman, R. D., Saha, T., Swamy, J. and Singh, D. 2023. Ex situ Conservation of Palms with Special Reference to Endemic and IUCN Red List Species in Acharya Jagadish Chandra Bose Indian Botanic Garden, Howrah, India. Journal of Experimental Agriculture International, 45(12), 223–234. <https://doi.org/10.9734/jeai/2023/v45i122283>~~
10. Bar-Ness, Y.D. 2013. Conservation of Indian Heritage Trees. INTACH.
11. Singh, J. N. 1979. The Great Banyan Tree (*Ficus benghalensis* L.), Indian Botanic Garden, Howrah-The Periodical Variation in its Soil Fertility Status. Nelumbo 21 (1-4), 22-27.

Formatted: Font: Italic

Formatted: Indent: Left: 0", Hanging: 0.49"

UNDER PEER REVIEW