

## INTERGRATED FISH ANDPOULTRY PRODUCTION COMBINED

**Comment [H1]:** This title needs to be improved. The context or the dependent variable is not coming out. It was out of reading the abstract that I was able to get the context of the research.

### Abstract

*A locally improvised poultry house built with pure local material called bamboo and covered with dried treated grasses (roof) known as sana which housed 300 laying birds was built near 20m X 20m earthen fish pond with an underground spring water as source of intake with a flow through system and stocked with an improved catfish *clariasgeripinus* juveniles 2000 in number. The floor of the poultry house was cemented to prevent leakages of water and permanently covered with thick nylon white rubber and layer were placed at high level above the floor on top of bamboo tree having gaps in between them. The usage of these gaps created was to allow layers (chicken) dropping directly on the rubber which eventually served as breeding places for maggot. The maggot was collected washed thoroughly and processed to serve as feed to the fishes. The processed maggot reduces about 40% of total cost of feed in raising the fishes. The birds produced 70 pieces of eggs per day 7½ crates and a crate is sold for N2,700 per crate. The fishes were harvested after six (6) months of stocking with a total piece of 1,680 with an average of 800g (1344kg) and is sold at N1500 per kg. The revenue realized took care of both capital and recurrent cost with a working force of three (3) people. The experiment contribute significantly ( $P < 0.05$ ) in reducing the level of poverty and also contribute to food security, transformation agenda of Federal government of Nigeria.*

**Comment [H2]:** There should be a first sentence telling the reader the aim of the research paper.

**Comment [H3]:** The author should have mentioned the p-value so that it is justified that it was less than 0.05.

**KEYWORDS:** Maggot, bamboo, *Clariasgeripinus*, Juvenile and treated grasses.

### 1.0 Introduction

Poultry is the foremost developing segment of the Agricultural sector in Nigeria, Africa and indeed globally. The increase in poultry production has enhanced the per capita availability of eggs and per kilogram of poultry meat per annum (Ajadi et al 2019).

In Nigeria, the average daily consumption of 54g protein with 6.5g from animal source falls below the recommended daily protein of 86g and 8.4g of protein from animal source (Abu, et.al., 2008) the reason being the exorbitant cost of fish, chicken and its products owing to high cost of feed which accounts for 60% - 80% total cost of production (Abu, 2008).

The integrated poultry-fish based farming provides enormous opportunities for improvisation of some of the necessary inputs especially feeds so that cost is saved and profit is improved as supplementary income for men and women empowerment and livelihood security to rural families.

The farming system also offers opportunities to these resource-poor families at meagre financial investments and addresses the issues of women empowerment and livelihood security at the grass root level. To encourage poultry entrepreneurship among women in rural households so as to support backyard or intensive broiler production and hatchery/small poultry feed mills in the rural areas and villages.

The basic reasons behind poor growth of poultry production in most of our villages and rural areas is the accessibility to basic poultry needs like good feed mills, feed ingredients and veterinary clinics and most importantly movement of youths from rural area to cities. (Lawal et al., 2013)

This type of system will bring a kind of recycling in Agriculture between fishery and poultry will create jobs for the rural people with minimum capital requirement to take off there by reducing poverty and also contribute to food security, transformation agenda of Federal government of Nigeria.

### 1.1 Justification

- i-Poultry **feaces** is produced in large quantity in every poultry farm and this can cause nuisance
- ii-Majority of the rural people are jobless
- iii-one of the major **inputes** in poultry production is land, which is readily available in the rural area

**Comment [H4]:** Replace this with manure

**Comment [H5]:** Correct this. Inputs not Inputes

### 1.2 Objective of the study

- i-Build both the poultry and fish pond on each other
- ii-Feed the fishes with poultry dropping

**Comment [H6]:** The objective is not written well. Let what you want to achieve come out well from the objective. It should be SMART.

## 2 Material and Methods

Site of experiment: The experiment was carried out **in** the Agricultural garden **if** the department of Agricultural Technology

**Comment [H7]:** Replace with at

**Comment [H8]:** Correct this. Of not if

Construction: A locally improvised poultry house built with pure local material called bamboo and covered with dried treated grasses (roof) known as sana which housed 100 laying birds was built near 20m X 20m earthen fish pond with an underground spring water as source of intake with a flow through system and stocked with an improved catfish *clariasgeripinus* juveniles 2000 in number. The floor of the poultry house was cemented to prevent leakages of water and permanently covered with thick nylon white rubber and layer were placed at high level above the floor on top of bamboo tree having gaps in between them. The usage of these gaps created was to allow layers (chicken) dropping directly on the rubber which eventually served as breeding places for maggot.

Sources of the juvenile : The juvenile was purchased from Ministry of Agriculture, fisheries section along yidi road in Ilorin.

Comment [H9]: Correct this to of

Sources of the layers : The point of layer was purchased from, Affcom farm Kulende junction , sango in Ilorin, the point of lay was allowed to start laying properly.

Comment [H10]: Correct this to junction

Sources of feed ingredients : The few feed ingredients that was used was purchased from Gbemisolafeedmill house, Offa garage, Ilorin

Egg collection : egg were collected everyday into crates directly around 4pm, no sorting was done and the eggs was sold the way it is picked.

Statistics analysis : Data like feed intake, number of eggs collected, equivalent amount of money for the eggs collected, all these was subjected to complete randomized design.

### 3 Results and Discussion

Every three hundred laying (200) birds will be fed by a bag of feed which is 25kg but the 300 birds that was used in the experiment, 1½ a bag will be fed to them on a daily basis, the table 1 below shows the amount of feed consumed when the maggot is supplemented and this accounts for about 40% reduction of the feed they were supposed to consumed a day and this has drastically reduced the amount of money for keeping the laying birds while the production rate is well above 65% or 70% in some situation, The fishes were harvested after six (6) months of stocking with a total piece of 1,680 with an average of 500g (1344kg) and is sold at N1500 per kg. The revenue realized took care of both capital and recurrent cost with a working force of three (3) people.

**Table 1 : Performance characteristics of poultry and Fish combined production**

Months	1	2	3	4	5	6
Feed intake	7.5	7.9	7.5	7.0	7.0	7.1
Egg prod	2130	2100	2070	2130	2190	2130
No of crates	2.3	2.3	2.3	2.3	2.4	2.3
Equiv. amt of money (N)	106,500	105,500	103,500	106,500	1680	106,500
SEM	0.1	0.4	0.1	0.1	0.2	0.1

#### 4 Conclusion and Recommendation

##### Conclusion

i- The cost of feed has been reduced by 40%

ii- Egg production rate is about 70%

iii- Sales of fishes is another source of income

**Comment [H11]:** Conclusion should be continuous prose and not numbered.

##### Recommendations

i- This system of farming could be adopted even in large scale production

ii- Maggot could also be used in poultry feed production because it is a good source of methionine.

**Comment [H12]:** Should be continuous prose and not numbered

##### References

Abu A.O Onifade, A.Abanikanda A.A, and Obineye, R.I. (2008): Status and Promotional Strategies for Rabbit Production in Nigeria. 9<sup>th</sup> World Rabbits Congress Varona, Italy

Adewoye, S.O. and Omotosho, J.S. (1997): Nutrient Composition of some freshwater fishes in Nigeria. *Biosci. Res. Commun.* 11 (4) 333-336.

Ajad M.I, Lawal W.S, Banjoko I;K and Ahmed S.A (2019) Comparative Proximate And Mineral Analysis Of Smoked Dried And Solar-Dried African Mud Catfish

Akinneye, J.O, Amoo, I.A, Arannilewa, S.T.(2007): Effect of drying method on the nutritional composition of three species of (*Bonga spp*, *Sardinella spp*. And *Heterotisniloticus*. J. fish. Int; 2; 99-103 biology journal of North America, 5(3): 109-117.

Bolawaet *al.* (2011). Proximate Composition Properties of Different Fish Species Obtained from Lagos,Nigeria. *Internet Journal of Food Safety*, **13**, 342-344.

Burgress, G.H.O (1975): "Increasing the direct consumption of fish. In: WW Pirie (Edu). Food Protein Sources. *International Biological Programme*4. Cambridge University Press, Cambridge, pp 187-200. Calabar, p: 183.

Cui, Y. and Wootton, R. J. (2011): Effects of ration, temperature and body size on the body composition, energy content and condition of the minnow, *Phoxinusphoxinus*(L). *Journal of Fish Biology*, **32**: 749-764.

Dawson, A.S. & Griman, A.S. (1980). Quantitative seasonal changes in the protein, lipid and energy content of carcass, ovaries and liver of adult plaice (*Pleuronectesplatena*) *Fisheries* 48: 297-302.

Lawal W.S, Oladunjoye I.O, Banjokol;K and Ahmed S.A 2013Performance characteristics, sensory evaluation of broiler birds fed *Jatropha caucas* (A tertiary education trust fund sponsored project of Kwara state Polytechnic, Ilorin Nigeria )