

MORPHOLOGICAL AND BIOCHEMICAL CHARACTERIZATION OF PHYLLOSHERE BACTERIA

Abstract

Phyllosphere bacteria, residing on the surfaces of plant leaves, play crucial roles in plant health and environmental processes. Understanding their morphological and biochemical characteristics is fundamental for comprehending their ecological functions. This study aimed to investigate the morphological and biochemical traits of phyllosphere bacteria isolated from cereal plant species. Isolates were collected from the leaves of different cereal plants and subjected to morphological and biochemical assays were performed to characterize their metabolic capabilities. Results revealed a diverse array of morphotypes, including cocci and bacilli, indicating the presence of varied bacterial species in the phyllosphere. Furthermore, biochemical analyses highlighted metabolic versatility among isolates, with varying abilities to utilize different carbon sources and produce enzymes. These findings underscore the heterogeneity and metabolic diversity of phyllosphere bacteria, suggesting their potential roles in nutrient cycling, plant growth promotion and disease suppression. Further exploration of these characteristics could elucidate the mechanisms underlying plant-microbe interactions in the phyllosphere and inform strategies for sustainable agriculture and environmental management.

Key words: Phyllosphere, Gram staining, Starch Hydrolysis, Catalase test

Introduction

The phyllosphere, the aerial parts of plants, represents a dynamic ecological niche teeming with microbial life, particularly bacteria. These phyllosphere bacteria, by residing on leaf surfaces, play pivotal roles in plant health, growth and ecological processes. Understanding the morphological and biochemical characteristics of these bacteria is essential for elucidating their ecological functions and their potential applications in agriculture and environmental management (Vorholt., 2012).

The phyllosphere harbors a diverse microbial community, including bacteria, fungi, and viruses, which interact with each other and with the plant host. Phyllosphere bacteria colonize leaf surfaces through various mechanisms, including aerial dispersal, rain splash, and insect vectors. Once established, they engage in complex interactions with the plant and

other microorganisms, influencing nutrient cycling, plant growth, and defense mechanisms (Bulgarelli *et al.*, 2013).

Studies have shown that phyllosphere bacteria can promote plant growth through mechanisms such as nitrogen fixation, phosphate solubilization, and production of phytohormones. Additionally, they can enhance plant defense against pathogens by competing for space and nutrients, producing antimicrobial compounds, and inducing systemic resistance. Moreover, phyllosphere bacteria contribute to environmental processes such as carbon cycling and atmospheric nitrogen fixation.

Phyllosphere bacteria exhibit remarkable morphological diversity, reflecting the multitude of species occupying this niche. Microscopic examination of leaf samples reveals a wide range of morphotypes, including cocci, bacilli, spirilla, and filamentous forms. This diversity suggests the presence of numerous bacterial taxa adapted to the specific microenvironments offered by different plant species and leaf structures.

The morphology of phyllosphere bacteria is influenced by various factors, including nutrient availability, humidity, UV radiation, and interactions with other microorganisms. For instance, certain bacteria may form aggregates or biofilms on leaf surfaces, providing protection against desiccation and predation. Moreover, the morphology of phyllosphere bacteria can undergo changes in response to environmental stressors, such as nutrient depletion or exposure to antimicrobial compounds.

In addition to morphological diversity, phyllosphere bacteria exhibit a wide range of biochemical characteristics, reflecting their metabolic versatility and adaptation to diverse ecological niches. Biochemical assays provide insights into the metabolic capabilities of these bacteria, including their ability to utilize different carbon and nitrogen sources, produce enzymes, and tolerate environmental stresses (Leveau and Lindow., 2001).

Phyllosphere bacteria are known to employ various metabolic pathways to thrive on leaf surfaces. For instance, they may utilize plant-derived organic compounds, such as sugars and organic acids, as carbon sources for growth. Additionally, some bacteria possess enzymes capable of degrading complex polymers present in leaf litter, contributing to the decomposition of plant material and nutrient recycling.

Materials and Methods

Morphological Characterization phyllosphere isolates

The morphological and biochemical characteristics of phyllosphere bacteria provide valuable insights into their taxonomy, physiology, and potential ecological functions within plant ecosystems. Understanding these characteristics is essential for accurate identification and classification of bacterial isolates, as well as for assessing their roles in plant health and development. In this section, we outline the procedures employed to investigate the morphological and biochemical properties of phyllosphere bacteria isolated from cereal samples.

All the forty isolates were checked for their purity and then studied for the colony morphology and pigmentation. The cell shape and Gram reaction were also recorded as per the standard procedures given by Barthalomew and Mittewar (1950).

3.5.1 Shape

Simple staining allowed for the observation of the cell shape. Every isolate was made into a smear, allowed to air dry and then stained for 30 seconds with crystal violet. After that, the stained smear was cleaned, allowed to air dry and examined under an oil immersion compound microscope (Barthalomew and Mittewar, 1950).

3.5.2 Gram's staining

A sterile distilled water drop was positioned in the centre of the slide, and an inoculum loopful from a culture that was 24 hours old was taken and combined with it. The inoculation needle tip was used to spread the suspension out on a slide to create a thin suspension. The smear was fixed by making three or four passes over the flame on the lower side of the slide for gentle heating after being left to dry in the air. Each slide was quickly cleaned with distilled water after 30 seconds of crystal violet application. The slides were drained and cleaned using distilled water, 95% ethyl alcohol, and water after being covered with Gram's iodine solution for 60 seconds. After using safranin for 30 seconds, the area was cleaned with distilled water and allowed to air dry. According to Aneja (2003), bacteria were classified as Gram-positive or purple and Gram-negative or pink.

3.5.3 Motility test

The sterile motility agar medium was used to stably inoculate each test culture. The cultures were then incubated for 48 hours at 37 °C to monitor bacterial motility.

3.5.4 Endospore staining

The organisms to be tested for endospores were prepared and heat-fixed into smears. A piece of absorbent paper that was cut to fit the slide was used to cover the smear, and it was set on a wire mesh on a ring stand. After saturating the paper with malachite green, the slide was heated until steam emerged from the top. To maintain the slide's steam for roughly three minutes, reheating was done as needed. Malachite green was gradually added to the paper as it started to dry in order to keep it moist and prevent a noticeable drop in temperature. Afterwards, the slide was carefully cleaned with tap water after the paper was taken off with tweezers. After draining and counterstaining the slide for 45 seconds with 0.5 percent safranin, the slide was cleaned with tap water, pat dried, and scrutinized. The spores will appear green, and the vegetative cells will appear red (Aneja, 2003; Harley and Prescott, 2002).

3.6 Biochemical characterization and tentative identification of phyllosphereisolates

Biochemical characterization of phyllosphere bacteria involves a series of tests aimed at identifying specific enzymatic activities and metabolic pathways exhibited by these microorganisms. The following are the key biochemical tests used for characterization:

3.6.1 Catalase test

A small amount of bacterial growth from each isolate is transferred onto a glass slide or into a test tube using a sterile loop or inoculation needle. A few drops of 3% hydrogen peroxide solution are added to each bacterial growth. Observations are made for the presence of bubbles, indicating positive catalase activity (Blazevic and Ederer, 1975).

3.6.2 Oxidase test (Cappuccino and Sherman, 1996)

A small amount of bacterial growth is transferred onto an oxidase test strip or disk. Color changes are observed according to the manufacturer's instructions within a specified time period, indicating positive or negative oxidase activity.

3.6.3 Starch Hydrolysis

Bacterial isolates are streaked onto starch agar plates. Plates are incubated at the appropriate temperature for the given bacteria. After incubation, plates are flooded with

iodine solution and observed for the presence of clear zones around bacterial growth, indicating starch hydrolysis.

3.6.4 Hydrogen Sulphide production

Bacterial isolates are inoculated into SIM agar tubes. Tubes are incubated at the appropriate temperature for the given bacteria. After incubation, tubes are observed for the presence of black precipitates, indicating positive hydrogen sulfide production. (Blazevic and Ederer, 1975).

3.6.5 Methyl red test (Seeley and Vandemark, 1981)

Bacterial isolates are cultured in MR-VP broth tubes. Methyl red indicator is added to the tubes. After incubation, tubes are observed for the development of a red color, indicating positive methyl red test.

3.6.6 Urease test

Bacterial isolates are streaked onto urea agar plates. Plates are incubated at the appropriate temperature for the given bacteria. After incubation, plates are observed for a pink color change, indicating positive urease activity. (James and Sherman, 1992).

3.6.7 Gelatin liquefaction

Bacterial isolates are inoculated into gelatin tubes. Tubes are incubated at the appropriate temperature for the given bacteria. After incubation, tubes are chilled to assess for solidification or liquefaction of gelatin. A liquid consistency indicates positive gelatin liquefaction. (Blazevic and Ederer, 1975).

3.6.8 Nitrate reduction test

The pre-sterilized nitrate broth was used to inoculate the isolates, and they were then incubated for 48 hours at 37°C. A porcelain dish was filled with one drop of culture, one drop of sulphuric acid, and three drops of the nitrate test reagent. The test will be deemed successful if the colour changes to red. (Beishir, 1987).

3.6.9 Indole production (Venant *et al.*, 2013)

Bacterial isolates are cultured in tryptone broth tubes. Kovac's reagent is added to the tubes. After incubation, tubes are observed for a red coloration in the reagent layer, indicating positive indole production.

3.6.10 Citrate Utilization

Bacterial isolates are streaked onto citrate agar slants. Slants are incubated at the appropriate temperature for the given bacteria. After incubation, slants are observed for growth and color change of the medium from green to blue, indicating positive citrate utilization.

3.6.11 Voges-Proskauer Reaction

Test tubes were filled with sterilized V-P broth (pH 6.5) and incubated for 3, 5, and 7 days. Following incubation, 3 ml of 40% (w/v) NaOH was mixed with cultures, and 0.5 mg of creatine was added. The test tubes were then checked for the production of acetyl methyl carbinol. At room temperature for half an hour, the tubes were checked to see if any red colour was produced.

Casein Hydrolysis (Smibert and Kreig, 1981)

Skim milk agar plates were streaked with the isolates, and they were then allowed to incubate at room temperature. The creation of clear zones surrounding the colonies, a sign of casein hydrolysis, was regarded as a positive reaction.

Result and Discussion

Morphological characterization of rhizosphere and phyllosphere isolates

Based on the morphological and microscopic characterization, the bacterial colonies were identified upto generic level. The bacterial isolates were examined for variations in colony morphology, shape, gram reaction, and motility. The morphological characteristics of all the forty phyllosphere were represented in the Table 1.

The bacteria that were isolated for the study were identified as *Bacillus subtilis* and *Bacillus pumilus* based on their morphological and biochemical characteristics (Awais *et al.*, 2007).

Colony Morphology

The colony morphology of all the 40 isolates varied from creamish to white, yellow, pinkish to red color and smooth circular small size to raised irregular large size colonies (Table 1) (Plate 1).

Meena *et al.* (2012) distinguished between the methylotrophic isolates using their colony morphology and growth characteristics. Vinod *et al.* (2017) used morphological and cultural characteristics, such as colony morphology and pigmentation, to isolate and identify 24 PSB from maize. Of these, eight were *Pseudomonas* and sixteen were *Bacillus*.

Shape

All the forty phyllosphere isolated bacteria were rod shaped and were represented in the Table 1. Dhruveet *al.* isolated 41 distinct phosphate-solubilizing bacteria in 2017.

Gram reaction and Motility

Among forty phyllosphere bacterial isolates, 19 isolates were Gram positive as they show crystal violet by gram reaction process includes CPRMS 4, CPRMS 5, CPRMS 7, CPRMS 8, CPRMS 10, CPRMS 11, CPRMS 14, CPRMS 15, CPRMS 19, CPRMS 21, CPRMS 23, CPRMS 25, CPRMS 27, CPRMS 29, CPRMS 31, CPRMS 32, CPRMS 34, CPRMS 38, CPRMS 40 and 21 of 40 isolates were confirmed as gram negative as they produce pink color on gram reaction *viz.* CPRMS 1, CPRMS 2, CPRMS 3, CPRMS 6, CPRMS 9, CPRMS 12, CPRMS 13, CPRMS 16, CPRMS 17, CPRMS 18, CPRMS 20, CPRMS 22, CPRMS 24, CPRMS 26, CPRMS 28, CPRMS 30, CPRMS 33, CPRMS 35, CPRMS 36, CPRMS 37, CPRMS 39 (Plate 2) and All the isolated rhizosphere bacteria were motile.

A study was carried out to isolate *Bacillus* species from the rhizosphere of soybean plants. According to Wahyudi *et al.* (2011), every single one of these isolates had positive Gram reaction and endospore appearance results.

Table 1: Morphological characterization of phyllosphere bacteria

Sl. No.	Isolates	Morphological characters				
		Colony morphology	Shape	Gram reaction	Spore formation	Motility
1	CPRMS 1	Creamy, white, raised	Rod	-ve	-ve	Motile
2	CPRMS 2	White, small, smooth	Rod	-ve	+ve	Motile
3	CPRMS 3	Irregular, flat, creamy whitish	Rod	-ve	-ve	Motile
4	CPRMS 4	Light cream, flat, irregular	Rod	+ve	+ve	Motile
5	CPRMS 5	Irregular, reddish, flat	Rod	-ve	-ve	Motile
6	CPRMS 6	Whitish, small, irregular	Rod	-ve	-ve	Motile
7	CPRMS 7	Irregular, small, light cream	Rod	+ve	-ve	Motile
8	CPRMS 8	Small, white, circular, flat	Rod	+ve	+ve	Motile
9	CPRMS 9	Small white, slimy, flat	Rod	-ve	-ve	Motile
10	CPRMS 10	Cream, small, regular	Rod	+ve	-ve	Motile
11	CPRMS 11	Creamy whitish, large, irregular	Rod	+ve	+ve	Motile
12	CPRMS 12	Whitish, large, irregular, flat	Rod	-ve	-ve	Motile
13	CPRMS 13	Small yellowish, raised	Rod	-ve	+ve	Motile
14	CPRMS 14	Irregular white smooth	Rod	+ve	+ve	Motile
15	CPRMS 15	Irregular, creamy, wrinkled	Rod	+ve	-ve	Motile
16	CPRMS 16	Large, whitish, flat	Rod	-ve	-ve	Motile
17	CPRMS 17	White, small, flat, irregular	Rod	-ve	+ve	Motile
18	CPRMS 18	Irregular, whitish, medium, flat	Rod	-ve	+ve	Motile
19	CPRMS 19	Small, white, flat	Rod	+ve	-ve	Motile
20	CPRMS 20	Irregular, white, smooth	Rod	-ve	-ve	Motile

Sl. No.	Isolates	Morphological characters				
		Colony morphology	Shape	Gram reaction	Spore formation	Motility
21	CPRMS 21	White, smooth, small, flat	Rod	+ve	+ve	Motile
22	CPRMS 22	White, large, irregular	Rod	-ve	-ve	Motile
23	CPRMS 23	Irregular, dull white, dry	Rod	+ve	-ve	Motile
24	CPRMS 24	Cream, small, regular	Rod	-ve	+ve	Motile
25	CPRMS 25	Small, orange, circular, white	Rod	+ve	+ve	Motile

26	CPRMS 26	Small, white, flat	Rod	-ve	-ve	Motile
27	CPRMS 27	Creamy white, raised	Rod	+ve	-ve	Motile
28	CPRMS 28	Light cream, regular, flat	Rod	-ve	+ve	Motile
29	CPRMS 29	Small yellowish, raised	Rod	+ve	-ve	Motile
30	CPRMS 30	White, slimy, flat	Rod	-ve	+ve	Motile
31	CPRMS 31	White, smooth, irregular	Rod	+ve	+ve	Motile
32	CPRMS 32	White, large, flat	Rod	+ve	-ve	Motile
33	CPRMS 33	Small, white irregular	Rod	-ve	+ve	Motile
34	CPRMS 34	Cream yellow, flat, circular	Rod	+ve	+ve	Motile
35	CPRMS 35	Red, raised, regular	Rod	-ve	-ve	Motile
36	CPRMS 36	Creamy, wrinkled, irregular	Rod	-ve	-ve	Motile
37	CPRMS 37	Whitish, medium, flat, irregular	Rod	-ve	+ve	Motile
38	CPRMS 38	Large, whitish, flat	Rod	+ve	-ve	Motile
39	CPRMS 39	Yellow, slimy, raised	Rod	-ve	+ve	Motile
40	CPRMS 40	Cream, irregular, flat	Rod	+ve	-ve	Motile




Plate 1: Morphological characterization of phyllosphere bacteria




Plate 2: Gram Staining of phyllosphere bacteria

Endospore staining

Among forty phyllosphere isolates 19 isolates produces endospores, whereas 21 isolates shows negative result for the endospore staining test. The production of endospores and a Gram-positive reaction characterize members of the genus *Bacillus*, which are commonly found in soil (Baumann, 2004).

Biochemical characterization and tentative identification of rhizosphere and phyllosphere isolates

All the forty phyllosphere isolates were undergone for the biochemical test with respect to chemical reaction. Based on the result of biochemical tests the isolates were tentatively identified up to generic level. The results of all the biochemical tests were described in the Table 2.

Catalase test

All the forty phyllosphere isolates showed positive result for the catalase test (Plate

3). It means all the isolates are aerobic in nature.

Oxidase test

Thirty of forty phyllosphere isolates were positive for oxidase test.

Starch hydrolysis

Thirty three isolates of forty rhizosphere strains and 28 isolates of forty phyllosphere strains showed positive result for the starch hydrolysis.

Hydrogen sulphide test

Eighteen of forty phyllosphere isolates were showed positive result for the hydrogen sulphide test.

Methyl red test

All the forty isolates showed positive results for the methyl red test (Plate 4).

Urease Test

Fifteen isolates of forty phyllosphere strains were positive for the urease test.

4.5.7 Gelatin Liquefaction

All the phyllosphere isolates showed positive result for the gelatin liquefaction test.

4.5.8 Nitrate reduction test

Thirty six of 40 rhizosphere isolates and 28 of 40 phyllosphere isolates showed positive result for the nitrate reduction test.

4.5.9 Indole test

None of the phyllosphere isolates showed positive result for the Indole test.

4.5.10 Citrate Utilization

Thirty eight of forty phyllosphere isolates were showed positive result for the citrate

utilization test (Plate 5).

4.5.11 Voges-Proskauer Reaction

All most all the phyllosphere isolates showed positive result for the Voges-Proskauer reaction.

4.5.12 Casein Hydrolysis

Among forty phyllosphereisolates 22 isolates showed positive result for the casein hydrolysis.

Mondal *et al.* (2016) obtained comparable outcomes by isolating and analyzing nine putative phosphate-solubilizing bacterial isolates using Pikovskaya's liquid medium. Biochemical tests revealed they tested positive for obligate aerobes, starch hydrolysis, and catalase.



Plate 3: Catalase test




Plate 4: Methyl red test




Plate 5: Citrate Utilization test

Table 2: Biochemical characterization and tentative identification of phyllospherebacteria from different cereals sample

Sl. No.	Isolates	Biochemical characterization												Tentative genus
		SH	CAT	HSP	OXD	CH	MR	VP	CU	UR	GL	NR	IN	
1	CPRMS 1	+	+	+	-	+	+	+	+	-	+	+	-	<i>Pseudomonasspp</i>
2	CPRMS 2	+	+	+	+	+	+	+	+	+	+	+	-	<i>Pseudomonasspp</i>
3	CPRMS 3	+	+	+	+	+	+	+	+	+	+	+	-	<i>Serratiaspp</i>
4	CPRMS 4	+	+	-	+	-	+	+	+	+	+	+	-	<i>Bacilluspp</i>
5	CPRMS 5	+	+	-	+	-	+	+	+	+	+	+	-	<i>Serratia marcesence</i>
6	CPRMS 6	+	+	+	-	+	+	+	+	-	+	+	-	<i>Pseudomonas spp</i>
7	CPRMS 7	+	+	+	-	+	+	+	+	-	+	+	-	<i>Bacilluspp</i>
8	CPRMS 8	+	+	-	+	-	+	+	+	+	+	+	-	<i>Bacilluspp</i>
9	CPRMS 9	+	+	+	+	+	+	+	+	+	+	+	-	<i>Serratiaspp</i>
10	CPRMS 10	+	+	+	-	+	+	+	+	-	+	+	-	<i>Bacilluspp</i>
11	CPRMS 11	+	+	+	-	+	+	+	+	-	+	+	-	<i>Bacilluspp</i>
12	CPRMS 12	+	+	+	-	+	+	+	+	-	+	+	-	<i>Pseudomonasspp</i>
13	CPRMS	+	+	-	+	-	+	+	+	+	+	+	-	<i>Serratiaspp</i>

	13													
14	CPRMS 14	+	+	-	+	-	+	+	+	+	+	+	-	<i>Bacillus</i> spp
15	CPRMS 15	+	+	-	+	-	+	+	+	+	+	+	-	<i>Bacillus</i> spp
16	CPRMS 16	+	+	-	+	-	+	+	+	+	+	+	-	<i>Pseudomonas</i> spp
17	CPRMS 17	+	+	+	+	+	+	+	+	+	+	+	-	<i>Methylobacterium</i> spp
18	CPRMS 18	+	+	+	+	+	+	+	+	+	+	+	-	<i>Pseudomonas</i> spp
19	CPRMS 19	+	+	+	-	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
20	CPRMS 20	+	+	-	-	-	+	+	+	-	+	+	-	<i>Pseudomonas</i> spp

Sl. No.	Isolates	Biochemical characterization												Tentative genus	
		SH	CAT	HS P	OX D	C H	M R	VP	CU	UR	GL	N R	IN		
21	CPRMS 21	+	+	+	-	+	+	+	+	+	+	+	+	-	<i>Bacillus</i> spp
22	CPRMS 22	+	+	+	+	+	+	+	+	+	+	+	+	-	<i>Pseudomonas</i> spp
23	CPRMS 23	+	+	+	+	+	+	+	+	+	+	+	+	-	<i>Bacillus</i> spp
24	CPRMS 24	+	+	+	+	-	+	+	+	+	+	+	+	-	<i>Serratias</i> spp
25	CPRMS 25	+	+	-	+	-	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
26	CPRMS 26	+	+	+	-	+	+	+	+	+	-	+	+	-	<i>Pseudomonas</i> spp
27	CPRMS 27	+	+	+	-	+	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
28	CPRMS 28	+	+	+	+	-	+	+	+	+	+	+	+	-	<i>Pseudomonas</i> spp
29	CPRMS 29	+	+	+	+	+	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
30	CPRMS 30	+	+	-	-	+	+	+	+	+	-	+	+	-	<i>Methylobacteriu</i> <i>m</i> spp
31	CPRMS 31	+	+	-	-	+	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
32	CPRMS 32	+	+	-	-	+	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp
33	CPRMS 33	+	+	+	+	-	+	+	+	+	-	+	+	-	<i>Pseudomonas</i> spp
34	CPRMS 34	+	+	-	+	-	+	+	+	+	+	+	+	-	<i>Bacillus</i> spp

35	CPRMS 35	+	+	+	+	-	+	+	+	-	+	+	-	<i>Methylobacteriu m radiotolerans</i>
36	CPRMS 36	+	+	+	+	-	+	+	+	-	+	+	-	<i>Pseudomonassp p</i>
37	CPRMS 37	+	+	-	+	+	+	+	+	+	+	+	-	<i>Pseudomonassp p</i>
38	CPRMS 38	+	+	-	+	+	+	+	+	+	+	+	-	<i>Bacilluspp</i>
39	CPRMS 39	+	+	+	-	+	+	+	+	+	+	+	-	<i>Serratiaspp</i>
40	CPRMS 40	+	+	+	-	-	+	+	+	+	+	+	-	<i>Bacilluspp</i>

Conclusion

In conclusion, our research on the morphological and biochemical characteristics of phyllosphere bacteria isolated from cereal samples has provided valuable insights into the diversity and metabolic potential of these microorganisms within plant ecosystems. Through a comprehensive analysis of colony morphology and biochemical tests, including catalase, oxidase, starch hydrolysis, gelatin liquefaction, hydrogen sulfide production, methyl red test, urease activity, indole production, and citrate utilization, we have elucidated the metabolic capabilities of the isolated bacteria. Our findings demonstrate the presence of diverse metabolic pathways among phyllosphere bacteria, reflecting their adaptation to the unique environmental conditions of plant surfaces. These metabolic traits play significant roles in nutrient cycling, plant-microbe interactions, and ecosystem functioning within the phyllosphere environment. Overall, our study contributes to the understanding of the phyllosphere microbiome and its significance in agricultural ecosystems. Future research may further explore the ecological roles and applications of phyllosphere bacteria, ultimately contributing to the development of sustainable agricultural practices.

References:

- Aneja, K. R., 2003, Experiments in microbiology, plant pathology and biotechnology. New Age International
- Awais, M., Shah, A. A., Hameed, A. and Hasan, F., 2007, Isolation, identification and optimization of bacitracin produced by *Bacillus* sp. *Pakistan J. Bot.*, 39(4): 1303.

- Barthalomew, J. W. and Mittewer, J., 1950, A simplified bacterial Strain. *Stain Tech.*, 25: 153.
- Bauman, R. W., 2004, *Microbiology*, Pearson Education Inc., San Francisco, CA.
- Beishir, L., 1987, *Microbiology in Practice: a self-instructional laboratory course*. 4: 120–130. New York: Harper and Row Publications.
- Blazevic, D. J. and Ederer, G. M., 1975, *Principles of biochemical tests in diagnostic microbiology*, Wiley and Company, New York., 13-45.
- Bulgarelli, D., Schlaeppi, K., Spaepen, S., van Themaat, E. V. L., & Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*, 64, 807-838.
- Cappuccino, J. G. and Sherman, N., 1996, *In Microbiology, A laboratory manual*. The Benjamin/Cummings Publishing Company Inc. (4th Edition), Melopark, California.
- Dhurve, N. G., Ingle, R. W., Lad, R. S. and Madavi, P. N., 2017, Characterization of phosphate solubilizing bacteria isolated from paddy rhizosphere of Vidarbha region. *Int. J. Chemi. Studies.*, 5(6): 24-30.
- Harley and Prescott., 2002, *Laboratory Exercises in Microbiology*, 58.
- James, C. R. and Sherman, N., 1992, *Microbiology and laboratory manual*, Rockland Community college, Suffern, New York, 3rd ed. The Benjamin/Cummings Publishing Co. Inc., Redwood city, California.
- Leveau, J. H. J., & Lindow, S. E. (2001). Appetite of an epiphyte: quantitative monitoring of bacterial sugar consumption in the phyllosphere. *Proceedings of the National Academy of Sciences*, 98(6), 3446-3453.
- Meena, K. K., Kumar, M., Kalyuzhnaya, M. G., Yandigeri, M. S., Singh, D. P., Saxena, A. K. and Arora, D. K., 2012, Epiphytic pink-pigmented methylotrophic bacteria enhance germination and seedling growth of wheat (*Triticum aestivum*) by producing phytohormone. *Antonie Van Leeuwenhoek.*, 101: 777-786.

- Mondal, D., Islam, M. S., Hoque, M. F., Hossain, M. K., Islam, M. K., Hossin, M. S. and Ahsan, S. M., 2016, Isolation and screening of potential phosphate solubilizing bacteria from tidal saline soils of Bangladesh. *J. Environ. Res.*, 4(3): 198-207.
- Seeley, H. W. and Vandemark, P. S., 1981, *Microbes in action—Laboratory manual for microbiology*, Freeman and Company, San Francisco, USA: 388.
- Smibert, R. M. and Kreig, N. R., 1981, General characterization. In : *Methodology for General Bacteriology*, Eds. P. Gerhardt, Academic Publisher, New York. 1: 400- 450.
- Venant, N., Pascal, K. and Ernest, S., 2013, Isolation of *Bacillus* strains from the rhizosphere of Tomato and their *in vitro* antagonistic effects against phytopathogenic fungi. *J. Microbiol.*, 2(3): 065-071.
- Vinod, B., Triveni, Subhash, R. and Sathyanarayana, 2017, Isolation and characterization of phosphate solubilizing bacteria from agriculture soil of Jaipur, Rajasthan. *Bull. Env. Pharmacol. Life Sci.*, 6(1): 194-200.
- Vorholt, J. A. (2012). Microbial life in the phyllosphere. *Nature Reviews Microbiology*, 10(12), 828-840.
- Wahyudi, A. T., Astuti, R. P., Widyawati, A., Meryandini, A. and Nawangsih, A. A., 2011, Characterization of *Bacillus* sp. strains isolated from rhizosphere of soybean plants for their use as potential plant growth for promoting rhizobacteria *chnol*, 11(32): 8085-8091.