

Characterization and Classifications of Soils of an Experimental farm in Tiruvannamalai District, Tamil Nadu

ABSTRACT

Characterization and classification of soil resources is essential scientifically to plan agricultural land use agroforestry, agri-horticulture and silvipastoral farming systems. In the present study, an attempt was made to characterize and classify the soils occurring on different topography in an experimental farm of Tiruvannamalai District of Tamil Nadu. Six representative pedons (covering all the soil types) very shallow to moderately deep, dark reddish brown to yellowish red (red soils) in colour with sub-angular blocky in structure, These soils were sandy loam to sandy clay loam in texture, slightly acidic to mildly alkaline (3.28-7.70) in reaction (non-saline soils) and had low to medium (1.2 to 6.4 g/kg) organic carbon content. Calcium and magnesium were the dominant exchangeable cations followed by sodium and potassium. Soils were grouped under Alfisols and, Inceptisols orders.

Keywords: Soil survey, soil characterization, soil classification, soil physical and chemical characteristics

1. INTRODUCTION

Soil is recognized as a base for every production system and knowledge of their properties, extent and spatial distribution is extremely important to maintain soil resources to sustain the ecosystem [1] and site-specific soil management practices. The systematic study of morphology, physicochemical characteristics and taxonomy of soils provides information on the nature and type of soils, their constraints, potentials, capabilities and their suitability for various uses [2,3]. The data generated through systematic soil resource inventory and characterization will help to formulate agro-interventions for management of these soils under existing delivery extension systems agriculture schemes, and value chains, so that new technologies could be adopted for sustainable production.

2. MATERIALS AND METHODS

2.1 Description of the study area

The study was carried out in soils of experimental farm of Agricultural college and Research Institute, Vazhavachanur, Tamil Nadu Agricultural university in Tiruvannamalai district. Geographically the study area is located between 12°4'15" N to 12°4'45" N Latitude and 78°59'0" E to 77°59'25" E Longitudes (Fig.1). The annual rainfall of the region is 759.4 mm. The mean maximum and minimum temperatures are 38°C and 21°C, respectively. Geologically it is underlain by granites. Geomorphologically, the Vazhavachanur farm is categorized as Pediplain.

2.2 Soil sampling methodology

A detailed soil survey was carried out using cadastral map of Vazhavachanur village. The soils belong to Kurumbalur series. Six typical pedons were selected based on landform and their soil morphological properties (Table 2). The horizon-wise soil samples were collected, processed and analysed using standard analytical methods and soils were classified [4].




Fig.1. Study area located in AC & RI, Vazhavachanurfarm of Tiruvannamalai District, Tamil Nadu, India

3. RESULTS AND DISCUSSION

3.1 Physiography and Soils

Physiographically, the area has been characterized into Pediplain shallow weathered and Pediplain moderately weathered which were further subdivided based on slope and erosion categories. Six soil profiles were dug in C, D, G, J, O and P blocks of AC&RI, Vazhavachanur farm based on the physiography of the soils. The soils of C, D and O Block are categorized as pediplain moderately weathered. The soils of G, J and P block belongs to pediplain shallow weathered.

3.2 Morphological features of soils

The soils of Pediplain shallow weathered (soils of G, J and P block) are moderately shallow, well drained, noncalcareous and severely eroded. The soils of nearly level pediplain moderately weathered (soils of C, D and O Block) are moderately deep, moderately well drained, non

calcareous and moderately eroded. The soils of the farm belong to the Kurumbalur soil series. Based on the properties, the soils were grouped under Inceptisol and Alfisol (Table 1).

3.3 Physical properties of soils

The coarse fragments of the pedons varied from 9.55 to 59.89 per cent (Table 1). The coarse fragments were mainly of quartz fragments. The irregular nature of gravels suggested that they were formed in situ. The surface soils possessed loamy sand to sandy clay loam texture and subsurface soils possessed sandy loam to sandy clay loam texture. The clay content of these soils ranged from 11.00 per cent in G block to 38.28 per cent in P Block. Unimodal distribution i.e., increase of clay upto certain depth and then decrease was observed in soils of O block. An increase in clay content was primarily due to illuviation. The clay content gradually increased with the depth in soils of C,D and J blocks. This may be due to combined effect of *insitu* clay formation and illuviation [5]. In P block there was no wide textural class variation. This could be ascribed to increased compaction and decreased aggregation caused by decreased OC and clogging of pores by dispersed clay in sub-soil layers [6].

Table 1. Morphological and physical properties of soils of AC & RI, Vazhavachanur farm

Horizon	Depth	Colour	Texture	Structure	Bulk density M gm ⁻³	Particle density M gm ⁻³	Gravel (%)	Clay (%)	Silt (%)	Sand (%)
C block- Loamy skeletal, Mixed, isohyperthermic, Typic Rhodustalfs										
Ap	0-35	5YR4/6	sl	2msbk	1.27	2.30	26.47	15.10	9.20	75.70
Bt1	35-60	2.5YR4/6	scl	3msbk	1.33	2.00	31.76	25.00	22.20	52.80
Bt2	60-80	2.5YR4/6	scl	2msbk	1.41	1.80	59.89	27.60	22.00	50.40
D Block –Loamy skeletal, Mixed, isohyperthermic, Typic Haplustalfs										
Ap	0-30	7.5YR 6/6	sl	2msbk	1.14	2.50	9.55	18.90	13.50	67.60
Bt1	30-50	5YR5/6	scl	2msbk	1.43	2.20	15.76	24.38	10.73	64.89
Bt2	50-60	5YR5/6	scl	2msbk	1.51	2.00	25.81	29.43	9.99	60.58
G Block –Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs										
Ap	0-20	5YR3/4	ls	1mgr	1.26	2.22	28.47	11.00	13.40	75.60
Bw	20-50	2.5YR3/6	scl	1msbk	1.42	2.00	43.35	37.28	4.82	57.90
J Block – Fine loamy, Mixed, isohyperthermic, Vertic Haplustepts										
Ap	0-20	10YR 3/2	scl	3msbk	1.46	2.16	9.68	19.20	15.68	65.12
B1	20-40	7.5YR3/4	sl	3msbk	1.22	2.05	24.68	14.20	12.26	73.54
O Block–Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs										
Ap	0-25	5YR 4/4	scl	2msbk	1.11	2.22	11.67	24.00	5.10	70.90
Bt1	25-40	2.5YR 4/8	scl	2msbk	1.25	2.17	15.93	26.80	7.00	66.20
Bt2	40-70	2.5YR 4/6	scl	2msbk	1.17	1.96	22.83	23.20	18.80	58.00
P Block –Fine loamy, Mixed, isohyperthermic, Typic Haplustepts										
Ap	0-30	5YR 4/3	scl	2msbk	1.33	1.81	13.74	38.28	8.25	53.47
B1	30-60	5YR 4/6	scl	3msbk	1.25	1.66	24.93	36.11	3.87	60.02

Soil texture : sl - Sandy loam; scl – sandy clay loam; ls - loamy sand; cl - Clay loam; sc – sandy clay; c - Clay
Soil structure : f- fine; m - Medium; c - coarse; 1 -weak; 2 - moderate; 3-strong;gr – granular; sbk - subangular blocky, abk -angular blocky

3.4 Chemical properties of soils

The soils are slightly acidic (pH 6.28) to mildly alkaline (pH 7.70) and this variation was attributed to the nature of parent material (Table 2). The electrical conductivity values (dS m⁻¹) varied from 0.01 in

C Block to 0.40 in J Block. The OC content ranged from 1.2 to 6.4 g/kg, and categorized as low to the medium. The surface horizons had higher than the sub-surface horizons because of more biomass addition [7]. The CaCO₃ ranged from 0.13 to 0.88 %. The cation exchange of the soils (cmolP⁽⁺⁾ kg⁻¹) varied from 6.44 in D Block to 27.57 in C block. The wide range of CEC is related to the amount and type of clay, and the of organic carbon content in these soils [8].The depth wise distribution of CEC had same trend as that of clay distribution. The exchangeable bases in the all the pedons were in the order of Ca⁺² > Mg⁺² > Na⁺ > K⁺[9].

Table 2. Chemical properties of soils of AC &RI, Vazhavachanur farm

Horizon	Depth	PH (1:2.5)	EC (dSm ⁻¹)	OC (g/kg)	CaCO ₃ (%)	CEC cmol (p ⁺) kg ⁻¹	Exchangeable Cations cmol (p ⁺) kg ⁻¹				BSP (%)	ESP (%)
							Ca	Mg	Na	K		
C block- Loamy skeletal, Mixed, isohyperthermic, Typic Rhodustalfs												
Ap	0-35	6.56	0.02	6.4	0.75	27.57	15.00	4.50	0.54	0.19	73.40	1.97
Bt1	35-60	6.28	0.01	5.6	0.25	16.92	8.00	4.00	0.22	0.16	73.15	1.28
Bt2	60-80	6.15	0.01	3.6	0.13	21.41	4.24	2.12	0.26	0.15	31.62	1.21
D Block –Loamy skeletal, Mixed, isohyperthermic, Typic Haplustalfs												
Ap	0-30	6.30	0.10	5.1	0.53	6.44	2.38	1.92	0.05	0.05	68.32	0.78
Bt1	30-50	6.60	0.10	4.8	0.35	13.35	2.12	1.06	0.44	0.28	29.14	3.26
Bt2	50-60	6.90	0.10	3.9	0.18	10.00	4.00	1.62	0.44	0.28	63.30	4.35
G Block –Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs												
Ap	0-20	6.90	0.05	4.8	0.42	9.25	5.00	2.50	0.51	0.03	86.96	5.52
Bw	20-50	7.10	0.02	4.5	0.28	15.62	10.80	0.42	0.68	0.02	76.31	4.35
J Block – Fine loamy, Mixed, isohyperthermic, VerticHaplustepts												
Ap	0-20	7.20	0.40	4.6	0.48	21.41	4.24	2.12	0.26	0.15	31.62	1.21
B1	20-40	6.90	0.30	3.0	0.41	14.72	7.50	3.00	0.22	0.19	74.11	1.48
O Block– Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs												
Ap	0-25	6.85	0.12	2.4	0.65	10.60	3.20	1.40	0.07	0.39	47.74	0.66
Bt1	25-40	6.27	0.17	4.5	0.45	9.80	2.40	2.40	0.01	0.27	51.84	0.10
Bt2	40-70	7.15	0.22	4.2	0.33	8.60	2.10	1.70	0.01	0.12	45.70	0.12
P Block – Fine loamy, Mixed, isohyperthermic, Typic Haplustepts												
Ap	0-30	6.40	0.20	2.9	0.88	16.19	10.05	4.38	0.36	0.11	92.07	2.41
B1	30-60	7.70	0.20	1.2	0.75	14.76	9.49	4.75	0.36	0.15	98.97	2.43

3.5 Classification of soils

Based on the morphological, physical and chemical properties of the soils were classified (4 Soil Survey Staff 2022). viz. Inceptisol and Alfisols. J Block and P block soils were grouped under Inceptisols based on the presence of Cambic subsurface horizon. J Block soils was placed under VerticHaplustepts due to the presence of Vertic properties. C Block soils were classified as Rhodustalfs because of occurrence of sub-horizons in the upper 100 cm of the argillic horizon or throughout the entire argillic horizon if less than 100 cm thick, more than 50 per cent colours that have hue of 2.5YR or redder and value, moist, of 3 or less. Similarly at sub group level, these soils did not exhibit any integration with other taxa nor deviated from central concept of Rhodustalfs are placed in Typic Rhodustalfs. C, D and G block soils were grouped under Alfisols since the illuviation of clay and the presence of base saturation more than 35% throughout the depth of the soil horizons and ustic moisture regime.

Table 3. Soil mapping units and taxonomic classification of AC &RI, Vazhavachanur farm

Name of the block-name of the series	Mapping unit	Taxonomic classification
C Block - Kurumbalur	$\frac{\text{Kbr-sl(g)-d}_4}{\text{A-e}_2}$	Loamy skeletal, Mixed, isohyperthermic, Typic Rhodustalfs
D Block- Kurumbalur	$\frac{\text{Kbr-sl(g)-d}_4}{\text{A-e}_2}$	Loamy skeletal, Mixed, isohyperthermic, Typic Haplustalfs
G Block- Kurumbalur	$\frac{\text{Kbr-ls-d}_3}{\text{B-e}_1}$	Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs
J Block- Kurumbalur	$\frac{\text{Kbr-scl-d}_3}{\text{B-e}_1}$	Fine loamy, Mixed, isohyperthermic, VerticHaplustepts
O Block- Kurumbalur	$\frac{\text{Kbr-scl-d}_4}{\text{B-e}_1}$	Fine loamy, Mixed, isohyperthermic, Typic Haplustalfs
P Block- Kurumbalur	$\frac{\text{Kbr-scl-d}_4}{\text{B-e}_1}$	Fine loamy, Mixed, isohyperthermic, Typic Haplustepts

4. CONCLUSION

The soils of the study area were slightly acidic to mildly alkaline inreaction (non saline) and had low to medium OC contentand low to medium CEC. These soils were classified asTypic Rhodustalfs, Typic Haplustalfs, Typic Haplustepts and VertifHaplustepts. The main constraints were depth, slope,excessive drainage and low water holding capacity in these soils.

REFERENCES

1. Sarmah T, Dutta S, Karmakar R.M, Tamuly D Dutta N. Characterization andclassification of some alluvium-derived rice and associated non-rice soils of Jorhat district ofAssam. Journal of the Indian Society of Soil Science.2019; 67: 379-388.
2. Jagdish Prasad, Ray SK, Gajbhiye KS Singh SR. Soils of Selsura research farm inWardha district, Maharashtra and their suitability for crops. Agropedology.2009;19:84-91.
3. Sashikala G, Naidu MVS, Ramana KV,Nagamadhuri KV, Reddy PKA, SudhakarP,. Krishna GT. Characterizationand classification of soils in semi-arid region ofTatrakallu village of Anantapuramu district inAndhra Pradesh. Journal of the Indian Societyof Soil Science. 2019; 67: 389-401.
4. Soil Survey Staff. Keys to Soil Taxonomy, 13thed. USDA-Natural Resources ConservationService 2022.

5. Vasundhara R, Srinivasan R, Dharumarajan S, Kalaiselvi B, Rajendra Hegde. Characterization and Classifications of Soils of Eastern Ghats Region of Karnataka. *Agropedology*. 2022; 32 (02): 200-210
6. Khan MAA, Kamalakar J. Physical, physicochemical and chemical properties of soils of newly established agro-biodiversity park of Acharya NG Ranga Agricultural University, Hyderabad, Andhra Pradesh. *International Journal of Farm Sciences*. 2012; 2: 102-116.
7. Vasundhara R, Prakash NB, Anil Kumar KS, Rajendra Hegde. Characterization and classification of areca nut-growing soils of Karnataka, India. *Journal of Plantation Crops*. 2020; 48: 91-102.
8. Sekhar C, Naidu MVS, Ramprakash T, Balaguravaiah D. Characterization and classification of soils in the central parts of Prakasam district in Andhra Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*. 2017; 6: 2699-2712.
9. Arunkumar V, Natarajan S, Sivasamy R. Soil resource mapping of Vellamadai Village, Coimbatore district fused (IRS 1C LISS III and PAN merged) space borne multispectral data. *Madras Agricultural Journal*. 2004; 91 (7-12): 399-405.

UNDER PEER REVIEW