

Original Research Article

Fertigation-Driven Optimization of Growth Characters in Banana (*Musa acuminata* AAA) cv. Red banana

ABSTRACT

The main aim of the study was to standardize the fertigation schedule and nutrient levels for growth enhancement in red banana. [The study was conducted at the Department of Fruit Science College of Agriculture, Vellayani, Thiruvananthapuram during 2021-2023.](#) The statistical design used was factorial [Randomized Block Design](#) replicated thrice with 12 plants per plot at a spacing of 2.1m x 2.1 m laid out with 12 treatment combinations of 4 different nutrient levels (60%, 80%, 100% and 120% RD of N and K) and 3 fertigation schedule (equal splits at weekly interval, at fortnightly interval, and according to the growth stage of the crop) and a control (KAU POP recommendation). Soil test based [on](#) fertilizer recommendation was derived based [on](#) KAU POP (200:200:400 g plant⁻¹ year⁻¹). Farm yard manure was [applied](#) as organic source at 10 kg plant⁻¹) and P (200 g plant⁻¹) to all treatments as basal. Water soluble fertilizers used included Urea as nitrogen source and Muriate of Potash as potassium source; Phosphorus was supplied through Rajphos. [Data was collected from pseudo stem height, girth, number of leaves, leaf index, leaf area, days to shooting, time taken from shooting to harvest and crop duration. Data was analyzed using analysis of variance.](#) The highest pseudostem height (310.53 cm) and girth (89.30 cm) was observed with the interactive effect of 120% of the recommended dose of N and K applied according to the crop growth stage) which was on par with F₂S₃ (80% of the recommended dose of N and K applied according to the crop growth stage) at flowering stage. The total functional leaf area was maximum [on](#) F₃S₃ at the flowering (23.86 m²) and harvest stage (21.50 m²). Data on days to shooting (363 days), shoot to harvest (89 days) and total crop duration (452 days) in red banana showed that the interaction effect F₂S₃ has taken the least number of days. [In conclusion](#) the growth characters were observed to be better at 80% recommended dose of N and K with split application according to the growth stage of the crop (F₂S₃).

Keywords: Red banana, fertigation, growth characters, urea, muriate of potash, rajphos,

1. INTRODUCTION

Banana is one of the most important fruit crops grown in India. India is the largest producer of banana in the world with a production of 30.81 million tonnes from an area of 8.84 million

hectares[1]. Among the banana varieties, Red banana or Chenkadali (fruit locally known as Kappa pazham) is one of the most relished and highly priced varieties of Kerala rich in antioxidants, beta carotene, vitamin C and vitamin B₆. This finicky-to-grow cultivar is extensively cultivated in southern parts of the state and bordering regions of Kanyakumari district in Tamil Nadu for its sweet flavour and creamy and distinctive purplish red peel colour.

Banana being a gross feeder requires high amount of nutrients for proper growth, development and optimum production. The unscientific crop management practices followed by farmers have led to poor utilization of nutrients resulting in low productivity. Fertigation has vast potential to improve nutrient use efficiency, saving labour towards [some cultural practices thereby](#) reducing the cost of production [and](#) maintaining soil health and meeting the specific nutritional requirements of the crop [2]. Fertilizer use efficiency in fertigation increases up to 67 per cent over conventional fertilizer application. Forty per cent saving in fertilizer usage can be achieved without any reduction in banana yield. Banana plants can effectively utilize the accurately-placed fertilizer in solution form in the active root zone area, resulting in vigorous growth, early flowering and early bunch development.

Knowledge about the precise dosage of fertilizers for application through fertigation is crucial for optimizing crop productivity. The success of any fertigation program predominantly hinges on the strategic scheduling of fertilization, as the frequency of application significantly influences the fulfilment of crop growth and development requirements. An accurate fertilizer recommendation for Red banana cultivation in Kerala is lacking. Despite this, numerous farmers are increasingly embracing its commercial cultivation. Growers face challenges such as extended crop duration, substantial fertilizer needs, and the elevated cost of production. Therefore, it is imperative to establish standardized nutrient requirements and a fertigation schedule for red banana. Addressing these considerations, the current study aims to standardize nutrient levels and a fertigation schedule to enhance the growth of red bananas under a fertigation system.

2. MATERIALS AND METHODS

The present study was carried out at the Department of Fruit Science, College of Agriculture, Vellayani, Thiruvananthapuram, Kerala during the period 2020-2023. The study employed a randomized block design, encompassing 12 treatment combinations and a control, each replicated three times. The combinations of fertigation levels (F) and split application of

fertilizers (S) and fertigation treatments were fixed according to the soil test based on N and K recommendation. The requirement of lime was calculated based on the initial soil status and the same was applied uniformly for all treatments. Soil test based fertilizer recommendation was derived based on 200:200:400 g plant⁻¹ year⁻¹ as per KAU POP (KAU, 2016). The fertigation levels included four levels of recommended dose of fertilizers (RDF) with respect to N and K viz., F₁ – 60 per cent RDF, F₂ – 80 per cent RDF, F₃ – 100 per cent RDF and F₄ – 120 per cent RDF. The split application included S₁ – Equal splits of N and K at weekly intervals up to 12 months, S₂ – Equal splits of N and K at fortnightly intervals up to 12 months, S₃ – split application of N and K in the proportion given in table no-1 and the control was 100% RDF with manual application of nutrients with conventional land management and drip irrigation system.

Comment [H1]: The referencing method is not unique

Table 1: Stages of growth and split application of N and K

Stages of growth	N(%)	N(g)	K(%)	K(g)
5 th to 12 th week	30	60	10	40
13 th to 25 th week	35	70	10	40
26 th to 37 th week	20	40	30	120
38 th to 49 th week	15	30	25	100
50 th to 52 nd week	-	-	25	100
Total	100	200	100	400

Farm yard manure @ 10 kg plant⁻¹ and P (200 g plant⁻¹) as rock phosphate was added given uniformly to all treatments as basal. Water soluble fertilizers used for the work included urea as nitrogen source and muriate of potash as potassium source.

Assessment of vegetative characters was conducted at five months after planting (MAP), nine MAP and at flowering. Pseudo stem height was measured from soil level to the base of the unopened leaf and expressed in centimeters, while girth of the pseudo stem was measured at 10 cm height of the pseudo stem from the ground level. The total number of fully opened functional leaves that appeared healthy on the plants were counted and recorded. The third fully opened leaf from the top (index leaf) was considered for measuring the leaf area (leaf area of index leaf = length of lamina x width of lamina x constant (0.8)) and was expressed in meter square. Days to shooting was calculated from planting to shooting. Time taken from shooting to harvest and crop duration was observed and expressed in days. [How did you analyze the collected data?](#)

3. RESULTS AND DISCUSSION

Pseudostem height (cm):

The effect of fertigation level and split application of fertilizers have shown significant effect on pseudostem height at 5 months after planting (MAP), 9 MAP, flowering stage and at the time of harvest (Table 2). Among the fertigation levels F_4 (120% of the recommended dose) recorded significantly higher values for plant height throughout the growth period and the lowest was recorded by F_1 (60% of the recommended dose) and in the split application level S_3 (split application according to growth stage) recorded the highest value. The interactive effect has shown that F_4S_3 recorded the highest value at 5 MAP (178.73 cm) which was at par with F_2S_3 (174 cm) and the lowest figure was recorded for F_1S_1 during all stages of growth. From the contrast analysis data it was found that the control was significantly different from the treatment combinations. This might be due to the fact that nutrients applied prior to the experiment through the soil may not have been available to the plants at different stages resulting in lower pseudostem height as reported by [3]. The steady increase in pseudostem height through fertigation could be best explained with the regular supply of plant nutrients and water which increased the availability of N, P and K in crop root zone ultimately leading to the enhanced uptake of these nutrients as reported by [4]. The results were similar to the findings byef [5] and [6].

Pseudostem Girth (cm):

Perusal of the data in Table 3, at 5 MAP revealed that pseudo stem girth (63.03 cm) was the highest for 75 % of the recommended dose of N and K through fertigation (F_3) along with split application according to the growth stage (S_3) which was at par with F_4S_3 (62.00 cm). Pseudostem girth at 9 MAP (77.27 cm) and flowering stage (89.30 cm) were the highest for F_4S_3 which was at par with F_3S_3 . This may be due to increased amount of nitrogen and potash in the soil and those supplied as a result of fertigation which contribute in the formation of complex nitrogenous substances such as proteins and amino acids which are the building blocks of tissues. Similar findings have also been reported [7]. The lowest value for pseudostem girth among the treatment combinations was recorded for F_1S_1 at 9MAP and flowering stage. The control significantly varied from the treatment combinations; this may be due to low uptake of nutrients by the plants through conventional method.

Total functional leaf area:

Effect of fertigation level and split application had significant effect on functional leaf area throughout the growth period of the crop (Table 4). The leaf area consistently increased in

the F₃ fertigation level and S₃ split application treatments. The interactive effect has shown that at 5 MAP the total functional leaf area was maximum for F₂S₃ (7.59 m²) and at flowering (23.86 m²). The lowest leaf area was recorded for F₁S₃ at 5 MAP, 9MAP. These results are in conformity with the findings by [8] and [9] who stated maximum leaf retention was at lowest fertigation level combined with consortium of fertilizer application. Higher frequency of irrigation and increased availability of soil moisture under subsurface drip fertigation might have led to effective absorption and utilization of available nutrients and better proliferation of roots resulting in quick canopy growth and physiological parameters [10]. According to [11], frequent application of nutrients through drip system improves the uptake of nutrients through two main mechanisms: i) continuous replenishment of nutrients in the depletion zone at the vicinity of root interface; and ii) enhanced transport of dissolved nutrients by mass flow, due to the higher water content in the medium.

Days to shooting and harvest

Results (Table 5) on days to shooting (325.33 days), shoot to harvest (89 days) and total crop duration (414.33 days) in red banana showed that the interaction effect of 80% of the recommended dose of N and K through drip irrigation (F₂) along with split application with regard to the crop growth stages (S₃) has taken the least number of days. The maximum number of days for shooting (455 days), shoot to harvest (106 days), and crop duration (561 days) were observed for F₁S₁ (i.e. 60% of the recommended dose of N and K through drip irrigation (F₁) along with weekly application of fertilizers (S₁). The early flowering in subsurface drip fertigation with fertilizers inoculated plants may be ascribed to easy uptake of nutrients and simultaneous transport of growth promoting substances like cytokinin to the auxiliary buds resulting in breakage of apical dominance [10]. Similar results were obtained by [12] who found that the drip irrigation minimized the days to harvest (398 days) as compared to surface method of irrigation (435 days). According to [13] fruit setting took place 28 days earlier and the average harvesting period was less by 32 days in drip irrigation system. Compared to soil application of fertilizers, low concentrations of fertilizers are applied in more number of splits in fertigation. This enables a precise and uniform application of nutrients to the wetted area, where the active roots are focused resulting in continuous supply of nutrients according to demand, reducing the losses and enhancing the availability [14].

Table 2: Effect of nutrient level and fertigation schedule on plant height (cm)

Treatments	5 MAP	9 MAP	At flowering
Fertigation level (F)			
F ₁	123.37	206.07	256.30

Formatted: Font: Bold

F ₂	143.51	229.52	281.50
F ₃	143.39	226.91	272.77
F ₄	155.89	242.48	286.53
SE m (±)	1.727	2.32	1.81
CD (0.05)	5.066	6.804	5.316
Split application (S)			
S ₁	130.04	211.87	261.55
S ₂	133.12	216.03	265.36
S ₃	161.47	250.84	295.91
SE m (±)	1.496	2.009	1.57
CD (0.05)	4.387	5.892	4.604
Interaction (F X S)			
F ₁ S ₁	120.00	200.20	251.00
F ₁ S ₂	125.00	210.00	257.87
F ₁ S ₃	125.13	208.00	260.03
F ₂ S ₁	126.00	207.07	265.00
F ₂ S ₂	130.53	211.50	269.23
F ₂ S ₃	174.00	270.00	310.27
F ₃ S ₁	130.16	210.20	253.00
F ₃ S ₂	132.00	214.33	260.23
F ₃ S ₃	168.00	256.20	305.07
F ₄ S ₁	144.00	230.00	277.20
F ₄ S ₂	144.93	228.30	274.10
F ₄ S ₃	178.73	269.16	308.30
SE m (±)	2.992	2.32	3.14
CD (0.05)	8.774	6.804	9.208
Treatment mean	141.54	226.20	274.27
Control mean	105.20	198.13	236.26
Control vs. treatments	S	S	S

Table 3: Effect of nutrient level and fertigation schedule on plant girth (cm)

Treatments	5 MAP	9 MAP	At flowering
Fertigation level (F)			
F ₁	44.08	61.43	70.28
F ₂	51.83	66.80	75.50
F ₃	56.35	67.50	79.97
F ₄	55.38	68.09	83.12
SE m (±)	1.488	1.444	1.285

CD (0.05)	4.364	4.236	3.769
Split application (S)			
S ₁	48.87	63.85	72.52
S ₂	49.34	62.31	75.87
S ₃	57.52	71.70	83.26
SE m (±)	1.289	1.251	1.113
CD (0.05)	3.78	3.669	3.264
Interaction (F X S)			
F ₁ S ₁	43.23	58.17	67.30
F ₁ S ₂	44.00	62.03	73.27
F ₁ S ₃	45.00	64.10	70.27
F ₂ S ₁	48.20	66.17	71.07
F ₂ S ₂	47.23	59.07	68.23
F ₂ S ₃	60.07	75.17	87.20
F ₃ S ₁	49.03	63.07	71.47
F ₃ S ₂	57.00	69.17	82.17
F ₃ S ₃	63.03	70.27	86.27
F ₄ S ₁	55.00	68.00	80.27
F ₄ S ₂	49.13	59.00	79.80
F ₄ S ₃	62.00	77.27	89.30
SE m (±)	2.577	2.502	2.226
CD (0.05)	4.363	7.337	6.529
Treatment mean	51.91	65.95	77.21
Control mean	41.00	58.00	63.23
Control Vs treatments	S	S	S

Table 4: Effect of nutrient level and fertigation schedule on total functional leaf area (m²)

Treatments	5 MAP	9 MAP	A _t flowering
Fertigation level (F)			
F ₁	2.33	5.60	7.02
F ₂	5.04	10.21	16.43
F ₃	5.89	10.29	20.05
F ₄	5.79	9.52	19.94
SE m (±)	0.02	0.02	0.017
CD (0.05)	0.06	0.059	0.05

Formatted Table

Split application (S)			
S ₁	4.26	8.52	14.75
S ₂	4.16	7.85	14.01
S ₃	5.86	10.34	18.82
SE m (±)	0.018	0.017	0.015
CD (0.05)	0.052	0.051	0.043
Interaction (F X S)			
F ₁ S ₁	2.71	6.05	7.30
F ₁ S ₂	2.23	4.68	6.49
F ₁ S ₃	2.04	6.05	7.25
F ₂ S ₁	3.80	9.60	13.13
F ₂ S ₂	3.72	9.27	14.03
F ₂ S ₃	7.59	11.75	22.13
F ₃ S ₁	4.87	9.63	16.51
F ₃ S ₂	5.73	9.83	19.77
F ₃ S ₃	7.05	11.40	23.86
F ₄ S ₁	5.64	8.76	22.03
F ₄ S ₂	4.96	7.64	15.76
F ₄ S ₃	6.77	12.14	22.03
SE m (±)	0.035	0.035	0.03
CD (0.05)	0.104	0.102	0.087
Treatment mean	4.75	8.90	15.80
Control mean	2.37	5.70	8.43
Control vs. treatments	S	S	S

Table 5: Effect of nutrient level and fertigation schedule on days to shooting, shoot to harvest duration and crop duration

	Days to shooting	Shoot to harvest duration	Crop duration
Fertigation level (F)			
F ₁	434.11	104.94	539.06
F ₂	384.11	96.78	480.89
F ₃	386.56	97.05	483.61
F ₄	389.83	97.44	487.28
SE m (±)	2.158	1.371	2.343
CD (0.05)	6.328	4.022	6.87

Split application (S)			
S ₁	417.79	99.79	517.58
S ₂	405.12	101.00	506.12
S ₃	373.04	96.37	469.42
SE m (±)	1.868	1.187	2.029
CD (0.05)	5.48	3.483	5.95
Interaction (F X S)			
F ₁ S ₁	455.00	106.00	561.00
F ₁ S ₂	422.67	105.00	527.67
F ₁ S ₃	424.67	103.83	528.50
F ₂ S ₁	413.33	100.00	513.33
F ₂ S ₂	413.67	101.33	515.00
F ₂ S ₃	325.33	<u>89.00</u>	414.33
F ₃ S ₁	413.17	<u>96.17</u>	509.33
F ₃ S ₂	412.33	<u>99.00</u>	511.33
F ₃ S ₃	334.17	<u>96.00</u>	430.17
F ₄ S ₁	389.67	<u>97.00</u>	486.67
F ₄ S ₂	371.83	<u>98.67</u>	470.50
F ₄ S ₃	408.00	<u>96.66</u>	504.67
SE m (±)	3.737	2.375	4.057
CD (0.05)	<u>10.96</u>	<u>5.16</u>	<u>11.90</u>
Treatment mean	398.65	<u>99.05</u>	497.70
Control mean	409.00	105.00	514.00
Treatment vs. Control	S	S	S

4. CONCLUSION

Assessment of the vegetative characters of red banana at 5 MAP, 9MAP and flowering stage has shown significant differences among the characters on application of different nutrient level and fertigation scheduling. The plant height, pseudo stem girth, total functional leaf area were observed to be the highest at 120% recommended dose of N and K with application of fertilizers according to growth stage of crop (F₄S₃) which was at par with F₂S₃ and F₃S₃. The days to shooting, shoot to harvest and total crop duration registered the least at 80% recommended dose of N and K with application of fertilizers according to the growth

Formatted: Line spacing: 1.5 lines

stage of the crop (F_2S_3). Thus overall growth characteristics proved better with interaction effect at 80% recommended dose of N and K along with split application according to the growth stage of the crop (F_2S_3).

REFERENCES

1. NHB [National Horticulture Board]. NHB home page[online]Available: 2020. [www.http://nhb.gov.in/statistics/areaproduction-statistics.html](http://nhb.gov.in/statistics/areaproduction-statistics.html) [14 February 2018]
2. Holder GD, Gimbs FA. Effects of nitrogen and irrigation on the growth and yield of banana. *Trop Agric. Israeli.* 1983; 68(4):331-334
3. Humajahan S, Patil D, Patil S, Venkateshalu RS, Ambika DS. Effect of fertigation on growth and yield of banana cv. Red banana (*Musa AAA*). *Int J Chem Studies.* 2018;6:2186–2189.
4. Basanta Singh, T, Patra SK, Chongtham Tania, Basudha Devi CH and Thokchom Narjit Singh. Effect of Drip Fertigation on the Plant Morphology and Crop Duration of Banana (cv. Martaman) in an Alluvial Soil. *Int. J. Curr. Microbiol. App. Sci.* 2018; 7(11): 3307-3315. doi: <https://doi.org/10.20546/ijcmas.2018.711.382>
5. Naresh Babu, Anamika Sharma and Shasher Singh. Effect of different nitrogen doses and their split application on growth, yield and quality of 'Jahajee' banana. *South Indian Hort.* 2004; 52:35-40.
6. Hedge DM. and Srinivas K. Effect of soil moisture stress on fruit growth and nutrient accumulation in banana cv. Robusta. *Fruits.* 1989; 44:135-138.
7. Santosh DT, and Tiwari KN. Estimation of water requirement of banana crop under drip irrigation with and without plastic mulch using dual crop coefficient approach. In *IOP Conference Series: Earth Environ. Sci.* 2019. 344: doi:10.1088/1755-315/344/1/012024
8. Senthilkumar M, Ganesh S, Srinivas K and Panneerselvam P. Enhancing Growth and Yield in Banana Cv. Robusta (AAA) through Fertigation with Microbial Consortium. *J Hort Sci.* 2013; 8(2): pp.240–245. doi: <https://doi.org/10.24154/jhs.v8i2.310>.

Formatted: Line spacing: 1.5 lines

9. Zhang X, Li H, Zhang T, Ma S. Effects of different water and nitrogen levels on nitrogen use efficiency under drip irrigation. *J. Irrig. Drain.* 2017; 37:45–50.
10. Mahendran PP, Yuraj M, Pareswari C, Gurusamy A and Krishnasamy S. Enchaining growth, yield and quality of banana through subsurface drip fertigation. *Int J Chem Env Biol Sci.* 2013; 1(2): 391- 394.
11. Sathya S, James Pitachi G, Indirani R and Kannathasan, M. Effect of fertigation on availability of nutrient (N, P & K) in soil - a review, *Agriculture Reviews.* 2008; 29(3): 214-219.
12. Sharma R. and Kispotta W. Study on Drip Irrigation in Banana - District of Kaushambi (U.P.). *J Business Manag.* 2016; 18(1):10-12.
13. Shashidhara KK, Bheemappa A, Hirevekanagoudar LV, Shashidhar KC. Benefits and constraints in adoption of drip irrigation among the plantation crop growers, *Karnataka J Agric Sci.* 2007; 20(1):82- 84.
14. Pintu RV. Precision farming in banana (*Musa AAB nendran*) for productivity enhancement. Ph. D. thesis, Kerala Agricultural University, Thrissur, 2017; 231p.