

Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* (Swinhoe) on Maize (*Zea mays* L.)

ABSTRACT: Spotted stem borer, *Chilo partellus* (Lepidoptera: Pyralidae) regarded as one of the most significant and prevalent insect pests of maize and its mystic aspect of feeding makes it challenging to control. Hence current study aims to evaluate comparative study of synthetic insecticides with biopesticides against *Chilo partellus*. Eight treatments viz., azadirachtin 10,000ppm, neem oil 3%, karanj oil 3%, neem oil 3% + imidacloprid 17.5%SL, karanj oil 3% + imidacloprid 17.5%SL, imidacloprid 17.5%SL, spinosad 45%SC and untreated control were tested under field condition in randomized block design with three replications. Among the treatments, lowest larval population of *Chilo partellus* per plant were recorded in karanj oil 3% + imidacloprid 17.5%SL (1.99) followed by neem oil 3% + imidacloprid 17.5%SL (2.42), imidacloprid 17.5%SL (2.71), spinosad 45%SC (3.17), neem oil 3% (3.73), karanj oil 3% (3.95) and azadirachtin 10,000ppm (4.44) were found superior over the untreated control (7.24). The highest yield was recorded in karanj oil 3% + imidacloprid 17.5%SL with 39.20 q/hectare and most economic treatment was imidacloprid 17.5%SL with highest cost benefit ratio (1:2.2). The lowest cost benefit ratio was recorded in untreated control (1:1.4).

Keywords: *Chilo partellus*, Maize stem borer, Management, Insecticides, Biopesticides, Efficacy

1. INTRODUCTION: Maize (*Zea mays* L.) was domesticated in Central America. It is the world's leading crop, widely cultivated as cereal grain. Globally, maize is known as queen of cereals because of its highest genetic yield potential (Directorate of Maize Research, 01-03-2024) that the crop is used for variety of purposes including foods and fodder using diverse plant parts, such as grain, leaves, stalks, tassels, and cobs (Singh *et al.*, 2006). It is an important source of vitamins, minerals, lipids, protein, starch, and fiber and also highly nutritious for cattle and poultry (Prasanna *et al.*, 2001). In terms of nutrition, maize grains include 4% oil, 70% carbohydrates, 2-3% crude fiber, and 10% protein in addition to vitamins A and E, riboflavin and nicotinic acid (Joshi 2015). Maize contributes over 20 per cent of food calories of Asian population (Shiferaw *et al.*, 2011). The global consumption

pattern of maize is, feed-61%, food-17% and industry-22%. It has attained a position of industrial crop globally as 83% of its production in the world is used in feed, starch and bio-fuel industries. Further, using maize directly or indirectly more than 3000 products are being made providing a wide opportunity for value addition. Because of its myriad uses, it is a prime driver of the global agricultural economy. With a yield of 1423 million T, it is grown on 188 million hectares of land in 170 countries across the globe. China has maximum area under maize followed by the USA, both together representing 39% of world maize area. Since 2005, India ranks 4th in terms of area with 9.89 million hectares land under maize. (Indian Institute of Maize Research, ICAR 2024). Production of Maize in the country during 2023-24 is estimated at record 224.82 lakh tones. (PJ TSAU, 2024).

Notable insect pests of the crop are *Helicoverpa armigera* (Hubner) and *Mythimna separata* (Walker). *Spodoptera frugiperda*, *Atherigona soccata* (Rondani), *Rhopalosiphum maidis* (Fitch), and *Chilo partellus* (Swinhoe) (Chouraddi *et al.*, 2017). Scientists have undertaken numerous investigations and come to the conclusion that the maize stem borer, *Chilo partellus* is a significant pest of pearl millet (*Pennisetum typhoideum* (Rich), sorghum (*Sorghum bicolor* L.), and maize (*Zea mays* L.) across Asia and Africa. (Panwar, 2005). It was also noticed infesting Rice (*Oryza sativa*), Sugarcane (*Saccharum officinarum*) and several millets and grasses. (Kauma *et al.*, 2008). *Chilo partellus*, a member of the Pyralidae family of insects, is a notable biotic barrier to maize cultivation worldwide. (Pingali., 2000). It is most significant pests in Asia and Africa. (Arabjafari and Jalali., 2007). Approximately 139 distinct insect pest species attack maize; among these, *Chilo partellus* is a major pest in various agroclimatic zones of India. (Shukla and Kumar, 2005). It is very difficult to control the *Chilo partellus*, because of cryptic and nocturnal habits of adult moths (Singh *et al.*, 2014).

Chilo partellus lay oval-shaped, creamy white eggs of around 0.8 mm long. It takes 4–8 days for the eggs to develop into larvae and takes 28–35 days for the larvae to develop into pupa. Larvae in their final instar measure 25–30 mm long, and their bodies have rows of dark spots. Pupae are long, cylindrical and dark brown colored, males are smaller than females. The pupal period is 5-12 days. The moths are pale brown colored with an approximate wingspan of 20-30 mm. These moths have 3–8 days lifespan. When they attain maturity, they mate and lay eggs. The life cycle of *Chilo partellus* takes 25–50 days to complete (Panchal and Kachole, 2013). Since maize has more sugars and amino acids than other gramineous hosts, it is more vulnerable to harm from stem borer infestation (Souza,

2002). It starts to infest the crop three to four weeks after sowing and continues to do so until the crop reaches maturity. (Sarup *et al.*, 1978) The leaf-eating and stem-tunneling activity of the larvae is the most common symptom of *Chilo partellus* damage to maize plants. In a natural field, the first signs of infestation are characteristic leaf lesions and scarification caused by the first and second instars of *Chilo partellus* (Sithole, 1990). After hatching, stem borer larvae move over the plant, gather in the funnel, and feed on the curled leaves for a few days before approaching the stalk and stem (Mushore, 2005). When the infestation is severe, the larvae, either in the leaf whorl or in the stem, can cut through the meristematic tissues; the central leaves dry up to produce the 'dead heart' symptom (Groote, 2002). Exit holes and tunnels in the main stem inhibit plant growth and promote bacterial and fungal diseases (Songa *et al.*, 2001). Dead hearts reduce translocation, ear damage, lodging, initial leaf senescence and in severe cases complete crop failure (Naz *et al.*, 2003, Gupta *et al.*, 2010). The yield losses exhibit significant regional variations, with a range of 25-40% depending on the pest population density and crop phenological stage of infection (Khan *et al.*, 2015).

2. MATERIALS AND METHODS

The study was conducted during Kharif, 2023 at Central Research farm, SHUATS, Prayagraj, Uttar Pradesh, India. Experimental design employed was a Randomized Block Design (RBD) consisting of eight different treatments including untreated control, each being replicated thrice with a plot size of was 2m×1m and treatments were assigned randomly. The treatments included azadirachtin 10,000ppm(1ml/l), neem oil 3% (30ml/l), karanj oil 3% (30ml/l), neem oil 3% + imidacloprid 17.5%SL (30ml/l+0.25ml/l), karanj oil 3% + imidacloprid 17.5%SL (30ml/l+0.25ml/l), imidacloprid 17.5%SL (0.5ml/l), spinosad 45%SC (0.5ml/l) and untreated control. Two sprays were carried out at an interval of 15 days. Five plants were randomly selected in each treatment and observations were taken one day before spraying and three, seven and fourteen-days after spraying. Insecticides and biopesticides were applied at their recommended doses at economic threshold level (ETL = 10% infestation). (Iqbal *et al.*, 2017). The healthy marketable yield from different treatments were collected separately and weighed. The treatment cost and common cost of cultivation per hectare was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price; while the net benefit was obtained by subtracting the total cost of plant protection from total income.

3. RESULTS AND DISCUSSION

3.1 Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* after first spray:

The data on the larval population of spotted stem borer on 3rd, 7th and 14th days after first spray revealed that all the treatments were significantly superior over control. Among the treatments, the lowest larval population of *Chilo partellus* was recorded in karanj oil 3% + imidacloprid 17.5%SL (2.48) followed by neem oil 3% + imidacloprid 17.5%SL (2.91), imidacloprid 17.5%SL (3.13), spinosad 45%SC (3.66), neem oil 3% (4.20), karanj oil 3% (4.44) and azadirachtin 10,000ppm (5.00) which were found to be less effective when comparing all the treatments, which were significantly superior over the control (6.93). Among all the treatments (karanj oil + imidacloprid, neem oil + imidacloprid, imidacloprid), (imidacloprid, spinosad), (spinosad, neem oil), (neem oil, karanj oil) and (karanj oil, azadirachtin) were found statistically significant at par with each other.

3.2 Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* after second spray:

The data on the larval population of spotted stem borer on 3rd, 7th and 14th DAS days after the second spray revealed that all the treatments were significantly superior over control. Among all the treatments lowest larval population of spotted stem borer was recorded in karanj oil 3% + imidacloprid 17.5%SL (1.50) followed by neem oil 3% + imidacloprid 17.5%SL (1.93), imidacloprid 17.5%SL (2.30), spinosad 45%SC (2.68), neem oil 3% (3.26), karanj oil 3% (3.46) and azadirachtin 10,000ppm (3.88) were found to be less effective when comparing all the treatments, which were significantly superior over the control (7.55). Among all the treatments (neem oil and karanj oil) were found statistically significant at par with each other.

3.3 Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* after first and second spray (Overall mean):

The data on the larval population of spotted stem borer on mean (3rd, 7th and 14th DAS) days after first and second spray revealed that all the treatments were significantly superior over control. Among all the treatments lowest larval population of spotted stem borer was recorded in karanj oil 3% + imidacloprid 17.5%SL (1.99) followed by neem oil 3% + imidacloprid 17.5%SL (2.42), imidacloprid 17.5%SL (2.71), spinosad 45%SC (3.17), neem oil 3% (3.73), karanj oil 3% (3.95) and azadirachtin 10,000ppm (4.44) is found to be least

effective then all the treatments and was significantly superior over the control (7.24). Among all the treatments (karanj oil + imidacloprid, neem oil + imidacloprid, imidacloprid), (neem oil + imidacloprid, imidacloprid, spinosad), (spinosad, neem oil, karanj oil) and (neem oil, karanj oil azadirachtin) were found statistically significant at par with each other.




Figure 1. Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* (Swinhoe) after first and second spray

3.4 Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* on maize yield and cost benefit ratio:

The highest yield was recorded in karanj oil 3% + imidacloprid 17.5%SL with 39.20 q/ha followed by neem oil 3% + imidacloprid 17.5%SL with 36.50 q/ha, imidacloprid 17.5%SL with 34.40 q/ha, spinosad 45%SC with 32.50 q/ha, neem oil 3% with 30.70 q/ha, karanj oil 3% with 29.85 q/ha, azadirachtin 10,000ppm with 27.40 q/ha and untreated control 19.35 q/ha were recorded which was least yield.

Among all the treatments studied, the best and most economic treatment was imidacloprid 17.5%SL with cost benefit ratio of 1:2.2, followed by karanj oil 3% + imidacloprid 17.5%SL (1:2.0), spinosad 45%SC (1:1.9), neem oil 3% + imidacloprid 17.5%SL (1:1.7), azadirachtin 10,000ppm (1:1.7), karanj oil 3% (1:1.6), neem oil 3% (1:1.5) and untreated control (1:1.4).

UNDER PEER REVIEW

Table 1. Comparative study of synthetic insecticides with biopesticides against spotted stem borer, *Chilo partellus* on Maize

S. No.	Treatments	Dosage	Larval Population of <i>chilo partellus</i> / 5 plants										Yield (q/ha)	C: B Ratio	
			First spray					Second spray							Overall mean
			1DBS	3DAS	7DAS	14DAS	Mean	3DAS	7DAS	14DAS	Mean				
T ₁	Azadirachtin (10,000 ppm)	1 ml/lit	5.80	5.60 ^b	4.40 ^b	5.00 ^b	5.00 ^b	4.13 ^b	3.60 ^b	3.93 ^b	3.88 ^b	4.44 ^b	27.40	1:1.7	
T ₂	Neem oil (3%)	30 ml/lit	5.93	5.06 ^c	3.26 ^d	4.26 ^c	4.20 ^{cd}	3.40 ^d	3.13 ^c	3.26 ^c	3.26 ^c	3.73 ^{bc}	30.70	1:1.5	
T ₃	Karanj oil (3%)	30 ml/lit	5.60	5.33 ^{bc}	3.66 ^c	4.33 ^c	4.44 ^{bc}	3.73 ^c	3.26 ^{bc}	3.40 ^c	3.46 ^c	3.95 ^{bc}	29.85	1:1.6	
T ₄	Neem oil (3%) + Imidacloprid (17.8SL)	30ml/lit + 0.25 ml/lit	5.80	3.86 ^e	2.13 ^e	2.73 ^e	2.91 ^f	2.20 ^f	1.66 ^e	1.93 ^f	1.93 ^f	2.42 ^{de}	36.50	1:1.7	
T ₅	Karanj oil (3%) + Imidacloprid (17.8SL)	30ml/lit + 0.25 ml/lit	6.06	3.40 ^f	1.73 ^f	2.33 ^f	2.48 ^f	1.66 ^g	1.33 ^e	1.53 ^g	1.50 ^g	1.99 ^e	39.20	1:2.0	
T ₆	Imidacloprid (17.8SL)	0.5 ml/lit	5.73	4.06 ^e	2.33 ^c	3.00 ^e	3.13 ^{ct}	2.46 ^f	2.13 ^d	2.33 ^e	2.30 ^e	2.71 ^{de}	34.40	1:2.2	
T ₇	Spinosad (45 SC)	0.5 ml/lit	5.73	4.46 ^d	3.00 ^d	3.53 ^d	3.66 ^{de}	2.86 ^e	2.46 ^d	2.73 ^d	2.68 ^d	3.17 ^{cd}	32.50	1:1.9	
T ₈	Control		6.06	6.60 ^a	6.93 ^a	7.26 ^a	6.93 ^a	7.40 ^a	7.60 ^a	7.66 ^a	7.55 ^a	7.24 ^a	19.35	1:1.4	
F-Test			NS	S	S	S	S	S	S	S	S	S			
S.Ed.(±)			NS	0.15	0.18	0.13	0.32	0.13	0.13	0.16	0.13	0.81			
CD (0.05)			NS	0.32	0.38	0.28	0.67	0.28	0.28	0.33	0.28	0.95			

DBS- Day before Spraying, DAS- Day after Spraying, NS- Non significant, S- Significant

4. CONCLUSION: *Chilo partellus*, a lepidopteran stem borer, damages the worldwide maize ecosystem economically. Due of its cryptic nature of feeding, it is challenging to control with synthetic insecticides. The excessive use of synthetic insecticides and its associated detrimental effects have been widely criticized. Insects quickly develop resistance to synthetic insecticides when they are frequently exposed to them but biopesticides contain a variety of naturally occurring active ingredients with unique modes of actions, such as antifeedant, repellent, oviposition deterrent and synergistic effects, making resistance difficult to develop. Therefore, when synthetic insecticides and biopesticides are used together, they have substantially greater efficacy than when they are applied separately because of their synergistic and complimentary effects to each other. In present study, karanj oil 3% + imidacloprid 17.5%SL followed by neem oil 3% + imidacloprid 17.5%SL shows highest efficacy with highest yield 39.20 and 36.50 q/hectare respectively against *Chilo partellus* but economically best treatment was imidacloprid (17.8SL) which shows highest cost benefit ratio (1:2.2). Combinations of synthetic insecticides with biopesticides have lower cost benefit ratio to some extent compared to synthetic insecticides because of the high dosages and expense of biopesticides but it considered as a feasible alternative for synthetic insecticides and reduces adverse acute and chronic effects in environment along with bioaccumulation and biomagnifications.

REFERENCES

- Agricultural market intelligence center, Professor Jayashankar Telangana State Agricultural University (PJTSAU), Maize outlook (2024).
- Arabjafari, K. H. and Jalali, S. K. (2007). Identification and analysis of host plant resistance in leading maize genotypes against spotted stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera:Pyralidae). *Pakistan Journal of Biological sciences*, **10**(11): 1885-1895
- Chouraddi, M. and Mallapur, C.P. (2017). Impacts of newer molecules of insecticides on management of maize stem borers. *Journal of Entomology and Zoology Studies*, **5**(4) 1424-1428
- Groote, H. (2002). Maize yield losses from stem borers in Kenya. *International Journal of Tropical Insect Science*, **22**:89-96.
- Directorate of Maize Research, ICAR, 01-03-2024

Gupta, S., Handore, K. and Pandey, I.P. (2010). Effect of insecticides against *Chilo partellus* (Swinhoe) damaging *Zea mays* (maize). *International Journal of Parasitology Research*, **2**(2): 4-7.

Indian Institute of Maize Research, (2024)

Iqbal J, Farooq SU, Alqarni AS, Ali H, Zeshan A, Ansari MJ. Management of maize stem borer (*Chilo partellus*) with insecticides of three formulations under field conditions. *Indian Journal of Agricultural Sciences*, 2017;87(12):1720- 1724.

Joshi, M. Text Book of Field Crops. New Delhi: PHI Learning Private Limited, 2015.

Kauma, T.M., Schulthess, F., LeRu, B.P., Mueke, J., Ogwang, J.A. and Omwega C.O (2008) Abundance and diversity of lepidopteran stemborers and their parasitoids on selected grasses in Uganda, *Crop Protection*. **27**: 505-513.

Khan, I.A., Khan, M., Akbar, R., Saeed, M., Farid, A., Ali., Ijaz. and Alam, M. (2015). Assessment of different control methods for the control of Maize Stem Borer, (*Chilo partellus*) Swinhoe in Maize crop. *Journal of Entomology and Zoology Studies*, **3**(4):327-330.

Mushore, K. (2005). Assessment of suitability of different populations of stemborer species for the development of *Cotesia flavipes* (Hymenoptera: Braconidae) and the Establishment of the latter in Zimbabwe, Thesis, University of Zimbabwe, p 14.

Naz, F., Faridullah, M. and Din, M. (2003). Insect pest of maize and their losses. *Asian Journal of Plant Science*, **2**(5): 412-414.

Panchal, B.M. and Kachole, M.S. (2013). Life cycle of *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae) on artificial diets. *International Journal of Plant and Environmental Science* **3**(4): 19-22.

Panwar, V.P.S. (2005). Management of Maize stalk borer *Chilo partellus* (Swinhoe) in maize. *Stresses on Maize in Tropics*. pp. 324-375

Pingali, P.I. (2000). World Maize Facts and Trends Meeting World Maize Need: Technological Opportunities and Priorities for the Public Sector. International Maize and Wheat Improvement Center, Mexico. D.F. 57p.

Prasanna, B.M., Vasal, S.K., Kassahun, B. and Singh, N. N. (2001). Quality protein maize. *Current Science*, **81**(10): 1308–1318.

- Sarup, P., Siddiqui, K.H. and Marwaha, K.K. (1978). Trends in maize pest management research in India together with bibliography. *Journal of Entomological Research*, **111**:19-68.
- Shiferaw, B., Prasanna, B., Hellin, J. and Banziger, M. (2011). Crops that feed the world. Past successes and future challenges to the role played by maize in global food security. *Food Security*, **3**, 307–327.
- Shukla, A and Kumar, A. (2005). Maize stem borer (*Chilo partellus* Swinhoe). A review of plant protection Bulletin, University of Agriculture and Technology, Udaipur, India
- Singh, S.K., Chauhan, S.K., Singh, R.P., Singh, R.K. and Singh, R.B. (2006). Studies on abiotic factors related to yield gaps across the environment in tropical maize (*Zea mays* L.). *Analysis of Agricultural Research*, **27**(4):373-376.
- Singh, A., Dwivedi, S.A., Sharma, R.C. and Mishra, P.K. (2014). Management of stem borer, *chilo partellus* swinhoe on maize crop *Society for Scientific Development in Agriculyure and Technology Progressive Research*, **9** (2): 284-286.
- Sithole, S.Z. (1990). Status and control of *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae) in southern Africa. *Insect Science and Its Application* **11**: 481-488.
- Souza, F.S., Rebeiro, R.H.E., Veloso, C.A.C. and Correan, L.A. (2002). Yielding and phenotypic stability of corn cultivars in three municipal districts of parastate Brazil *Pesquisa-Agrepercuria Brasileira*, **37**: 1269-1274
- Songa, J.M., Guofa, Z. and Overhalt, W. A. (2001). Relationships of stem borer damage and plant physical conditions to maize yield in a semi-arid zone of Eastern Kenya. *International Journal of Tropical Insect Science*, **21**(3): 224-43.