

ASSESSMENT OF PLANT DIVERSITY AND SOIL QUALITY MANAGEMENT IN FOREST LAND OF WEST GODAVARI DISTRICT, ANDHRA PRADESH, INDIA

Abstract

The field experiment was set in home-garden areas under forest land of eight small towns viz. Borampalem, Venkatapuram, Gundugolanukunta, Ramannapalem, Abdullapuram, Tirumaladevpetta, Venkatakrishnapuram, Appalarajugudem in West Godavari, Andhra Pradesh during November to March in 2021–22 to assess the Plant and Tree Diversity and Soil Quality parameters. The plant/tree species *Phoenix sylvestris*, reported the highest frequency (245.00), dominance (3.27), and basal area (7542.96 cm²). *Plumbago zeylanica* had the highest relative density (6.96) and relative abundance (6.81) among all the shrub species. *Plumbago zeylanica* reported the highest Important value index (IVI) 17.10 while Herb species with highest density was found in *Centella asiatica* (0.95), frequency (35), and highest Important value index (IVI) 14.60. Soil quality parameters reflect that at the depth 0-15 cm the average value of pH is found to be 7.52, organic carbon is 0.56%, the amount of available nitrogen is 213.78 kg/ha, amount of available phosphorus is 41.25 kg/ha and available potassium is 38.22 kg/ha. At the depth of 15-30 cm the average value of pH is found to be 6.61, organic carbon is 0.39%, the amount of available nitrogen is 189.51 kg/ha, amount of available phosphorus is 34.75 kg/ha and available potassium is 31.37 kg/h.

Keywords: Agro-forestry models, Forest land, Plant and Tree diversity, Soil quality, Relative Density, West Godavari District.

INTRODUCTION:

Since trees make up a significant portion of the vegetation, it is vital to continuously monitor and manage them to ensure that the diversity of species and habitats is maintained through successional processes (Turner 1987; Attua and Pabi 2013). Diversity in tree species is essential to tropical forest biodiversity (Evariste et al., 2010) and a significant component of forest ecosystem diversity (Rennolls and Laumonier, 2000; Tchouto et al., 2006). To track the dynamics of tropical forests over time and evaluate the impact of disturbance and climate change on plant demography, tree census plots have been set up in a variety of forest types (Condit et al 1996; Laurance et al 2004; Mohandass and Davidar 2009). The diversity of tree species affects the climate, stand structure, species composition, and geomorphology of the forest. Understanding the structure of forest stands is crucial for understanding forest ecosystems, since it plays a significant role in the biodiversity of stands (Ozcelik 2009). Our capacity to maximize biodiversity conservation as a result of deforestation and degradation will be strengthened by an immediate inventory of tree species that offers information on diversity (Baraloto et al 2013). Plant species diversity is complex in nature, and its structure and composition vary depending on climatic conditions and topography (Raturi 2012). Tropical forests are the most complex of all terrestrial ecosystems and the most heavily exploited ecosystems in the biosphere (Bahuguna 1999). It only accounts for 7% of the earth's land surface yet is

home to more than half of the world's plant and animal variety (Wilson 1988). Despite their direct benefits to human survival, they are disappearing at a rate of 0.8 to 2% every year (May & Stumpf 2000, Sagar *et al.* 2003), and dry deciduous forests in particular are the most damaged and least protected ecosystems on the planet (Murphy & Lugo 1986). Biodiversity is still declining despite a national policy focused at preserving and enhancing nature. The rapid decline of tropical forests worldwide has been accelerated by habitat destruction, deforestation, human settlements, globalization, agricultural expansion, and other development-related infrastructure over the last century. These factors have adversely affected biodiversity, climate change, ecological services, soil productivity, and the livelihoods of both forest dwellers and those living in rural areas (Myers 1992, Raghubanshi & Tripathi 2009). Considering the importance of plant diversity and their role in improvement of soil quality parameters, present study was conducted and findings are presented in this paper.

MATERIALS AND METHODS

Experimental areas are situated on the southeast coast of the nation, alongside the Bay of Bengal, is Andhra Pradesh, the eighth largest state in the nation. The State's geographical area is 1,62,968 square kilometers, or 4.96% of the nation's total area. One of Andhra Pradesh's most significant districts, the West Godavari, is located between 16° 15' 00" and 17° 30' 00" of north latitude and between 80° 50' 00" and 81° 55' 00" of east longitude. It covers an area of 8506 sq. km with a coastline that is roughly 23.00 km.




Fig1: Experimental areas of West Godavari district, Andhra Pradesh

Analysis of the plant and tree diversity and Analysis of data

Quadrat methods used for collection of data for analysis of parameter of plant and tree diversity. In this species are enumerated within random quadrat of different size as per on the size and nature of plant community. 10m, x10m., 5mX5m and 1mx1m for trees, shrubs and herbs respectively were laid out at selective sites. Nearly all types of vegetation can be quantified using the quadrat approach to determine the plant community. The numbers and names of each plant recorded when the sample pits are created. The frequency, density, and dominance of the flora were then determined and evaluated by formula suggested by Curtish and McIntosh 1950. The important value index, relative density, relative dominance, and relative frequency were analysed using formula suggested by Philips 1959. The following formulas were used to determine frequency, density, and dominance.

$$\text{Density} = \frac{\text{Total number of individuals of the species}}{\text{Total number of sample units studied}}$$

$$\text{Frequency} = \frac{\text{Number of sample plots in which species occurred}}{\text{Total number of sample plots studied}} \times 100$$

$$\text{dominance} = \frac{\text{Total cover basal area}}{\text{Total area sampled}} \times 100$$

Basal area:

The area that a tree's base occupies is thought to be a reliable measure of the size, weight, or volume of the tree. The formula was used to determine the basal area.

$$BA = Cbh^2/4\pi,$$

Where, Cbh diameter measured at breast height

Important Value Index

A measure of a species' dominance in a particular study area is called an important value. Foresters use it as a common tool to inventory a forest. Instead of counting every tree in a forest, foresters typically identify key locations inside the forest and sample a predetermined area surrounding those locations. Relative frequency, relative density, and relative dominance are the three types of data that are gathered; each of these values is expressed as a percent and spans from 0 to 100. The sum of these three measurements is the significant value, which can be anywhere between 0 and 300.

The value of the Important Value Index (IVI) was 300. It covers three essential components.

$$\text{Relative Density} = \frac{\text{Total number of individuals of the species}}{\text{Total number of sample units studied}} \times 100$$

$$\text{Relative Frequency} = \frac{\text{Number of sample plots in which species occurred}}{\text{Total number of sample plots studied}} \times 100$$

$$\text{Relative dominance} = \frac{\text{Total cover basal area}}{\text{Total area sampled}} \times 100$$

Simpson's index (1949): It is calculated by a particular formula, which examines the species dominance of a specific area of study:

$$Cd = \pi(n_i/N)^2$$

n_i = Total number of the individuals in each species

N = Total number of the individuals of all species

Shannon–wiener index (1963): Whenever assessing the species diversity within a community, the Shannon-Weiner index (H) is often utilised. Shannon's Index takes into consideration the evenness and richness of the species that are there. It's calculated by using this formula.

$$H = -\sum (n_i/N) \log(n_i/N)$$

n_i = The overall number of individuals in the collective that belong in the i^{th} species

Soil sample analysis

Eight locations in the West Godavari District provided soil samples, which have been gathered for physical-chemical research at depths of 0–15 cm and 15–30 cm. After the samples were correctly mixed, let to air dry, crushed, and sieved through a 2-mm screen to remove any remaining foreign material, the samples were stored in preparation for their physical and chemical analysis. The mechanical characteristic of the soil is determined by using Bouyoucos Hydrometer method (1965) and physical property is done by cylindrical methods (Muthuvel *et al.* 1992). Soil parameter analysed for pH and EC (dSm⁻² at 25° C or mmho/m at 25° C) % organic carbon, available Nitrogen (N) kg/ha, Phosphorous (P₂O₅) kg/ha, Potassium (K₂O) kg/ha).

Chart 1. Physio–Chemical properties of the Soil

S.No.	Parameter	Method	Reference
1	pH	Digital pH meter	Jackson, 1973
2	EC	Method no.4 USDA Handbook no.16	Richards, 1954
3	Organic Carbon	Walkley and Blackman Method	Walkley and Black 1947
4	Available Nitrogen.	Alkaline pomegranate Method	Subbiah and Asija, 1954
5	Available Phosphorous	Olsen's Colorimetric Method	Olsen <i>et al.</i> 1994
6	Available Potassium	Flame Photometric Method	Toth and Prince, 1949

S.NO	Nameofplants	Total no. of individuals	Density	Frequency	BasalArea (cm ²)	Abundanc e	RelativeDe nsity	RelativeFreq uency	RelativeDom inance	IVI
------	--------------	--------------------------	---------	-----------	------------------------------	------------	------------------	--------------------	--------------------	-----

RESULTSANDDISCUSSION

Table 1 shows that 239 individuals in all, representing 41 tree species across 35 groups, were taken into consideration. Among all the species, *Phoenix sylvestris* (49) followed by *Salix babylonica*(14), *Tamarindus indica*(10), *Lanneacoromandelica*(8) reported the maximum number of individuals while *Azadirachta indica* (2), *Moringa oleifera* (3), reported with the least number of individuals was *Terminalia elliptica* reported the smallest basal area (314.16 cm²), while *Phoenix sylvestris* recorded the highest basal area (7542.96). *Phoenix sylvestris* was found to have the highest density (2.45) *Sterculia urens* has the smallest density (0.1). *Phoenix sylvestris* had the highest frequency (245.00), while *Sterculia urens* had the lowest frequency (10.00). *Phoenix sylvestris* has the highest Dominance (3.27), while *Sterculia urens* had the lowest (0.25). The relative density was highest in *Phoenix sylvestris* (20.33) followed by *Salix babylonica*(5.81) whereas the lowest was *Moringa oleifera* (0.83). The relative frequency was found to be the highest in *Phoenix sylvestris*(20.50) whereas the least was reported in *Moringa oleifera* (0.84). The species with the highest Important value index (IVI) was *Phoenix sylvestris* (50.25) whereas the least was *Terminalia elliptica*(2.89).

1.	<i>Azadirachta indica</i>	2	0.2	20	1385.44	1.33	1.66	1.67	1.73	5.06
2.	<i>Bombax cieba</i>	7	0.35	35	804.25	2.33	2.90	2.93	1.00	6.84
3.	<i>Ficus benghalensis</i>	6	0.3	30	1590.43	2.00	2.49	2.51	1.99	6.99
4.	<i>Mangifera indica</i>	4	0.2	20	490.87	1.33	1.66	1.67	0.61	3.95
5.	<i>Psidium guajava</i>	6	0.3	30	706.86	2.00	2.49	2.51	0.88	5.88
6.	<i>Prosopis cineraria</i>	3	0.15	15	490.87	1.00	1.24	1.26	0.61	3.11
7.	<i>Ficus religiosa</i>	6	0.3	30	1256.64	3.00	2.49	2.51	1.57	6.57
8.	<i>Tamarindus indica</i>	10	0.5	50	1134.11	2.00	4.15	4.18	1.42	9.75
9.	<i>Eucalyptus globulus</i>	3	0.15	15	3318.31	1.50	1.24	1.26	4.14	6.64
10.	<i>Elaeis guineensis</i>	7	0.35	35	4417.86	1.75	2.90	2.93	5.52	11.35
11.	<i>Coccoloba nucifera</i>	4	0.2	20	1963.50	2.00	1.66	1.67	2.45	5.79
12.	<i>Phoenix sylvestris</i>	49	2.45	245	7542.96	3.27	20.3	20.50	9.42	50.25
13.	<i>Jasminum spp</i>	7	0.35	35	1590.43	1.75	2.90	2.93	1.99	7.82
14.	<i>Moringa oleifera</i>	2	0.1	10	1256.64	1.00	0.83	0.84	1.57	3.24
15.	<i>Indian mahogany</i>	4	0.2	20	3848.45	1.33	1.66	1.67	4.81	8.14
16.	<i>Terminalia arjuna</i>	4	0.2	20	1963.50	2.00	1.66	1.67	2.45	5.79
17.	<i>Shorea robusta</i>	4	0.2	20	4071.50	1.33	1.66	1.67	5.08	8.42
18.	<i>Delonix regia</i>	3	0.15	15	3318.31	1.50	1.24	1.26	4.14	6.64
19.	<i>Saraca asoca</i>	3	0.15	15	5026.55	1.50	1.24	1.26	6.28	8.78
20.	<i>Murrayakoenigii3</i>	6	0.3	20	1134.11	1.50	2.49	1.67	1.42	5.58
21.	<i>Leucaena leucocephala</i>	3	0.15	15	804.25	1.00	1.24	1.26	1.00	3.50
22.	<i>Terminalia catappa</i>	3	0.15	15	1385.44	1.00	1.24	1.26	1.73	4.23
23.	<i>Carica papaya</i>	6	0.3	30	1590.43	1.50	2.49	2.51	1.99	6.99
24.	<i>Anacardium occidentale</i>	4	0.2	20	1809.56	1.33	1.66	1.67	2.26	5.59
25.	<i>Acacia nilotica</i>	6	0.3	30	1075.21	1.50	2.49	2.51	1.34	6.34
26.	<i>Dalbergia latifolia</i>	3	0.15	15	2922.47	1.50	1.24	1.26	3.65	6.15
27.	<i>Phyllanthus emblica</i>	5	0.25	25	2290.22	1.25	2.07	2.09	2.86	7.03
28.	<i>Pongamia pinnata</i>	4	0.2	20	1590.43	2.00	1.66	1.67	1.99	5.32
29.	<i>Salix babylonica</i>	14	0.7	70	2827.43	2.33	5.81	5.86	3.53	15.20
30.	<i>Pterocarpus</i>	5	0.25	25	3848.45	1.25	2.07	2.09	4.81	8.97
31.	<i>Terminalia bellirica</i>	4	0.2	20	706.86	1.33	1.66	1.67	0.88	4.22
32.	<i>Wrightia tinctoria</i>	3	0.15	15	660.52	1.33	1.24	1.26	0.82	3.32
33.	<i>Sterculia aurens</i>	4	0.2	20	1963.50	0.25	1.66	1.67	2.45	5.79
34.	<i>Madhuca longifolia</i>	5	0.25	25	1452.20	0.29	2.07	2.09	1.81	5.98
35.	<i>Bauhinia racemosa</i>	6	0.3	30	1590.43	1.20	2.49	2.51	1.99	6.99
36.	<i>Senegalia chundra</i>	3	0.15	15	1075.21	1.00	1.24	1.26	1.34	3.84
37.	<i>Boswellia ovalifoliolata</i>	3	0.15	15	660.52	1.00	1.24	1.26	0.82	3.32
38.	<i>Lannea coromandelica</i>	8	0.4	40	1017.88	1.33	3.32	3.35	1.27	7.94
39.	<i>Alstonia scholaris</i>	4	0.2	20	1734.94	2.00	1.66	1.67	2.17	5.50
40.	<i>Erythrina suberosa</i>	3	0.15	15	1452.20	1.50	1.24	1.26	1.81	4.31
41.	<i>Terminalia elliptica</i>	3	0.15	15	314.16	1.00	1.24	1.26	0.39	2.89
	Total	239	12.05	1195	80083.9	61.7	100	100	100	300

Table 1 Phyto-sociological analysis of trees in 8 villages of West Godavari District, at Andhra

Table 2 shows that a total number of 230 individuals belonging to 30 shrubs species and 19 families

reported in study areas. *Plumbago zeylanica* had the highest relative density (6.96) while *Carissa spinarum*, *Maytenusemarginata* had the least (1.30). The relative frequency was found to be highest in *Clerodendrum phlomidis*(7.50), while it was lowest in *Carissa spinarum*, *Maytenusemarginata*, *Calotropis gigantea*(1.67). The relative abundance was highest in *Plumbago zeylanica* (6.81) while it was lowest in *Phyllanthus reticulatus*(2.38). *Plumbago zeylanica* reported the highest IVI(17.10), followed by *Clerodendrum phlomidis*(16.24) whereas *Carissa spinarum* reported the least Important value index (IVI)(5.53).

Table 2 Phyto-sociological analysis of Shrubs in 8 villages of West Godavari District at Andhra Pradesh

S.NO	Name of plants	Total no of Individual	Density	Frequency	Basal Area	Abundance	Relative Density	Relative frequency	Relative dominance	IVI
1.	<i>Memecylonumbellatum</i>	14	0.7	35	283.5	2.0	6.09	5.83	3.41	15.3
2.	<i>Phyllanthusreticulatus</i>	7	0.35	25	153.9	1.4	3.04	4.17	2.38	9.59
3.	<i>Piperhooglandii</i>	10	0.5	25	254.5	2.0	4.35	4.17	3.41	11.9
4.	<i>Plumbagozeylanica</i>	16	0.8	20	113.1	4	6.96	3.33	6.81	17.1
5.	<i>Ricinuscommunis</i>	8	0.4	20	314.2	2.0	3.48	3.33	3.41	10.2
6.	<i>Sennaalata</i>	7	0.35	15	415.5	2.3	3.04	2.50	3.97	9.52
7.	<i>Vitexnegundo</i>	10	0.5	25	113.1	2	4.35	4.17	3.41	11.9
8.	<i>Woodfordiafruticosa</i>	7	0.35	15	572.6	2.3	3.04	2.50	3.97	9.52
9.	<i>Maytenusemarginata</i>	3	0.15	10	201.1	1.5	1.30	1.67	2.55	5.53
10.	<i>Acaciafarnesiana</i>	7	0.35	20	346.4	1.75	3.04	3.33	2.98	9.36
11.	<i>Azimatetracantha</i>	8	0.4	15	314.2	2.7	3.48	2.50	4.54	10.5
12.	<i>Balaniesroxburghii</i>	6	0.3	15	254.5	2	2.61	2.50	3.41	8.51
13.	<i>Bambusavulgaris</i>	6	0.3	20	452.4	1.5	2.61	3.33	2.55	8.50
14.	<i>Calotropisgigantea</i>	5	0.25	10	380.1	2.5	2.17	1.67	4.26	8.10
15.	<i>Canthiumparviflorum</i>	6	0.3	20	490.9	1.5	2.61	3.33	2.55	8.50
16.	<i>Capparisaphylla</i>	8	0.4	20	283.5	2.0	3.48	3.33	3.41	10.2
17.	<i>Cappariszeylanica</i>	6	0.3	20	346.4	1.5	2.61	3.33	2.55	8.50
18.	<i>Carissaspinarum</i>	3	0.15	10	415.5	1.5	1.30	1.67	2.55	5.53
19.	<i>Catunaregamspinosa</i>	7	0.35	20	283.5	1.75	3.04	3.33	2.98	9.36
20.	<i>Clerodendrumphlomidis</i>	14	0.7	45	113.1	1.6	6.09	7.50	2.65	16.2
21.	<i>Dichrostachyscinerea</i>	5	0.25	10	452.4	2.50	2.17	1.67	4.26	8.10
22.	<i>Dodonaeviscosa</i>	6	0.3	20	201.1	1.5	2.61	3.33	2.55	8.50
23.	<i>Flacourtiaindica</i>	9	0.45	20	490.9	2.25	3.91	3.33	3.83	11.0
24.	<i>Gmelinaasiatica</i>	5	0.25	15	201.1	1.7	2.17	2.50	2.84	7.51
25.	<i>Helicteresisora</i>	12	0.6	35	572.6	1.7	5.22	5.83	2.92	13.9
26.	<i>Ixoraarborea</i>	7	0.35	20	346.4	1.8	3.04	3.33	2.98	9.36
27.	<i>Jatrophaurcas</i>	7	0.35	20	254.5	1.75	3.04	3.33	2.98	9.36

28.	<i>Justicia adhatoda</i>	6	0.3	15	254.5	2.0	2.61	2.50	3.41	8.51
29.	<i>Lantana camara</i>	6	0.3	15	254.5	2	2.61	2.50	3.41	8.51
30.	<i>Lawsonia inermis</i>	9	0.45	25	314.2	1.8	3.91	4.17	3.07	11.1
31.	Total	230	11.5	60	9443.6	58.7	100	100	100	300

Table 3 shows that a total number of 321 individuals belonging to 35 herb species and 25 families. *Centella asiatica* had the highest number of individuals (19) while the least was *Tephrosia purpurea* (2). *Centella asiatica* had the highest density (0.95) and *Tephrosia purpurea* had the lowest density (0.1). *Centella asiatica* (35) had the highest frequency, while *Commelinabenghalensis* had the lowest. *Withania somnifera*, *Tridax procumbens* (10). *Centella asiatica* reported the highest IVI (14.60) followed by *Indigofera trita* (12.80) whereas *Tephrosia purpurea* reported with the least Important value index (IVI) (3.35).

Table 3. Phytosociological analyses of Herbs in 8 villages of West Godavari District of Andhra Pradesh

S.NO	Name of plants	Total no. of individual	Density	Frequency	Abundance	Relative Density	Relative Frequency	Relative Dominance	IVI
1.	<i>Cassitoria</i>	8	0.4	15	2.7	2.49	2.46	2.89	7.85
2.	<i>Centella asiatica</i>	19	0.95	35	2.7	5.92	5.74	2.95	14.6
3.	<i>Cheilocostus speciosus</i>	9	0.45	15	3	2.80	2.46	3.26	8.52
4.	<i>Cleome viscosa</i>	11	0.55	20	2.75	3.43	3.28	2.98	9.69
5.	<i>Commelinabenghalensis</i>	5	0.25	10	2.5	1.56	1.64	2.71	5.91
6.	<i>Cyperus rotundus</i>	9	0.45	15	3	2.80	2.46	3.26	8.52
7.	<i>Dendrophthoe falcata</i>	10	0.5	20	2.5	3.12	3.28	2.71	9.11
8.	<i>Elytraria acaulis</i>	12	0.6	20	3	3.74	3.28	3.26	10.2
9.	<i>Hygrophila auriculata</i>	9	0.45	15	3	2.80	2.46	3.26	8.52
10.	<i>Hyptis suaveolens</i>	12	0.6	20	3	3.74	3.28	3.26	10.2
11.	<i>Indigofera trita</i>	16	0.8	30	2.7	4.98	4.92	2.89	12.8
12.	<i>Martynia annua</i>	8	0.4	15	2.7	2.49	2.46	2.89	7.85
13.	<i>Pentanema indicum</i>	7	0.35	20	1.75	2.18	3.28	1.90	7.36
14.	<i>Phyllanthus floridus</i>	12	0.6	20	3	3.74	3.28	3.26	10.2
15.	<i>Phyllanthus amarus</i>	6	0.3	15	2	1.87	2.46	2.17	6.50
16.	<i>Rauwolfia serpentina</i>	7	0.35	15	2.3	2.18	2.46	2.53	7.17
17.	<i>Senna obtusifolia</i>	3	0.15	10	1.5	0.93	1.64	1.63	4.20
18.	<i>Senna occidentalis</i>	11	0.55	20	2.75	3.43	3.28	2.98	9.69
19.	<i>Sennatoria</i>	3	0.15	10	1.5	0.93	1.64	1.63	4.20
20.	<i>Sesamum alatum</i>	7	0.35	15	2.3	2.18	2.46	2.53	7.17
21.	<i>Tephrosia purpurea</i>	2	0.1	10	1	0.62	1.64	1.09	3.35
22.	<i>Tridax procumbens</i>	7	0.35	10	3.5	2.18	1.64	3.80	7.62
23.	<i>Vandatesella</i>	13	0.65	30	2.17	4.05	4.92	2.35	11.3
24.	<i>Withania somnifera</i>	9	0.45	10	4.5	2.80	1.64	4.88	9.33
25.	<i>Acalypha indica</i>	13	0.65	30	2.17	4.05	4.92	2.35	11.3

26.	<i>Achyranthusaspera</i>	11	0.55	15	3.7	3.43	2.46	3.98	9.87
27.	<i>Acoruscalamus</i>	10	0.5	20	2.5	3.12	3.28	2.71	9.11
28.	<i>Aervalanata</i>	5	0.25	15	1.67	1.56	2.46	1.81	5.83
29.	<i>Agavaamericana</i>	11	0.55	20	2.75	3.43	3.28	2.98	9.69
30.	<i>Anisomelesindica</i>	8	0.4	15	2.7	2.49	2.46	2.89	7.85
31.	<i>Argemonemexicana</i>	4	0.2	15	1.3	1.25	2.46	1.45	5.15
32.	<i>Bacopamonneri</i>	10	0.5	15	3.3	3.12	2.46	3.62	9.19
33.	<i>Biophyumsensitivum</i>	13	0.65	20	3.25	4.05	3.28	3.53	10.8
34.	<i>Byttneriaberbaceae</i>	12	0.6	15	4	3.74	2.46	4.34	10.5
35.	<i>Xanthiumstrumarium</i>	9	0.45	15	3	2.80	2.46	3.26	8.52
36.	Total	321	16.0	610	92.1	100	100	100	300

Soil nutrient status

Table 4 Revealed that the chemical characteristics of various soil samples collected from the studied area is reported in the table1 and 2, respectively, at a depth of 0–15 and 15–30 cm. The chemical parameters are studied by the help of suitable methods. At the depth of 0-15 cm: The mean value of the organic carbon is (0.56%) which means the soil is comparatively fertile and good for the cultivation. The organic carbon is greater in the top soil due to continue mixing of leaf litters and high value of vegetation decomposition. The primary nutrient for vegetation growth and an important factor that determines the quality of the soil is nitrogen. The mean value of available Nitrogen in the soil is (213.78 kg/ha) which means that the nitrogen content in the soil is good enough. The mean value for the available Phosphorous is (41.25 kg/ha) it is quite low for that reason recommendable amount of SSP is to be used for the phosphorous. The mean value of the available potassium is (38.22 kg/ha) Potassium availability is generally low. The pH of study areas is slightly alkaline.

Table 4 Nutrient status of the soil sample at 0-15cm in WestGodavariDistrict of Andhra Pradesh

Name of the village	pH	EC(Ds/m)	Organic carbon (%)	N(kg/ha)	P(kg/ha)	K(kg/ha)
Borampallem	7.62	0.81	0.87	245.6	96	55.1
Venkatapuram	7.43	0.23	0.76	232.2	19	50
Gundugolanukunta	7.03	0.34	0.23	223.2	53	40.1
Ramannapalem	7.90	0.75	0.45	212.1	33	35.1
Abdullapuram	8.00	0.45	0.20	232.1	34	40.2
Tirumaladevipeta	7.24	0.29	0.81	152.1	20	35.1
Venkata krishnapuram	7.36	0.62	0.35	222.9	45	20.2
Appalarajugudem	7.62	0.55	0.81	190.1	30	30.0
Average	7.52	0.50	0.56	213.7	41.2	38.2

Soil analysis at the deep of 15-30cm

Table 5 revealed that as the organic carbon content in the deeper layer of soil (0.39%) was found to be lesser than top soil. Nitrogen content decrease with increases in soil depth and also due to decrease in the organic carbon the mean value of nitrogen at 15 – 30cm is (189.51 kg/ha). The same trends of decrease in available Phosphorous and Potassium (34.75 P kg/ha and 31.37 K kg/ha, respectively) was recorded with in soil depth. The decreased availability of organic matter in the soil's deeper layers may be the cause of its nutritional loss.

Table 5. Nutrient status of the soil sample at 15-30cm

Name of the village	pH	EC(Ds/m)	Organic carbon (%)	N(kg/ha)	P(kg/ha)	K(kg/ha)
Borampallem	6.36	0.7	0.80	230	85	49
Venkatapuram	7.2	0.15	0.61	220.1	15	45
Gundugolanukunta	6.7	0.3	0.18	190.1	50	30
Ramannapalem	6.24	0.71	0.39	180.2	31	25
Abdullapuram	6.66	0.42	0.10	200.2	25	35
Tirumaladevipeta	6.75	0.25	0.62	142.1	10	30
Venkata krishnapuram	6.5	0.59	0.32	170.9	40	15
Appalarajugudem	6.51	0.42	0.12	182.5	22	22
Avg	6.61	0.44	0.39	189.51	34.75	31.37




Fig.2pH of soil sample at 0-15cm and 15-30cm




Fig.3Electrical conductivity of soil sample at 0-15cm and 15-30cm




Fig.4 Organic carbon of soil sample at 0-15cm and 15-30cm



Fig.5 Nitrogen of soil sample at 0-15cm and 15-30cm




Fig.6 Phosphorus of soil sample at 0-15cm and 15-30cm




Fig.7. Potassium of soil sample at 0-15cm and 15-30cm

Conclusion:

Present research work concludes that villages of West Godavari district are mainly composed of 41 tree species belonging to 35 families. *Phoenix sylvestris* (7542.96) is the largest populous species with highest basal area, the highest density, frequency (245.00), and the highest dominance (3.27). Shrub species *Plumbago zeylanica* had the highest relative density (6.96), relative abundance (6.81) and the highest Important value index (17.10). Herbs species with highest density was found for *Centella asiatica* (0.95), highest frequency (35) and highest IVI (14.60). The diversity of the study regions shows the disturbance due to soil quality variation, environmental factors and human intervention. As the soil nutrient status shows the remarkable low content of phosphorous and potassium. On the basis of questionnaire survey, it can be concluded that the compensation of diversity losses due to human activity in the study region can be achieved by adopting various Agro-forestry model by the farmers especially home garden.

Reference

Attua, E.M. & Pabi, O. 2013. Tree species composition, richness and diversity in the northern forest- savanna ecotone of Ghana. *Journal of Applied Biosciences*, 69: 5437-5448.

Armenteras, D., Rodríguez, N., & Retana, J. (2009). Are conservation strategies effective in avoiding the deforestation of the Colombian Guyana Shield. *Biological Conservation*, 142, 1411–1419.

Bouyoucos, 1962 Hydrometer method improved for making particle size analysis of soils *Agron. J.*, 54 (1962), pp. 464-465

Baraloto C, Paine CET, Poorter L, Beauchene J, Bonal D, *et al.* (2010) Decoupled leaf and stem economics in rain forest trees. *Ecol Lett* 13: 1338–1347.

V.K. BAHUGUNA And A. UPADHYAY Forest fires in India: policy initiatives for community participation *The International Forestry Review* Vol, pp. 122-127 (6 pages)

Chittibabu, C.V., & Parthasarathy, N. (2000). Attenuated tree species diversity in human-impacted tropical evergreen forest sites at Kolli hills, Eastern Ghats, India. *Biodiversity and Conservation*, 9, 1493–1519.

Cintra, R., Ximenes, A.C., Gondim, F.R., & Kropf, M.S. (2005). Forest spatial heterogeneity and palm richness, abundance and community composition in Terra Firme Forest, Central Amazon. *Revista Brazilian Botany*, 28(1), 75–84.

Choudary 2018 Sustainable intensification influences soil quality, biota, and productivity in cereal-based agroecosystems *Applied Soil Ecology*, 126 (2018), pp. 189-198.

Condit R, Hubbell SP, Lafrankie JV, Sukumar R, Manokaran N, Foster RB, Ashton PS. Species-area and species-individual relationships for tropical trees: a comparison of three 50-ha plots. *Journal of Ecology* 1996;84:549-62.

Condit, R., Ashton, P.S., Baker, P., Bunyavejehwin, S., Gunatilleke, S., Gunatilleke, N., Yamakura, T. (2000).

- Spatial patterns in the distribution of tropical tree species. *Science*, 288, 1414–1418.
- Dierick, D., & Hölscher, D. (2009). Species-specific tree water use characteristics in reforestation stands in the Philippines. *Agricultural and Forest Meteorology*, 149, 1317–1326.
- Evariste, W. T., Hong, S. Q., Jie, K., & Tongxi, W. A. N. G. (2010). Design and simulation of Gaussian shaping amplifier made only with CMOS FET for FEE of particle detector, *Nuclear Science and Techniques* Vol.21, No 5, October 2010 P.312-315
- Galley, R.E. (2014). Ecology of tropical rain forests. In R.K. Monson (Ed.), *Ecology and the environment* New York, NY: *The Plant Sciences Springer*. 8, 247–272
- Gandhi, D.S., & Sundarapandian, S. (2014). Inventory of trees in tropical dry deciduous forests of Tiruvannamalai district, Tamil Nadu, India. *Biodiversitas, Journal of Biological Diversity*, 15(2), 169–179.
- Gonmadje, C.F., Doumenge, C., McKey, D., Tchouto, G.P.M., Sunderland, T.C.H., Malinga, M.P.B., & Sonke, B. (2011). Tree diversity and conservation value of Ngovayang's lowland forests, Cameroon. *Biodiversity Conservation*, 20, 2627–2648.
- Gopalakrishna, S.P., Kaonga, M.L., Somashekar, R.K., Suresh, H.S., & Suresh, R. (2015). Tree diversity in the tropical dry forest of Bannerghatta National Park in Eastern Ghats, southern India. *European Journal of Ecology*, 1(2), 12–27.
- Gordon, J.E., & Newton, A.C. (2006). The potential misapplication of rapid plant diversity assessment in tropical conservation. *Journal for Nature Conservation*, 14, 117–126.
- Gunaga, S., Rajeswari, N., & Vasudeva, R. (2013). Tree diversity and disturbance of Kaan forests: Relics of a community protected climax vegetation in the central Western Ghats. *Tropical Ecology*, 54, 117–131.
- Hitimana, J., Kiyiapi, J.L., & Njunge, J.T. (2004). Forest structure characteristics in disturbed and undisturbed sites of Mt. Elgon moist low montane forest, Western Kenya. *Forest Ecology and Management*, 194, 269–291.
- Huang, W., Pohjonen, V., Johansson, V., Nashanda, M., Katigula, M.I.L., & Luukkanen, O. (2003). Species diversity, forest structure and species composition in Tanzanian tropical forests. *Forest Ecology and Management*, 173, 111–124.

J. T. Curtis and R. P. McIntosh the Interrelations of Certain Analytic and Synthetic Phytosociological Characters. *Ecology* Vol. 31, No. 3 (Jul., 1950), pp. 434-455 (22 pages)

Jackson. 1969. Soil Chemical Analysis: Advanced Course: a Manual of Methods Useful for Instruction and Research in Soil Chemistry, Physical Chemistry of Soils, Soil Fertility, and Soil Genesis. Quantitative determination of quartz in soils, sediments, and rocks by pyrosulfate fusion and hydrofluosilicic acid treatment. *Soil Sci.* Appears in 7 books from 1896-2005

- Kumar, J.I.N., Kumar, R.N., Bhoi, R.K., & Sajish, P.R. (2010). Tree species diversity and soil nutrient status in three sites of tropical dry deciduous forest of western India. *Tropical Ecology*, 51, 273–279.
- Kumar 2009 The effects of disturbance on forest structure and diversity at different altitudes in Garhwal Himalaya *Chi Journal Ecol*, 28 (3), 424-432.
- Lalfakawma, S.U.K., Roy, S., Vanlalhrithpuia, K., & Vanlalhluna, P.C. (2009). Community composition and tree population structure in undisturbed and distributed tropical semi-evergreen forest stands of north-east India. *Applied Ecology environmental research*, 7, 303-318.

Laurance WF, Oliveira AA, Laurance SG, Condit R, Nascimento HEM, Sanchez-Thorin AC, Lovejoy TE, Andrade A, D'Angelo S, Ribeiro JE. Pervasive alteration of tree communities in undisturbed Amazonian forests. *Nature* 2004;428(6979):171-5

May RM, Stumpf MPH (2000). Species-area relationships in tropical forests. *Science* 290(5499):2084-2086

Myers N (1992) Population/environment linkages: discontinuities ahead. *Ambio* 21: 116–118.

Majumdar, K., Shankar, U., & Datta, B.K. (2012). Tree species diversity and stand structure along major community types in lowland primary and secondary moist deciduous forests in Tripura, Northeast India. *Journal of Forestry Research*, 23(4), 553–568.

Mani, S., & Parthasarathy, N. (2006). Tree diversity and stand structure in inland and coastal tropical dry evergreen forests of peninsular India. *Current Science*, 90, 1238–1246.

Margules, C.R., Pressey, R.L., & Williams, P.H. (2002). Representing biodiversity: Data and procedures for identifying priority areas for conservation. *Journal of Biosciences*, 27(4), 309–326.

Mohandass, D., & Davidar, P. (2009). Floristic structure and diversity of a tropical montane evergreen forest (shoal) of the Nilgiri Mountains, southern India. *Tropical Ecology*, 50(2), 219–229.

P G Murphy, and A E Lugo ECOLOGY OF TROPICAL DRY FOREST Vol. 17:67-88 (Volume publication date November 1986)

Naidu, M.T., & Kumar, O.A. (2015). Tree species diversity in the Eastern Ghats of northern Andhra Pradesh, India. *Journal of Threatened Taxa*, 7(8), 7443-7459.

Nath, P.C., Arunachalam, S., Khan, M.L., Arunachalam, K., & Barbhuiya, A.N. (2005). Vegetation analysis and tree population structure of tropical wet evergreen forests in and around Namdapha National Park, northeast India. *Biodiversity Conservation*, 14, 2109–2135.

Nagarjuna, A. 2009. Vertical gradient and resource partitioning of rainfall. Theoretical and Applied Climatology, migratory birds on Barringtonia tree in Nelapattu 92(1-2): 31-45. bird sanctuary. *World J. Zoology*, 4(3): 223-224.

Nath 2018 Impact of land-use changes on the storage of soil organic carbon in active and recalcitrant pools in a humid tropical region of India. *Sci. Total Environ.*, 624, 908-917.

Ozcelik B, Lee JH, Min DB (2003) Effects of light, oxygen, and pH on the absorbance of 2,2-diphenyl-1-picrylhydrazyl (DPPH). *J Food Sci* 68(2):487. doi:10.1111/j.1365-2621.2003.tb05699.x

Olson, D. R. (1994). *The world on paper: The conceptual and cognitive implications of writing and reading*. Cambridge University Press.

Pandey, S.K. and Shukla, R.P. (2003). Diversity and richness of plants in Darjeeling Himalaya with an eye on Gaddikhana forestry beat, Szechal East Zone Forest Range, Darjeeling. *Indian Journal of Forestry*, 35: 39-44.

Panda, P.C., Mahapatra, A.K., Acharya, P.K., & Debata, A.K. (2013). Plant diversity in tropical deciduous forests of Eastern Ghats, India: A landscape level assessment. *International Journal of Biodiversity Conservation*, 5, 625–639.

Pandey S.K and Shukla, R.P. (2005). Plant community and diversity patterns within the forested landscape of north-eastern U.P. *Indian Forester*, 131; 1217-1226.

Parthasarathy, N. (2001). Changes in forest composition and structure in three sites of tropical evergreen forests around Sengaltheri, Western Ghats. *Current Sciences*, 80, 389–393.

Phillips, O., Martinez, R.V., Vargas, P.N., Monteagudo, A.L., Zans, M.E.C., & Sanchez, W.G. (2003). Efficient plot-based floristic assessment of tropical forests. *Journal of Tropical Ecology*, 19, 629–645.

Pragasam, L.A., & Parthasarathy, N. (2010). Landscape-level tree diversity assessment in tropical forests of southern Eastern Ghats, India. *Flora-Morphology, Distribution, Functional Ecology of Plants*, 205, 728–737.

Premavani, D., Naidu, M.T., Kumar, O.A., & Venkaiah, M. (2017). Diversity and distribution of tree species in tropical forests of Northcentral Eastern Ghats, India. *Asian Journal of Forestry*, 1(1), 27–32.

Parthasarathy N (1986) Studies on the vascular flora, structure and nutrient cycling in Kalakad Reserve Forest, Western Ghats, Tamil Nadu. PhD Thesis, University of Madras, Madras, 313.

Parthasarathy N (1988) A phytogeographic analysis of the flora of Kalakad reserve forest Western Ghats. *Journal of the Indian Botanical Society*, 67: 342–345.

Phillips, E.A. (1959) *Methods of Vegetation Study*. Holt, Rinehart and Winston, New York, 107 p.

Ramesh P, Panwar NR, Singh AB, Ramana S, Rao AS (2009) Impact of organic-manure combinations on the productivity and soil quality in different cropping systems in Central India.

Raturi (2012) [169] for temperate mixed forests and sub-tropical forests. Biodiversity can also vary based on biogeography, habitat, and disturbance (Sagar et al., 2003 [176])

Rennolls, K Laumonier, Diversity structure analysis at two sites in the tropical rain forest of Sumatra *Journal of Tropical EcologySpecies*, Volume 16, Issue 2, March 2000, Pages 253-270.

Sagar R, Raghubanshi AS, Singh JS (2003). Tree species composition, dispersion and diversity along a disturbance gradient in a dry tropical forest region of India. *Forest Ecology and Management* 186(1-3):61-71.

Raghubanshi AS & Tripathi A (2009) Effect of disturbance, habitat fragmentation and alien invasive plants on floral diversity in dry tropical forest of Vindhyan Highlands: a review. *Tropical Ecology* 50(1): 57–69.

Richards, L.A. (1954) *Diagnosis and Improvement of Saline Alkali Soils*, Agriculture, 160, Handbook 60. US Department of Agriculture, Washington DC.

Swamy PS (2000) Forest ecosystem structure and composition along an altitudinal gradient in the Western Ghats, South India. *Journal of Tropical Forest Science* 12: 104–123.

Shukla R.P. and Ramakrishnan P.S. 1986. Architecture and growth strategies of tropical trees in relation to successional status. *Journal of Ecology*, 74: 33–46.

Sharma KL, Mandal B (2009) Soil quality and its relevance for sustainable agriculture. *Journal Indian Soc SoilSci*, 57:572–586.

Subbiah, B.V. and Asija, G.L. (1956) A Rapid Procedure for the Estimation of Available Nitrogen in Soils. *Current Science*, 25, 259-260.

Turner, M.G. 1987. *Landscape Heterogeneity and Disturbance*. Springer-Verlag, New York, 240pp.

Tchouto, M.G.P., Diversity patterns in the flora of the Campo-Ma'an rain forest, Cameroon: Do tree species tell it all? *Biodiversity and Conservation* Volume 15, Issue 4, April 2006, Pages 1353-1374

Wilson, E. O. (1988b). The current state of biological diversity. In E. O. Wilson & F. M. Peter (Eds.), *Biodiversity* (pp. 3–18). National Academy Press.

Utkarsh G et al (1998) On the patterns of tree diversity in the Western Ghats of India. *Curr Sci*.75: 594–603.

Vasu D, Singh SK, Ray SK, Duraisami VP, Tiwary P, Chandran P, Nimkar AM, Anantwar SG (2016) Soil quality index as a tool to evaluate crop productivity in semi-arid Deccan plateau India. *Geoderma*, 282:70–79.

TOTH, S. J.; PRINCE, A. L. ESTIMATION OF CATION-EXCHANGE CAPACITY AND EXCHANGEABLE Ca, K, AND Na CONTENTS OF SOILS BY FLAME PHOTOMETER TECHNIQUES *Soil Science* 67(6):p 439-446, June 1949.

Walkley, A. & Black I.A., 1934. An examination of the Degtjareff Method for Determining Soil Organic Matter, and a proposed Modification of the Chromic Acid Titration Method. *Soil Science*, 37(1): 29-38