

Yield maximization of rice (*Oryza sativa*) through integrated nutrient management

Abstract:

This experiment was conducted on a permanent layout at the west byde of BRRI farm, Gazipur during T. Aman 2019 to Boro 2021-22 to find out the suitable management practices for yield maximization of rice and soil health. Seven treatments in Randomized Complete Block (RCB) Design with three replications were imposed and each treatment was assigned in 5m × 4m sized plot. The treatments combinations were T₁= Absolute Control (No nutrient supply), T₂ = BRRI dose N-P-K-S@83-17-53-12 kg ha⁻¹ in T. Aman and 138-21-75-21 kg ha⁻¹ in Boro season, T₃= Soil Test Based (STB) Fertilizer Dose N-P-K-S @ 67-10-40-10 kg ha⁻¹ in T. Aman and 134-16-75-10 kg ha⁻¹ in Boro, T₄ = STB dose + 1 t ha⁻¹Cowdung, T₅ = STB dose + 1 t ha⁻¹ Poultry manure, T₆ = STB dose + 1 t ha⁻¹ Vermicompost and T₇ = STB dose + 0.33 t ha⁻¹ CD + 0.33 t ha⁻¹ PM + 0.33 t ha⁻¹ VC. Thirty-days old seedling of BRRI dhan87 in T. Aman and forty-two days old seedling of BRRI dhan89 in Boro season were transplanted at 20 cm × 20 cm spacing in both seasons. All manures, soil and plant samples analysis were done by the help of soil Science Division BRRI, Gazipur. Initial soil (0-15 cm depth) properties were: soil texture, clay loam; pH, 6.94; organic Carbon, 1.59%; Nitrogen, 0.18%; Phosphorus, 21.88 ppm and Potassium, 0.19meq/100g soil. Thirty days old seedling of BRRI dhan87 in T. Aman and 42 days old seedling of BRRI dhan89 in Boro season were transplanted at 20cm x 20cm spacing. Grain yield, tiller number, panicle number, plant height and grain number were significantly affected by the different integrated nutrient management during both T. Aman and Boro season. Grain yield, tiller number, panicle number, plant height and grain number were significantly affected by the different nutrient management in both T. Aman and Boro season. Poultry manure related treatments and BRRI recommended dose performed better than the others in all the parameter except 1000-grain weight. On the other hand, Absolute Control (No nutrient supply) produced the lowest result. Every parameter, Poultry manure treatments have been performed the best. STB dose with one t ha⁻¹ poultry manure is better for maximization of rice yield.

Keywords: Cowdung, Poultry Manure, Vermicompost, Yield maximization, Soil fertility, Rice (*Oryza sativa*)

Introduction:

Rice is the staple food for the 170 million people of the country and is grown on around 10.5 million ha [5]. It's an indispensable food for more than 50% of the world's population. Its requirement is increasing rapidly for fulfilling the demand of ever growing human population. To meet the rising demand, different approaches are being practiced such as maximizing rice production with the application of different fertilizers [2], the integrated use of organic fertilizer

Comment [U1]: Please elaborate more what the meaning of best. Usually, best if we compare in the treatments. And also please elaborate how much the yield when you mentioned maximization

with inorganic fertilizer. The global annual production of rice is about 496.40 million tons from 162.06 million hectares of land in 2019–2020 [3]. Globally, agriculture is becoming challenging because of climate change and soil health degradation. Organic matter (OM) is a vital component to keep soil alive and productive and for providing better ecosystem services [10,8]. Along with global agriculture, Bangladesh also faces the challenge of producing crops from its limited land resources to meet the huge demand from its population. Moreover, every year, cultivated land is being reduced due to human settlement and rapid urbanization and industrialization. Therefore, the intensification of land with modern crop varieties for increased production is crucial [11]. Although modern rice varieties have the highest yield, there is still a serious yield gap. In Bangladesh, the lack of judicious nutrient management is considered as the major causes of yield gap in rice production. An imbalanced use of inorganic fertilizers and pesticides without organic fertilizers has led to deterioration of soil health and crop yield loss which has become a concern. More dependency on chemical fertilizers and imbalanced nutrient management practices has impaired the productivity of soils in many Asian countries including Bangladesh. Increased the cropping intensity and regular cultivation of high-yielding rice varieties has increased the removal of nitrogen (N), phosphorus (P), potassium (K) and other macro and micro-nutrients from the soils in Bangladesh [14].

Soil OM content is reported to be declining, which is considered as one of the most serious threats in Bangladesh agriculture [11]. The integration of all organic, natural and inorganic sources of nutrients is an efficient and environmentally friendly technology of crop production which is known as integrated nutrient management (INM). Management practices such as diversified cropping systems and the application of different organic wastes and optimum fertilization, are believed to offer high potential for increasing the nutrient-use efficiency, carbon © levels in soils, and the crop yield [11,12,10,15]. In addition, the release pattern of inorganic fertilizers compared to organic ones are higher [11]. On the other hand, organic fertilizers are decomposed slowly and nutrients are available for a longer period of time which helps to maintain soil nutrient status. However, most of the soils of Bangladesh have an OM content that is less than 1.5% and, in many cases, it is less than 1% [4]. Hence, a large amount of N fertilizer is used to increase soil fertility and crop productivity. Excess N may enhance mineralization of OM, which may decrease soil C content and increase in carbon dioxide (CO₂) emission. Cowdung (Cd) Poultry manure (PM) and vermicompost (VC) are the potential organic sources of soil organic C and plant nutrients [16,13]. Many reports showed that Cd, PM and VC are the source of N, P, K, Ca, Mg, and S that can improve the soil fertility and plant can uptake more of these nutrients [11,16,9]. As Cowdung increases the fertility of the soil by increasing the nutrients in it. VC is rich in microbial activity and contains antagonistic organisms to control plant pathogens, it is also an effective biocontrol agent [7]. Jorgensen et al. [6] reported that adding cowdung, PM and VC in particular increases the contents of organic carbon (OC) and of microbial biomass, the cation exchange capacity, and the biological activities of soils. The above

Comment [U2]: If you have update data such as 2023 at least 2022

findings suggest that organic matter along with inorganic fertilizers can improve soil fertility as well as soil organic C than when applied alone. The soils of Bangladesh have low reserves of C and plant nutrients due to increasing cropping intensity, higher rates of decomposition of organic matter under prevailing hot and humid climate, use of lesser quantity of organic manure, and little or no use of green manure [11]. It is predicted that the crop production of Bangladesh would be tremendously vulnerable to climate change and, therefore, the food security will be at risk. The addition of C-enriched materials such as cowdung, PM and VC improves soil physical, chemical, and biological characteristics. The improvement of soil properties favors the development of the crop root system, elongating it both at the surface level and in the deep soil, which ultimately helps to accumulate more C in the soil. The continuous application of organic manure builds up the soil C content and ensures an adequate N supply. The soil organic C pool in agricultural land is capable of enhancing agricultural sustainability and serving as a potential sink of atmospheric CO₂. The objectives of this study were to find out the suitable integrated nutrient management practice for yield maximization of rice and soil health.

Materials and Methods:

This experiment was initiated on a permanent layout at the west byde of BRRRI farm, Gazipur during T. Aman 2019 to Boro 2021-22 to find out the suitable management practice for yield maximization of rice and soil health. Seven treatments in Randomized Complete Block (RCB) Design with three replications were imposed and each treatment was assigned in 5m × 4m sized plot. The treatments combinations were T₁= Absolute Control (No nutrient supply), T₂ = BRRRI dose N-P-K-S@83-17-53-12 kg ha⁻¹ in T. Aman and 138-21-75-21 kg ha⁻¹ in Boro season, T₃= Soil Test Based (STB) Fertilizer Dose N-P-K-S @ 67-10-40-10 kg ha⁻¹ in T. Aman and 134-16-75-10 kg ha⁻¹ in Boro, T₄ = STB dose + 1 t ha⁻¹ Cowdung, T₅ = STB dose + 1 t ha⁻¹ Poultry manure, T₆ = STB dose + 1 t ha⁻¹ Vermicompost and T₇ = STB dose + 0.33 t ha⁻¹ CD + 0.33 t ha⁻¹ PM + 0.33 t ha⁻¹ VC. Thirty-days old seedling of BRRRI dhan87 in T. Aman and forty-two days old seedling of BRRRI dhan89 in Boro season were transplanted at 20 cm × 20 cm spacing in both seasons. All manures, soil and plant samples analysis were done by the help of soil Science Division BRRRI, Gazipur. Initial soil (0-15 cm depth) properties were: soil texture, clay loam; pH, 6.94; organic Carbon, 1.59%; Nitrogen, 0.18%; Phosphorus, 21.88 ppm and Potassium, 0.19meq/100g soil. Thirty days old seedling of BRRRI dhan87 in T. Aman and 42 days old seedling of BRRRI dhan89 in Boro season were transplanted at 20cm x 20cm spacing. The flooded water level at 5-7 cm depth was maintained during rice cultivation, and drained out the water 21 days before rice harvesting. Yield and Yield components were collected at harvesting time. Collected data were statistically analyzed using a standard statistical procedure (Crop stat 7.2).

Results and Discussion:

Grain yield, tiller number, panicle number, plant height and Grain number were significantly affected by the different nutrient management in both T. Aman and Boro season. Poultry

Comment [U3]: To compare between treatments, do you use statistical analysis?

manure related treatments and BRRRI recommended dose performed better than the others in all the parameter except 1000-grain weight. On the other hand, Absolute Control (No nutrient supply) produced the lowest result. The details have discussed below.

Plant height:

Different nutrient management have significant effects in rice plant height. In T. Aman season 2019 and 2021, the tallest rice plant (126.27 cm) was found in the STB dose + 1 t/ha Poultry manure management. The smallest rice plant (114.40 cm) was found in the Absolute control plot (Table 1).

In T. Aman season 2020, the tallest rice plant (129.64 cm) was found in the BRRRI recommended dose, which is statistically similar with other nutrient management doses except absolute control. The smallest rice plant (124.87 cm) was found in the Absolute control plot.

And in T. Aman season 2021, the tallest rice plant (125.69 cm) was found in the STB dose with 1 t ha⁻¹ Poultry manure, which is statistically similar with other nutrient management doses except absolute control. The smallest rice plant (121.27 cm) was found in the absolute control plot (Table 1).

During Boro season 2019-20, 2020-21 and 2021-22 STB dose, BRRRI dose, STB dose + 1 t/ha Poultry litter, STB dose + 1t/ha cowdung, STB dose + 1t/ha Vermicompost and STB dose + 0.33 t/ha CD + 0.33 t/ha PM + 0.33 t/ha VC used plot gave almost similar plant height which was statistically significant from control plot. STB dose provides the tallest plant (101.99 cm) whereas control plot gave the smallest plant (96.40 cm) in 2019-20.

STB dose with 1 t/ha, poultry litter provides the tallest plant (105.03 cm) whereas control plot gave the smallest plant (96.98 cm) in 2020-21.

STB dose with 1 t ha⁻¹ poultry manure showed the tallest plant (103.82 cm) whereas control plot gave the smallest plant (94.37 cm) in 2021-22. (Table 1)

Tiller number:

Tiller production varies significantly among the different nutrient management in T. Aman season 2019-21. STB dose + 1t/ha poultry litter plot produced statistically highest tiller number followed by others treatment. In T.Aman 2019, STB dose + 1t/ha poultry litter gave the highest number of tiller (314 tiller m⁻²) whereas control plot gave the lowest number of tiller (230 tiller m⁻²) among all the treatments.

In T. Aman season 2020, STB dose + 1t/ha poultry litter gave the highest number of tiller (296 tiller m⁻²) whereas control plot gave the lowest number of tiller (204 tiller m⁻²) among all the treatments.

In T. Aman season 2021, STB dose + 1 t ha⁻¹ poultry manure gave the highest number of tiller (286 tiller m⁻²) and the second highest STB dose + 0.33 t ha⁻¹ CD + 0.33 t ha⁻¹ PM + 0.33 t ha⁻¹ VC (273 tiller m⁻²) whereas control plot gave the lowest number of tiller (207 tiller m⁻²) among all the treatments (Table 1).

Tiller production varies significantly among the different nutrient management in Boro season 2019-20, STB dose + 1t/ha poultry litter plot produced statistically highest tiller number (257 tiller m⁻²) followed by others treatment. whereas control plot gave the lowest number of tiller (202 tiller m⁻²) among all the treatments.

In Boro season 2020-21, STB dose + 1t/ha poultry litter gave the highest number of tiller (283 tiller m⁻²) whereas control plot gave the lowest number of tiller (210 tiller m⁻²) among all the treatments

In Boro season 2021-22, STB dose + 1 t ha⁻¹poultry manure plot produced statistically the highest tiller number (275 tiller m⁻²) followed by others treatment whereas control plot gave the lowest number of tiller (202 tiller m⁻²) among all the treatments (Table 1).

Panicle number:

Panicle production was significantly affected by all the nutrient management during T. Aman season 2019-21. Here BRRI fertilizer dose and STB dose + poultry litter used plot gave statistically similar panicle number. In T. Aman 2019, the highest number of panicle (303 panicle m⁻²)found in STB dose + 1t/ha poultry manure. The lowest number of panicle (223 panicle m⁻²) among all the treatments was observed in control plot.

In T.Aman 2020, the highest number of panicle (274 panicle m⁻²)found in STB dose + 1t/ha poultry manure and the lowest number of panicle (180-panicle m⁻²) among all the treatments was observed in control plot

During T. Aman season 2021,the highest number of panicle (271 panicle m⁻²) was found in STB dose + 1 t ha⁻¹ poultry manure used plot. The lowest number of panicle (188-panicle m⁻²) among all the treatments was observed in absolute control plot (Table 1).

Panicle production was significantly affected by all the nutrient management during Boro season 2019-20, 2020-21 and 2021-22. Here STB dose + poultry litter used plot gave statistically highest panicle number. The lowest number of panicle among all the treatments was observed in control plot (Table 1).

Grain number and grain weight:

During T. Aman season 2019, STB dose + 1 t/ha Poultry litter provides the highest number of grain per panicle (117 grain panicle⁻¹) whereas control plot gave the lowest number of grain (98 grain panicle⁻¹). And there was no significant difference among the treatments in case grain weight.

In T.Aman 2020, STB dose + 1 t/ha Poultry litter and STB dose + 1 t/ha vermicompost plot provide the highest number of grain per panicle (104 grain panicle⁻¹) whereas control plot gave the lowest number of grain (90 grain panicle⁻¹). In addition, there was no significant difference among the treatments in case grain weight.

In T. Aman season 2021, the all-nutrient management plot except absolute control plot gave almost similar number of grains per panicle. STB dose + 1 t ha⁻¹ Poultry manure and STB dose + 1 t ha⁻¹ vermicompost plot gave the highest number of grains per panicle (102 grain panicle⁻¹)

whereas control plot gave the lowest number of grain (95 grain panicle⁻¹). In addition, there was no significant difference among the treatments in case grain weight (Table 1).

During Boro season 2019-20, STB dose + 1 t/ha Poultry litter provides the highest number of grain per panicle (158 grain panicle⁻¹) whereas control plot gave the lowest number of grain (110 grain panicle⁻¹).

In Boro season 2020-21, STB dose + 1 t/ha Poultry litter provides the highest number of grain per panicle (152 grain panicle⁻¹) whereas control plot gave the lowest number of grain (110 grain panicle⁻¹). On the other hand, there was no significant difference among the treatments in case grain weight.

In Boro season 2021-22, STB dose + 1 t ha⁻¹ Poultry manure, BRRRI dose, STB dose + 1 ha⁻¹ cowdung STB dose + 1 t ha⁻¹ VC, STB dose + 0.33 t ha⁻¹CD + 0.33 t ha⁻¹PM + 0.33 t ha⁻¹VC and STB dose used plot gave almost similar number of grains per panicle which was statistically significant from control plot. STB dose + 1 t ha⁻¹ Poultry manure provides the highest number of grains per panicle (139.77 grain panicle⁻¹) whereas control plot gave the lowest number of grain (108.02 grain panicle⁻¹). On the other hand, there was no significant difference among the treatments in case of grain weight (Table 1).

Grain yield:

During T. Aman season 2019-21 Grain yield was significantly affected by different nutrient management practices. In 2019, STB dose + 1 t/ha poultry manure (5.52 t ha⁻¹) and BRRRI dose (5.12 t ha⁻¹) gave the highest and statistically similar grain yield. The lowest yield was observed in control plot (3.90 t ha⁻¹).

During T. Aman season 2020, STB dose + 1 t/ha poultry manure (6.53 t ha⁻¹) and STB dose + 1 t/ha Vermicompost (6.11 t ha⁻¹) gave the highest grain yield and the lowest yield was observed in control plot (3.40 t ha⁻¹).

In T. Aman 2021, STB dose + 1 t ha⁻¹ poultry manure (5.81 t ha⁻¹) and BRRRI recommended dose (5.59 t ha⁻¹) produced the highest grain yield and the lowest yield observed in absolute control plot (3.23 t ha⁻¹) (Table 1).

During Boro season 2019-20, 2020-21 and 2021-22, grain yield was significantly affected by different nutrient management practices. In 2019-20, STB dose + 1 t/ha poultry manure (8.16 t ha⁻¹) and BRRRI dose (7.90 t ha⁻¹). The lowest yield was observed in control plot (3.16 t ha⁻¹).

In 2020-21, STB dose + 1 t/ha poultry manure (8.90 t ha⁻¹), BRRRI dose (8.52 t ha⁻¹) and STB dose + 1 t/ha VC (8.49 t ha⁻¹) gave the highest and statistically similar grain yield and the lowest yield was observed in control plot (4.48 t ha⁻¹).

In Boro season 2021-22, STB dose + 1 t ha⁻¹ poultry manure gave the highest grain yield (7.02 t ha⁻¹). The lowest yield was observed in control plot (4.16 t ha⁻¹) (Table 1).

Table 1. Yield and yield components of rice under different fertilizer management in T. Aman 2019-2021 and Boro 2019-2022

Treatments	Plant height (cm)	Tiller m ⁻² (no.)	Paniclem ⁻² (no.)	Grain panicle ⁻¹ (no.)	1000- grain wt. (g)	Grain yield (t ha ⁻¹)
T. Aman (BRRI dhan87), 2019						
T ₁	114.40	230	223f	98	24.10	3.90
T ₂	125.33	302	292	110	23.67	5.12
T ₃	122.60	252	241	99	24.17	4.34
T ₄	125.53	294	283	101	24.43	4.98
T ₅	126.27	314	303	117	24.27	5.52
T ₆	123.90	264	255	112	24.40	4.56
T ₇	124.43	279	268	99	23.98	4.67
LSD at 5% level	2.22	10.45	11.02	10.12	0.84	0.42
CV %	2.98	9.55	8.46	8.90	3.48	9.55
Boro (BRRI dhan89), 2019-20						
T ₁	96.40	202	198	110	24.59	3.16
T ₂	100.54	253	244	154	24.99	7.90
T ₃	101.99	245	238	139	24.73	6.32
T ₄	99.22	249	242	153	24.63	7.36
T ₅	101.10	257	254	158	24.87	8.16
T ₆	98.98	247	238	141	24.67	6.54
T ₇	99.55	248	240	149	24.97	7.00
LSD at 5% level	2.25	7.45	8.92	9.12	0.75	0.48
CV %	3.28	7.98	7.66	7.09	3.32	7.67
T. Aman (BRRI dhan87), 2020						
T ₁	124.87	204	180	90	22.10	3.40
T ₂	129.64	278	257	102	22.57	5.89
T ₃	126.87	263	246	97	22.34	5.46
T ₄	128.30	281	267	96	22.47	5.75
T ₅	129.23	296	274	104	22.52	6.53
T ₆	128.57	285	266	104	22.43	6.11
T ₇	128.57	278	255	98	22.37	5.64
LSD at 5% level	3.32	13.50	12.23	8.89	0.68	0.48
CV %	3.77	10.12	9.54	9.12	5.09	10.95
Boro (BRRI dhan89), 2020-21						
T ₁	96.98	210	190	110	22.28	4.48

T ₂	104.53	275	258	148	22.82	8.52
T ₃	104.63	258	242	146	22.83	7.89
T ₄	105.03	265	248	145	22.90	8.10
T ₅	104.55	283	263	152	22.88	8.90
T ₆	104.40	269	257	147	22.85	8.49
T ₇	104.60	260	242	147	22.81	7.95
LSD at 5% level	2.84	7.45	8.92	9.12	0.67	0.46
CV %	3.28	7.98	7.66	7.09	3.32	9.80
T. Aman (BRRl dhan87), 2021						
T ₁	121.27	207	188	95	22.25	3.23
T ₂	125.43	266	251	101	22.55	5.59
T ₃	122.20	258	242	98	22.12	5.20
T ₄	120.27	271	259	99	22.68	5.42
T ₅	125.69	286	271	102	22.56	5.81
T ₆	123.33	266	257	102	22.32	5.39
T ₇	122.37	273	254	97	22.38	5.34
LSD at 5% level	3.33	10.95	7.58	4.31	0.65	0.21
CV %	2.17	8.92	8.51	9.05	3.14	10.20
Boro (BRRl dhan89), 2021-22						
T ₁	94.37	202	182	108.02	22.30	4.16
T ₂	103.27	274	246	139.54	22.46	6.78
T ₃	101.60	258	237	139.28	22.47	6.37
T ₄	101.33	270	247	138.98	22.49	6.48
T ₅	103.82	275	250	139.77	22.56	7.02
T ₆	102.93	268	246	139.28	22.53	6.65
T ₇	102.17	264	240	140.15	22.57	6.47
LSD at 5% level	6.11	9.24	10.82	2.97	0.16	0.16
CV %	2.36	8.14	7.75	4.20	2.57	8.87

T₁ = Absolute control

T₂ = BRRR recommended dose N-P-K-S@83-17-53-12 kg ha⁻¹

T₃ = Soil Test Based (STB) Fertilizer dose N-P-K-S@67-10-40-10 kg ha⁻¹

T₄ = STB dose + 1 t ha⁻¹ cowdung

T₅ = STB dose + 1 t ha⁻¹ poultry manure

T₆ = STB dose + 1 t ha⁻¹ vermicompost and

T₇ = STB dose + 0.33 t ha⁻¹ CD + 0.33t ha⁻¹ PM + 0.33 t ha⁻¹ VC.

Initial soil (0-15 cm depth) properties:

Initial soil (0-15cm depth) properties of this experimental field were: soil texture, clay loam; pH, 6.94; organic Carbon, 1.59%; Nitrogen, 0.18%; Phosphorus, 21.88 ppm and Potassium, 0.19meq/100g soil. (Table 4)

Table 2. Initial soil (0-15 cm depth) properties were-

Soil texture	pH	% OC	% Total N	Available P (ppm)	Available K (meq/100g soil)
Clay loam	6.94	1.59	0.18	21.88	0.19

Chemical properties of soil after harvest:

The highest % OC and available K (meq/100g soil) were found from the treatment T₄ = STB dose + 1 t ha⁻¹cowdung application plot. The highest % total N was found from the treatment T₃ = Soil Test Based (STB) Fertilizer dose N-P-K-S@67-10-40-10 kg ha⁻¹ application plot. The highest available P (ppm) and available S (ppm) were found from the treatment T₅ = STB dose + 1 t ha⁻¹ poultry manureapplication plot. (Table 5)

Table 3. Chemical properties of soil after rice harvest under different fertilizer management in T. Aman and Boro seasons

Treatment	% OC	% Total N	Available P (ppm)	Available K (meq/100g soil)	Available S (ppm)
T1	1.160	0.117	19.150	0.134	26.035
T2	1.210	0.110	17.933	0.130	23.263
T3	1.240	0.131	21.200	0.137	26.223
T4	1.280	0.126	21.300	0.147	29.425
T5	1.210	0.128	23.833	0.141	29.948
T6	1.170	0.124	22.100	0.133	24.435
T7	1.190	0.112	20.750	0.128	24.835

Conclusions

Grain yield, tiller number, panicle number, plant height and grain number were significantly affected by the different integrated nutrient management practices during both T. Aman and Boro season. Every parameter, Poultry manure treatments have been performed the best. This study indicates STB dose with one t ha⁻¹ Poultry manure is better for maximization of rice yield.

References

1. Alam, M.A.; Rahman, M.M.; Biswas, J.C.; Akhter, S.; Maniruzzaman, M.; Choudhury, A.K.; Jahan, M.A.H.S.; Miah, M.M.U.; Sen, R.; Kamal, M.Z.U.; et al. Nitrogen transformation and carbon sequestration in wetland paddy field of Bangladesh. *Paddy Water Environ.* 2019, 17, 677–688.
2. Cassman KG, Peng S, Olk DC, Ladha JK, Reichardt W, Dobermann A, Singh U (2018) Opportunities for increased nitrogen-use efficiency from resource management in irrigated rice systems. *Field Crop Res* 56(1):7-39. [https://doi.org/10.1016/S0378-4290\(97\)00140-8](https://doi.org/10.1016/S0378-4290(97)00140-8)
3. FAO. FAOSTAT, Food and Agriculture Organization, Rome. Available online: <http://www.fao.org/faostat/en/#home> (accessed on 10 October 2021).
4. FRG. Fertilizer Recommendation Guide; Bangladesh Agricultural Research Council (BARC): Dhaka, Bangladesh, 2018.
5. Islam, M.M.; Urmi, T.A.; Rana, M.S.; Alam, M.S.; Haque, M.M. Green manuring effects on crop morpho-physiological characters, rice yield and soil properties. *Physiol. Mol. Biol. Plants* 2019, 25, 303–312.
6. Jorgensen, R.G.; Meyer, B.; Roden, A.; Wittke, B. Microbial activity and biomass in mixture treatments of soil and biogenic municipal refuse compost. *Biol. Fert. Soils* 1996, 23, 43–49.
7. Joshi, R.; Singh, J.; Vig, A.P. Vermicompost as an effective organic fertilizer and biocontrol agent: Effect on growth, yield and quality of plants. *Rev. Environ. Sci. Biotechnol.* 2015, 14, 137–159.
8. Liu, Y.; Ge, T.; van Groenigen, K.J.; Yang, Y.; Wang, P.; Cheng, K.; Zhu, Z.; Wang, J.; Li, Y.; Guggenberger, G.; et al. Rice paddy soils are a quantitatively important carbon store according to a global synthesis. *Commun. Earth Environ.* 2021, 2, 1–9.
9. Mahmud, M.; Abdullah, R.; Yaacob, J.S. Effect of vermicompost on growth, plant nutrient uptake and bioactivity of ex vitro pineapple (*Ananas comosus* var. MD2). *Agronomy* 2020, 10, 1333.

- 10.** Menšík, L.; Hlisnikovský, L.; Kunzová, E. The state of the soil organic matter and nutrients in the long-term field experiments with application of organic and mineral fertilizers in different soil-climate conditions in the view of expecting climate change. In *Organic Fertilizers-History, Production and Applications*; IntechOpen: London, UK, 2019.
- 11.** Rahman, M.M.; Alam, M.S.; Kamal, M.Z.U.; Rahman, G.M. Organic sources and tillage practices for soil management. In *Resources Use Efficiency in Agriculture*; Springer: Berlin/Heidelberg, Germany, 2020; pp. 283–328.
- 12.** Rahman, M.M. Nutrient-use and carbon-sequestration efficiencies in soils from different organic wastes in rice and tomato cultivation. *Commun. Soil Sci. Plant Anal.* 2013, *44*, 1457–1471.
- 13.** Rahman, F.; Rahman, M.M.; Rahman, G.K.M.M.; Saleque, M.A.; Hossain, A.T.M.S.; Miah, M.G. Effect of organic and inorganic fertilizers and rice straw on carbon sequestration and soil fertility under a rice-rice cropping pattern. *Carbon Manag.* 2016, *7*, 41–53.
- 14.** Saleque, M.A.; Abedin, M.J.; Bhuiyan, N.I.; Saman, S.K.; Panaullah, G.M. Long-term effects of inorganic and organic fertilizer sources on yield and nutrient accumulation of low land rice. *Field Crops Res.* 2004, *86*, 53–65.
- 15.** Sultana, M.; Jahiruddin, M.; Islam, M.R.; Rahman, M.M.; Abedin, M.A.; Solaiman, Z.M. Nutrient enriched municipal solid waste compost increases yield, nutrient content and balance in rice. *Sustainability* 2021, *13*, 1047.
- 16.** Suthar, S. Impact of vermicompost and composted farmyard manure on growth and yield of garlic (*Allium stivum* L.) field crop. *Int. J. Plant Prod.* 2009, *3*, 27–38. *Plants* 2022, *11*, 138 16 of 17