

The association between dietary intake and lifestyle pattern of diabetes patients in Manipur

ABSTRACT

The aim of the study was to examine the association between dietary intake and lifestyle pattern of diabetes patients in Manipur. Tribal and non tribal respondents between 45-64 years of age were randomly selected from the Regional Institute of Medical Sciences (RIMS) hospital Manipur. Total number of 200 study subjects, 100 tribal and 100 non tribal subjects constituted the study. The information were collected on socio-demographic profile, clinical, anthropometric measurements and dietary intakes of the patients by using 24-hour recall method. The data were analysed and tabulated using statistical tools such as frequency, percentage, means, standard deviation, MS Excel and Pearson's correlation test. The results have shown that alcohol, triglycerides, energy, oils and sugar consumption were significantly correlated with the prevalence of diabetes. The low calcium intake among non tribal individuals reveals a significant association with FBG (Fasting Blood Glucose) test results, as quoted low calcium level increases type II diabetes. While low calcium intake among the non tribal community showed significant relation to the PP (Post Prandial) test results, i.e. increase in PP. Tribal energy intake and non tribal protein intake were significantly related with HbA1c (Glycated Haemoglobin) test.

Out of which 100 tribal and 100 non tribal respondents were selected

Keywords: Diabetes, Tribal, Non tribal, Demographic profile, Lifestyle pattern and Clinical parameters.

Comment [d1]: 'were' randomly selected

Comment [d2]: Capitalize the first letter "Hospital"

Comment [d3]: Include "number of"

Comment [d4]: Format comma

Comment [d5]: Delete "s"

Comment [d6]: ...analyzed

Comment [d7]: ...rearrange

Comment [d8]: Rearrange

Comment [d9]: Rearrange

1. INTRODUCTION

Diabetes is a chronic disease that occurs either when the pancreas does not produce enough insulin or when the body cannot effectively use the insulin it produces (WHO, 2023). Its rising incidence in emerging nations like India is directly related to ageing populations, urbanized lifestyles and population structure (Ramachandran, 2001). Approximately 422 million individuals globally suffer from diabetes, with the majority residing in low and middle-income nations. The disease is directly responsible for 1.5 million deaths annually. Over the past few decades, there has been a steady rise in both the number of cases and the incidence of diabetes (WHO, 2023). According to NFHS 2019-21 reports, India had the proportion of women who were requiring medication or having high blood glucose levels (above 140 mg/dl) which varied across different religious groups: Hindus (13.3%), Muslims (13.8%), Christians (19.2%), Sikhs (14.6%), Buddhists (17.5%), Jains (9.5%), and others (10.3%). For men, the percentages were Hindus (15.5%), Muslims (15.5%), Christians (20.4%), Sikhs (14.2%), Buddhists (11.6%), Jains (17.3%), and others (13.5%) (James *et al.*, 2022). For Manipur, the percentage of women and men with blood glucose levels higher than 140 mg/dl or taking medicine to lower their blood glucose level were 13.6% and 16.5% respectively. This rate for women and men for schedule tribe accounts 6.4% and 9.9% respectively (Lhungdimet *et al.*, 2021).

Genetic and environmental variables, such as family history, older age, obesity, and physical inactivity, are risk factors for this condition. Long-term obesity and a sedentary lifestyle may also have an impact on the likelihood of developing type 2 diabetes Bakr *et al.*, (2015). Delay in the onset of T2DM can be made by the use of healthy diet, having regular physical activity, and also through the avoidance of tobacco (WHO, 2023). T2DM and its complications constitute a major public health problem worldwide, affecting almost all populations in both developed and developing countries, with high rates of diabetes-related morbidity and mortality (Wu *et al.*, 2014).

Comment [d10]: Cite

Comment [d11]: Remove the bracket

Comment [d12]: Use APA REFERENCES

Comment [d13]: Move this statement to the beginning of your introduction

2. MATERIAL AND METHODS

2.1 Study Area and Selection of Respondents

This cross-sectional study was carried out in the Manipur NCD (Non Communicable Disease) unit of RIMS Hospital. Patients with diabetes from the non-tribal Meiteis group as well as the tribal Nagas and Kukis group were chosen. The required sample sizes were selected from the hospital randomly. Total 200 study sample were selected. Out of which 100 tribal and 100 non tribal respondents were selected. Fifty from each gender. The chosen age range falls into late middle age (45–64 years old).

Comment [d14]: Write in full first

Comment [d15]: How did you calculate your sample size and what method of sample selection did you use

Please, indicate

Comment [d16]: The research design should be 'qualitative design' not cross-sectional

2.2 Methods of Data Collection

The main instruments used in the study were pre-tested and pre-designed interview schedules. Utilizing the 24-hour recall method, Informations were gathered on the patients' nutritional intakes, anthropometric measurements, clinical data, and sociodemographic profile. In anthropometric measurements are weight, height, and waist-hip ratio were taken. The formula kg/m^2 , where m^2 is the square of a person's height in meters, is used to calculate a person's BMI (Body Mass Index). Utilizing the WHO classification, the BMI was evaluated. Measurements of height, weight were obtained using a digital scale. The patients' height was determined using an anthropometer rod. Updated blood glucose monitoring and cholesterol levels records were taken from the patients.

Comment [d17]: Your statistical analysis should be in line with the interview method of data analysis

Please, fine-tune your research design and instruments used and data analysis. They must correlate

2.3 Statistical Analysis

The data were analysed and tabulated using statistical tools such as frequency, percentage, means, standard deviation, MS Excel and Pearson's correlation test.

3. Results and Discussion:

Table shows the association between the dietary intake and lifestyle pattern of diabetes patients in Manipur which describe thoroughly about demographic factors like age, occupation, qualification, marital status, total annual income and addiction pattern. Tables 1 show the demographic factors of the diabetic patients from tribal and non tribal communities. A majority from both groups falls within the age group of 55-64 years and annual income falls under 2 lakhs per annum. Service was the main occupation for male and majority of females were under housewives category. Male under service category and female under housewife category in both communities exhibited a higher prevalence of diabetes. Respondents who got higher education had more incidence of diabetes in both tribal and non tribal communities. Probably, the reason behind this is highly paid services because of good education level. More than 90% of the respondents were married. Among the surveyed population, tribal males (68%) exhibited the highest alcohol consumption as compared to others. Tobacco uses was prevalent among tribal males 70% and tribal females at 66%. In comparison non tribal males reported a lower tobacco usage at 50%, while non tribal females had the highest rate at 72%. A similar study conducted by Jayawardena *et al.*, (2014) and Wimmer *et al.*, (2023) revealed that old aged people and individuals with higher income were more prevalent to diabetes. A study conducted by Begum *et al.* (2004) found that housewives were more diabetic than others. A study by Kim *et al.*, (2021) noticed that more than 50% male diabetic patients used tobacco and consumed alcohol. According to Onalan and Gozel (2020), smoking cigarettes raises the incidence of diabetes.

Comment [d18]: Your methodology is not full enough

Description of your instrument
Population size
Sample size determination
Sampling methods,
Ethical consideration with approved protocol/reference number

Comment [d19]: Insert the table number

**Table 1. Distribution of study subjects based on socio-demographic details:
N= (Tribal=100, Non tribal=100), (Male=50, Female=50)**

Demographic factors	Tribal F (%)		Non tribal F (%)	
	Male	Female	Male	Female
1. Age Group				
45-54	14 (28)	18 (36)	15 (30)	16 (32)
55-64	36 (72)	32 (64)	35 (70)	34 (68)

2. Occupation				
Housewife	-	22 (44)	-	28 (56)
Farmer	9 (18)	1 (2)	12 (24)	-
Wage earner	1 (2)	3 (6)	4 (8)	2 (4)
Service	29 (58)	13 (26)	22 (44)	11 (22)
Business	11 (22)	11 (22)	12 (24)	9 (18)
3. Qualification				
Illiterate	2 (4)	2 (4)	3 (6)	6 (12)
Can read & write	1 (2)	3 (6)	4 (8)	2 (4)
Middle School	4 (8)	8 (16)	4 (8)	6 (12)
High School	10 (20)	10 (20)	7 (14)	15 (30)
Higher Secondary	14 (28)	11 (22)	13 (26)	7 (14)
Under graduate	15 (30)	16 (32)	15 (30)	12 (24)
Post graduate	4 (8)	-	4 (8)	2 (4)
4. Marital status				
Married	43 (86)	32 (64)	49 (98)	37 (74)
Unmarried	2 (4)	3 (6)	-	1 (2)
Widow	-	15 (30)	-	12 (24)
Widower	5 (10)	-	1 (2)	-
5. Total annual income of family (in rupees)				
<50,000	-	-	1 (2)	1 (2)
50,001 – 1,00,000	2 (4)	4 (8)	-	5 (10)
1,00,001 – 1,50,000	1 (2)	2 (4)	5 (10)	5 (10)
1,50,001 – 2,00,000	3 (6)	2 (4)	3 (6)	8 (16)
>2,00,000	44 (88)	42 (84)	41 (82)	31 (62)
6. Addiction Pattern				
Alcohol				
Yes	34 (68)	8 (16)	16 (32)	-
No	16 (32)	42 (84)	34 (68)	50 (100)
Tobacco				
Yes	35 (70)	33 (66)	25 (50)	36 (72)
No	15 (30)	17 (34)	25(50)	14 (28)

F= Frequency

Table 2 represents the association of demographic profile, lifestyle pattern and clinical parameters. The mean of tribal and non tribal population were compared height, weight, BMI, increased alcohol intake and higher triglyceride levels showed significance positive correlation among both tribal and non tribal population by using Pearson's correlation test. Additionally, nutrient such as mean energy intake, mean carbohydrate intake and mean protein (high) intake, exhibited significant results for both the communities. Moreover, the consumption of specific food categories such as cereals and millets, oils (excess) and sugar also demonstrated significant relationship across both demographic groups. The age groups of 50-69 years were significant for high chances of being diabetic as stated in a study conducted by Mathur *et al.*, (2022). In similar findings, high BMI (Motovuet *et al.*, 2017) and high triglyceride levels (Jayakumar *et al.*, 2020) are significantly correlated with the prevalence of diabetes. Likewise, discovered that over consumption of alcohol (Kim *et al.*, 2021), over intake of energy (Zujko *et al.*, 2020) and protein (Shahar *et al.*, 2005) were significantly correlated with the occurrence of diabetes. Similarly to the above finding (Sowmya *et al.*, 2016) revealed that excess refined cereal intake were significant with increased in diabetes. Bowen *et al.*, (2011) and Verma *et al.*, (2019) stated that excessive consumption of oils and occurrence rate of diabetes were significant.

**Table 2. Profiling of study subjects based on demographic, lifestyle, and clinical parameters:
N= (Tribal=100, Non tribal=100), (Male=50, Female=50)**

Variables	Tribal Diabetic (Mean ± SD)	Non tribal Diabetic (Mean ± SD)	p-value
Age	56.48 ± 5.40	57.2 ± 5.98	0.02*

Weight (Kg)	65.23 ± 8.46	64.3 ± 10.14	0.00*
Height (cm)	159.01 ± 9.14	167.45 ± 7.78	0.02*
BMI (kg/m ²)	25.40 ± 3.58	25.43 ± 4.61	0.03*
WHtR	1.02 ± 0.07	1.03 ± 0.07	0.29
Tobacco Consumption	1.32 ± 0.46	1.39 ± 0.49	0.27
Alcohol Consumption	1.58 ± 0.49	1.84 ± 0.36	0.01*
FBG	181.47 ± 47.26	164.45 ± 31.77	0.58
PP	260.46 ± 77.71	283.63 ± 81.87	0.55
HbA1c	9.39 ± 2.45	8.40 ± 1.86	0.96
Triglyceride (TG)	188.98 ± 74.79	156.24 ± 93.75	0.04*
Total Cholesterol (TC)	221.89 ± 73.43	185.64 ± 48.75	0.55
LDL	102.21 ± 37.32	93.40 ± 34.71	0.65
HDL	43.15 ± 10.65	50.33 ± 17.71	0.98
Energy (kcal)	2090.39 ± 411.02	2109.52 ± 458.14	0.00*
Carbohydrate (g/d)	262.31 ± 58.68	259.58 ± 73.82	0.00*
Protein (g/d)	95.39 ± 26.90	70.10 ± 18.35	0.04*
Total Fat (g/d)	75.06 ± 25.22	87.13 ± 29.51	0.34
Dietary Fibre (mg/d)	24.32 ± 7.10	23.83 ± 6.88	0.85
Cereals & Millets (g/d)	237.1 ± 64.98	226.7 ± 78.16	0.00*
Pulses & legumes (g/d)	15.3 ± 18.17	25 ± 23.67	0.86
Fruits and vegetables (g/d)	435.44 ± 132.86	325.89 ± 114.39	0.85
Dairy products (g/d)	140.56 ± 93.58	134.3 ± 100.00	0.10
Fish, egg and meats (g/d)	278.93 ± 135.26	138.95 ± 87.30	0.32
Oils (ml/d)	43.78 ± 20.98	19.49 ± 24.88	0.03*
Sugar (g/d)	6.17 ± 2.15	6.22 ± 2.64	0.03*

*Significant at the 0.05 level

BMI=Body Mass Index, WHtR=Waist Hip Ratio, FBG=Fasting Blood Glucose, PP=Post Prandial, HbA1c=GlycatedHaemoglobin, LDL=Low Density Lipoprotein and HDL=High Density Lipoprotein.

Table 3 demonstrates about the comparison between nutrients intake of both communities with respect to blood glucose monitoring (FBG, PP and HbA1c) and showed significant results in terms of energy, protein and calcium between them. The energy intake and lower calcium intake among non tribal individuals demonstrate a significant association with FBG test results. Among all nutrients only calcium intake among non tribal community shows significant effect on the PP test results. Further, energy intake of tribal and protein intake non tribal communities exhibited significant relationship with HbA1c test result. Similarly, (Ha *et al.*, 2019) and (Alami *et al.*, 2022) concludes that high energy intake is directly proportionate to increase in FPG test result. Another study conducted by Takeda *et al.*, (2021) and Kang and Kim (2012) also reported significant between high HbA1c levels and energy intake. Hajhashemyet *al.*, (2022) found that there was an inverse association between calcium intake type 2 diabetes mellitus.

Comment [d20]: It advisable you put the result in parenthesis in front of each variable of nutrient and blood glucose, most especially the significant figures

Table 3. Distribution of study subjects by comparing nutrients intake and blood glucose monitoring:

Nutrients	FBG (p-value)		PP (p-value)		HbA1c (p-value)	
	Tribal	Non Tribal	Tribal	Non Tribal	Tribal	Non Tribal
Energy (kcal)	0.76	0.00*	0.63	0.52	0.00*	0.04
Carbohydrate (g/d)	0.86	0.99	0.94	0.75	0.82	0.25
Protein (g/d)	0.31	0.32	0.30	0.19	0.32	0.01*
Fat (g/d)	0.26	0.21	0.30	0.17	0.62	0.12
Iron (mg/d)	0.29	0.71	0.74	0.60	0.07	0.76
Calcium (mg/d)	0.13	0.02*	0.33	0.03*	0.43	0.62
Dietary fibre (mg/d)	0.23	0.84	0.16	0.60	0.83	0.92

*Significant at the 0.05 level

4. CONCLUSION:

Based on the findings, it is evident that both tribal and non tribal communities exhibit significant association with demographic parameters, lifestyle patterns, and clinical findings. Age, weight, height, BMI, alcohol consumption and triglyceride levels emerged as key factors influencing diabetes in both groups. Additionally, nutrient intake, including energy, carbohydrate and protein, and specific food consumption such as cereals and millets, oils, and sugar, displayed correlation with diabetes. Specifically, increase energy and low calcium intake among non-tribal individuals is significantly correlated with the results of the FBG test. Low calcium intake among the non-tribal community was inversely correlated with the results of the PP test. And tribal energy intake and non-tribal protein intake were significantly correlated with the results of the HbA1c test. These findings emphasized the need for understanding and addressing diverse demographic and lifestyle factors to effectively mitigate the burden of diabetes in both tribal and non tribal populations. Further research and targeted public health strategies are required to address the complex interaction of factors contributing to diabetes prevalence and management within this community.

5. REFERENCES:

1. Alami, F., Mohseni, K.G., Ahmadzadeh, M., Vahid, F., Gholamalizadeh, M., Masoumivand, M. et al. The association between fasting blood sugar and index of nutritional quality in adult women. *Frontiers in Nutrition*. 2022;doi.org/10.3389/fnut.2022.883672.
2. Bakr, E.S.H. Nutritional Assessment of Type II Diabetic Patients. *Pakistan Journal of Nutrition*. 2015; 14(6):308-315.
3. Begum, F.A.A., Azad, A.K., Alim, M.A. and Ekram, A.S. Nutritional Status of Diabetic Patients Attending to a District Level Diabetic Center. *Journal of Teachers Association*. 2004; 17(2):89-92.
4. Bowen, L., Ebrahim, S., Stavola, B.D., Ness, A., Kinra, S., Bharathi, A.V. et al. Dietary Intake and Rural-Urban Migration in India: A Cross-Sectional Study. *Public Library of Science*. 2011; DOI: [10.1371/journal.pone.0014822](https://doi.org/10.1371/journal.pone.0014822).
5. Ha, N.T., Phuong, N.T. and Ha, L.T.T. How dietary intake of type 2 diabetes mellitus outpatients affects their fasting blood glucose levels? *AIMS Public Health*. 2019; doi: [10.3934/publichealth.2019.4.424](https://doi.org/10.3934/publichealth.2019.4.424).
6. Hajhashemy, Z., Rouhani, P. and Saneei, P. (2022). Dietary calcium intake in relation to type-2 diabetes and hyperglycemia in adults: A systematic review and dose-response meta-analysis of epidemiologic studies. *Scientific Reports*. 2022; Available: <https://doi.org/10.1038/s41598-022-05144-8>.
7. James, K.S., Singh, S.K., Lungdim, H., Shekhar, C., Dwivedi, L.K., Pedgaonkar, S. et al. National Family Health Survey (NFHS-5), 2019-21: India. Mumbai: International Institute for Population Sciences. 2022; 1:505-510.8. Jayawardena, R., Thennakoon, S., Byrne, N., Soares, M., Katulanda, P. and Hills, A. Energy and nutrient intakes among Sri Lankan adults. *International Archives of Medicine*. 2014; doi: [10.1186/1755-7682-7-34](https://doi.org/10.1186/1755-7682-7-34).
9. Kang, H.M. and Kim, D.J. Total Energy Intake May Be More Associated with Glycemic Control Compared to Each Proportion of Macronutrients in the Korean Diabetic Population. *Diabetes & Metabolism Journal*. 2012; 36(4):300-306.
10. Kim, S.W., Kim, H.J., Min, K., Lee, H., Lee, S.H., Kim, S. et al. The relationship between smoking cigarettes and metabolic syndrome: A cross-sectional study with non-single residents of Seoul under 40 years old. *Public Library of Science*. 2021; doi: [10.1371/journal.pone.0256257](https://doi.org/10.1371/journal.pone.0256257).
11. Lungdim, H., Singh, S.K., Pedgaonkar, S. and Gupta, J. (2021). National Family Health Survey (NFHS-5), India, 2019-21: Manipur. Mumbai: International Institute for Population Sciences. 2021; 1:158-159.

Comment [d21]: No need of numbering APA format of referencing

Arrange alphabetically and indent where necessary

12. Mathur, P., Leburu, S. and Kulothungan, V. Prevalence, awareness, treatment and control of diabetes in India from the countrywide national NCD monitoring survey. *Frontiers in Public Health*. 2022; doi: [10.3389/fpubh.2022.748157](https://doi.org/10.3389/fpubh.2022.748157).

13. Matovu, N., Matovu, F.K., Sseguya, W. and Tushemerirwe, F. Association of dietary intake and BMI among newly diagnosed type 2 diabetes patients attending diabetic clinics in Kampala. *B.M.C. Nutrition*. 2017; DOI: [10.1186/s40795-017-0141-7](https://doi.org/10.1186/s40795-017-0141-7).

14. Onalan, E. and Gozel, N. The association between the prevalence of cigarette smoking and complications in patients with type 2 diabetes. *Progress in Nutrition*. 2020; DOI: [10.23751/pn.v22i2.8155](https://doi.org/10.23751/pn.v22i2.8155).

15. Ramachandran, A., Snehalatha, C., Kapur, A., Vijay, V., Mohan, V., Das, A.K., et al. High prevalence of diabetes and impaired glucose tolerance in India: National Urban Diabetes Survey. *Diabetologia*. 2021; doi: [10.1007/s001250100627](https://doi.org/10.1007/s001250100627).

16. Shahar, D., Shai, I., Vardi, H., Shahar, A., Drora Fraser, D. Diet and eating habits in high and low socioeconomic groups. *Nutrition*. 2005; Available: <https://doi.org/10.1016/j.nut.2004.09.018>.

17. Sowmya, N., Lakshmi Priya, N., Arumugam, K., Venkatachalam, S., Vijayalakshmi, P., Ruchi, V. et al. Comparison of dietary profile of a rural south Indian population with the current dietary recommendations for prevention of non-communicable diseases (CURES 147). *Indian Journal of Medical Research*. 2016; 144(1):112–119.

18. Takeda, Y., Fujihara, K., Nedachi, R., Ikeda, I., Morikawa, S.Y., Hatta, M. et al. Comparing Associations of Dietary Energy Density and Energy Intake, Macronutrients with Obesity in Patients with Type 2 Diabetes (JDDM 63). *Nutrients*. 2021; DOI: [10.3390/nu13093167](https://doi.org/10.3390/nu13093167).

19. Verma, K., Gupta, A., Alisha and Gupta, P. (2019). Assessment of nutritional status and dietary pattern of diabetes patients in Allahabad City. *Journal of Pharmaceutical Innovation*. 2019; 8(3):402-408.

20. WHO. (2023). Diabetes. <https://www.who.int/news-room/fact-sheets/detail/diabetes#:~:text=Overview,hormone%20that%20regulates%20blood%20glucose>. Accessed on 10 Jan. 2024.

Comment [d22]:

21. Wimmer, R., Audétat, A., Binggeli, J., Schuetz, P. and Braun, N.K. Association of Sociodemographic, Socioeconomic and Lifestyle Characteristics with Low Protein and Energy Intake in the Healthy Swiss Population. *Nutrients*. 2023; doi: [10.3390/nu15092200](https://doi.org/10.3390/nu15092200).

22. Wu, Y., Ding, Y., Tanaka, Y and Zhang, W. Risk Factors Contributing to Type 2 Diabetes and Recent Advances in the Treatment and Prevention. *International Journal of Medical Sciences*. 2014; 11(11): 1185–1200.

23. Zujko, M.E., Waśkiewicz, A., Drygas, W., Mikołajczyk, A.C., Zujko, K., Szcześniewska, D. et al. Dietary Habits and Dietary Antioxidant Intake Are Related to Socioeconomic Status in Polish Adults: A Nationwide Study. *Nutrients*. 2020; doi: [10.3390/nu12020518](https://doi.org/10.3390/nu12020518).