

Biochar in Agriculture: Enhancing Crop Productivity and Disease Resistance

ABSTRACT

Biochar, derived through pyrolysis, presents a promising solution to the challenges faced in sustainable agriculture. This review delves into the diverse advantages of employing biochar to enhance crop yields while promoting environmental responsibility. Its cost-effectiveness and eco-friendly nature not only enrich soil fertility but also contribute to carbon capture, aiding in the fight against climate change. Additionally, while its effectiveness in disease control may vary, biochar shows potential in bolstering crops against environmental pressures. By altering soil characteristics, it encourages the growth of beneficial microbes and improves nutrient availability, ultimately supporting plant vitality. Moreover, integrating biochar into agricultural systems may prompt biochemical and physiological changes that activate plant defences against pathogens. This study thoroughly assesses biochar's impacts on soil health, crop output, and disease prevention, emphasizing its crucial role in advancing sustainable farming practices. Embracing biochar as a strategic resource offers great potential for cultivating resilient and environmentally friendly farming methods, marking a significant step towards sustainable crop disease management.

Keywords: Biochar, Disease suppression, Soil health, Sustainable agriculture, Crop yield

1. INTRODUCTION

The significant growth in the global population poses significant challenges to both food security and climate change. Projections suggest that the world's population will continue to rise, with estimates ranging from 9.4 to 10.1 billion people by 2050 [1]. With agricultural land becoming increasingly scarce, there is a pressing need for sustainable approaches to address imminent food shortages. The most effective solution is to enhance food productivity within existing constraints [2]. The primary reasons for decreased crop yields can be categorized into two main groups: biotic factors, which include pests, pathogens, and herbivores, and abiotic factors such as salinity, alkalinity, and chilling injuries. Statistics reveal that plant pathogens are responsible for a 26% reduction in yield, with soil-borne pathogens alone contributing to 10-20% of this loss [3, 4].

Traditional agricultural methods rely on agrochemical application, soil disinfection techniques, and the cultivation of resistant crop varieties to manage pests. However, pesticides pose significant environmental risks by harming beneficial soil microbes, polluting aquatic ecosystems, and leading to the accumulation of toxins in the food chain through water and soil pollution [5, 6]. Ongoing efforts

advocate for the widespread adoption of long-term sustainable organic farming as a preferable alternative to conventional agricultural practices.

Biochar, as defined by Lehmann et al. [7], is a carbon-rich substance derived from biomass through pyrolysis. Pyrolysis entails the thermal breakdown of organic matter in a high-temperature, oxygen-deprived setting, as elucidated by Lehmann and Joseph [8]. This process occurs within a reactor and leads to the conversion of organic material into diverse solid, liquid, and gaseous by-products. The solid fraction, encompassing fixed carbon and other inorganic components, is recognized as biochar, as outlined by Gheorghe et al. [9] and Yaman [10].

It exhibits potential in effectively combating various disease-causing agents such as fungi, bacteria, nematodes, and insect pests. Beyond its numerous beneficial attributes, biochar also improves soil quality and contributes to carbon storage [11]. Biochar resembles charcoal and is obtained by subjecting plant and animal residues to controlled, low-oxygen conditions via a process termed thermo chemical conversion [12, 13]. Various techniques, including pyrolysis, torrefaction, gasification, flash carbonization, and hydrothermal carbonization, are employed in biochar production. Among these methods, pyrolysis is commonly preferred as the most viable option.

In the Amazon regions, there's a notable presence of black carbon known as Terra Preta, which boasts high levels of carbon, nitrogen, phosphorus, potassium, and magnesium, creating a fertile soil conducive to agricultural productivity [14]. Recent research has highlighted various benefits of biochar use. For example, sugarcane bagasse biochar (SBB) has been utilized to counteract the negative effects of polystyrene micro plastics in rice ecosystems, resulting in reduced methane emissions and improved yield and biomass production [15]. Similarly, SBB has been applied to enhance the growth of corn and rice in soils contaminated with lead and cadmium, leading to increased shoot and root dry weights and demonstrating its effectiveness in remediating polluted soils [16, 17].

The remarkable porosity of biochar creates a favourable environment for microorganisms [14], influencing the structure and function of microbial communities and promoting the use of aromatic carbon sources. Generally, five mechanisms have been proposed to explain biochar's disease-suppressing properties, including enhancing systemic resistance, altering soil physicochemical characteristics, boosting populations of beneficial rhizospheric microorganisms, and releasing fungitoxic and allelopathic compounds, either directly or indirectly. Moreover, both systemic acquired resistance (SAR) and induced systemic resistance (ISR) pathways have been observed to contribute to biochar's disease suppression mechanism [18, 19, 20]. The aim of this review is to offer a comprehensive overview of the significance of biochar in promoting sustainable agriculture, particularly focusing on its impact on soil health, disease management, and the mechanisms through which it enhances plant resilience.

2. BIOCHAR IN AGRICULTURE

Biochar plays a pivotal role in agriculture, offering advantageous attributes that positively influence soil health, microbial ecosystems, and plant development. Its porous structure and inherent oxidizing abilities facilitate the generation of beneficial organic compounds, rendering it versatile for applications like water purification, pest management, and fertilizer administration. Moreover, its nutrient richness and retention capabilities make it an effective soil amendment, while its protective mechanisms shield plants from pathogens and aid in environmental rehabilitation.

The properties of biochar are intricately linked to the pyrolysis process, which encompasses variables such as heating rate, temperature, duration of exposure, and post-production treatments. Managing these factors allows for tailoring biochar to specific requirements. Various feedstocks and pyrolysis temperatures influence the physical characteristics of biochar; for instance, biochar derived from rice and wheat exhibits higher density compared to that from maize and pearl millet at 400°C. Lower pyrolysis temperatures produce biochar with reduced surface area and ash content, while higher temperatures result in biochar with increased surface area and ash content.

Additionally, the chemical composition of biochar is affected by temperature variations. Pyrolysis at lower temperatures (below 400°C) yields biochar with elevated levels of nitrogen, sulfur, potassium, and phosphorus, along with lower carbon content and pH. Conversely, higher temperatures (ranging from 600 to 900°C) generate biochar with decreased nitrogen, sulfur, phosphorus, and potassium content, but with higher carbon content and pH. Furthermore, the choice of raw material also influences the chemical properties of biochar; for instance, biochar derived from rice residue exhibits the highest Cation Exchange Capacity (CEC) [21].

2.1. Enhancing Agriculture through Biochar: A Beneficial Influence

While conventional chemical approaches are commonly utilized for managing plant diseases, they carry notable drawbacks. Pesticide usage results in the accumulation of harmful chemical residues, fosters pesticide resistance in pathogens, and diminishes natural enemies of pests, posing health hazards to farmers and environmental risks such as groundwater pollution [22]. Conversely, biochar offers a sustainable alternative in agriculture. Serving as a soil enhancer, biochar contributes to mitigating climate change by reducing emissions of greenhouse gases and promoting carbon storage owing to its resistance to microbial degradation [23].

Research indicates that the application of biochar, either alone or in combination with fertilizers, enhances crop productivity by approximately 10% [24]. This enhancement arises from various mechanisms: biochar's alkaline properties elevate soil pH, its high capacity for retaining water and nutrients benefits crops, it absorbs and neutralizes harmful substances, and it encourages the proliferation of beneficial microorganisms, all of which support plant well-being [25]. Moreover,

biochar contributes to the management of plant diseases by eliciting systemic defense reactions, improving soil characteristics, and maintaining a balanced microbial ecosystem [26].

2.2. Transformative Effects of Biochar on Soil Chemical Properties

Biochar is becoming increasingly popular in agriculture due to its ability to improve essential soil chemical properties such as cation exchange capacity (CEC) and neutralizing soil acidity. This enhancement is attributed to the presence of carboxyl functional groups on the surface of biochar, which contribute to increased alkalinity and CEC. Factors influencing the alkalinity of biochar include the composition of alkaline ash, choice of raw material, and pyrolysis temperature [27, 28]. The main mechanism behind the increase in soil pH buffering capacity is the release of cations resulting from the protonation of carboxyl groups on the surface of biochar and the dissolution of carbonates [29].

Application of biochar improves soil nutrient retention by raising levels of NH_4^+ and Ca^{2+} . Furthermore, it reduces leaching of nitrogen, phosphorus, and potassium, as well as nitrous oxide emissions. It also decreases inorganic nitrogen while increasing organic nitrogen, which stimulates root activity for enhanced nitrogen uptake by plants [30]. The nutrient content of biochar varies depending on the raw material and pyrolysis process, with options such as sludge or manure biochar containing fertilizer elements [31]. Biochar is considered chemically stable, with long-lasting effects due to its dense structure and gradual release of organic carbon as carbon dioxide [32, 33].

2.3. Transformative Effects of Biochar on Soil Physical Properties

Biochar is widely recognized as a potent soil amendment essential for regulating pH levels, improving soil density and thermal properties, stimulating plant growth, enhancing soil structure and fertility, boosting nutrient availability, and augmenting water retention capacity [34]. Its ability to decrease soil density enhances water retention by improving aeration and reducing compaction. Various factors, such as the source material and pyrolysis temperature used in biochar production, as well as the soil type, influence its impact on soil water retention. Notably, while biochar can enhance water retention in sandy and loamy soils, its efficacy may be diminished in clay soils [35]. The influence of biochar on soil physical properties plays a crucial role in shaping microbial communities in the rhizosphere [36, 37].

2.4. Influencing Soil Microbial Communities: The Effect of Biochar

Soil microorganisms play a crucial role in maintaining soil health by facilitating the decomposition of organic matter, cycling nutrients, controlling diseases, and promoting plant growth [38, 39]. While biochar generally resists microbial decomposition, there may be some initial breakdown due to both chemical reactions and microbial activity [40]. The degradation of biochar provides organic nutrients for beneficial microbes such as mycorrhizal fungi and bacteria [41], resulting in an increase in

beneficial bacteria and a decrease in harmful fungi [42]. Application of biochar stimulates root growth, fosters microbial growth, and enhances nutrient cycling efficiency [43].

Biochar also influences soil pH, which in turn affects critical microbial populations involved in nutrient cycling and plant disease control, notably increasing the abundance of Actinobacteria and Proteobacteria [44, 45]. The Proteobacteria phylum aids in plant pathogen control by fostering the growth of antagonistic bacteria through nitrogen, carbon, and sulfur cycling [46]. Additionally, it supports the growth of nitrogen-fixing bacteria, thereby increasing biological nitrogen fixation in plants via root nodule formation [47].

The application of biochar enriches the diversity of beneficial microbial communities such as plant growth-promoting rhizobacteria (PGPR), which in turn assists in the cycling of soil organic matter and its availability to plants [48]. Incorporating biochar alters soil microbiota, impacting nitrogen availability by reducing nitrogen mineralization and disrupting the balance between nitrogen oxidizers and reducers [49].

3. BIOCHAR AS A POTENTIAL WEAPON AGAINST PLANT PATHOGENS

In 1847, Allen's report was among the earliest recognitions of biochar's potential in addressing plant diseases such as mildew and rust [50]. Substantial evidence suggests that biochar significantly reduces the occurrence and severity of diseases globally [51]. Biochar provides carbon, supporting the proliferation of lignin-degrading microbes like Dothideomycetes and Actinobacteriales. By promoting the growth of gamma and beta-proteobacteria, it aids in combating fungal pathogens. The application of biochar stimulates the development of beneficial microbes such as *Pseudomonas*, *Serratia*, and *Stenotrophomonas*, which play a crucial role in suppressing fungal pathogens [52].

3.1. Biochar for the Control of Plant Pathogenic Oomycetes

Employing a 5% (vol/vol) concentration of biochar derived from pine plant material initiates the activation of plant defenses, inducing systemic resistance. This effect has been observed in *Quercus rubra*, countering *Phytophthora cinnamomi*, and in *Acer rubrum*, combating *Phytophthora actorum*. This mechanism helps mitigate disease progression and alleviates physiological stress, as observed by Zwart and Kim [53]. Conversely, the application of softwood biochar resulted in increased root colonization by *Pythium multivium* in sweet pepper, lettuce, basil, and geranium. Nonetheless, there were no apparent adverse effects on the root system or hindrances in plant growth, as documented by Gravel et al. [54].

3.2. Biochar for the Control of Plant Pathogenic Fungi

The use of hardwood biochar resulted in an increase in the populations of beneficial organisms like arbuscular mycorrhizal fungi (AMFs), as demonstrated by Elmer [55]. Moreover, AMFs have been

acknowledged for their role in stimulating systemic resistance, as highlighted in a study by Poveda et al.[56]. These findings strongly suggest that biochar exerts its beneficial effects by altering soil characteristics or nutrient availability and promoting microbial communities in the soil, as discussed in research by Elmer[55] and Elmer and Pignatello[57]. Similar results were observed against pathogens like *F. solani* and *Ilyonectriadestructans* in *Panax ginseng* plants cultivated in soil enriched with rice husk biochar, attributed to changes in the microbial population in the rhizosphere, as demonstrated by Eo et al.[58].

Likewise, incorporating green waste biochar with compost (ranging from 0 to 3% concentration by weight) was found to mitigate *F. oxysporum* sp. *lycopersici*-induced wilt in tomatoes, linked to an increase in beneficial microorganisms. These microorganisms could potentially act through direct antagonistic effects or indirectly by inducing systemic resistance in plants, as suggested by Akhter et al.[59]. The positive effect of biochar derived from beech wood chips or garden waste in combating *F. oxysporum* sp. *lycopersici* in tomato plants likely arises from its ability to alter substances released by plant roots, which could play a crucial role in plant response to disease pressure, as proposed by Akhter et al.[60]. Additionally, biochar enhances the plant's ability to resist foliar plant pathogenic fungi. The primary mechanism identified for this enhanced resistance involves the systemic activation of defense mechanisms, as observed by Poveda et al.[56]. *Botrytis cinerea* is a versatile fungal pathogen responsible for necrotic disease in numerous host plants found across various plant species.

Utilizing biochar derived from various sources, such as holm oak for strawberry fields and eucalyptus wood chips for tomato cultivation, enhances the diversity of bacteria in the rhizosphere. This activation of systemic resistance against pathogens like *B. cinerea* results in a decrease in disease incidence[61]. Additionally, the application of biochar from greenhouse waste in tomato plants triggers early- and late-acting defense responses against *B. cinerea* by stimulating gene expression associated with jasmonic acid and ethylene responses. This process also increases the accumulation of active oxygen species like H_2O_2 , which are crucial for resistance[62].

Similarly, the incorporation of citrus wood biochar at varying percentages in tomato and pepper cultivation effectively reduces diseases caused by *B. cinerea* and *Leveillulataurica* (powdery mildew) by inducing systemic resistance[51]. Pepper plant waste biochar also demonstrates effectiveness against fungal pathogens such as *Colletotrichum acutatum* and *Podosphaera aphanis*[18], which cause anthracnose and powdery mildew in strawberries, respectively[63, 64]. In the *Rosellinia necatrix*–*Persea americana* pathosystem, the addition of biochar notably decreases both the severity and occurrence of the disease[65].

There is a strong correlation between biochar and the effectiveness of biocontrol fungus *Trichoderma harzianum* against *Macrophomina phaseolina*[66]. Combining biochar with *Bacillus subtilis* helps mitigate Fusarium wilt in radishes[67]. The potential for disease suppression varies

depending on factors such as the diversity of raw materials, production methods, application rates, and soil types [68]. For instance, corn stover biochar effectively manages soybean root rot caused by *Fusariumvirguliforme*, while sawdust and poultry waste biochar significantly reduce maize ear rot caused by *Fusariumverticilloides* [69, 70].

3.3. Biochar for the Control of Plant Pathogenic Bacteria

Recent research has investigated the use of biochar in addressing bacterial plant diseases, particularly focusing on combating bacterial wilt, a significant concern in global vegetable cultivation, attributed to *Ralstoniasolanacearum* [71]. The protective effects of biochar result from its capacity to improve soil physical and chemical properties, as well as promote an increased presence of beneficial bacteria and actinomycetes near plant roots. Consequently, this reduces the swarming capability and ability to colonize roots by *R. solanacearum*, as evidenced in studies conducted by Lu et al. [72], and Gu et al. [73].

3.4. Mechanism of plant disease control by application of biochar

Disease management can arise from both direct and indirect factors. The soil environment and the microbiome in the root zone are closely linked, and even small alterations in soil characteristics can affect the makeup of microorganisms surrounding the roots. It's crucial to note that while biochar's direct interactions may have a lesser impact on foliar pathogens, its effects on pathogens inhabiting the soil are typically more significant [74]. Table 1 lists the application of biochar and its presumed mechanism of action against different pathogens.

3.4.1. Induction of systemic plant defense

When biochar is administered away from the site of infection in foliar diseases, it initiates resistance, indicating systemic activation. This phenomenon may be attributed to chemical triggers, phytotoxic compounds, and the assistance of beneficial microorganisms. Various factors such as volatile compounds, minerals, and free radicals induce resistance in plants. Both systemic acquired resistance (SAR) and induced systemic resistance (ISR) pathways contribute to this process by prompting a rapid increase in lipid peroxidation and the production of superoxide. Consequently, free radicals are generated and polyunsaturated fatty acids are altered, ultimately suppressing diseases through systemic resistance [75, 76].

The application of biochar resulted in reduced disease severity in pepper and tomato plants infected with powdery mildew and graymold caused by *L. taurica* and *B. cinerea*, respectively. This decrease was attributed to the activation of induced systemic resistance [51]. Similarly, biochar application lowered the incidence of foliar diseases in strawberries caused by *P. aphanis* and *B. cinerea*, activating systemic resistance against these pathogens [18].

3.4.2. Inhibition of growth and development of pathogens

Biomass pyrolysis involves significant chemical transformations, breaking down O-alkyl carbons associated with carbohydrates while simultaneously producing aliphatic and aromatic carbon compounds [77]. Furthermore, this process can yield organic substances possessing strong antifungal properties. The resultant biochar comprises 29 primary compounds, including fatty acids, dicarboxylic acids, carboxylic acid derivatives, phenolic compounds, and tri-terpene acids. However, it is essential to recognize that the direct toxicity of biochar to pathogens is unlikely the sole reason for disease suppression. Various other mechanisms and factors are likely at play in achieving this outcome [78].

UNDER PEER REVIEW

Table 1 Biochar origin, application and putative mode of action against various plant pathogens

Pathogens	Pathosystem			Biochar		
	Species	Crop	Mode of action	Origin	Application rate	Reference
Bacteria	<i>Ralstoniasolanacearum</i>	Tomato	The soil bacteria and actinomycetes have experienced a rise in their population density	Wheat straw, Peanut shell	2% (Wt/Wt)	[72]
		Tobacco	Enhanced soil physicochemical characteristics have led to a greater abundance of bacteria in the rhizosphere	Rice straw	3 tons/ha	[11]
		Tomato	Decreased motility of pathogens and colonization in the rhizosphere have been observed	Pinewood	3% (Wt/Wt)	[73]
		Brinjal	Plants exhibit increased levels of phenolics, flavonoids, and peroxidase content	Leaf waste	6% (Wt/Wt)	[79]
	<i>Xanthomonascampestris</i> pv. <i>vesicatoria</i>	Chilli	Boosted synthesis of phenols, catalase, peroxidase, and flavonoids has been observed	Rice straw, Fly ash	3% (Wt/Wt)	[80]
Fungi	<i>Botrytis cinerea</i> , <i>Leveillulataurica</i>	Tomato, Pepper	Induced Systemic Resistance (ISR)	Citrus Wood	3 & 1% (Wt/Wt)	[51]
	<i>Podosphaeraaphanis</i>	Strawberry	Induced Systemic Resistance (ISR)	Pepper plant	3% (Wt/Wt)	[18]
	<i>Botrytis cinerea</i>	Tomato	Induced Systemic Resistance (ISR)	Greenhouse waste	3% (Wt/Wt)	[62]
	<i>Botrytis cinerea</i>	Strawberry	Rise in the diversity of bacteria within the rhizosphere	Holm Oak	3% (Wt/Wt)	[81]
	<i>Fusariumoxysporum</i> f.sp . <i>asparagi</i>	Asparagus	Root colonization by arbuscular mycorrhizal fungi (AMFs) has shown an increase	Hardwood Biomass	3% (Wt/Wt)	[55]
	<i>Botrytis cinerea</i>	Tomato	Increase in the variety and alterations in the metabolic capabilities of the microbiome in the rhizosphere	Eucalyptus wood chips	1 % (Wt/Wt)	[61]
	<i>Fusariumsacchari</i> , <i>Fusarium verticillioides</i> , <i>Fusariumproliferatum</i>	Sugarcane	Changes in the soil microbial community have resulted in increased diversity and beneficial modifications in soil functions	Palm residues	15tons/ ha	[82]

4. CHALLENGES AND CONSIDERATIONS IN BIOCHAR INTEGRATION

Contemporary literature extensively discusses the advantages of integrating biochar into sustainable agriculture. Nonetheless, it's imperative to acknowledge the potential drawbacks of biochar on soil health. Future research endeavors should prioritize identifying limitations that hinder its capacity to promote sustainable crop cultivation and combat climate change. The production of organic pollutants such as polycyclic aromatic hydrocarbons, furans, and dioxins during biochar pyrolysis poses risks to human health [83, 84]. Furthermore, biochar may serve as a reservoir for heavy metals like copper, cadmium, and nickel. For instance, the introduction of biochar could substantially elevate copper and arsenic levels in the soil, adversely affecting both soil quality and plant vitality [85, 86]. Aging biochar, influenced by biological organisms and environmental conditions, diminishes its ability to absorb heavy metals and detrimentally impacts soil-dwelling organisms [87]. Several studies have observed an increase in weed proliferation following the application of biochar, with weed growth rising by as much as 200% in lentil cultivation [88]. Research indicates that the benefits of biochar vary depending on soil composition, plant species, and specific plant components. For example, applying biochar promoted the vegetative growth of tomatoes without affecting fruit yield [89]. Further investigation and attention are warranted to understand the limitations associated with biochar utilization and devise strategies to mitigate them for the advancement of sustainable agricultural practices.

5. CONCLUSION

Biochar emerges as a promising solution for enhancing crop yields and safeguarding crops in a manner that is both environmentally friendly and sustainable. It offers a cost-effective strategy that aligns well with conservation initiatives. Furthermore, biochar contributes to carbon capture, thereby improving crop productivity and resilience to various environmental pressures. Produced via pyrolysis, it represents an energy-efficient process. While the direct relationship between biochar application and disease suppression may vary, it nonetheless supports soil health by promoting microbial growth and supplying essential nutrients to plants. Disease management could also involve biochemical or physiological shifts that activate systemic resistance. Consequently, integrating biochar into soil has the potential to significantly influence the evolution of sustainable agricultural practices.

6. FUTURE PROSPECTS

Future research is crucial to uncover the metabolites involved in the interaction between pathogens and host plants within soil treated with biochar. Further investigations are needed to elucidate the disease-suppressing capabilities and novel mechanisms at play. The impact of different biochar types

on diverse pathogens remains unexplored, and additional studies are required to determine methods for promoting beneficial soil microbes without inadvertently increasing pathogen populations and virulence. A deeper comprehension of biochar's chemical and physical properties and their effects on pathogenic microbes, as well as the signalling pathways implicated in inducing disease resistance, is imperative. While most research has been conducted in controlled environments, long-term field studies are essential to assess the practicality of biochar in combating plant diseases. Many farmers lack awareness of the myriad benefits of biochar, underscoring the importance of offering training and workshops to disseminate knowledge about its promising long-term advantages in organic agriculture and establish it as a viable method for disease control in the future.

References

1. United Nations. World population prospects: The 2019 revision. Department of Economic and Social Affairs, Population Division, United Nations. Online Edition. Rev.1. 2019. <https://population.un.org/wpp/Download/Standard/Population/>.
2. Giller KE, Delaune T, Silva JV, Descheemaeker K, van de Ven G, Schut AG, et al. The future of farming: Who will produce our food?. *Food Security*. 2021;13(5):1073-99. <https://doi.org/10.1007/s12571-021-01184-6>
3. Deutsch CA, Tewksbury JJ, Tigchelaar M, Battisti DS, Merrill SC, Huey RB, et al. Increase in crop losses to insect pests in a warming climate. *Science*. 2018;361(6405):916-9. <https://doi.org/10.1126/science.aat3466>
4. Roopa KP, Gadag AS. Management of Soil-Borne Diseases of Plants Through Some Cultural Practices and Actinobacteria. *Plant Health Under Biotic Stress: Volume 1: Organic Strategies*. 2019;129-45. https://doi.org/10.1007/978-981-13-6043-5_7
5. Egamberdieva D, Jabborova D, Hashem A. *Pseudomonas* induces salinity tolerance in cotton (*Gossypiumhirsutum*) and resistance to *Fusarium* root rot through the modulation of indole-3-acetic acid. *Saudi journal of biological sciences*. 2015;22(6):773-9. <https://doi.org/10.1016/j.sjbs.2015.04.019>
6. Jaiswal AK, Graber ER, Elad Y, Frenkel O. Biochar as a management tool for soilborne diseases affecting early stage nursery seedling production. *Crop protection*. 2019;120:34-42. <https://doi.org/10.1016/j.cropro.2019.02.014>
7. Lehmann J, Gaunt J, Rondon M. Bio-char sequestration in terrestrial ecosystems—a review. *Mitigation and adaptation strategies for global change*. 2006;11:403-27. <https://doi.org/10.1007/s11027-005-9006-5>
8. Lehmann J, Joseph S. Biochar for environmental management: an introduction. In *Biochar for environmental management* 2015 (pp. 1-13). Routledge. <https://doi.org/10.4324/9780203762264>

9. Gheorghe C, Marculescu C, Badea A, Dinca C, Apostol T. Effect of pyrolysis conditions on bio-char production from biomass. In Proceedings of the 3rd WSEAS Int. Conf. on RENEWABLE ENERGY SOURCES 2009 Jul 1 (pp. 239-241). Tenerife, Canary Islands Spain: University of La Laguna.
10. Yaman S. Pyrolysis of biomass to produce fuels and chemical feedstocks. Energy conversion and management. 2004;45(5):651-71. [https://doi.org/10.1016/S0196-8904\(03\)00177-8](https://doi.org/10.1016/S0196-8904(03)00177-8)
11. Zhang C, Lin Y, Tian X, Xu Q, Chen Z, Lin W. Tobacco bacterial wilt suppression with biochar soil addition associates to improved soil physiochemical properties and increased rhizosphere bacteria abundance. Applied Soil Ecology. 2017;112:90-6. <https://doi.org/10.1016/j.apsoil.2016.12.005>
12. Lehmann J, Kern D, German L, McCANN J, Martins GC, Moreira A. Soil fertility and production potential. Amazonian Dark Earths: Origin Properties Management. 2003;105-24. https://doi.org/10.1007/1-4020-2597-1_6
13. Novak JM, Busscher WJ, Watts DW, Amonette JE, Ippolito JA, Lima IM, et al. Biochars impact on soil-moisture storage in an ultisol and two aridisols. Soil Science. 2012;177(5):310-20. <https://doi.org/10.1097/SS.0b013e31824e5593>
14. Barrow CJ. Biochar: potential for countering land degradation and for improving agriculture. Applied Geography. 2012;34:21-8. <https://doi.org/10.1016/j.apgeog.2011.09.008>
15. Rassaei F. Rice yield and carbon dioxide emissions in a paddy soil: a comparison of biochar and polystyrene microplastics. Environmental Progress and Sustainable Energy. 2024;43(1):e14217. <https://doi.org/10.1002/ep.14217>
16. Rassaei F. Biochar effects on rice paddy cadmium contaminated calcareous clay soil: A study on adsorption kinetics and cadmium uptake. Paddy and Water Environment. 2023a;21(3):389-400. <https://doi.org/10.1007/s10333-023-00937-7>
17. Rassaei F. Sugarcane bagasse biochar affects corn (*Zea mays* L.) growth in cadmium and lead-contaminated calcareous clay soil. Arabian journal of Geosciences. 2023b;16(3):181. <https://doi.org/10.1007/s12517-023-11225-3>
18. Meller Harel Y, Elad Y, Rav-David D, Borenstein M, Shulchani R, Lew B, et al. Biochar mediates systemic response of strawberry to foliar fungal pathogens. Plant and Soil. 2012;357:245-57. <https://doi.org/10.1007/s11104-012-1129-3>
19. Bonanomi G, D'Ascoli R, Scotti R, Gaglione SA, Caceres MG, Sultana S, et al. Soil quality recovery and crop yield enhancement by combined application of compost and wood to vegetables grown under plastic tunnels. Agriculture, Ecosystems and Environment. 2014;192:1-7. <https://doi.org/10.1016/j.agee.2014.03.029>
20. Bonanomi G, Ippolito F, Scala F. A "black" future for plant pathology? Biochar as a new soil amendment for controlling plant diseases. Journal of Plant Pathology. 2015;97(2). <https://doi.org/10.4454/jpp.v97i2.3381>

21. Naeem MA, Khalid M, Aon M, Abbas G, Tahir M, Amjad M, et al. Effect of wheat and rice straw biochar produced at different temperatures on maize growth and nutrient dynamics of a calcareous soil. *Archives of Agronomy and Soil Science*. 2017;63(14):2048-61. <https://doi.org/10.1080/03650340.2017.1325468>
22. Ali S, Ullah MI, Sajjad A, Shakeel Q, Hussain A. Environmental and health effects of pesticide residues. *Sustainable Agriculture Reviews 48: Pesticide Occurrence, Analysis and Remediation Vol. 2 Analysis*. 2021;311-36. https://doi.org/10.1007/978-3-030-54719-6_8
23. Laird DA. The charcoal vision: a win–win–win scenario for simultaneously producing bioenergy, permanently sequestering carbon, while improving soil and water quality. *Agronomy journal*. 2008;100(1):178-81. <https://doi.org/10.2134/agronj2007.0161>
24. Jeffery S, Verheijen FG, van der Velde M, Bastos AC. A quantitative review of the effects of biochar application to soils on crop productivity using meta-analysis. *Agriculture, ecosystems and environment*. 2011;144(1):175-87. <https://doi.org/10.1016/j.agee.2011.08.015>
25. Lehmann J, Rillig MC, Thies J, Masiello CA, Hockaday WC, Crowley D. Biochar effects on soil biota—a review. *Soil biology and biochemistry*. 2011;43(9):1812-36. <https://doi.org/10.1016/j.soilbio.2011.04.022>
26. Poveda J, Martinez-Gomez A, Fenoll C, Escobar C. The use of biochar for plant pathogen control. *Phytopathology*. 2021;111(9):1490-9. <https://doi.org/10.1094/PHYTO-06-20-0248-RVW>
27. Al-Wabel MI, Al-Omran A, El-Naggar AH, Nadeem M, Usman AR. Pyrolysis temperature induced changes in characteristics and chemical composition of biochar produced from conocarpus wastes. *Bioresource technology*. 2013;131:374-9. <https://doi.org/10.1016/j.biortech.2012.12.165>
28. Han L, Sun K, Yang Y, Xia X, Li F, Yang Z, et al. Biochar's stability and effect on the content, composition and turnover of soil organic carbon. *Geoderma*. 2020;364:114184. <https://doi.org/10.1016/j.geoderma.2020.114184>
29. Shi RY, Hong ZN, Li JY, Jiang J, Baquy MA, Xu RK, et al. Mechanisms for increasing the pH buffering capacity of an acidic Ultisol by crop residue-derived biochars. *Journal of Agricultural and food chemistry*. 2017;65(37):8111-9. <https://doi.org/10.1021/acs.jafc.7b02266>
30. Xia H, Riaz M, Zhang M, Liu B, El-Desouki Z, Jiang C. Biochar increases nitrogen use efficiency of maize by relieving aluminium toxicity and improving soil quality in acidic soil. *Ecotoxicology and environmental safety*. 2020;196:110531. <https://doi.org/10.1016/j.ecoenv.2020.110531>
31. Brassard P, Godbout S, Raghavan V. Soil biochar amendment as a climate change mitigation tool: key parameters and mechanisms involved. *Journal of environmental management*. 2016;181:484-97. <https://doi.org/10.1016/j.jenvman.2016.06.063>

32. Liu G, Xie M, Zhang S. Effect of organic fraction of municipal solid waste (OFMSW)-based biochar on organic carbon mineralization in a dry land soil. *Journal of Material Cycles and Waste Management*. 2017;19:473-82. <https://doi.org/10.1007/s10163-015-0447-y>
33. Luo C, Yang J, Chen W, Han F. Effect of biochar on soil properties on the Loess Plateau: Results from field experiments. *Geoderma*. 2020;369:114323. <https://doi.org/10.1016/j.geoderma.2020.114323>
34. Blanco-Canqui H. Biochar and soil physical properties. *Soil Science Society of America Journal*. 2017;81(4):687-711. <https://doi.org/10.2136/sssaj2017.01.0017>
35. Razzaghi F, Obour PB, Arthur E. Does biochar improve soil water retention? A systematic review and meta-analysis. *Geoderma*. 2020;361:114055. <https://doi.org/10.1016/j.geoderma.2019.114055>
36. Barros JA, Medeiros EV, Notaro KA, Moraes WS, Silva JM, Nascimento TC, et al. Different cover promote sandy soil suppressiveness to root rot disease of cassava caused by *Fusariumsolani*. *African Journal of Microbiology Research*. 2014;8(10):967-73. <https://doi.org/10.5897/AJMR2014.6607>
37. de Medeiros EV, dos Santos Moraes MD, da Costa DP, Duda GP, de Oliveira JB, Araujo da Silva JS, et al. Effect of biochar and inoculation with *Trichodermaaureoviride* on melon growth and sandy Entisol quality. *Australian Journal of Crop Science*. 2020;14(6):971-7. <https://doi.org/10.21475/ajcs.20.14.06.p2302>
38. Barrios E. Soil biota, ecosystem services and land productivity. *Ecological economics*. 2007;64(2):269-85. <https://doi.org/10.1016/j.ecolecon.2007.03.004>
39. Coban O, De Deyn GB, van der Ploeg M. Soil microbiota as game-changers in restoration of degraded lands. *Science*. 2022;375(6584):abe0725. <https://doi.org/10.1126/science.abe0725>
40. Lehmann J. Bio- energy in the black. *Frontiers in Ecology and the Environment*. 2007;5(7):381-7. <https://doi.org/10.1890/1540-9295>
41. Tryon EH. Effect of charcoal on certain physical, chemical, and biological properties of forest soils. *Ecological Monographs*. 1948;18(1):81-115. <https://doi.org/10.2307/1948629>
42. Liang B, Lehmann J, Sohi SP, Thies JE, O'Neill B, Trujillo L, et al. Black carbon affects the cycling of non-black carbon in soil. *Organic Geochemistry*. 2010;41(2):206-13. <https://doi.org/10.1016/j.orggeochem.2009.09.007>
43. Wang M, Sun Y, Gu Z, Wang R, Sun G, Zhu C, et al. Nitrate protects cucumber plants against *Fusariumoxysporum* by regulating citrate exudation. *Plant and Cell Physiology*. 2016;57(9):2001-12. <https://doi.org/10.1093/pcp/pcw124>
44. Xu N, Tan G, Wang H, Gai X. Effect of biochar additions to soil on nitrogen leaching, microbial biomass and bacterial community structure. *European journal of soil biology*. 2016;74:1-8. <https://doi.org/10.1016/j.ejsobi.2016.02.004>

45. Ali N, Khan S, Yao H, Wang J. Biochars reduced the bioaccessibility and (bio) uptake of organochlorine pesticides and changed the microbial community dynamics in agricultural soils. *Chemosphere*. 2019;224:805-15. <https://doi.org/10.1016/j.chemosphere.2019.02.163>
46. Igalavithana AD, Kim KH, Jung JM, Heo HS, Kwon EE, Tack FM, et al. Effect of biochar pyrolyzed in N₂ and CO₂, and feedstock on microbial community in metal (loid) contaminated soils. *Environment international*. 2019;126:791-801. <https://doi.org/10.1016/j.envint.2019.02.061>
47. Soothar MK, Hamani AK, Sardar MF, Sootahar MK, Rahim R, Abubakar SA, et al. Maize (*Zea mays* L.) seedlings rhizosphere microbial community as responded to acidic biochar amendment under saline conditions. *Frontiers in Microbiology*. 2021;12:789235. <https://doi.org/10.3389/fmicb.2021.789235>
48. Kavitha B, Reddy PV, Kim B, Lee SS, Pandey SK, Kim KH. Benefits and limitations of biochar amendment in agricultural soils: A review. *Journal of environmental management*. 2018;227:146-54. <https://doi.org/10.1016/j.jenvman.2018.08.082>
49. Brtnicky M, Datta R, Holatko J, Bielska L, Gusiatin ZM, Kucerik J, et al. A critical review of the possible adverse effects of biochar in the soil environment. *Science of the Total Environment*. 2021;796:148756. <https://doi.org/10.1016/j.scitotenv.2021.148756>
50. Allen RL. A brief compend of American agriculture. CM Saxton, 1847.
51. Elad Y, David DR, Harel YM, Borenshtein M, Kalifa HB, Silber A, et al. Induction of systemic resistance in plants by biochar, a soil-applied carbon sequestering agent. *Phytopathology*. 2010;100(9):913-21. <https://doi.org/10.1094/PHYTO-100-9-0913>
52. Sadegh-Zadeh F, Parichehreh M, Jalili B, Bahmanyar MA. Rehabilitation of calcareous saline-sodic soil by means of biochars and acidified biochars. *Land Degradation and Development*. 2018;29(10):3262-71. <https://doi.org/10.1002/ldr.3079>
53. Zwart DC, Kim SH. Biochar amendment increases resistance to stem lesions caused by *Phytophthora* spp. in tree seedlings. *HortScience*. 2012;47(12):1736-40. <https://doi.org/10.21273/HORTSCI.47.12.1736>
54. Gravel V, Dorais M, Menard C. Organic potted plants amended with biochar: its effect on growth and *Pythium* colonization. *Canadian Journal of Plant Science*. 2013;93(6):1217-27. <https://doi.org/10.4141/cjps2013-315>
55. Elmer WH. Effect of leaf mold mulch, biochar, and earthworms on mycorrhizal colonization and yield of asparagus affected by *Fusarium* crown and root rot. *Plant disease*. 2016;100(12):2507-12. <https://doi.org/10.1094/PDIS-10-15-1196-RE>
56. Poveda J, Abril-Urias P, Escobar C. Biological control of plant-parasitic nematodes by filamentous fungi inducers of resistance: *Trichoderma*, mycorrhizal and endophytic fungi. *Frontiers in Microbiology*. 2020;11:530260. <https://doi.org/10.3389/fmicb.2020.00992>

57. Elmer WH, Pignatello JJ. Effect of biochar amendments on mycorrhizal associations and *Fusarium* crown and root rot of asparagus in replant soils. *Plant Disease*. 2011;95(8):960-6. <https://doi.org/10.1094/PDIS-10-10-0741>
58. Eo J, Park KC, Kim MH, Kwon SI, Song YJ. Effects of rice husk and rice husk biochar on root rot disease of ginseng (*Panax ginseng*) and on soil organisms. *Biological Agriculture and Horticulture*. 2018;34(1):27-39. <https://doi.org/10.1080/01448765.2017.1363660>
59. Akhter A, Hage-Ahmed K, Soja G, Steinkellner S. Potential of *Fusarium* wilt-inducing chlamydospores, in vitro behaviour in root exudates and physiology of tomato in biochar and compost amended soil. *Plant and Soil*. 2016;406:425-40. <https://doi.org/10.1007/s11104-016-2948-4>
60. Akhter A, Hage-Ahmed K, Soja G, Steinkellner S. Compost and biochar alter mycorrhization, tomato root exudation, and development of *Fusariumoxysporum* f. sp. *lycopersici*. *Frontiers in Plant Science*. 2015;6:146969. <https://doi.org/10.3389/fpls.2015.00529>
61. Kolton M, Graber ER, Tsehansky L, Elad Y, Cytryn E. Biochar- stimulated plant performance is strongly linked to microbial diversity and metabolic potential in the rhizosphere. *New Phytologist*. 2017;213(3):1393-404. <https://doi.org/10.1111/nph.14253>
62. Mehari ZH, Elad Y, Rav-David D, Graber ER, Meller Harel Y. Induced systemic resistance in tomato (*Solanumlycopersicum*) against *Botrytis cinerea* by biochar amendment involves jasmonic acid signaling. *Plant and Soil*. 2015;395:31-44. <https://doi.org/10.1007/s11104-015-2445-1>
63. Carisse O, Morissette-Thomas V, Van der Heyden H. Lagged association between powdery mildew leaf severity, airborne inoculum, weather, and crop losses in strawberry. *Phytopathology*. 2013;103(8):811-21. <https://doi.org/10.1094/PHYTO-11-12-0300-R>
64. Dinler H, Benlioglu S. The possibility to control diseases caused by *Colletotrichum* species in strawberries. *International Journal of Agriculture Forestry and Life Sciences*. 2019;3(2):362-70.
65. Vida C, de Vicente A, Cazorla FM. The role of organic amendments to soil for crop protection: induction of suppression of soilborne pathogens. *Annals of Applied Biology*. 2020;176(1):1-5. <https://doi.org/10.1111/aab.12555>
66. Araujo AS, Blum LE, Figueiredo CC. Biochar and *Trichodermaharzianum* for the Control of *Macrophominaphaseolina*. *Brazilian Archives of Biology and Technology*. 2019;62:e19180259. <https://doi.org/10.1590/1678-4324-2019180259>
67. Chen W, Wu Z, Liu C, Zhang Z, Liu X. Biochar combined with *Bacillus subtilis* SL-44 as an eco-friendly strategy to improve soil fertility, reduce *Fusarium* wilt, and promote radish growth. *Ecotoxicology and Environmental Safety*. 2023;251:114509. <https://doi.org/10.1016/j.ecoenv.2023.114509>

68. Gao Y, Lu Y, Lin W, Tian J, Cai K. Biochar suppresses bacterial wilt of tomato by improving soil chemical properties and shifting soil microbial community. *Microorganisms*. 2019;7(12):676. <https://doi.org/10.3390/microorganisms7120676>
69. Rogovska N, Laird D, Leandro L, Aller D. Biochar effect on severity of soybean root disease caused by *Fusariumvirguliforme*. *Plant and soil*. 2017;413:111-26. <https://doi.org/10.1007/s11104-016-3086-8>
70. Akanmu AO, Sobowale AA, Abiala MA, Olawuyi OJ, Odebode AC. Efficacy of biochar in the management of *Fusariumverticillioides*Sacc. causing ear rot in *Zea mays* L. *Biotechnology Reports*. 2020;26:e00474. <https://doi.org/10.1016/j.btre.2020.e00474>
71. Choudhary DK, Nabi SU, Dar MS, Khan KA. *Ralstoniasolanacearum*: A wide spread and global bacterial plant wilt pathogen. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(2):85-90.
72. Lu Y, Rao S, Huang F, Cai Y, Wang G, Cai K. Effects of biochar amendment on tomato bacterial wilt resistance and soil microbial amount and activity. *International Journal of Agronomy*. 2016;2016. <https://doi.org/10.1155/2016/2938282>
73. Gu Y, Hou Y, Huang D, Hao Z, Wang X, Wei Z, et al. Application of biochar reduces *Ralstoniasolanacearum* infection via effects on pathogen chemotaxis, swarming motility, and root exudate adsorption. *Plant and Soil*. 2017;415:269-81. <https://doi.org/10.1007/s11104-016-3159-8>
74. Graber ER, Elad Y. Biochar impact on plant resistance to disease. *Biochar and soil biota*. 2013;278. <https://doi.org/10.1201/b14585-3>
75. Formela M, Samardakiewicz S, Marczak L, Nowak W, Narozna D, Bednarski W, et al. Effects of endogenous signals and *Fusariumoxysporum* on the mechanism regulating genistein synthesis and accumulation in yellow lupine and their impact on plant cell cytoskeleton. *Molecules*. 2014;19(9):13392-421. <https://doi.org/10.3390/molecules190913392>
76. Abd-Allah EF, Abeer H, Alqarawi AA, Hend AA. Alleviation of adverse impact of cadmium stress in sunflower (*Helianthus annuus* L.) by arbuscular mycorrhizal fungi. *Pakistan Journal of Botany*. 2015;47(2):785-95.
77. Chen D, Cen K, Zhuang X, Gan Z, Zhou J, Zhang Y, et al. Insight into biomass pyrolysis mechanism based on cellulose, hemicellulose, and lignin: Evolution of volatiles and kinetics, elucidation of reaction pathways, and characterization of gas, biochar and bio-oil. *Combustion and Flame*. 2022;242:112142. <https://doi.org/10.1016/j.combustflame.2022.112142>
78. Torikai K, Yoshida S, Takahashi H. Effects of temperature, atmosphere and pH on the generation of smoke compounds during tobacco pyrolysis. *Food and chemical toxicology*. 2004;42(9):1409-17. <https://doi.org/10.1016/j.fct.2004.04.002>

79. Ahmad CA, Akhter A, Haider MS, Abbas MT, Hashem A, Avila-Quezada GD, et al. Demonstration of the synergistic effect of biochar and *Trichoderma harzianum* on the development of *Ralstoniasolanacearum* in eggplant. *Frontiers in Microbiology*. 2024;15:1360703. <https://doi.org/10.3389/fmicb.2024.1360703>
80. Akhter A, Asif M, Anwar W, Aftab Z, Khurshid M, Hashem A, et al. Biochemical and physiological elucidation of biochar induced defense response activation against bacterial leaf spot in chilies. *Applied Ecology and Environmental Research*. 2023;21(4). https://doi.org/10.15666/aeer/2104_30573074
81. De Tender CA, Debode J, Vandecasteele B, D'Hose T, Cremelie P, Haegeman A, et al. Biological, physicochemical and plant health responses in lettuce and strawberry in soil or peat amended with biochar. *Applied Soil Ecology*. 2016;107:1–12. <https://doi.org/10.1016/j.apsoil.2016.05.001>
82. Ahmad S, Zhai X, Wang M, Shi Y, Chen Y, Liang Q, et al. Biochar amendments improve soil functionalities, microbial community and reduce Pokkahboeng disease of sugarcane. *Chemical and Biological Technologies in Agriculture*. 2024;11(1):28. <https://doi.org/10.1186/s40538-024-00546-4>
83. Garcia-Perez M, Metcalf J. The formation of polyaromatic hydrocarbons and dioxins during pyrolysis: A review of the literature with descriptions of biomass composition, fast pyrolysis technologies and thermochemical reactions. Washington State University Extension, Pullman, Washington, 2008.
84. Hale SE, Lehmann J, Rutherford D, Zimmerman AR, Bachmann RT, Shitumbanuma V, et al. Quantifying the total and bioavailable polycyclic aromatic hydrocarbons and dioxins in biochars. *Environmental science and technology*. 2012;46(5):2830-8. <https://doi.org/10.1021/es203984k>
85. Beesley L, Moreno-Jimenez E, Gomez-Eyles JL. Effects of biochar and greenwaste compost amendments on mobility, bioavailability and toxicity of inorganic and organic contaminants in a multi-element polluted soil. *Environmental pollution*. 2010;158(6):2282-7. <https://doi.org/10.1016/j.envpol.2010.02.003>
86. Chen D, Liu X, Bian R, Cheng K, Zhang X, Zheng J, et al. Effects of biochar on availability and plant uptake of heavy metals—A meta-analysis. *Journal of Environmental Management*. 2018;222:76-85. <https://doi.org/10.1016/j.jenvman.2018.05.004>
87. Hale S, Hanley K, Lehmann J, Zimmerman A, Cornelissen G. Effects of chemical, biological, and physical aging as well as soil addition on the sorption of pyrene to activated carbon and biochar. *Environmental science and technology*. 2011;45(24):10445-53. <https://doi.org/10.1021/es202970x>
88. Safaei Khorram M, Fatemi A, Khan MA, Kiefer R, Jafarnia S. Potential risk of weed outbreak by increasing biochar's application rates in slow-growth legume, lentil (*Lens*

culinaris Medik.). Journal of the Science of Food and Agriculture. 2018;98(6):2080-8. <https://doi.org/10.1002/jsfa.8689>

89. Vaccari FP, Maienza A, Miglietta F, Baronti S, Di Lonardo S, Giagnoni L, et al. Biochar stimulates plant growth but not fruit yield of processing tomato in a fertile soil. Agriculture, Ecosystems and Environment. 2015;207:163-70. <https://doi.org/10.1016/j.agee.2015.04.015>

UNDER PEER REVIEW