

# SIGNALLING: GROWTH PROMOTERS AND INHIBITORS

## ABSTRACT

Plant growth and development are intricately regulated by a multitude of signaling pathways that respond to internal and external cues. Central to these regulatory mechanisms are the signaling pathways of growth promoters and inhibitors. Various growth-promoting hormones, including auxins, cytokinins, gibberellins, and brassinosteroids, stimulate cell division, elongation, and differentiation, playing vital roles in shaping plant architecture and influencing various developmental processes. The pathways of these growth promoters involve specific receptors and transcription factors that orchestrate growth in response to dynamic signals. In contrast, growth inhibitors, represented by hormones like abscisic acid, ethylene, jasmonic acid, and salicylic acid, act as crucial checkpoints that restrict growth during stress, pathogen attacks, or other adverse conditions. These signaling pathways trigger growth inhibition and activate defense mechanisms, enabling plants to adapt to challenging environments. Balancing the interplay between growth promoters and inhibitors allows plants to allocate resources efficiently and ensure their survival under changing circumstances. Understanding these signaling pathways is not only fundamental to plant biology but also essential for applications in agriculture, horticulture, and environmental management. This abstract provides a concise overview of these essential processes. This knowledge enables us to manipulate plant growth and development for improved crop yields, disease resistance, and stress tolerance. Ongoing research continues to unravel the complexities of these pathways, offering valuable insights into how plants delicately balance growth promotion and inhibition to thrive in an ever-evolving world.

*Keywords:* Auxin, Brassinosteroids, Cytokinin, Ethylene, Gibberellin, Salicylic acid.

## 1. INTRODUCTION

Plant growth and development are tightly controlled by a diverse set of signalling molecules, which are including regulators and inhibitors. These chemicals are critical in coordinating many parts of plant life, including germinating and cell elongation through blooming, fruit development, and stress response. Plants can adapt and grow in ever-changing settings due to the dynamic interaction of these chemicals. Plant growth hormones, commonly referred to as phytohormones, are chemical messengers that have a significant impact on several developmental processes within plants. These hormones are synthesized in certain tissues and delivered to specific locations to influence growth and development. Auxins, gibberellins, cytokinin's, abscisic acid, ethylene, and brassinosteroids are some of the primary plant growth hormones, each with its own particular role in forming the plant's architecture. Plant growth inhibitors, on the other hand, serve to restrict or inhibit specific components of plant development, as the name implies. While this may appear to be counterproductive, inhibitors are necessary for preserving plant equilibrium. They, for example, aid in the prevention of excessive growth and the regulation of responses to environmental stimuli. Abscisic acid is a well-known inhibitor during drought circumstances, Jasmonic acid in response to herbivory or pathogen assault, and salicylic acid during specific stress reactions. In plants the roles and quantities of these growth hormones and inhibitors are carefully regulated to respond to the plant's individual demands, developmental stage, and environmental difficulties. This sophisticated regulatory network lets plants adapt to light, gravity, temperature, and other environmental conditions, guaranteeing their survival and successful reproduction. Recognizing the application of hormones that stimulate plant growth and inhibitors is critical for horticulturalists, agriculturists, and other professionals who want to boost the productivity of crops, boost stress tolerance, and regulate plant development for a variety of reasons. It demonstrates the plant kingdom's extraordinary intricacy and flexibility, as these molecular messengers choreograph a complex pattern of development and evolution in order to maintain the plant's general well-being. "In actuality, these hormones exert many of their effects via a complicated chain of interactions with other hormones and signaling molecules. PGRs, also known


as biostimulants or bio inhibitors, function inside plant cells to stimulate or inhibit certain enzymes or enzyme systems, therefore aiding in the regulation of plant metabolism”[1]. “Botanical hormones are biological chemical messengers that regulate the maturation and growth of plants as well as reactions to stresses caused by abiotic and biotic factors, allowing plant homeostasis and adaptability to environmental changes”[2]. “Temperature is an environmental component that has a significant impact on the plant's development program. It is crucial in modulating the degree of dormancy experienced by the seeds”[3]. Plant hormones have powerful, seemingly independent effects on plant development as well as growth. Auxin stimulates growth in general; cytokinin's delay leaf senescence and impact cell division; abscisic acid causes bud hibernation and promotes stomatal closure; gibberellins aid in the sprouting of seeds.

## 2.CROSSTALK IN PHYTOHORMONAL SIGNALING IN PLANT GROWTH AND DEVELOPMENT:

### 2.1 signaling of auxin:

Initially, auxin signaling molecules were explored by studying the effects of exogenous administration of hormone combinations on plant proliferation or morphology [4].“Hormones, as endogenous variables, are critical in controlling developmental processes.Hormonal control of development is a complicated process including interactions between numerous hormones at the transcriptional, translational, and cellular levels”[5].“Furthermore, auxin is a key player in the


**Fig: 1. hormonal crosstalk controlling the proliferation of plants and development, with auxin at the centre.**



current hormonal crosstalk that influences several developmental phases”[6,5,7] Fig 1.It regulates cell division, elongation, and programmed cell death, hence influencing embryonic and post-embryonic development [8]

“Auxin's impact is cell type dependent: at the same dose, indole-3-acetic acid (IAA) boosts hypocotyl elongation, suppresses main root development, initiates lateral root initiation, and stimulates root hair creation”[2].“Auxin and cytokinin have been demonstrated to have both synergistic and antagonistic effects on shoot and root growth”[6,9].“Auxin and ethylene are two other hormones that interact in a traditional and complicated way. In regulating various developmental processes such as apical hook formation, root and shoot extension, root and shoot gravitational attraction, lateral root formation, root hair initiation and elongation, hypocotyl phototropism, and leaf abscission, they exhibit both synergistic and antagonistic interactions”[10].“Gibberellin (GA) and auxin have been shown to regulate pea stem elongation and senescence by[6] populus lateral root development, and Arabidopsis apical dominance, inflorescence, and root development”[11]. “Brassinolide (BR) combines with auxin to control cell elongation in roots, shoots, and tropisms”[12].

65 “Jasmonic acid (JA) and auxin work together to govern root growth, lateral root development, and flowering, but also  
 66 work against each other to control coleoptile elongation”[5]. “According to recent research, ABA and auxin interact to  
 67 regulate lateral root formation”[13]. “Auxin is distinctive among plant hormones in that it can travel both long and small  
 68 distances. Short-distance movement is slower, happens cell-to-cell, and is controlled by influx and exhaust carrier  
 69 proteins”[10]. “Auxin distribution in the root is more complicated, having two different polarity”[14]. “Through the outer  
 70 layers of root cells, IAA flows in the opposite way towards the shoot. IAA shootward migration is controlled by AUX1,  
 71 PIN2, and ABCB4. Auxin and cytokinin (CK) are phytohormones that regulate important cell specification processes,  
 72 drive growth, and are present during embryogenesis and establishing seedlings”[15,16]. There are several sites where  
 73 the mechanisms of auxin and cytokinin networks interact. In addition to the inter-hormonal interconnections revealed,  
 74 both hormones control their own metabolism and perception, which adds to the pathways' complexity. Many critical  
 75 elements of plant growth rely on interactions between auxins and cytokinin (e.g., shoot branching, the tip of the shoot  
 76 meristem, the root transition zone, lateral root initiation, and the root meristem[17] Fig 2.



77 **Fig:2. auxin interacts with cytokinin in various development stages.**

78 The auxin and cytokinin pathways interact in a variety of ways[17]. [9] proposed a model in which the size of the  
 79 meristem of the root is determined by enmity between auxin and cytokinin in the vascular evolution zone; this  
 80 antagonistic effect is mediated by battling legislation of SHY2, which in turn interferes with cytokinin biosynthesis while  
 81 suppressing auxin transport and signalling (Fig. 2)demonstrated that auxin initiates the activation of two type A ARRs,  
 82 ARR7 and ARR15, blocking cytokinin signalling [18] (Fig. 2)).The Aux/IAA SHORT HYPOCOTYL2 (SHY2) gene, a  
 83 regulator of auxin signalling, plays an important role in meristem growth and development [19],[20]show in this work  
 84 that ethylene-auxin cross-talk requires a dynamic relationship among ERF.B3 and IAA27. The evidence they provide to  
 85 support this is that overexpression of an important inhibitor form of ERF.B3 results in modified auxin sensitivity, root  
 86 development, and impaired chlorophyll accumulation, similar to the phenotypes of lines with downregulated IAA27  
 87 expression.


88 **2.2Signalling of Gibberellin:**

89 Gibberellins have a significant impact on germination, the advancement of plants in general (mostly through cell  
 90 expansion), floral development, and blooming duration. They regulate seed germination, leaf growth, stem elongation,  
 91 and blooming. The manner of GA activity in planta is yet unknown, as it is influenced by various both positive and  
 92 negative synergistic relationships with other domestic and environmental signals[21] (Fig. 3).Several genes that encode  
 93 GA signalling components involved in the initiation of seed germination have also been identified[22]. RGL2 inhibits GA  
 94 responses that predominantly affect how quickly seeds germinate[23]. SPY, another Arabidopsis gene, is a negative  
 95 control of GA signalling. However, unlike RGL2, it also governs all other GA-related developmental processes. SPY is  
 96 also a cytokinin signalling pathway activator[24].

97 GAs controls a variety of developmental activities in plants throughout their life cycle, from allowing seeds to germinate  
 98 through the broadening of leaves, stem elongation, flower induction, and seed formation[25]. Because GA interactions  
 99 with other hormones include components of the GA biosynthesis and reaction networks[25].of several developmental  
 100 processes. Unlike GA, which promotes germination, development, and blooming, ABA suppresses these activities.  
 101 Furthermore, the antagonistic interaction and ratio among these two hormones control the transition from  
 102 morphogenesis to seed viability. Several mechanisms have been discovered to underpin this antagonistic interplay in  
 103 several developmental phases (Fig 3).GA and auxin functions overlap in terms of cell growth and tissue differentiation  
 104 control. Auxin influences both GA signalling and GA biosynthesis (Fig. 3). GA promotion of root elongation in

105  
106  
107  
108  
109  
110  
111

Arabidopsis has been demonstrated to need auxin. The ablation of the shoot apex, which is a significant source of auxin, decreased GA-induced root growth, and this effect was restored by auxin administration. Ethylene suppresses proliferation in a GA-antagonistic way has demonstrated that the DELLA proteins mediate at least some of the inhibitory impact of ethylene on development and it cooperates with GA in this respect[26]. In Arabidopsis, GA increases seedling root elongation, which is blocked by ethylene. In water-soluble rice and *R. palustris*, submergence stimulates ethylene and GA production, and GA promotes ethanol-induced internode extension[27]. In Arabidopsis, GAs are intrinsic regulators of hypocotyl development via cellular elongation via DELLA protein



112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140

degradation.

**Fig.3. throughout the plant's life cycle, ga interacts both positively and adversely with other plant hormones.**

Ethylene inhibits GA production or decreases GA responses by stabilising DELLA. GA and ethylene interact in a network of positive and negative interactions. Ethylene inhibits GA production or decreases GA responses by stabilising DELLA. GA stimulates ethylene responses in seedlings cultivated in the dark and in the light (apical hook development in the dark and the elongation of the hypocotyl in the light). In deepwater rice and *R. palustris*, submergence encourages ethylene and GA production, and GA enhances ethylene-induced internode elongation.

### 2.3 Signalling of Cytokinin's:

Cytokinin's are a kind of plant hormone that is involved in many aspects of plant development, including organ formation, apical dominance, and leaf senescence [17]. Cell division, dominance of the apical layer, chloroplast biogenesis, nutrient mobilisation, leaf senescence, vascular differentiation, thereby photomorphogenic development, shoot differentiation, and anthocyanin production are all regulated by cytokinin's (CKs)[28,8]. The modulation of the type A ARR's ARR5, ARR7, and ARR15, which play a role in freezing tolerance, is influenced by cytokinin and ethylene signalling. Cytokinin also boosts ethylene production by enhancing the stability of ACS proteins, which catalyse the rate-limiting step in ethylene biosynthesis. CKs are frequently regarded as ABA inhibitors and auxin antagonists/synergists in a variety of plant activities [29]. Strigolactones, which were discovered in *Arabidopsis thaliana*, *Pisum sativa*, and other plants as branching mutants, are assumed to be synthesised mostly in the roots and carried upwards in the xylem to restrict bud expansion [30,31]. The mechanics of cytokinin transport are poorly understood, while possible efflux and influx transporters have been found[32]. A His-Asp phosphorelay comparable to that seen in bacterial two-component signalling systems, which are the primary means by which bacteria perceive and respond to environmental stimuli, is involved in the cytokinin signal transduction pathway [33]. Increased strigolactone levels in pea and *Arabidopsis* result in lower cytokinin levels in the xylem sap, but not in the shoot. Cytokinin interacts with strigolactones to control the formation of axillary buds. In addition to blocking the transcription factor BRC1, which operates downstream of strigolactones, cytokinin's hinder auxin-mediated regulation of the strigolactone biosynthetic gene MAX4. Strigolactones, in turn, suppress cytokinin production via IPT1. Cytokinin degradation is mediated by enzymes encoded by the CYTOKININ OXIDASE (CKX) gene family. The cytokinin signalling pathway interacts with ABA, controlling salinity and drought response. ABA inhibits multiple CKX genes, while AHK2 and AHK3 inhibit several ABA responsive genes. Cytokinin is a phytohormone that influences several agriculturally significant activities, including growth, nutritional responses, and exposure to both abiotic and biotic stresses. Plant cytokinin levels are controlled by

141 manufacturing and inactivation mechanisms. Cytokinin's are recognized by membrane-localized histidine kinase  
142 receptors and transduced to the nucleus via a His-Asp phosphorelay to promote the expression of a family of  
143 transcription factors[34].Cytokinins are plant hormones that play a crucial role in regulating various aspects of plant  
144 growth and development, including cell division, shoot initiation, leaf senescence, and more. The signaling pathway of  
145 cytokinins involves a complex network of components. Cytokinins are perceived by membrane-bound receptor proteins  
146 called cytokinin receptors. In *Arabidopsis thaliana*, there are three major cytokinin receptors: AHK2 (ARABIDOPSIS  
147 HISTIDINE KINASE 2), AHK3 (ARABIDOPSIS HISTIDINE KINASE 3), and AHK4/CRE1 (CYTOKININ RESPONSE 1).  
148 These receptors have histidine kinase activity and are involved in cytokinin perception[35]. Upon cytokinin binding to  
149 the receptors, a phosphorylation cascade involving histidine phosphotransfer proteins (AHPs) is initiated. AHPs transfer  
150 phosphate groups between proteins in the pathway[36].Cytokinins activate a family of transcription factors known as  
151 type-B *Arabidopsis* response regulators (ARRs) through phosphorylation by AHPs. These type-B ARR then control the  
152 expression of target genes involved in various cytokinin responses[37].Cytokinin signaling leads to a range of  
153 physiological responses, including enhanced cell division, shoot branching, delayed leaf senescence, and more, which  
154 contribute to plant growth and development[28].The cytokinin signalling pathway is tightly regulated by negative  
155 feedback loops, including the type-A ARR that act as negative regulators of cytokinin signalling[36].Understanding the  
156 cytokinin signalling pathway is crucial for elucidating the mechanisms by which cytokinins control plant growth and  
157 development.

#### 158 **2.4 Signalling of Brassinosteroid (BRs) :**

159 Brassinosteroids (BRs) are a type of polyhydroxylated steroidal hormone that plays important roles in plant growth and  
160 development, including cell elongation, cell division, senescence, vascular differentiation, reproduction,  
161 photomorphogenesis, and stress responses[38]. Brassinosteroids have been detected in nearly all plant tissues, with  
162 the largest concentrations observed in seeds, pollen, and young developing tissues [7].Brassinosteroids have strong  
163 impacts on overall plant growth via cell [39].Recent detailed analyses revealed that BRs are involved in many aspects  
164 of root development, including meristem size maintenance, root hair formation, lateral root initiation, gravitropic  
165 response, mycorrhiza formation, and nodulation in legume species[40].In comparison to the auxin route,  
166 Brassinosteroid perception is a more traditional signalling paradigm. Exogenous Brassinosteroid application, as well as  
167 brassinosteroid-deficient and brassinosteroid-insensitive mutants, have all resulted in a thorough functional model of  
168 brassinosteroid activity in plants Fig 4.Brassinosteroids (BRs) are a class of plant hormones that play essential roles in  
169 various aspects of plant growth and development. They are involved in processes such as cell elongation, cell division,  
170 vascular differentiation, and responses to environmental stimuli. The first step in the BR signalling pathway involves the  
171 perception of brassinosteroids by a cell surface receptor. The primary receptor for BRs in plants is the receptor kinase  
172 complex, which consists of the brassinosteroid insensitive 1 (BRI1) and BRI1-associated kinase 1 (BAK1). BRI1 is a  
173 leucine-rich repeat receptor kinase, and BAK1 is a co-receptor that interacts with BRI1 upon BR binding[41]. Upon  
174 binding of BRs to the BRI1-BAK1 receptor complex, a signal is transduced into the cell. This involves a cascade of  
175 phosphorylation events, leading to the activation of downstream components. The key transcription factors in the BR  
176 signalling pathway are the brassinazole-resistant 1 (BZR1) and BRI1-EMS-suppressor 1 (BES1) proteins.Activated  
177 BZR1 and BES1 transcription factors regulate gene expression by binding to specific target genes. They control the  
178 expression of genes involved in cell expansion, differentiation, and other processes, leading to the physiological  
179 responses associated with BRs.The changes in gene expression triggered by the activation of BZR1 and BES1 lead to  
180 various physiological responses. These responses can include increased cell elongation, root growth, and resistance to  
181 stress conditions, among other effects. It's important to note that the BR signalling pathway is highly interconnected  
182 with other signalling pathways, allowing plants to integrate multiple signals and respond appropriately to environmental  
183 conditions.

184




Fig:4. Brassinosteroids have a role in plant development throughout the life cycle.

### 2.5 Signalling of Salicylic acid (SA):

Salicylic acid (SA) is a phytohormone that acts as a signal molecule in the immune and plant development pathways [42,43,44]. SA is produced in plants by two separate processes: the isochlorismate (IC) and the phenylalanine (PA) ammonia-lyase (PAL) pathways[45,46]. Plants with changed endogenous SA levels exhibit aberrant growth characteristics[44,47]. A high amount of endogenous SA causes growth retardation. Many Arabidopsis mutants with over accumulating SA, such as has accelerated cell death 6 (acd6), permanent expresser of PR gene 5 (cpr5), and the SAP and MIZ domain protein gene 1 (siz1), present the dwarf plant morphology with a shorter stem, smaller leaves, and/or floral organs compared to the wild type[48,49,50]. The Arabidopsis SA-deficient mutant sid2 and the SA-depleted NahG transgenic line, on the other hand, display increased leaf biomass as compared to WT [51].

It is worth mentioning that the effects of SA may vary depending on the plant type or organ. Aim1 rice mutants with lower endogenous SA levels, for example, have weaker seedling and spore-producing roots than the original wild type[52]. It has been shown that overexpression of NahG can decrease the cell multiplication and expansion defects produced by siz1[50], suggesting that SA suppresses these two processes in the cells. However, SA buildup in a different situation might have a distinct effect on the development of cells control. [53] discovered that elevated SA levels stimulate cell proliferation and growth in acd6-1 leaves. SA may influence plant growth by either negatively or positively influencing cell division and expansion. Some evidence suggests that SA has a deleterious influence on the two cellular processes in Arabidopsis leaves. For example, SA-deficient NahG transgenic plants grew faster than WT plants and had a 1.7-fold increase in leaf rosette biomass at the early stage of propagation[51]. This higher growth effect on NahG transgenic plants was more noticeable at low temperatures and was caused by greater increase in cell size of rosette leaves.

SA signalling during plant immunological responses has been extensively studied, and NPR1 has been identified as a critical player in this process. NPR1 also plays a role in SA-mediated growth regulation by controlling cell division and expansion [53,54,55]. Arabidopsis npr1-1 mutant leaves contained fewer cells and more DNA, showing that NPR1 promotes cell division and inhibits endoreduplication in leaves[53]. [54] discovered that the xs2 mutant accumulated high SA levels and inhibited cell growth, creating smaller cells than the WT. Interestingly, the xs2 mutant's severe impairment in cell size was repaired in the xs2 npr1 double mutant. These findings suggest that the inhibition of cell growth in xs2 was mediated by an NPR1-dependent signalling pathway. Furthermore, the SA-over accumulating Arabidopsis mutant cad1 boosted QC cell division, which was rescued by mutations in SID2 or NPR1, demonstrating that the cad1 mutant promotes QC cell division via an NPR1-dependent SA signalling pathway [55]. To govern plant and organ growth, SA interacts with other hormones involved in the regulation of cell division and expansion, such as auxin, GA, and ethylene (ET) [56,57,58,47]. Auxin is a significant growth hormone that regulates these cellular activities, particularly in roots [59,60,61,62], and SA can influence root development by influencing auxin accumulation and transit.[63] shown that exogenous SA has a concentration-dependent effect on the root tip meristem in Arabidopsis.


### 2.6 Signalling of Abscisic acid (ABA):

222 Because of their regulatory action on abscission and dormancy, these were previously known as Dormin or Abscisin.  
 223 This hormone is widely distributed in higher plants and can be detected in a variety of organs and tissues (both old and  
 224 young). ABA causes abscission of the leaves of many plants as well as the fruits of some plant species. ABA appears  
 225 to be an intrinsic component that induces dormancy in at least some temperature zone woody plant buds. ABA also  
 226 slows or delays seed germination. ABA suppresses the growth of several plant tissues and organs, including leaves,  
 227 coleoptiles, stems, hypocotyls, and roots. It accelerates senescence by causing leaf abscission, degeneration of  
 228 excised leaves, and accelerating chlorophyll breakdown. Recent research has shown that ABA accelerates leaf  
 229 senescence and abscission in the absence of ethylene [64,65].

230 Several ABA auxotrophic maize mutants, termed vp2, vp5, vp7, and vp9, have been found by genetic screening of  
 231 viviparous mutants[66]. Plastids catalyse the conversion of zeaxanthin (C40) to xanthoxin (C15)(Fig5).Catabolism of  
 232 ABA occurs via the conversion of ABA to phaseic acid (PA), which is catalysed by a P450 encoded by CYP707As[67].  
 233 PA is subsequently catalysed to dihydrophaseic acid (DPA) and DPA4ODglucoside (DPAG) by PA reductase (PAR),  
 234 ABH2, and glycosyltransferase (GT) [68]. PA has been shown to preferentially activate a subset of ABA receptor PYLs  
 235 (Fig 5)[68].Abscisic acid occurs naturally in plants in both anionic (ABA) and protonated (ABAH) forms. ABA can diffuse  
 236 passively via the plasma membrane, and ABA diffusion decreases with cytoplasmic alkalization, which rises during  
 237 osmotic stressors[67,70].When plants are exposed to a variety of environmental circumstances, they rapidly acquire  
 238 ABA.

239 The maintenance of a basal level of free ABA consistent with different tissues in diverse settings is critical to the overall  
 240 plant's appropriate growth and development states. As a result, ABA catabolism is strictly regulated by both ABA  
 241 conjugation and catalytic hydroxylation. A UDPglucosyltransferase (UGT) encoded by UGT71C5 can glycosylate ABA.  
 242 ABAGE, or ABA glucose ester, is an inactive version of ABA [71]. When the environment changes, AtBG1 and AtBG2  
 243 encode glucosidases that rapidly convert ABAGE to active ABA, which is released from the endoplasmic reticulum and  
 244 vacuole [23,52].By quickly activating and inactivating ABA, the conjugation cycle established by glucosyltransferase  
 245 and glucosidase allows plants to phenotypically adjust to their environment via ABA mediated responses. Catabolism of  
 246 ABA occurs via the conversion of ABA to phaseic acid (PA), which is catalysed by a P450 encoded by CYP707As  
 247 (Kushiro et al. 2004). PA is subsequently catalysed to dihydrophaseic acid (DPA) and DPA4ODglucoside (DPAG) by  
 248 PA reductase (PAR), ABH2, and glycosyltransferase (GT) [68]. PA has been shown to preferentially activate a subset  
 249 of ABA receptor PYLs (Fig 5) [68].

250



## Fig:5. plant abscisic acid (aba) the process of biosynthesis breakdown, and (de)conjugation.

Pathogen infection, for example, can influence ABA homeostasis in host plants. Some biotrophic infections, such as wheat rust fungus, can induce ABA elevations that lead to increased cytoplasmic sugar accumulation by increasing TaSTP6 expression. In Arabidopsis, when infected with the necrotrophic disease *Botrytis cinerea*, the transcription factor WRKY33 increases ABA production by upregulating the transcription of NCED3 and NCED5 [20]. The tomato NAC transcription factor LeJA2 (for jasmonic acid 2) stimulates the production of ABA, which can inhibit pathogen entrance through stomata. Hormone crosstalk also has a role in ABA homeostasis. Auxin and GA, for example, influence endogenous ABA levels in forest strawberry (*Fragaria vesca*) through influencing the regulatory loops of FveCYP707As and FveNCEDs [71]. Furthermore, JA accumulation is essential for ABA accumulation in Arabidopsis roots following dehydration treatment [72].

Future research would offer insight on the relationship between global ABA content regulation and different biological processes. All of these systems work together to fine-tune the amount of ABA at various developmental stages and in response to diverse environmental variables. More signalling factors that function upstream of ABA accumulation, as well as the processes that coordinate the dynamic control of ABA content and biological activity, remain unknown. Protein phosphorylation and dephosphorylation are critical for maintaining an adequate balance of ABA-mediated growth regulation depending on the plant's environmental situation [73,74]. Protein phosphatases can interact with and inhibit SnRK2s, SnRK1s, SnRK3s, and even mammalian AMPKs, all of which play important roles in abiotic stress, ABA, and energy signalling [75,73].

### 2.7 Signalling of Ethylene:

Ethylene is a gaseous plant hormone that exerts a profound influence on various aspects of plant growth, development, and responses to environmental stimuli. It is often referred to as the "ripening hormone" because of its central role in fruit ripening. However, ethylene's significance extends far beyond fruit ripening, affecting processes ranging from seed germination to leaf senescence. This article introduces ethylene, summarizing its biosynthesis, signaling pathway, and diverse physiological effects, with references to key studies that have advanced our understanding of this essential plant hormone.

#### 2.7.1 Biosynthesis of Ethylene:

Ethylene is synthesized from the amino acid methionine through a series of enzymatic reactions. The two key enzymes in ethylene biosynthesis are ACC synthase (ACS), which converts methionine to 1-aminocyclopropane-1-carboxylic acid (ACC), and ACC oxidase (ACO), which transforms ACC into ethylene. This biosynthesis pathway is tightly regulated and can be influenced by various internal and external factors [76].

#### 2.7.2 Ethylene Receptors

Ethylene signals are perceived by membrane-bound receptors, such as Ethylene Response Sensor 1 (ERS1) and Ethylene Response Sensor 2 (ERS2), in association with Constitutive Triple Response 1 (CTR1). Upon ethylene binding, the receptors inactivate CTR1, initiating a signaling cascade. This cascade ultimately leads to the activation of transcription factors, such as Ethylene-Insensitive 3 (EIN3) and Ethylene-Insensitive 3-Like 1 (EIL1), which regulate the expression of ethylene-responsive genes [77].

#### 2.7.3 Physiological Effects of Ethylene:

Ethylene exerts diverse effects on plant growth and development. It is responsible for the "triple response" in seedlings, which includes a thickened stem, an exaggerated apical hook, and inhibited root elongation. Ethylene also plays a critical role in fruit ripening, leaf and flower senescence, abiotic and biotic stress responses, apical dominance, root development, and abscission, among other processes [78]. Understanding the multifaceted roles of ethylene in plant physiology is essential for crop management, post-harvest handling, and horticultural practices. Moreover, ethylene's significance in stress responses and environmental adaptations makes it a subject of ongoing research in the field of plant biology. Ethylene is a gaseous plant hormone that plays a significant role in regulating various aspects of plant growth and development, as well as responses to environmental cues. Here is an overview of ethylene, its biosynthesis, and its effects on plants:

Ethylene Biosynthesis: (Fig 6)

1. Methionine Conversion: Ethylene is synthesized from the amino acid methionine. The first step involves the conversion of methionine to S-adenosylmethionine (SAM) by the enzyme methionine adenosyltransferase.
2. SAM to ACC: SAM is then converted to 1-aminocyclopropane-1-carboxylic acid (ACC) by ACC synthase (ACS).
3. ACC to Ethylene: ACC is subsequently converted to ethylene through the action of ACC oxidase (ACO).

#### 2.7.3 Ethylene Receptors:

Ethylene signals are perceived by a family of membrane-bound receptors, primarily Ethylene Response Sensor 1 (ERS1) and Ethylene Response Sensor 2 (ERS2), which function in association with Constitutive Triple Response 1 (CTR1).


In the presence of ethylene, the receptors are activated, leading to inactivation of CTR1 through phosphorylation. Inactivation of CTR1 allows for the de-repression of Ethylene-Insensitive 2 (EIN2). Active EIN2 acts as a positive regulator of ethylene signaling, promoting the cleavage and nuclear translocation of the Ethylene-Insensitive 3 (EIN3)

and Ethylene-Insensitive 3-Like 1 (EIL1) transcription factors. EIN3 and EIL1 subsequently activate the expression of ethylene-responsive genes.

#### 2.7.4 Effects of Ethylene:

Ethylene induces the "triple response" in seedlings, which includes a thickening and shortening of the stem, an exaggerated curvature of the apical hook, and inhibition of root elongation. Ethylene is well-known for its role in fruit ripening. It promotes the ripening process and can be used for the controlled ripening of fruits. Ethylene also accelerates the senescence of leaves and flowers, leading to their wilting and decay. Ethylene is involved in responses to various stresses, such as pathogen infection and environmental stressors like drought and flooding. Ethylene is implicated in apical dominance, regulating the growth of lateral branches and buds. Ethylene can affect root development, promoting root hair elongation and controlling root architecture. Ethylene triggers leaf and fruit abscission, allowing for the shedding of older leaves and ripe fruit.

Understanding the role of ethylene in plant physiology is essential for various applications in agriculture, horticulture, and the mitigation of crop losses. Ethylene is also a key target for research in the field of plant biology and stress responses. Ethylene is a versatile plant hormone with a wide range of applications in agriculture, horticulture, and post-harvest management. Its effects on plant growth and development make it a valuable tool for various purposes. Ethylene is widely used to initiate and control the ripening of climacteric fruits, such as tomatoes, bananas, apples, and



pears. Ethylene gas is applied to unripe fruit to trigger the ripening process, allowing for better timing of harvest and distribution. Ethylene is involved in the regulation of flower senescence. Controlling ethylene exposure can extend the vase life of cut flowers, making them last longer in floral arrangements.

**Fig:6. pictural representation of ethylene biosynthesis**

Ethylene promotes seed germination in some plant species. Pre-treating seeds with ethylene gas can enhance germination rates and uniformity. Ethylene plays a role in apical dominance, influencing the growth of lateral branches and buds. Controlling ethylene levels can affect branching patterns in plants. Ethylene can stimulate the formation of root hairs and influence root architecture. It is used in some cases to enhance root development in crops. Ethylene triggers the abscission of leaves and fruits. This property is harnessed for defoliation in cotton farming and for the controlled shedding of ripe fruit. Ethylene is involved in plant responses to various stresses, such as drought, flooding, and pathogen attack. Understanding ethylene signaling can help in developing stress-tolerant crops. Ethylene can affect the flavor and aroma of certain fruits, influencing their taste and scent. Ethylene is used in post-harvest management to regulate fruit ripening and control storage conditions. It can extend the shelf life of some fruits and vegetables when managed properly. Ethylene is a valuable tool for researchers studying plant biology and physiology. It can be applied to investigate the effects of ethylene on various aspects of plant growth and development. Ethylene interacts with other plant hormones, such as auxins and cytokinins, to regulate plant responses to different stimuli. Understanding these interactions is essential for plant manipulation. Ethylene management is also important in enclosed environments, such as greenhouses and storage facilities, to prevent unwanted effects on plant growth and maintain optimal conditions for crops. Ethylene has industrial applications outside of agriculture, including its use in the production of plastics, chemicals, and fuel. The controlled application of ethylene is a valuable strategy in modern agriculture and horticulture, enabling growers to optimize crop yield, quality, and post-harvest handling. Its diverse effects on plant growth and development make it a versatile tool for various purposes in the plant science and agricultural industries. The signaling pathway of ethylene in plants involves a complex network of components that regulate various physiological responses. Ethylene signals are perceived by a family of membrane-bound receptors, including Ethylene Response Sensor 1 (ERS1) and Ethylene Response Sensor 2 (ERS2). These receptors are part of a family of ethylene receptors that can

351 form heterodimers with Constitutive Triple Response 1 (CTR1), an ethylene signaling negative regulator[79]. In the  
352 absence of ethylene, CTR1 functions as a negative regulator of ethylene signaling. However, when ethylene binds to  
353 the receptors, CTR1 is inactivated, leading to the de-repression of Ethylene-Insensitive 2 (EIN2). EIN2 acts as a  
354 positive regulator of ethylene signaling and serves as a critical node in the pathway[80].Active EIN2 promotes the  
355 cleavage and nuclear translocation of the transcription factors Ethylene-Insensitive 3 (EIN3)and Ethylene-Insensitive 3-  
356 Like 1 (EIL1). These transcription factors regulate the expression of ethylene-responsive genes involved in various  
357 physiological processes, including those associated with growth, development, and stress responses[81]. The  
358 activation of ethylene-responsive genes leads to a wide range of physiological responses, including the "triple  
359 response" in seedlings, which includes a thickening and shortening of the stem, an exaggerated curvature of the apical  
360 hook, and inhibition of root elongation. Ethylene also plays a key role in processes like fruit ripening, leaf and flower  
361 senescence, and responses to abiotic and biotic stresses[82]. The ethylene signaling pathway is a complex and finely  
362 regulated system that governs various aspects of plant growth and development, as well as responses to  
363 environmental cues. The references cited above provide insights into the molecular and biochemical aspects of  
364 ethylene signaling in plants. For more detailed and up-to-date information, consult recent research articles and reviews  
365 in the field. The signaling pathway of ethylene in plants is a well-studied process that involves a complex network of  
366 components. Ethylene signals are perceived by a family of membrane-bound receptors known as ethylene receptors. In  
367 *Arabidopsis thaliana*, the receptors include Ethylene Response Sensor 1 (ERS1) and Ethylene Response Sensor 2  
368 (ERS2). These receptors are part of the family of ethylene receptors responsible for sensing ethylene gas[83].In the  
369 absence of ethylene, the receptors are bound to Constitutive Triple Response 1 (CTR1), which functions as a negative  
370 regulator of ethylene signaling. When ethylene is present, it binds to the receptors, leading to the inactivation of CTR1  
371 and allowing Ethylene-Insensitive 2 (EIN2) to be de-repressed[84]. EIN2 acts as a positive regulator of ethylene  
372 signaling and plays a pivotal role in the pathway. It promotes the cleavage and nuclear translocation of the transcription  
373 factors Ethylene-Insensitive 3 (EIN3) and Ethylene-Insensitive 3-Like 1 (EIL1). These transcription factors, in turn,  
374 regulate the expression of ethylene-responsive genes that control various physiological processes[85]. The activation  
375 of ethylene-responsive genes results in a broad spectrum of physiological responses in plants, including the "triple  
376 response" in seedlings, control of fruit ripening, leaf and flower senescence, and responses to abiotic and biotic  
377 stresses[82].The ethylene signaling pathway is a crucial regulatory system governing various aspects of plant growth  
378 and development. The references provided here offer insights into the molecular and biochemical components of the  
379 ethylene signaling pathway.

### 380 **3.CONCLUSION**

381 The signaling pathways of growth promoters and inhibitors play pivotal roles in regulating plant growth and  
382 development. Auxins, such as indole-3-acetic acid (IAA), are essential growth promoters that regulate cell elongation,  
383 apical dominance, phototropism, and root development. The auxin signaling pathway primarily involves the transport  
384 protein PIN1, which regulates auxin distribution, and the auxin response factors (ARFs) that control gene expression.  
385 Cytokinins are hormones that promote cell division and shoot initiation. The signaling pathway of cytokinins includes  
386 receptor kinases and transcription factors like type-B *Arabidopsis* response regulators (ARRs).In conclusion, the  
387 signaling pathways of growth promoters and inhibitors are intricate and interconnected, allowing plants to fine-tune their  
388 growth and development in response to internal and external cues. These pathways enable plants to adapt to changing  
389 environmental conditions, optimize resource allocation, and defend against stress and pathogens. A comprehensive  
390 understanding of these signaling pathways is crucial for agricultural and horticultural practices, as well as for advancing  
391 our knowledge of plant biology and stress responses. Ongoing research continues to uncover the complexities of these  
392 pathways, providing insights into how plants balance growth promotion and inhibition to thrive in diverse environments.  
393 The intricate signaling pathways of growth promoters and inhibitors are pivotal mechanisms that govern the dynamic  
394 and adaptable nature of plant growth and development. Growth inhibitors, including abscisic acid, ethylene, jasmonic  
395 acid, and salicylic acid, serve as crucial regulators that restrain growth under unfavorable conditions or in response to  
396 stress. These pathways are activated in times of stress, pathogen attack, or other challenges, leading to growth  
397 inhibition and the activation of defense mechanisms. By balancing growth promotion and inhibition, plants can  
398 strategically allocate resources and adapt to changing circumstances to ensure their survival. Understanding the  
399 complexities of these signaling pathways is not only fundamental for plant biology but also critical for various  
400 applications, including agriculture, horticulture, and environmental management. It allows us to manipulate plant growth  
401 and development for improved crop production, disease resistance, and stress tolerance. Ongoing research continues  
402 to unravel the intricate details of these pathways, offering new insights into how plants achieve the delicate balance  
403 between promoting growth and limiting it when necessary. This knowledge is invaluable for addressing the growing  
404 challenges of food security, environmental sustainability, and climate resilience in the face of a change.

### 406 **ACKNOWLEDGEMENTS**

407 We express our sincere gratitude to all authors contributed to the completion of this paper.

408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467

## AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## REFERENCES

- [1] Bisht, T. S., Rawat, L., Chakraborty, B., & Yadav, V. (2018). A Recent Advances in Use of Plant Growth Regulators (PGRs) in Fruit Crops-A Review.
- [2] Yeshitila, M., & Gobena, A. (2020). Review on phytohormones signaling cross-talk: To control plant growth and development. *International Journal of Advanced Research in Biological Sciences*, 7(3), 54-57.
- [3] Koornneef, M., Bentsink, L., & Hilhorst, H. (2002). Seed dormancy and germination. *Current opinion in plant biology*, 5(1), 33-36.
- [4] Skoog, F. and Miller, C.O. 1957. Chemical regulation of growth and organ formation in plant Tissues cultured in vitro. *Symp.Soc. Exp. Biol.* 11: 118–131.
- [5] Chandler, J.W.; Werr, W. 2009 . Cytokinin–auxin crosstalk in cell type specification. *Trends Plant Sci. Chem.* 276, 26405–26410.
- [6] Swarup R, Parry G, Graham N, Allen T, Bennett M 2002. Auxin cross-talk: integration of signalling pathways to control plant development. *Plant Mol Biol* 49: 411–426.
- [7] Stephen Depuydt and Christian S. Hardtke 2011. Hormone Signalling Crosstalk in Plant Growth Regulation *Current Biology* 21, R365–R373.
- [8] Davies, P. J. 2004. *Plant Hormones: Biosynthesis, Signal Transduction Action*: (Kluwer Academic).
- [9] Dello iolo, R., et al. 2008 . A genetic framework for the control of cell division and differentiation in the root meristem. *Science*. 322, 1380–1384 dependent auxin distribution. *Plant Cell* 19: 2197– 2212.
- [10] Muday GK, Rahman A, Binder B 2012. Auxin and ethylene: collaborators or competitors *Trends Plant Sci*, in press.
- [11] Silverstone AL, Chang C, Krol E, Sun TP 1997 .Developmental regulation of the gibberellin's biosynthetic gene GA1 in *Arabidopsis thaliana* . *Plant J* 12: 9–19.
- [12] Hardtke CS, Dorcey E, Osmont KS, Sibout R 2007. Phytohormone collaboration: zooming in on auxin-brassinosteroid interactions. *Trends Cell Biol* 17: 485–492.
- [13] Shkolnik-Inbar D, Bar-Zvi D 2010. ABI4 mediates abscisic acid and cytokinin inhibition of lateral root formation by reducing polar auxin transport in *Arabidopsis*. *Plant Cell* 22: 3560–3573.
- [14] Abidur Rahman 2013. Auxin: a regulator of cold stress response *Physiologia Plantarum* 147: 28–35.
- [15] Bennett T, Scheres B. 2010. Root development—two meristems for the price of one? *Current Topics in Developmental Biology* 91, 67–102.
- [16] Perilli S, Moubayidin L, Sabatini S. 2010. The molecular basis of cytokinin function. *Current Opinion in Plant Biology* 13, 21–26.
- [17] Sedeer El-Showk, Raili Ruonala and Ykä Helariutta 2013 . cytokinin signalling and crosstalk *The Company of Biologists Ltd Crossing paths: Development* 140, 1373-1383.
- [18] Zhang,W.,To,J. P. C.,Cheng,C.-Y.,Eric Schaller,G.andKieber,J. J. 2011. Type-A response regulators are required for proper root apical meristem function through post- transcriptional regulation of PIN auxin efflux carriers. *Plant J*.68, 1-10.
- [19] Elena Pacifici , Laura Polverari\* and Sabrina Sabatini 2015 . Plant hormone cross-talk: the pivot of root growth *Journal of Experimental Botany*, Vol. 66, No. 4 pp. 1113–1121, 2015 doi:10.1093/jxb/eru534.
- [20] Liu M, Chen Y, Chen Y, Shin J-H, Mila I, Audran C, Zouine M, Pirrello J, Bouzayen M. 2018.The tomato Ethylene Response Factor Sl- ERF.B3integrates ethylene and auxin signaling via direct regulation of Sl-Aux/IAA27. *New Phytologist* 219: 631–64.
- [21] Nemhauser JL, Hong FX, Chory J 2006. Different plant hormones regulate similar processes through largely nonoverlapping transcriptional responses. *Cell* 126: 467–475.
- [22] Tyler L, Thomas SG, Hu J, Dill A, Alonso JM, Ecker JR, Sun TP. 2004. Della proteins and gibberellin regulated seed germination and floral development in *Arabidopsis*. *Plant Physiology* 135, 1008–1019.
- [23] Lee KH, Piao HL, Kim HY, Choi SM, Jiang F, Hartung W, Hwang I, Kwak JM, Lee IJ, Hwang I (2006) Activation of glucosidase via stress-induced polymerization rapidly increases active pools of abscisic acid. *Cell* 126: 1109–1120.
- [24] Greenboim-Wainberg Y, Maymon I, Borochoy R, Alvarez J, Olszewski N, Ori N, Eshed Y, Weiss D. 2005. Crosstalk between gibberellin and cytokinin: the *Arabidopsis* GA response inhibitor SPINDLY plays a positive role in cytokinin signaling. *The Plant Cell* 17, 92–102.
- [25] Sun TP, Gubler F 2004 .Molecular mechanism of gibberellin signaling in plants. *Annu Rev Plant Biol* 55: 197–223
- [26] Achard P, Gong F, Cheminant S, Alioua M, Hedden P, Genschik P 2008. The cold-inducible CBF1 factor-dependent signaling pathway modulates the accumulation of the growth repressing DELLA proteins via its effect on gibberellin metabolism. *Plant Cell* ; 20:2117-29.

- 468 [27] Benschop JJ, Bou J, Peeters AJM, Wagemaker N, Gu "hl K, Ward D, Hedden P, Moritz T, Voeselek LACJ 2006.  
469 Long-term submergence-induced elongation in *Rumex palustris* requires abscisic acid-dependent biosynthesis of  
470 gibberellin. *Plant Physiol* 141: 1644–1652,
- 471 [28] Mok DW, Mok MC 2001. Cytokinin metabolism and action. *Annu Rev Plant Physiol Plant Mol Biol* 52:89-118.
- 472 [29] POSPÍŠILOVÁ J 2003. Interaction of cytokinins and abscisic acid during regulation of stomatal opening in bean  
473 leaves. *Photosynthetica* 41: 49.
- 474 [30] Beveridge CA. 2006. Axillary bud outgrowth: sending a message. *Current Opinion in Plant Biology* 9, 35–40.
- 475 [31] Kohlen W, Charnikhova T, Liu Q, Bours R, Domagalska MA, Beguerie S, Verstappen F, Leyser O, Bouwmeester  
476 H, RuyterSpira C. 2011. Strigolactones are transported through the xylem and play a key role in shoot architectural  
477 response to phosphate deficiency in nonarbuscular mycorrhizal host *Arabidopsis*. *Plant Physiology* 155, 974–987
- 478 [32] Duran-Medina, Y., Diaz-Ramirez, D. and Marsch-Martinez, N. (2017). Cytokinin on the Move. *Front. Plant Sci.* 8,  
479 146.
- 480 [33] Cheung, J. and Hendrickson, W. A. (2010). Sensor domains of two-component regulatory systems. *Curr. Opin.*  
481 *Microbiol.* 13, 116-123.
- 482 [34] Kieber, J. J., & Schaller, G. E. (2018). Cytokinin signaling in plant development. *Development*, 145(4), dev149344.
- 483 [35] Inoue, T., Higuchi, M., Hashimoto, Y., Seki, M., Kobayashi, M., Kato, T., ... & Kakimoto, T. (2001). Identification of  
484 CRE1 as a cytokinin receptor from *Arabidopsis*. *Nature*, 409(6823), 1060-1063.
- 485 [36] J. P., Haberer, G., Ferreira, F. J., Deruère, J., Mason, M. G., Schaller, G. E., ... & Kieber, J. J. (2004). Type-A  
486 *Arabidopsis* response regulators are partially redundant negative regulators of cytokinin signaling. *The Plant Cell*,  
487 16(3), 658-671.
- 488 [37] Rashotte, A. M., Carson, S. D., To, J. P., & Kieber, J. J. (2003). Expression profiling of cytokinin action in  
489 *Arabidopsis*. *Plant Physiology*, 132(4), 1998-2011.
- 490 [38] Clouse, S. D., Sasse, J. M. (1998). BRASSINOSTEROIDS: Essential Regulators of Plant Growth and  
491 Development. *Annual Review of Plant Physiology and Plant Molecular Biology*, 49(1), 427-451.
- 492 [39] Pacifici, E., Polverari, L., and Sabatini, S. 2015. Plant hormone cross-talk: the pivot of root growth. *J. Exp. Bot.*  
493 66:1113–1121.
- 494 [40] Wei Z. and Li J. 2016. Brassinosteroids Regulate Root Growth, Development, and Symbiosis. *Mol. Plant.* 86–100.
- 495 [41] Wang, X., Chory, J. (2006). Brassinosteroids regulate dissociation of BK11, a negative regulator of BRI1 signaling,  
496 from the plasma membrane. *Science*, 313(5790), 1118-1122.
- 497 [42] Hayat, S., Ali, B., and Ahmad, A. (2007). "Salicylic acid: biosynthesis, metabolism and physiological role in plants,"  
498 in *SALICYLIC ACID: A Plant Hormone*. eds. S. Hayat and A. Ahmad (Netherlands: Springer), 1–14.
- 499 [43] Vlot, A. C., Dempsey, D. A., and Klessig, D. F. (2009). Salicylic acid, a multifaceted hormone to combat disease.  
500 *Annu. Rev. Phytopathol.* 47, 177–206. doi: 10.1146/annurev.phyto.050908.135202.
- 501 [44] Rivas-San Vicente, M., and Plasencia, J. (2011). Salicylic acid beyond defence: its role in plant growth and  
502 development. *J. Exp. Bot.* 62, 3321–3338. doi: 10.1093/jxb/err031.
- 503 [45] Serino, L., Reimann, C., Baur, H., Beyeler, M., Visca, P., and Haas, D. (1995). Structural genes for salicylate  
504 biosynthesis from chorismate in *Pseudomonas aeruginosa*. *Mol. Gen. Genet.* 249, 217–228. doi:  
505 10.1007/bf00290369.
- 506 [46] Zhang, Y. L., and Li, X. (2019). Salicylic acid: biosynthesis, perception, and contributions to plant immunity. *Curr.*  
507 *Opin. Plant Biol.* 50, 29–36. doi: 10.1016/j.pbi.2019.02.004.
- 508 [47] Pokotylo, I., Hodges, M., Kravets, V., and Ruelland, E. (2021). A ménage à trois: salicylic acid, growth inhibition,  
509 and immunity. *Trends Plant Sci.* doi: 10.1016/j.tplants.2021.11.008 [Epub ahead of Print].
- 510 [48] Bowling, S. A., Clarke, J. D., Liu, Y., Klessig, D. F., and Dong, X. (1997). The *cpr5* mutant of *Arabidopsis*  
511 expresses both NPR1-dependent and NPR1-independent resistance. *Plant Cell* 9, 1573–1584. doi:  
512 10.1105/tpc.9.9.1573.
- 513 [49] Rate, D., Cuenca, J., Bowman, G., Guttman, D., and Greenberg, J. (1999). The gain-of-function *Arabidopsis* *acd6*  
514 mutant reveals novel regulation and function of the salicylic acid signaling pathway in controlling cell death,  
515 defenses, and cell growth. *Plant Cell* 11, 1695–1708. doi: 10.1105/tpc.11.9.1695.
- 516 [50] Miura, K., Lee, J., Miura, T., and Hasegawa, P. M. (2010). *SIZ1* controls cell growth and plant development in  
517 *Arabidopsis* through salicylic acid. *Plant Cell Physiol.* 51, 103–113. doi: 10.1093/pcp/pcp171
- 518 [51] Abreu, M. E., and Munné-Bosch, S. (2009). Salicylic acid deficiency in *NahG* transgenic lines and *sid2* mutants  
519 increases seed yield in the annual plant *Arabidopsis thaliana*. *J. Exp. Bot.* 60, 1261–1271. doi: 10.1093/  
520 jxb/ern363.
- 521 [52] Xu, L., Zhao, H. Y., Ruan, W. Y., Deng, M. J., Wang, F., Peng, J. R., et al. (2017). *ABNORMAL INFLORESCENCE*  
522 *MERISTEM1* functions in salicylic acid biosynthesis to maintain proper reactive oxygen species levels for root  
523 *MERISTEM* activity in rice. *Plant Cell* 29, 560–574. doi: 10.1105/tpc.16.00665.
- 524 [53] Vanacker, H., Lu, H., Rate, D. N., and Greenberg, J. T. (2001). A role for salicylic acid and NPR1 in regulating cell  
525 growth in *Arabidopsis*. *Plant J.* 28, 209–216. doi: 10.1046/j.1365-313X.2001.01158.

- 526 [54] Fujikura, U., Kazune, E., Horiguchi, G., Seo, M., Yuri, K., Yuji, K., et al. (2020). Suppression of class I  
527 compensated cell enlargement by *xs2* mutation is mediated by salicylic acid signaling. *PLoS Genet.* 16:e1008873.  
528 doi: 10.1371/journal.pgen.1008873
- 529 [55] Wang, Z., Rong, D., Chen, D., Xiao, Y., Liu, R., Wu, S., et al. (2021). Salicylic acid promotes quiescent center cell  
530 division through ROS accumulation and down-regulation of *PLT1*, *PLT2*, and *WOX5*. *J. Integr. Plant Biol.* 63, 583–  
531 596. doi: 10.1111/jipb.13020
- 532 [56] Ari, Y., Sam, F., Siddiqui, H., Bajguz, A., and Hayat, S. (2020). Salicylic acid in relation to other phytohormones in  
533 plant: A study towards physiology and signal transduction under challenging environment. *Environ. Exp. Bot.*  
534 175:104040. doi: 10.1016/j.envexpbot.2020.104040.
- 535 [57] Emamverdian, A., Ding, Y., and Mokhberdoran, F. (2020). The role of salicylic acid and gibberellin signaling in  
536 plant responses to abiotic stress with an emphasis on heavy metals. *Plant Signal. Behav.* 15:1777372. doi:  
537 10.1080/15592324.2020.1777372.
- 538 [58] Mazzoni-Putman, S. M., Brumos, J., Zhao, C., Alonso, J. M., and Stepanova, N. A. (2021). Auxin interactions with  
539 other hormones in plant development. *Cold Spring Harb. Perspect. Biol.* 13:a039990. doi:  
540 10.1101/cshperspect.a039990
- 541 [59] Perrot-Rechenmann, C. (2010). Cellular responses to auxin: division versus expansion. *Cold Spring Harb.*  
542 *Perspect. Biol.* 2:a001446. doi: 10.1101/cshperspect.a001446.
- 543 [60] Barbeza, E., Dünserb, K., Gaidoraa, A., Lendla, T., and Buscha, W. (2017). Auxin steers root cell expansion via  
544 apoplastic pH regulation in *Arabidopsis thaliana*. *P. Natl. Acad. Sci. U. S. A.* 114, E4884–E4893. doi:  
545 10.1073/pnas.1613499114
- 546 [61] Huang, R. F., Zheng, R., He, J., Zhou, Z. M., Wang, J. C., Xiong, Y., et al. (2019). Noncanonical auxin signaling  
547 regulates cell division pattern during lateral root development. *P. Natl. Acad. Sci. U. S. A.* 116, 21285–21290. doi:  
548 10.1073/pnas.1910916116.
- 549 [62] Seo, D. H., Jeong, H., Choi, Y. D., and Jang, G. (2021). Auxin controls the division of root endodermal cells. *Plant*  
550 *Physiol.* 187, 1577–1586. doi: 10.1093/plphys/kiab341
- 551 [63] Pasternak, T., Groot, E. P., Kazantsev, F. V., Teale, W., Omelyanchuk, N., Kovrizhnykh, V., et al. (2019). Salicylic  
552 acid affects root meristem patterning via auxin distribution in a concentration-dependent manner. *Plant Physiol.*  
553 180, 1725–1739. doi: 10.1104/pp.19.00130.
- 554 [64] Ogawa M, Hanada A, Yamauchi Y, Kuwahara A, Kamiya Y, Yamaguchi S (2003) Gibberellin biosynthesis and  
555 response during *Arabidopsis* seed germination. *Plant Cell* 15: 1591–1604.
- 556 [65] Zhao Y, Zhang Z, Gao J, Wang P, Hu T, Wang Z, Hou YJ, Wan Y, Liu W, Xie S, Lu T, Xue L, Liu Y, Macho AP, Tao  
557 WA, Bressan RA, Zhu JK (2018) *Arabidopsis* duodecuple mutant of *PYL* ABA receptors reveals *PYL* repression of  
558 ABA-independent *SnRK2* activity. *Cell Rep* 23: 3340–3351.
- 559 [66] Reid JB (1990) Phytohormone mutants in plant research. *J Plant Growth Regul* 9: 97.
- 560 [67] Kushihiro T, Okamoto M, Nakabayashi K, Yamagishi K, Kitamura S, Asami T, Hirai N, Koshihara T, Kamiya Y,  
561 Nambara E (2004) The *Arabidopsis* cytochrome P450 *CYP707A* encodes ABA 8'-hydroxylases: Key enzymes in  
562 ABA catabolism. *EMBO J* 23: 1647–1656.
- 563 [68] Weng JK, Ye M, Li B, Noel JP (2016) Co-evolution of hormone metabolism and signaling networks expands plant  
564 adaptive plasticity. *Cell* 166: 881–893.
- 565 [69] Wilkinson S, Davies WJ (1997) Xylem sap pH increase: A drought signal received at the apoplastic face of the  
566 guard cell that involves the suppression of saturable abscisic acid uptake by the epidermal symplast. *Plant*  
567 *Physiol* 113: 559–573
- 568 [70] Karuppanapandian T, Geilfus CM, Muhling KH, Novak O, Gloser V (2017) Early changes of the pH of the apoplast  
569 are different in leaves, stem and roots of *Vicia faba* L. under declining water availability. *Plant Sci* 255: 51–58.
- 570 [71] Liao X, Li M, Liu B, Yan M, Yu X, Zi H, Liu R, Yamamuro C (2018) Interlinked regulatory loops of ABA catabolism  
571 and biosynthesis coordinate fruit growth and ripening in woodland strawberry. *Proc Natl Acad Sci USA* 115:  
572 E11542–E11550
- 573 [72] de Ollas C, Arbona V, Gómez-Cadenas A (2015) Jasmonoyl isoleucine accumulation is needed for abscisic acid  
574 build-up in roots of *Arabidopsis* under water stress conditions. *Plant Cell Environ* 38: 2157–2170
- 575 [73] Zhu JK (2016) Abiotic stress signaling and responses in plants. *Cell* 167: 313–324.
- 576 [74] Yang Y, Guo Y (2018) Elucidating the molecular mechanisms mediating plant salt-stress responses. *New Phytol*  
577 217: 523–539.
- 578 [75] Sanders MJ, Grondin PO, Hegarty BD, Snowden MA, Carling D (2007) Investigating the mechanism for AMP  
579 activation of the AMP-activated protein kinase cascade. *Biochem J* 403: 139–148
- 580 [76] Yang, S. F., & Hoffman, N. E. (1984). Ethylene biosynthesis and its regulation in higher plants. *Annual Review of*  
581 *Plant Physiology*, 35(1), 155-189.
- 582 [77] Johnson, P. R., & Ecker, J. R. (1998). The ethylene gas signal transduction pathway: A molecular perspective.  
583 *Annual Review of Genetics*, 32(1), 227-254.
- 584 [78] Abeles, F. B., Morgan, P. W., & Saltveit, M. E. (1992). *Ethylene in Plant Biology* (2nd ed.). Academic  
585 Press.

- 586 [79] Hua, J., Chang, C., Sun, Q., & Meyerowitz, E. M. (1995). Ethylene insensitivity conferred by Arabidopsis ERS  
587 gene. *Science*, 269(5231), 1712-1714.
- 588 [80] Qiao, H., Chang, K. N., Yazaki, J., Ecker, J. R. (2009). Interplay between ethylene, ETP1/ETP2 F-box proteins,  
589 and degradation of EIN2 triggers ethylene responses in Arabidopsis. *Genes & Development*, 23(4), 512-521.
- 590 [81] Solano, R., Stepanova, A., Chao, Q., & Ecker, J. R. (1998). Nuclear events in ethylene signaling: A transcriptional  
591 cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes &  
592 Development*, 12(23), 3703-3714.
- 593 [82] Guo, H., Ecker, J. R. (2003). Plant responses to ethylene gas are mediated by SCF(EBF1/EBF2)-dependent  
594 proteolysis of EIN3 transcription factor. *Cell*, 115(6), 667-677.
- 595 [83] Chang, C., Kwok, S. F., Bleecker, A. B., & Meyerowitz, E. M. (1993). Arabidopsis ethylene-response gene ETR1:  
596 similarity of product to two-component regulators. *Science*, 262(5133), 539-544
- 597 [84] Kieber, J. J., Rothenberg, M., Roman, G., Feldmann, K. A., & Ecker, J. R. (1993). CTR1, a negative regulator of  
598 the ethylene response pathway in Arabidopsis, encodes a member of the raf family of protein kinases. *Cell*, 72(3),  
599 427-441.
- 600 [85] Ju, C., Yoon, G. M., Shemansky, J. M., Lin, D. Y., Ying, Z. I., Chang, J., ... & Ecker, J. R. (2012). CTR1  
601 phosphorylates the central regulator EIN2 to control ethylene hormone signaling from the ER membrane to the  
602 nucleus in Arabidopsis. *Proceedings of the National Academy of Sciences*, 109(47), 19486-19491.
- 603