

Original Research Article

EFFECT OF RATE OF FOLIAR FERTILIZER AND LEVEL OF IRRIGATION WATER APPLICATION ON CARROT PRODUCTIVITY

Comment [WU1]: Similarity index report????

Comment [WU2]: In what agro ecological condition???

Comment [WU3]: Name of country???

ABSTRACT

Carrot (*Daucuscarota L.*) productivity is still low in Tanzania due to inadequate management. There is dearth of information from the literature on the best management practices with respect to irrigation water (IW) and fertilizer levels for optimal carrot production. Consequently, farmers have limited access to means of boosting carrot productivity levels. Thus, a study was conducted to evaluate the effect of various levels of irrigation water application for carrots using drip irrigation and their interaction with foliar fertilizer on the growth, yield, and sugar content of carrots under a semi-arid climate in Morogoro region, Tanzania. Crop water requirement (ET_c) was calculated using established procedures upon which water application levels were based. Foliar fertilizer was applied at levels of 1 (F_1), 1.5 (F_2), and 2 (F_3) g/L/plant, while irrigation water levels applied were 60 (I_1), 80 (I_2), and 100 (I_3) percent of ET_c . The experiment was a 3x3 factorial with treatment combinations F_1I_1 , F_1I_2 , F_1I_3 , F_2I_1 , F_2I_2 , F_2I_3 , F_3I_1 , F_3I_2 , and F_3I_3 conducted over two seasons during 2020/2021 and 2021/2022. The highest yield of 30.9 t ha⁻¹ of carrots was obtained under 100% of ET_c IW application, and the lowest yield of 15.1 t ha⁻¹ was obtained under the 60% of ET_c IW application. Carrots under a deficit irrigation water level of 20%, i.e., 80% of ET_c IW were found to have the highest content of sugar. On the other hand, the highest yield of 26.1 t ha⁻¹ of carrots was obtained under the 2 g/L/plant fertilizer rate, and the lowest yield of 17.9 t ha⁻¹ of carrots was obtained under the 1 g/L/plant fertilizer rate. Carrots under the 1.5 g/L/plant fertilizer rate were found to have the highest content of sugar. Therefore, for optimum growth and yield, full irrigation in combination with foliar fertilizer application at a rate of 2 g/L/plant is recommended for carrots under Morogoro conditions. However, for high sugar content, deficit irrigation at 80% of ET_c in combination with foliar fertilizer application at a rate of 1.5 g/L/plant is recommended.

Key words: Carrots, foliar fertilizer, drip irrigation, sugar content, carrot yield, carrot growth.

1. INTRODUCTION

Carrots (*Daucus carota L.*) are one of the important and major root vegetables used for different purposes in the daily human diet [26]. It is rich in beta-carotene, which enhances resistance to blood and eye diseases, and a large amount of carrots in the diet has a favorable effect on the nitrogen balance [11,33].

Due to the importance of the crop for human existence, increased production to maximize yield is inevitable. However, in most parts of sub-Saharan Africa (SSA), carrot yield has persistently been low due to limited information on appropriate agronomic factors such as irrigation water level [23] and rate of fertilizer application [8]. Elsewhere, carrot yield has reportedly been increased under improved management. [21] reported that carrot growth, quality, and yield were positively affected by a 75% water application level in Germany. On the other hand, [17] reported that the foliar application of potassium at a rate of 3 ml/L/plant significantly enhanced the growth, yield, and root quality of carrots in Egypt. In Tanzania, information on the best management practices in managing irrigation water (IW) and fertilizer levels for optimal carrot production is limited. Therefore, farmers have limited access to means of boosting carrot productivity levels.

In addition, interaction between fertilizer and irrigation water application plays an important role in the production of carrots [13]. According to [24], various studies have been conducted, but these studies have mainly concentrated on carrot production and productivity. As a result, it remains unclear as to which fertilizer rate in combination with irrigation water level produces the best quantity and quality of carrots. For instance, [36] reported that the growth, yield, and quality parameters of carrots are significantly enhanced by foliar application of potassium under full irrigation, while [6] reported that application of foliar fertilizer in combination with a 75% water application level positively affected the growth of carrots. These studies point to the need for full investigation on the interactions between irrigation levels and the rates of fertilizer application and their effects on the growth, yield, and quality of carrots. There is a need to provide Tanzanian farmers with the information to know at what rate of fertilizer application and irrigation level the yield, quality, and growth of carrots would be enhanced as the increase in demand dominates the current production. According to [30], in Tanzania, the average per capita

daily carrot consumption is approximately 50 g, while the recommended daily consumption is at least 80 g [42]. This indicates that the consumption of carrots is still below the recommended level, therefore prompting increased production under irrigation and fertilization systems.

Comment [WU4]: Where is research gap in your study???

Comment [WU5]: Hypothesis of the research???

The study was therefore aimed at assessing (1) the effect of foliar fertilizer rates on the carrot growth, yield, and quality parameters; (2) the effect of irrigation water levels on the carrot growth, yield, and quality parameters; and (3) the interaction of water application and foliar fertilizer levels on the carrot growth, yield, and quality parameters.

2. MATERIALS AND METHODS

2.1 Description of the study area

The field experiment was conducted at the School of Engineering and Technology (SoET) research field (grid zone 350 213.19 E, 9 243 788.94 N, 37M; 512 m above sea level) as shown in Figure 1. The SoET is part of the Sokoine University of Agriculture (SUA) and is situated 3 km from the town center within Morogoro Municipality, Tanzania.




Figure 1: Map showing the location of the study site

2.2 Experimental Design

A factorial arrangement of treatments was laid out in a split-plot design. Two factors, factor 1 being water application regime, which was at 3 levels (100, 80, and 60% of ET_c), and factor 2, being foliar fertilizer application regime, which was at 3 levels (1, 1.5, and 2 g/L/plant), were investigated in a 3 x 3 factorial with three replications. There were a total of 9 treatment combinations (plots) for each replication. Each subplot measured 1.5 m x 2 m in size and was separated from the next by a 0.5 m buffer zone.

Drip irrigation system was used for the irrigation of carrots, with gate valves installed at the head of each lateral feeding the whole plot. The discharge of one emitter corresponding to full irrigation, i.e., 100% of ET_c (I_3) application level, was 2 litres per hour (L/h). The discharges of the other emitters corresponding to water application levels of 80 and 60% of ET_c (I_2 and I_1) were attained based on the 2 L/h discharge by measuring the discharge from an emitter using graduated cylinders and adjusting the valves. The measurements were taken at three positions (at the beginning, middle, and end) of the lateral. Thinning has been done two weeks after the emergence of the crop to attain the required spacing of 10 cm between plants, which gave 45 plants per sub-plot with a row spacing of 20 cm. No control for zero irrigation was applied because the experiment took place during the dry season. The applied water depth was 475.83 mm (I_3), 380.67 mm (I_2), and 285.498 mm (I_1) in season one, while in season two it was 233.64 mm (I_3), 186.92 mm (I_2), and 140.19 mm (I_1).

2.3 Crop water requirements

The reference crop evapotranspiration (ET_o) was determined using the Penman-Monteith equation [5] with the aid of INSTAT plus (v3.6) software [41].

The crop water requirement was determined as follows:

$$ET_c = ET_o \times K_c \quad (1)$$

Where,

ET_c : Crop evapotranspiration (mm day^{-1})

ET_o : Reference evapotranspiration (mm day^{-1})

K_c : Crop coefficient

Source: [15].




Figure 2: Mean monthly climatic data during the experiment

2.4 Crop growth parameters

The plant growth parameters that were collected include plant height, the number of leaves per plant, and the length of leaves. From each treatment, the height of 5 tagged carrot plants was measured using a ruler from the ground level to the top of the root shoot. The obtained measurements were recorded in cm. On the same plants, the length of leaves was measured from the bulb neck to the tip of the leaf using a ruler. The number of fully expanded leaves was counted and recorded as the number of leaves per plant.

2.5 Yield parameter

Yield data was obtained after harvesting. The same carrots that had been used to take data for the plant growth parameters were also used for the determination of fresh root weight by using a weighing scale balance. The obtained weights were recorded in tons/ha.

2.6 Root growth parameters

The same carrots that had been used to take data for the plant growth parameters were also used for the determination of the root growth and quality of carrots. The root length was determined using a vernier caliper starting from the shoulder to the end of the tap root. The shoulder and core diameter of roots were measured at 0.5 cm from the top of the shoulder using a vernier caliper, and measurements were recorded in mm.

2.7 Quality parameter

The total soluble solids, i.e., sugar level in carrots were measured using a hand-held refractometer (0–30% Brix) and recorded in percentage.

2.8 Statistical analysis

Analysis of variance (ANOVA) on the data collected was done using Genstat statistical software. The mean separation was carried out using Duncan's multiple range test (DMRT) at the 0.05 probability level [12].

Comment [WU6]: Edaphic, soil & environmental conditions are missing????

3. RESULTS AND DISCUSSION

3.1 Soil and irrigation water analysis

Tables 1 and 2 show the results of soil and irrigation water tests in the study area. According to the classification suggested by [19], the soil texture was sandy clay. Sandy clay soil is suitable for vegetable production but needs regular watering and fertilizing to ensure healthy development [27]. According to [34], the sodium adsorption ratio (SAR) was low. Irrigated water with low SAR is safe with regard to sodicity hazard [43].

Table 1: Physical and chemical properties of the soil in the study area

Soil properties	Values	Description
pH 1:2.5 in H ₂ O	6.12	Medium acidic soil
Total Nitrogen (%)	0.11	Low
Extractable Phosphorus (mg kg ⁻¹)	0.28	Low
EC _e (dS/m)	0.16	Normal
Potassium (Cmol kg ⁻¹)	0.208	Medium
Particle size distribution (%)		
Clay	39.76	Sandy clay
Sand	54.56	
Silt	5.68	

Table 2: Irrigation water quality

Parameters	Results	Description
Ph	7.8	Normal
Sodium (me L ⁻¹)	9.95	Slight to moderate
EC _w (dSm ⁻¹)	1.013	Medium
Phosphorus (mg L ⁻¹)	0.04	Non-problem
SAR (meq L ⁻¹)	1.33	Low
Bicarbonate (meq L ⁻¹)	0.38	Good
Carbonate (meq L ⁻¹)	1.38	Medium
Nitrogen (mg L ⁻¹)	0.03	Normal

3.2 Effect of different irrigation water levels on the growth, yield, and quality of carrots

3.2.1 Plant height

Carrots irrigated at 100% water application level produced the most plant height in season one (43.27 cm) and season two (40.92 cm) compared to 60% and 80% water application levels, while 60% water application level resulted in the least plant height in the first season (37.4 cm) as well as the second season (37.42 cm) (Table 3). However, during the first season, the plant heights for 80% and 100% water application levels were not significantly different at the 5% level, but both differed significantly ($p < 0.05$) from the 60% water application level. In the second season, plant height differed significantly ($p < 0.05$) among the three water application levels.

It is evident that the plant height increased gradually as irrigation water levels increased up to full irrigation. This could be attributed to the favorable soil moisture for proper plant growth associated with the rapid increase and expansion of plant cells that play vital roles in the biological and physiological processes of carrot plants, resulting in an increase in the plant height. This is in agreement with the findings by [29], who reported that the height of carrots increased with increasing levels of irrigation water.

Table 3: Effect of different water application levels on the growth, quality, and yield of carrots

Water level	Plant height (cm)	Number of leaves	Leaf length (cm)	Fresh root weight(t/ha)	Root length (cm)	Root core diameter(mm)	Root shoulder diameter (mm)	TSS (%)
First season								
60%	37.4a	9.6 a	10.66a	16.7 a	14.25a	13.19 a	23.77 a	8.15a
80%	41.54 b	9.9 a	10.76a	21.9 b	16.07b	16.53 b	26.82 b	8.5a
100%	43.27b	10.6 b	11.34a	30.9 c	17.07 b	18.83 c	29.08 c	8.11a
LSD (5%)	2.185	0.366	0.716	1.3	1.008	1.728	1.816	0.499
Second season								
60%	37.4 a	8.7 a	9.53 a	15.1 a	13.82 a	12.8 a	23.39 a	6.38 a
80%	38.83 b	9.7 b	10.06 b	21.2 b	15.36 b	14.03 b	25.90 b	7.67b
100%	40.92 c	10.4 c	10.54 c	28.4 c	16.94 c	15.98 c	27.13 b	6.34 a
LSD (5%)	0.606	0.378	0.325	1.19	1.02	1.025	2.073	1.107

TSS: Total Soluble Solids, LSD: Least Significance Difference, DMRT: Duncan's Multiple Range Test.

Means followed by the same letter (s) in the same column are not significantly different according to DMRT.

3.2.2 Number of leaves

Irrigating carrots at 100% water application level produced the highest number of leaves in the first season (10.6) as well as in the second season (10.4) compared with the other treatments. The lowest number of leaves per plant was obtained from the 60% water application level in both seasons (Table 3). The number of leaves per plant was not significantly different ($p > 0.05$) between the 60% and 80% water application levels, but both differed significantly ($p < 0.05$) from the 100% water application level in the first season. However, in the second season, the number of leaves per plant differed significantly ($p < 0.05$) among the three water application levels.

From the results, it is evident that water application levels of 60 and 80% decreased the number of leaves per plant. This could be due to the effect of water deficit which decreased the photosynthetic capacity that led to decreased leaf stomata conductance due to stomata closure as pointed out by [38].

3.2.3 Leaf length

Comment [WU7]: Need more discussion

Results in Table 3 show that irrigating carrots at 100% water application level resulted in the maximum length of leaves in season one (11.34 cm) as well as in season two (10.54 cm). Likewise, the 60% water application level recorded the minimum length of leaves in the first season (10.66 cm) as well as in the second season (9.53 cm). Further, the leaf lengths were not significantly different ($p > 0.05$) among the three water application levels in the first season. However, in the second season, the three water application levels differed significantly ($p < 0.05$) in relation to leaf length.

A similar explanation to that advanced for the number of leaves could apply to the leaf length as these are very much related phenomena [22].

3.2.4 Fresh root weight

The highest fresh root weight in the first season (30.9 t ha⁻¹) as well as in the second season (28.4 t ha⁻¹) was recorded under the 100% water application level, while the lowest fresh root weight in the first season (16.7 t ha⁻¹) as well as in the second season (15.1 t ha⁻¹) was obtained under the 60% water application level. Fresh root weight was significantly different ($p < 0.05$) among the three water application levels in both seasons (Table 3).

The reduced fresh root weight under the 60% water application level could be due to the soil water stress that caused the carrot plants to absorb inadequate nutrients essential for root growth. In connection with this, [28] reported that drought stress throughout the entire growth period of carrots caused a more severe reduction in the root production.

3.2.5 Root length

Full irrigation resulted in the maximum length of roots in season one (17.07 cm) as well as in season two (16.94 cm), while the minimum root length in season one (14.25 cm) as well as in season two (13.82 cm) was recorded under the 60% water application level. Further, the root length was not significantly different ($p > 0.05$) between the 80 and 100% water application levels, but both differed significantly ($p < 0.05$) from that under the 60% water application level in season one. However, in season two, the root length differed significantly ($p < 0.05$) among the three water application levels (Table 3).

The increased root length due to higher water application levels could be due to availability of sufficient moisture which helped in rapid cell elongation leading to longer root formation. This is consistent with findings by [1] and [2], who reported that root length of carrots was higher with higher amount of water level.

3.2.6 Root core and shoulder diameters

The maximum root core diameter of 18.83 mm for season one and 15.98 mm for season two were obtained under the 100% water application level, while the minimum root core diameter of 13.19 mm in the first season and 12.8 mm in the second season were obtained under the 60% water application level. Likewise, the maximum root shoulder diameter in the first season (29.08 mm) and in the second season (27.13 mm) were obtained under the 100% water application level and the minimum root shoulder diameter in the first season (23.77 mm) and in the second season (23.39 mm) were obtained under the 60% water application level (Table 3). Further, the root core diameter differed significantly ($p < 0.05$) among the three water application levels in the first as well as the second season. On the other hand, the root shoulder diameter followed the same trend as that of the root core diameter in the first season in terms of statistical significance but during the second season the root shoulder diameter under the 80 and 100% water application levels did not differ significantly ($p > 0.05$).

Deficit irrigation (60 and 80% water application levels) produced the lowest values in root core and shoulder diameters for carrots. This could be due to the effect of prolonged water stress, with consequent reduction in overall growth. In confirmation of this, [37] reported that water deficits cause water stress in plants, prevent plant and root growth, and reduce the absorbing areas and capacities of plant roots.

3.2.7 Total soluble solids (TSS)

As shown in Table 4, the highest TSS in the first season (8.5%) and the second season (7.67%) were obtained under the medium water application level (80% water application level), while the lowest TSS in the first season (8.11%) and the second season (6.34%) were obtained under full irrigation. Further, the TSS were not significantly influenced by water application levels in the first season. However, in the second season, the TSS under the 60 and 100% water application levels differed significantly ($p < 0.05$) from those under the 80% water application level.

The abundance of TSS content in the roots of the lesser irrigated carrot plants is more of a physiological characteristic that could be explained by plant photosynthate redistribution into the roots as a result of the water deficit, which resulted in lower water content but increased sugar content. A similar observation has also been made by [20], who reported that increased moisture has a negative influence on TSS content on carrot roots.

3.3 Effects of Foliar Fertilizer on the Growth, Yield, and Quality of Carrots

3.3.1 Plant height

Spraying foliar fertilizer at a rate of 2 g/L/plant resulted in the most plant height in season one (43.27 cm) and season two (41.14 cm), while the foliar fertilizer rate of 1 g/L/plant resulted in the least plant height in the first season (37.55 cm) as well as the second season (36.18 cm). The plant heights under 1.5 and 2 g/L/plant fertilizer levels were not significantly different at the 5% level in the first season, but both differed significantly ($p < 0.05$) from the 1 g/L/plant fertilizer level. However, in the second season, plant height differed significantly ($p < 0.05$) among the three fertilizer levels. The results (Table 4) show that carrot height consistently increased with fertilizer rate. The application of higher fertilizer rates has the potential to fulfill the requirements of the plant for nutrients than lower fertilizer rates. This observation was also made by [9], who reported that the highest level of foliar feeding produced the most carrot height compared to the lowest fertilizer level. In their study, [4] reported that foliar application of up to 1.5 g/L/plant fertilizer significantly enhanced carrot growth by increasing carrot height. A study by [17] reported that spraying carrot plants with foliar fertilizer at a rate of 3 ml/L/plant produced the highest values of plant height compared to control treatment (without foliar application) in two seasons.

3.3.2 Number of leaves

As shown in Table 4, the number of leaves per plant followed the same trend as that of plant height in the two seasons in terms of statistical significance. Spraying foliar fertilizer at a rate of 2g/L/plant produced the highest number of leaves per plant in the first season (10.4) as well as in the second season (10.08) compared with the other treatments. The lowest number of leaves per plant was obtained from the 1 g/L/plant fertilizer rate in both seasons.

In agreement with this, [16] stated that the use of high foliar feeding rates improved the number of leaves on carrots compared to low fertilizer rates. [9] reported that among the different fertilizer rates (0.5, 1, 1.5 and 2 ml/L/plant), 2 ml/L/plant produced the highest number of leaves.

Table 4: Effect of different rates of foliar fertilizers on the growth and yield parameters of carrots

Fertilizer level	Plant height (cm)	Number of leaves	Leaf length (cm)	Fresh root weight(t/ha)	Root length (cm)	Root core diameter(mm)	Root shoulder diameter (mm)	TSS (%)
First season								
1	37.55 a	9.4 a	10.47 a	19.8 a	15.53 a	13.09 a	22.6 a	8.27b
1.5	41.39 b	10.2 b	10.92 ab	23.6 ab	15.71 a	17.28 b	28.15 b	8.88 c
2	43.27 b	10.4 b	11.37 b	26.1 b	16.16 a	18.18 b	28.92 b	7.6 a
LSD(0.05)	2.185	0.366	0.716	1.3	1.008	1.728	1.816	0.499
Second season								
1	36.18 a	9.2 a	9.45 a	17.9 a	15 a	11.49 a	20.38 a	6.36a
1.5	39.85 b	9.7 b	10.29 b	22.9 b	15.42 a	15.11 b	27.53 b	8.25b
2	41.14 c	10.08 c	10.39 b	24.03 b	15.7 a	16.21 c	28.51 b	5.78 a
LSD(0.05)	0.606	0.378	0.325	1.19	1.02	1.025	2.073	1.107

3.3.3 Leaf length

Spraying foliar fertilizer at a rate of 2 g/L/plant resulted in the maximum length of leaves in season one (11.37 cm) as well as in season two (10.39 cm). Likewise, the 1 g/L/plant fertilizer rate recorded the minimum length of leaves in the first season (10.47 cm) as well as in the second season (9.45 cm). Further, the leaf lengths were not significantly different ($p>0.05$) between the 1 and 1.5 g/L/plant fertilizer rates in the first season as was the case for the 1.5 and 2 g/L/plant fertilizer rates but the 1 g/L/plant fertilizer rate was significantly different ($p<0.05$) from the 2 g/L/plant fertilizer level. In the second season, the leaf lengths under the 1.5 and 2 g/L/plant fertilizer rates were not significantly different ($p>0.05$) but both differed significantly ($p<0.05$) from the 1 g/L/plant fertilizer rate (Table 4).

Foliar nutrition using a high fertilizer rate can eliminate problems like fixation and immobilization of nutrients by penetrating the stomata of the leaf, entering the cells rapidly and fulfilling the nutrient demand of the growing plant, resulting in enhanced length of the leaf [14]. This is in agreement with the findings of [6], who reported that an increased application rate of foliar feeding to carrots enhanced the length of the leaves.

3.3.4 Fresh root weight

The highest fresh root weight in the first season (26.1 t/ha) as well as in the second season (24.03 t/ha) were recorded under the 2 g/L/plant fertilizer level, while the lowest fresh root weight in the first season (19.8 t/ha) as well as in the second season (17.9 t/ha) were obtained under the 1 g/L/plant fertilizer level. Fresh root weight followed the same trend as that for leaf length in terms of statistical significance in both seasons (Table 4). The fresh root weight consistently increased with increasing fertilizer rates. In their study, [40] reported that higher fertilizer rates directly increased the root fresh weight by producing heavier roots than the lower rates.

3.3.5 Root length

Results (Table 4) indicate that the root length was not significantly influenced by the rate of fertilizer application in both seasons. Nevertheless, the highest rate of 2 g/L/plant resulted in the maximum length of root in both seasons, while the minimum root length was recorded under the 1 g/L/plant fertilizer rate in both seasons. This positive effect on the root length due to increased fertilizer levels could be attributed to the favorable fertilizer nutrients for proper plant growth associated with rapid increase and expansion of plant cells that play vital roles in the biological and physiological processes of carrot plants, resulting in an increase of the root length. A similar observation was also made by [8], who reported that higher fertilizer rates produced longer roots than the lowest fertilizer rates.

3.3.6 Root shoulder and core diameters

Results (Table 4) indicate that the root shoulder diameter did not differ significantly ($p > 0.05$) between the 1.5 and 2 g/L/plant fertilizer levels in the first as well as the second seasons but both differed significantly ($p < 0.05$) from that under the 1 g/L/plant fertilizer level. On the other hand,

the root core diameter did not differ significantly ($p>0.05$) between the 1.5 and 2 g/L/plant fertilizer levels in the first season, but during the second season, the root core diameter differed significantly ($p<0.05$) among the three fertilizer levels. The maximum root core diameter of 18.18 cm for season one and 16.21 mm for season two was obtained under the 2 g/L/plant fertilizer level, while the minimum root core diameter of 13.09 mm in the first season and 11.49 mm in the second season was obtained under the 1 g/L/plant fertilizer level. Likewise, the maximum root shoulder diameter in the first season (28.92 mm) and the second season (28.51 mm) was obtained under the 2 g/l/plant fertilizer level. The minimum root shoulder diameter in the first season (22.6 mm) and the second season (20.23 mm) was obtained under the 1 g/l/plant fertilizer level. These results are in agreement with those of [9]. Other studies [7, 17, 39] reported that higher fertilizer rates directly increased the root core and shoulder diameters than the lower fertilizer rates.

3.3.7 Total soluble solids (TSS)

The highest TSS in the first season (8.88%) and the second season (8.25%) were obtained under the 1.5 g/L/plant fertilizer level, while the lowest TSS in the first season (7.6%) and the second season (5.78%) were obtained under the 2 g/L/plant fertilizer level. Further, the TSS were significantly influenced by fertilizer levels in season one. However, TSS under the 1 g/L/plant fertilizer level did not differ significantly ($p>0.05$) from that under the 2 g/L/plant fertilizer level during the second season (Table 6). The highest fertilizer rate (2 g/L/plant) resulted in decreased sugar levels in the carrots. This is an intriguing result as it tends to suggest that there is an optimal level of fertilization for TSS accumulation. In confirmation of this, [18, 31, 32] reported that less fertilized treatments had a greater tendency to increase sugar content in carrots than the most fertilized treatments.

3.4 Effect of the Interaction between Irrigation Water and Foliar Fertilizer Levels on the Growth, Yield, and Quality of Carrots

3.4.1 Plant height

The interaction between full irrigation \times 2 g/L/plant fertilizer level produced the highest plant height in the first season (46.43 cm) as well as in the second season (43.08 cm) compared to other interactions, while 60% water application level \times 1 g/L/plant fertilizer rate resulted in the

least plant height in the first season (33.83 cm) as well as in the second season (34.49 cm). Further, the plant height under the 60% water application level \times 1 g/L/plant fertilizer rate differed significantly ($p < 0.05$) with other combinations in both seasons (Table 5). The height of carrots increased consistently with fertilizer and irrigation water increments, up to 2 g/L/plant in combination with full irrigation. This could mean that the nutrients from irrigation water (full irrigation) in combination with the nutrients from foliar feeding enhanced photosynthesis which resulted in better plant growth. The findings are in harmony with those by [10, 35], who reported that the increase in irrigation water and fertilizer levels resulted in increased carrot height.

Table 5: Effects of the interaction between water application and foliar fertilizer levels on the plant height

Fertilizer levels	1 g	1.5 g	2 g
Water levels	Plant height		
	First season		
60%	33.83 a	38.49 b	39.87 bc
80%	39.11 b	42 bc	43.50 cd
100%	39.7 bc	43.69 cd	46.43 d
LSD (5%)	3.784		
	Second season		
60%	34.49 a	38.29 c	39.5 d
80%	35.74 b	39.91 de	40.85 ef
100%	38.31 c	41.37 f	43.08 g
LSD (5%)	1.105		

3.4.2 Number of leaves

Results (Table 6) show that the number of leaves per plant under the full irrigation \times 2 g/L/plant fertilizer level differed significantly ($p < 0.05$) with other combinations in the first season. However, in the second season, the 60% water application level \times 1 g/L/plant fertilizer level differed significantly ($p < 0.05$) with other combinations. Application of full irrigation in combination with 2 g/L/plant fertilizer rate produced the highest number of leaves in season one (11.27) as well as in season two (10.51), while the lowest number of leaves in season one (8.93)

as well as in season two (8.16) was obtained under the 60% water application level × 1 g/L/plant fertilizer level. A similar explanation as that advanced for the case of plant height for the full irrigation × the 2 g/L/plant fertilizer rate combination would seem to be the case for the number of leaves produced [3].

Table 6: Effects of the interaction between water application and foliar fertilizer levels on the number of leaves

Fertilizer levels	1 g	1.5 g	2 g
Water levels	Number of leaves		
	First season		
60%	8.93 a	9.93 bc	9.97 bc
80%	9.47 ab	10.17 bc	10.14 bc
100%	10.03 bc	10.5 c	11.27 d
LSD (5%)	0.635		
	Second season		
60%	8.16 a	8.88 b	9.31 bc
80%	9.16 bc	9.75 cd	10.43 de
100%	10.36 de	10.43 de	10.51 e
LSD (5%)	0.65		

3.4.3 Leaf length

The maximum leaf lengths in season one (11.56 cm) as well as in season two (11.25 cm) were obtained under the full irrigation in combination with the 2 g/L/plant fertilizer rate. Likewise, the minimum leaf lengths in season one (9.72 cm) as well as in season two (9 cm) were obtained under the 60% water application level × 1 g/L/plant fertilizer rate (Table 7). The leaf lengths were not significantly different ($p > 0.05$) among various interactions in season one. However, in season two, the full irrigation × 2 g/L/plant fertilizer level differed significantly ($p < 0.05$) from other the combinations. A similar explanation as that given for the plant height and number of leaves could be advanced for the case of the length of leaves [25].

Table 7: Effects of the interaction between water application and foliar fertilizer levels on the leaf length of carrots

Fertilizer levels	1 g	1.5 g	2 g
Water levels	Leaf length		
	First season		
60%	9.72 a	10.85 ab	11.41 b
80%	10.48ab	10.66 ab	11.14 b
100%	11.22ab	11.24 ab	11.56 b
LSD (5%)	1.241		
	Second season		
60%	9 a	10.09 cde	9.5 abc
80%	9.47 ab	10.28 de	10.42 de
100%	9.87 bed	10.51 e	11.25 f
LSD (5%)	0.563		

Conclusion

This study was conducted to evaluate the effects of different water application levels in combination with foliar fertilizer rates on the growth, sugar content, and yield of carrots under drip irrigation. Spraying foliar fertilizer at a rate of 2 g/L/plant enhanced the growth and yield of carrots compared to other fertilizer levels in two seasons. However, 1.5 g/L/plant fertilizer level resulted in the highest sugar content in the roots. On the other hand, full irrigation increased carrot production in both seasons compared to other water application levels. However, carrots under an IW level of 80% were found to have the highest content of sugar. Therefore, for optimum growth and yield, full irrigation and foliar fertilizer application at a rate of 2 g/L/plant is recommended for carrots under Morogoro conditions. On the other hand, for high sugar content, deficit irrigation at 80% of ET_c in combination with foliar fertilizer application at a rate of 1.5 g/L/plant is recommended.

Comment [WU8]: Recommendation about future study???

Comment [WU9]: Contribution to knowledge???

References

1. Afrin A, Islam MA, Hossain M, Hafiz MH. Growth and yield of carrot influenced by organic and inorganic fertilizers with irrigation interval. *Journal of Bangladesh Agricultural University*. 2019; 17(3): 338–343.
2. Ahmad Z, ali N, Ahmad M, Ahmad S. Yield and economics of carrot production in organic farming. *Pakistan Journal of Agriculture*. 2005; 22(7): 12-15.
3. Alam M, Mallik S, Costa D. Effect of irrigation on the growth and yield of carrot in hill valley. *Bangladesh Journal of Agricultural Research*. 1970; 35(2): 323–329.
4. Alhariri A, Boras M. Responses of seed germination and yield related traits to seed pretreatment and foliar spray of humic and amino acids compounds in carrot. *International Journal of Chemical Studies*. 2020; 8(4): 26–30.
5. Allen RG, Pereira LS, Raes D, Smith M. Crop evapotranspiration, guidelines for computing crop water requirements. Irrigation and drainage paper 56. FAO: Rome. 1998; 301-302.
6. Anub RR. Growth and yield of carrots as affected by different irrigation frequency, organic soil amendments and inorganic fertilizer. *International Journal of Humanities and Social Sciences*. 2019; 11(4): 9–19.
7. Arshad B, Khan AA, Babar MM, Sarki A. Effect of different levels of water soluble NPK fertilizers on the growth and yield of carrot by using drip irrigation. *International Journal for Research in Applied Science and Engineering Technology*. 2015; 3(5): 708–714.
8. Badr MA, Helmy YI. The influence of foliar application of potassium on yield and quality of carrot plants grown under sandy soil conditions. *Australian Journal of Basic and Applied Science*. 2011; 5(3): 171–174.
9. Badr M, Abouhoussein S, Helmy Y. The influence of foliar application of potassium on yield and quality of carrot plants grown under sandy soil conditions. *Australian Journal of Basic and Applied Sciences*. 2010; 4(1): 123-126.
10. Batra BR. Effect of different levels of irrigation and fertilization on growth and yield of carrot for root production. *Vegetable Science*. 1990; 17(2): 127–139.
11. Bjarnadottir A. Nutrition facts and health benefits. *Journal of Food Sciences*. 2015; 3(7): 12-15.
12. David, DB. Multiple range and multiple f tests. *International Biometric Society*. 2013; 11(1): 1–42.

Comment [WU10]: Tally each reference with text???

Comment [WU11]: Reference format??

13. Dauda SN, Ajayi FA, Ndor E. Growth and yield of water melon as affected by poultry manure application. *Electronic Journal of Environmental, Agricultural and Food Chemistry*.2009;8(4): 305–311.
14. Devi ND. Responds of vegetable crops to foliar feeding of water soluble fertilizer.*Brazilian Journal of Soil Science*. 2016;11(1): 242–251.
15. Doorenbos J, Pruitt WO. Crop water requirements.Irrigation and Drainage paper.FAO., Sub-regional office for East and Southern Africa. 1977; 35-54.
16. El-HelalyM. Effect of foliar application of humic and fulvic acids on yield and its components of some carrotCultivars. *Journal of Horticultural Science & Ornamental Plants*.2018;10(3): 159–166.
17. El-nasr A, Ibrahim EA. Effect of different potassium fertilizer rates and foliar application with some sources of potassium on growth, yield and quality of carrot plants.*Journal of Plant Production*. 2011;2(4): 559–569.
18. Evers AM. Effects of different fertilization practices on the quality of stored carrot.*Agricultural and Food Science*. 1989; 61(2): 123–134.
19. FAO. guidelines for soil profile description soil development and conservation services. Land and Water Division.(2nd Edition). FAO: Rome.1977; 96–97.
20. Fikselová M, Mareček J, MellenM. Carotenes content in carrot roots as affected by cultivation and storage. *Vegetable Crops Research Bulletin*.2010. 73(1): 47–54.
21. GutezeitB. Yield and quality of carrots as affected by soil moisture and N-fertilization. *Journal of Horticultural Science and Biotechnology*. 2001;76(6): 732–738.
22. Jahan I, Hossain M, Karim M. Effect of salinity stress on plant growth and root yield of carrots. *Progressive Agriculture*.2020; 30(3): 263–274.
23. Joslanny HV, CatarinyCAI, Elis M, Laylton AS, Gustavo H, Pedro HFF.irrigation levels and soil covers in carrot crop. *Journal of Agriculture and water resources*. 2020; 5(4): 12-15.
24. Kifle M, Gebremicael TG, Girmay A, GebremedihinT. Effect of surge flow and alternate irrigation on the irrigation efficiency and water productivity of onion in the semi-arid areas of North Ethiopia.*Agricultural Water Management*. 2017;187(5): 69–76.
25. Kelly TM. Nutrient management of vegetable and row crops.*Journal of Food Sciences*.2015;9 (3): 6-8.

26. Kwabena A. Influence of grass cutter, chicken manure and NPK amendments on soil characteristics, growth, and yield response of carrotstthesis. MSc level, university of Education, Winneba.2011;73- 76 pp
27. Landon, JR. Booker tropical soil manual. Paperback edition. Longman scientific and technical copublish: USA. 1991; 96–98.
28. Li P, He S, Yang N, Xiang G. Groundwater quality assessment for domestic and agricultural purposes in Yan’an City, northwest China: implications to sustainable groundwater quality management on the loss Plateau. *Environmental Earth Sciences*.2018;77(23): 1–16.
29. Ludong DPM. Effects of irrigation rate on the growth, yield, nutritive value, and water use efficiency of Carrot and Broccoli. Published dissertation for award of MSc. degree at Curtin University of Technology, Australia.2008; 87pp.
30. Lukmanji Z, Hertzmark E. Mlingi N, Assey V. Tanzania food composition tables. Muhimbili University of Allied Science,Tanzania.2008; 259pp.
31. Mbatha AN. Influence of organic manure on the yield and quality of vegetables. *Japanese Journal of Soil Science and Plant Nutrition*. 1936;10(1): 37–46.
32. Noella J, Umuhoza K, Sylvestre H, Philippe S. Nutritional quality of carrots influenced by farm yard manure. *World Journal of Agricultural Sciences*.2014; 2(7):102–107.
33. Pant B, ManandharS. Vitro propagation of carrots. *Scientific World*.2007;5(5): 51–53.
34. Park DM, White SA, Mccarty LB, MenchykN. Interpreting irrigation water quality reports. 2015. Sited visited on 24/08/2021.
35. Prabhakar M, Srinivas K, Hegde DM. Effect of irrigation regimes and nitrogen fertilisation on growth, yield, N uptake, and water use of carrots. *Gartenbauwissenschaft*.1991; 56(5): 206-209.
36. Quezada C, Fischer S, Campos J, Ardiles D. Water requirements and water use efficiency of carrot under drip irrigation in haploxerand soil. *Soil Science and Plant Nutrients*. 2006; 11(1): 16 – 28.
37. Reid JB, Gillespie RN. Yield and quality responses of carrots to water deficits. *New Zealand Journal of Crop and Horticultural Science*.2017;45(4): 299–312.
38. Sato T, Abdalla OS, Oweis TY, SakurataniT. Effect of supplemental irrigation on leaf stomatal conductance of field grown wheat in northern Syria. *Agric. Water Manage*.

2006;85(6): 105-112.

39. Sharangi AB, Paria NC. Growth, yield, and qualitative responses by carrot to varying levels of nitrogen and potassium. Hort. J. 1995;8(2): 161–164.
40. Shibairo SI, Upadhyaya MK, Toivonen PMA. Potassium nutrition and postharvest moisture loss in carrots. Journal of Horticultural Science and Biotechnology. 2016; 73(6): 862-866.
41. Stern R, Rijks D, Dale I, Knock J. INSTAT climatic guide. Statistical service centre: Washington. 2006; 247–281.
42. WHO. Region Report. Arusha region, Food Security and Cooperatives. Dar es Salaam. 2003; 512 pp.
43. Zaman M, Shahid SA, Heng L. Guideline for salinity assessment, mitigation and adaptation using nuclear and related techniques. Springer nature: Switzerland. 2018; <https://doi.org/10.1007/978-3-319-96190-3>.