

## **“Comparative efficacy of some insecticides and their combination against okra shoot and fruit borer, *Earias vitella* (Fabricius)”**

**ABSTRACT:** The research was conducted on okra crop to check the efficacy of some insecticides and their combinations against okra shoot and fruit borer, *Earias vittella* (Fab.). The study was carried out during *kharif* season 2023 at Central Research Farm of Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj (U.P.) with eight treatments including untreated check which were replicated thrice. All the treatments were found to be superior over untreated control, Neem oil 3%, Spinosad 45 SC, Cypermethrin 25 EC, Profenophos 50 EC, Spinosad 45SC + Neem Oil 3%, Cypermethrin 25EC + Neem oil 3%, Profenofos 50 EC + Cypermethrin 25 EC were tested to compare the efficacy against *Earias vitella* and their influences on yield of okra. Here the best and the most economic treatment was Profenofos 50 EC + Cypermethrin 25 EC followed by Cypermethrin 25EC + Neem oil 3%, Spinosad 45SC + Neem Oil 3%, Spinosad 45 SC, Cypermethrin 25 EC, Profenophos 50 EC and the least effective treatment was Neem oil 3%. The highest yield among the treatments was noticed in Profenofos 50 EC + Cypermethrin 25 EC (140quintal/hac.) and the lowest in Neem oil 3% (75quintal/hac.) with cost benefit ratio 1:4.7 and 1:2.0, respectively.

---

**Keywords:** Okra shoot and fruit borer, *Earias vitella*, insecticides, chemicals, combinations, management.

**1. INTRODUCTION:** An important vegetable crop, okra (*Abelmoschus esculentus* L.) is a member of the Malvaceae family. (Singh *et al.*, 2008) cultivated in tropical, subtropical and mild temperate parts of the world. Okra locally known as Bhendi and Lady's finger worldwide. It is generally a self-pollinating crop belonging to the Malvaceae (Sekyere *et al.*, 2011). Okra is referred to as the "Vegetable's Queen." It is prized for the soft green fruits it bears. It can be prepared in a number of ways and is a key ingredient in a variety of recipes (Janu and Kumar 2022). Before sugarcane juice is made into brown sugar and jiggery, the root and stems are used to clarify it. (Baloch *et al.*, 1990). It has good nutritional value particularly the high content of vitamin C (13mg/100g), fat (0.2g/100g), carbohydrate (64g/100g), iron (1.5mg/100g), moisture (89.6g/100g), protein (1.9g/100g), fibre

(1.2g/100g), calories 35 (kcal/100g), and other minerals (**Fageria et al., 2012**). The total area and production under okra in the world is reported to be 1.26 million ha and 22.29 million tonnes, respectively. With an area of 1148 thousand hectares, an annual production of 6346 million tonnes, and a productivity of 11.9 million tonnes/ha, India leads the world in okra production, accounting for 5784 thousand tonnes (72% of total global production). Gujarat is the state that produces the most okra, with an estimated 1019.42 thousand tons produced annually. (**National Horticulture Board, 2021-22**). The several insect pests of okra, some are serious pests that frequently arise; for example, the borer has been recorded to inflict damage to okra shoots of 24.6 to 26.0 percent and losses to fruits of 40 to 100 percent. (**Pareek et al., 2003 and Shinde et al., 2005**). The fruit borer, *Earias vittella*, which feeds on okra shoots, lays light greenish blue eggs with longitudinal ridges. The completely developed larvae (caterpillar) had a length of 1.64 cm and was orange, black, and green in color. The larvae have a pale-yellow ventral side and a brownish dorsal side with white stripes. Its pupa is a rough, bluntly rounded, gray, inverted boat-shaped cocoon that forms on the stem or fruits and is chocolate brown in color. Its adults have tiny forewings that measure 1.25 cm wide. The adult borer's head and thorax are ochreous white, while its forewings are pale white with a green spot in the middle that resembles a wedge, and its hind wings are a silvery, creamy white color. The middle of the forewing of *Earias vittella* moths has a slender, light green longitudinal band. The main distinction between a male and a female is that the male has thick hair at the anal end of the body, while the female is larger than the male and has a V-shaped anal region. (**Aman et al., 2022**). During the night, the female moth deposits 200–400 eggs by herself on the delicate leaves, bracts, and flower buds of okra plants. Incubation period of eggs are 3-4 days and the caterpillar passes through 6 stages, becoming full grown in 10-16 days. The moth emerges about 8–14 days during the summer and 18–23 days during the winter after pupating on plants or on the ground among fallen leaves. It takes 17–29 days to complete the lifecycle. In a year, multiple overlapping generations are completed. (**Kaveri and Kumar 2020**).

**2. MATERIALS AND METHODS:** The spraying was done after the population level reaches to ETL. The ETL level of the pest is 5% of shoot damage and 10% fruit damage (**Shirale et al., 2012**). The observations were recorded one day before spraying, 7<sup>th</sup> and 14<sup>th</sup> days after spraying. The assessment of the shoot damage was done by calculating the number of damaged shoots and total numbers of healthy shoots observe from five randomly selected

plants per plot and expressed in percentage. The percent of fruit damage was assessed at each picking by counting the total number of affect fruits from each plot.

### **Shoot Infestation**

The total number of shoots, as well as the number of infested shoots, was observed and recorded at weekly intervals from five selected plants in each plot. Shoot infestation was determined by percent using the following formula-

%Shoot infestation =

$$\frac{\text{Number of infested shoots}}{\text{Number of total shoots}} \times 100$$

### **Fruit Infestation**

The percentage of fruit infestation determined using the following formula:

%Fruit infestation =

$$\frac{\text{Number of infested fruit}}{\text{Number of total fruits}} \times 100$$

**Choudhury et al., (2021)**

The healthy marketable yield from different treatments were collected separately and weighed. There were two sprays throughout the research period and the treatment cost and common cost of cultivation per hectare was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price; while the net benefit was obtained by subtracting the total cost of plant protection from total income. The C: B was calculated by following formula -

$$\text{Cost Benefit Ratio} = \frac{\text{Gross return}}{\text{Total cost of cultivation}}$$

**(Nalini and Kumar, 2016)**

## **3. RESULTS AND DISUSSION**

**3.1 Effect of some insecticides and their combination against Okra shoot and fruit borer, *Earias vitella* during *kharif* season (2023) (First spray).**


The data of overall mean (7DAS and 14 DAS) resulted that all the treatments were found to reduce the shoot infestation significantly over control and lowest shoot damage was recorded in Profenofos 50 EC + Cypermethrin 25 EC (4.63%) followed by Cypermethrin 25EC + Neem oil 3% (7.22%), Spinosad 45SC + Neem Oil 3% (8.11%), Spinosad 45 SC (9.40%), cypermethrin 25 EC (12.24%), profenophos 50 EC (15.21%). The least effective treatment was Neem oil 3% (17.43%) compared to all the treatments and maximum damage of shoot is found in untreated plot (27.66%).

### **3.2 Effect of some insecticides and their combination against Okra shoot and fruit borer, *Earias vitella* during kharif season (2023) (second spray).**

The data of overall mean (7DAS and 14 DAS) of second spray indicated that among all the treatments lowest fruit damage was recorded in Profenofos 50 EC + Cypermethrin 25 EC (3.05%) followed by Cypermethrin 25EC + Neem oil 3% (4.29%), Spinosad 45SC + Neem Oil 3% (5.54%), Spinosad 45 SC (5.95%), cypermethrin 25 EC (7.70%), profenophos 50 EC (8.56%). The least effective treatment was Neem oil 3% (9.38%) compared to all the treatments and maximum damage of fruit is found in untreated plot (22.04%).

### **3.3 Effect of some insecticides and their combination against okra shoot and fruit borer, *Earias vitella* during kharif season (2023) (Overall mean of first and second spray).**

The results of Profenofos 50 EC + Cypermethrin 25 EC percent mean infestation of first and second spray (3.84%) are closely confirming with the finding of **Pardeshi et al. (2010)** (6.47%) was superior against *E. vittella*. Whereas **Padwal et al. (2013)** (6.8%) and **Nalini and Kumar (2016)** (8.91%) found Cypermethrin 25EC + Neem oil 3% (5.75%) next superior, followed by Spinosad 45SC + Neem Oil 3% (6.82%), **Rawat et al. (2020)** (6.70%). Spinosad 45 SC (7.67%), were next superior treatment supported by **Janu and Kumar (2022)** (12.02%) and **Rajput and Tayde (2017)** (10.46%). Cypermethrin 25 EC (9.97%) were closely related to **Neelima et al. (2021)** (10.29%) and **Manikanta and Kumar (2022)** (11.48%) followed by Profenophos 50 EC (11.88%) supported by **Patta et al. (2019)** (24.03%), **Madhuri and Kumar (2022)** (18.33%), **Kumar and Singh (2022)** (13.78%) and Neem oil 3% (13.40%) was least effective among all the treatments supported by **Naik et al. (2022)** (13.45%) and **Pachole et al. (2017)** (11.85%).



**Figure:1 Effect of some insecticides and their combination against okra shoot and fruit borer, *Earias vitella* during *kharif* season (2023) (Overall mean of first and second spray).**

**Effect on yield** –The data revealed that T<sub>7</sub> Profenofos 50 EC + Cypermethrin 25 EC, T<sub>6</sub> Cypermethrin 25EC + Neem oil 3%, T<sub>5</sub> Spinosad 45SC + Neem Oil 3% and T<sub>2</sub> Spinosad 45 SC were found significantly superior over rest of the treatments and recorded 140 quintal/hac.,137quintal/hac.,132quintal/hac and 105quintal/hac yield of healthy fruits respectively. The remaining treatments viz., T<sub>3</sub> cypermethrin 25 EC, T<sub>4</sub> profenophos 50 +EC, T<sub>1</sub> Neem oil 3%were found least effective with low yield 95quintal/hac., 86quintal/hac., 70quintal/hac. respectively.

**Cost benefit ratio**-When cost benefit ratio worked out among the treatments studied, the best and most economic treatment was found to be (T<sub>7</sub>) Profenofos 50 EC + Cypermethrin 25 EC

with C:B ratio (1:4.7) which was highest among all the treatments, the result is in closely agreement with **Pardeshi et al. (2010)**. (T<sub>6</sub>) Cypermethrin 25EC + Neem oil 3% also reported profitable yield and cost benefit ratio (1:3.6) these findings are supported by **Padwal et al. (2013)** and **Nalini and Kumar (2016)**. The cost benefit ratio of (T<sub>5</sub>) Spinosad 45SC + Neem Oil 3% (1:3.4) these results were to the findings reported by **Rawat et al. (2020)** and, the next best cost benefit ratio obtained in the treatment (T<sub>2</sub>) Spinosad 45 SC (1:3.3) was supported by **Devi et al. (2023)** and **Janu and Kumar (2022)**. The results of cost benefit ratio obtained in the treatment (T<sub>3</sub>) Cypermethrin 25 EC was (1:3.2) similar to **Naik et al. (2022)** and **Sureshsing and Tayde (2017)**. Treatments (T<sub>4</sub>) Profenophos 50 EC and (T<sub>1</sub>) Neem oil 3% showed least cost benefit ratio (1:2.9) and (1:2.0) respectively, the result is in closely agreement with **Madhuri and Kumar (2022)**, **Kumar and Singh (2022)** and **Rani and Kumar (2022)**, **Naik et al. (2022)**.

**Table 1 Effect of some insecticides and their combination against okra shoot and fruit borer, *Earias vitella* after first and second spray**

S.No.	Treatments	Dosage	Percent infestation of shoot and fruit borer								Overall mean	Yield (q/hac.)	C: B
			1 <sup>st</sup> spray (shoot infestation)				2 <sup>nd</sup> spray (fruit infestation)						
			1DBS	7DAS	14DAS	Mean	1DBS	7DAS	14DAS	Mean			
T <sub>1</sub>	Neem oil 3 %	30 ml lit <sup>-1</sup>	20.84	16.77 <sup>b</sup>	18.09 <sup>b</sup>	17.43 <sup>b</sup>	17.50 <sup>ab</sup>	8.40 <sup>b</sup>	10.37 <sup>b</sup>	9.33 <sup>b</sup>	13.40 <sup>b</sup>	75	1:2.0
T <sub>2</sub>	Spinosad 45 SC	0.5 ml lit <sup>-1</sup>	19.59	8.47 <sup>de</sup>	10.34 <sup>de</sup>	9.40 <sup>e</sup>	13.22 <sup>d</sup>	4.56 <sup>d</sup>	7.24 <sup>de</sup>	5.95 <sup>e</sup>	7.67 <sup>cd</sup>	105	1:3.3
T <sub>3</sub>	Cypermethrin 25 EC	0.5 ml lit <sup>-1</sup>	18.84	11.36 <sup>cd</sup>	13.12 <sup>cd</sup>	12.24 <sup>d</sup>	15.04 <sup>c</sup>	6.74 <sup>c</sup>	8.66 <sup>cd</sup>	7.70 <sup>d</sup>	9.97 <sup>bc</sup>	95	1:3.2
T <sub>4</sub>	Profenofos 50 EC	1.5 ml lit <sup>-1</sup>	19.38	14.21 <sup>bc</sup>	16.22 <sup>bc</sup>	15.21 <sup>c</sup>	16.24 <sup>bc</sup>	7.56 <sup>bc</sup>	9.57 <sup>bc</sup>	8.56 <sup>c</sup>	11.88 <sup>b</sup>	86	1:2.9
T <sub>5</sub>	Spinosad 45 SC +Neem oil 3 %	0.25 ml lit <sup>-1</sup> + 30 ml lit <sup>-1</sup>	19.06	7.01 <sup>e</sup>	9.21 <sup>ef</sup>	8.11 <sup>f</sup>	12.73 <sup>d</sup>	4.60 <sup>d</sup>	6.48 <sup>ef</sup>	5.54 <sup>e</sup>	6.82 <sup>cde</sup>	132	1:3.4
T <sub>6</sub>	Cypermethrin 25 EC +Neem oil 3%	0.25 ml lit <sup>-1</sup> + 30 ml lit <sup>-1</sup>	18.39	6.59 <sup>ef</sup>	7.85 <sup>ef</sup>	7.22 <sup>g</sup>	11.96 <sup>d</sup>	3.32 <sup>e</sup>	5.26 <sup>fg</sup>	4.29 <sup>f</sup>	5.75 <sup>de</sup>	137	1:3.6
T <sub>7</sub>	Profenophos50 EC + Cypermethrin 25 EC	1.5 ml lit <sup>-1</sup>	14.33	3.34 <sup>f</sup>	5.92 <sup>f</sup>	4.63 <sup>h</sup>	9.76 <sup>e</sup>	2.05 <sup>f</sup>	4.05 <sup>g</sup>	3.05 <sup>g</sup>	3.84 <sup>e</sup>	140	1:4.7
T <sub>0</sub>	Untreated check	-	15.89	26.30 <sup>a</sup>	29.02 <sup>a</sup>	27.66 <sup>a</sup>	18.76 <sup>a</sup>	21.51 <sup>a</sup>	22.58 <sup>a</sup>	22.04 <sup>a</sup>	24.85 <sup>a</sup>	42	1:1.5
F-Test			NS	S	S	S	S	S	S	S	S		
S. Ed (±)			-----	3.58	3.65	0.88	1.61	1.21	1.53	0.68	3.69		
C.D.			3.15	1.61	1.70	3.48	0.75	0.57	0.70	2.81	3.24		

DBS- Day before Spraying, DAS- Day after Spraying, NS- Non significant, S- Significant.

**4. CONCLUSION:** *Earias vittella* is a major pest of okra with high damage potential which makes its control indispensable, so research was conducted to manage *E. vittella* using various chemical insecticides, botanicals and their combinations belonging to different groups is in vogue for suppression of this pest but only partial control of pest could be achieved by using chemicals and botanicals alone. So results of this study had indicated that all the treatments are significantly superior over the untreated control, but the combinations of the insecticides had shown the highest efficacy against *E. vittella* in case of infestation as well as yield. However, among different tested insecticides combination of profenofos + cypermethrin shows best result 3.84 percent infestation reduction with 140q/hac yield and 1:4.7 cost benefit ratio followed by Cypermethrin 25EC + Neem oil 3% (5.75%) whereas the least effective treatment was Neem oil 3%(13.40%).The sequence of treatment according to their reduction in mean percent infestation found T<sub>7</sub> profenofos + cypermethrin <T<sub>6</sub> Cypermethrin 25EC + Neem oil 3% <T<sub>5</sub> Spinosad 45SC + Neem Oil 3% <T<sub>2</sub> Spinosad 45 SC <T<sub>3</sub> cypermethrin 25 EC <T<sub>4</sub> profenophos 50 EC <T<sub>1</sub> Neem oil 3% <T<sub>0</sub> untreated check. The combinations of Botanicals and Chemical insecticides are also showing good results in suppressing pest population and can be a part of integrated pest management as an effective tool under chemical control. Hence, it is suggested that the effective insecticides may be alternated in harmony with the existing Integrated pest management programs in order to avoid the problems associated with insecticidal resistance, pest resurgence etc.

#### REFERENCES

- Aman, A. S., Kumar, A., Mishra, P. K., Kumar, P. and Chowdhary, A. (2022). Integrated Approach for Management of Okra Shoot and Fruit Borer (*Earias vittella* Fab.) *The Agriculture Magazine*, 1(2): 2583-1755.
- Anonymous 2021-22 National Horticulture Board.
- Baloch, A.F., Qayyum, S. M. and Baloch, M.A. (1990) Growth and yield performance of okra (*Abelmoschus esculentus* L) *Cultivars Gomal University Journal Research*,10:191.
- Choudhury, M. A. R., Mondal, M. F., Khan, A. U., Hossain, M. S., Azad, M. O. K., Prodhan, M. D. H. and Naznin, M. T. (2021). Evaluation of biological approaches for

controlling shoot and fruit borer (*Earias vittella* F.) of okra grown in peri-urban area in Bangladesh. Horticulture, 7(1), 7.

**Fageria, M.S., Choudhary, B.R., Dhaka and R.S. (2012).** Vegetable Crops Production Technology. Text Book ISBN 978-81-272- 0702.

**Janu, R. and Kumar, A. (2022).** Field efficacy of selected insecticides against okra shoot and fruit borer (*Earias vittella* (Fab.)). *The Pharma innovation journal*; **11**(4): 1549 – 1551.

**Kaveri, G. and Kumar A. (2020)** Field efficacy of certain biopesticides against okra shoot and fruit borer, *Earias vittella* (Fabricius) on okra, *Abelmoschus esculentus* (Linn.) Moench *Journal of Entomology and Zoology Studies*; **8**(6): 1279-1281

**Kumar, R. and Singh P.P. (2022).** Field efficacy of insecticides against okra shoot and fruit borer *Earias vitella* (F.) *Indian Journal of Entomology* **84**(1): 145-148.

**Madhuri, K. and Kumar, A. (2022).** To study the field efficacy of certain chemicals and neem oil against shoot and fruit borer *Earias vittella* (Fabricius) on okra. *The Pharma Innovation Journal*; **11**(4): 1575-1578.

**Manikanta, S.E.N. and Kumar, A. (2022)** Efficacy of certain chemicals and essential oils against okra shoot and fruit borer [*Earias vittella* (Fabricius)] *The Pharma Innovation Journal*: **11**(4): 1385-1389

**Naik, V. A. K., Tayde, A. R. and Chandra, A. S. (2022).** Efficacy of Selected Chemical Insecticides against Shoot and Fruitborer [*Earias vittella* (Fabricius)] on Okra [*Abelmoschus esculentus* (L.) Moench.]. *International Journal of Plant & Soil Science* **34**(23): 656-661.

**Nalini, C. and Kumar, A. (2016).** Population Dynamics and Comparative Efficacy of Certain Chemicals and Biopesticides against Okra Shoot and Fruit Borer (*Earias Vitella*); *An International Quarterly Journal of Life Sciences.*; **11**(3):1589-1592.

**Neelima, K., Kakde, A., Thakur, S. and Venugopal, U. (2021).** Efficacy of selected botanicals against shoot and fruit borer (*Earias vittella* Fab.) on okra [*Abelmoschus esculentus* (L.) Moench]. *The Pharma Innovation Journal*; **10**(11): 2131-2134.

- Pachole, S.H., Thakur, S. and Simon, S. (2017)** Comparative bioefficacy of selected chemical insecticides and bio-rationals against shoot and fruit borer [*Earias vittella* (Fabricius)] on okra *Abelmoschus esculentus* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*. **6** (5): 1493-1495.
- Padwal, K. G. and Kumar, A. (2013)** Efficacy of plant products and combinations with cypermethrin in management of *Earias vittella* of Okra. *Annuals Plant Protection Sciences* **22** (1): 73-75.
- Pardeshi, A.M., Bharodia, R.K., Jethva, D.M., Rathod, R.T. and Patel, P.V. (2010)** Bioefficacy of chemical insecticides against of chemical insecticides against *Earias vittella* (Fabricius) on okra. *Agricultural Science Digest*, **31** (1): 25 – 29.
- Pareek, B.L., and Bhargava, M.C. (2003)** Estimation of avoidable losses in vegetable crops caused by borers under semi-arid conditions of Rajasthan. *Insect Environment*; **9**:59- 60.
- Patta, S., Sharma, A.K., Pachori, R. and Mishra, Y.K. (2019).** Bio-efficacy of newer and bio-rational insecticides against shoot and fruit borer (*Earias vittella* Fabricius) of okra. *International Journal of Chemical Studies*; **7**(5): 3402-3404.
- Rajput, G. S. and Tayde, A. (2017).** Population dynamics and comparative efficacy of certain novel insecticides, botanicals and bioagents, against shoot and fruit borer (*Earias vitella* Fabricius) of okra crop [*Abelmoschus esculentus* (L.) Moench]. *Journal of Entomology and Zoology Studies*; **5**(4): 1667-1670.
- Rani, K. and Kumar, A. (2022).** Field efficacy of different chemicals against shoot and fruit borer [*Earias vittella* (Fabricius)] of okra [*Abelmoschus esculentus* (L.) Moench]. *The Pharma Innovation Journal*; **11**(4): 1603-1607.
- Rawat, J. K., Navneet and Reddy, D.P. (2020).** To evaluate comparative efficacy of spinosad individual and in combination with botanicals against okra shoot and fruit borer (*Earias vitella* Fabricius). *Journal of Entomology and Zoology Studies*; **8**(3): 189-192.

- Sekyere, D.O., Akromah, R., Nyamah, E.Y., Brenya, E. and Yeboah, S., (2011),** Characterization of okra (*Abelmoschus spp.* L.) germplasm based on morphological characters in Ghana, *Journal of Plant Breeding and Crop Science*. **3**(13), 367-378.
- Shirale, D., Patil, M., Zehr, U. and Parimi, S. (2012).** Newer insecticides for the management of brinjal fruit and shoot borer, *Leucinodes orbonelis*. *Indian Journal of Plant Protection*, **40**(4): 273-275.
- Singh, S., Choudhary, D. P., Sharma, H. C., Mahla, R. S., Mathur, Y. S., and Ahuja, D. B. (2008).** Effect of insecticidal modules against jassid and shoot and fruit borer in okra. *Indian journal of Entomology*, **70**(3): 197-199.
- Sureshsing, M. C. and Tayde, A. R. (2017).** Efficacy of certain biorationals against shoot and fruit borer (*Leucinodes orbonalis* Guenee) of brinjal (*Solanum melongena* L.). *Journal of Pharmacognosy and Phytochemistry*; **6**(4): 1857-1859.