

Original Research Article

INFLUENCE OF VARIOUS ORGANIC MANURES ON THE GROWTH AND YIELD OF RADISH AT DIFFERENT GROWTH STAGES (*Raphanus sativus* L.)

ABSTRACT

Add brain storming sentence/problem statement A field experiment was conducted to evaluate the influence of different liquid organic manures on growth and yield of radish (*Raphanus sativus*) at organic farming block of Zonal Agricultural Research Station, V. C. Farm, Mandya during late Kharif 2020. The experiment was laid in Randomized Complete Block Design with nine treatments and three replications. The experiment consisted of nine treatments of three different levels of nitrogen was applied through FYM and different times of application of Panchagavya and Jeevamrutha. A significant variation was observed among the treatments. The 75% RDN through FYM + two time application of Panchagavya significantly increased the growth and yield attributes at 15, 30, 45 DAS and at harvest viz., plant height (8.39, 18.25, 28.16 and 33.06 cm), number of leaves per plant (6.22, 9.11, 16.04 and 18.43), leaf area index (0.08, 0.46, 0.70 and 1.01), root diameter (5.43cm at harvest), fresh shoot weight (8.53, 51.31, 115.29 and 132.43 g plant⁻¹), fresh root weight (2.99, 61.19, 161.08 and 213.08 g plant⁻¹), dry shoot weight (1.11, 10.76, 22.01 and 30.49 g plant⁻¹), dry root weight (0.90, 13.13, 26.74 and 43.28 g plant⁻¹), root yield (38.38 t ha⁻¹) and shoot yield (30.26 t ha⁻¹). In total, the results suggested that 75 per cent recommended dosage of nitrogen was applied through FYM with two times application of panchagavya is proved to be conducive for radish cultivation.

Formatted: Highlight

KEYWORDS: Panchagavya, Jeevamrutha, Growth, Yield and Radish

Formatted: Highlight

Panchagavya, Jeevamrutha, Growth, Yield and Radish

1 INTRODUCTION

Radish (*Raphanus sativus* L.) is one of important root crop of family Brassicaceae. It is a popular root vegetable in both tropical and temperate regions of world. Radish is grown for its tender root which is consumed either cooked or raw. Leaves and pods of some cultivars can be boiled and eaten as a vegetable. Radish root develops from both the primary root and the hypocotyls (Rawat and Pant.,Pant, 2021).It's a popular vegetable crop that's grown all across the globe. The major radish growing states in India are West Bengal, Haryana, Bihar, Assam, Punjab, Odisha, Chattisgarh, Madhya Pradesh, Uttar Pradesh, Karnataka and Maharashtra. In India, it was grown in an area of 205 thousand hectares during 2019-20 with an annual production of 3170 thousand metric tonnes. Radish is a good source of vitamin- A and C and also rich source of minerals like calcium, potassium and phosphorus. Radish contains glucose as the major sugar and smaller quantities of fructose and sucrose. Pink-skinned radishes are generally richer in ascorbic acid than white-skinned ones (Sadhu, 1986). It has diuretic, refreshing and cooling properties. It is also used for headache, neurological, sleeplessness and chronic diarrhoea. The roots are also useful in urinary complaints, piles, liver and gall bladder complaints (Hadley, 1993). The leaves of radish are good source for extraction of protein on a commercial scale and radish seeds are potential source of nondrying fatty oil suitable for soap making illuminating and edible purposes.

Formatted: Highlight

Comment [A1]: Too old

Comment [A2]: Too old

Growth, yield and quality of radish greatly depend on soil, climatic conditions and nutrient management. It is well documented fact that the increasing use of chemical fertilizers to increase vegetable production has been widely recognized but continuous and non judiciousnon-judicious use of inorganic inputs or commercial fertilizers have an adverse effect on soil health, environment, beneficial soil microorganisms, human being and leaving residual toxicity in the food products whereby reduces the quality of root and shoot (Mali *et al.*, 2018)-. In organic farming only organic manures or natural inputs like weeds, leaf litter and crop residues *etc.*, available on the farm are used. Thus, it reduces the cost of production against chemical inputs. Farm yard manure being a bulky organic material reduces the soil compaction and improves the aeration in addition to the supply of essential plant nutrients (Source). Apart from using conventional farm-based products there is an increasing demand for improvised materials like Panchagavya, Jeevamrutha and other liquid organic manures which mainly enrich the soil with indigenous microorganisms. These liquid formulations also contain macro nutrients, micro nutrients, many vitamins, essential amino acids, growth promoting factors like IAA, GA and thus

Formatted: Highlight

Formatted: Highlight

helps in improving the growth, yield and quality of the crops (Amalraj *et al.*, 2013).. Keeping this in view the present experiment was conducted to study the influence of various organic manures on the growth and yield of radish.

2 MATERIALS AND METHODS

A field experiment was conducted during late *Kharif* 2020 at organic farming block of Zonal Agricultural Research Station (ZARS) V. C. Farm, Mandya, University of Agricultural Sciences, Bangalore. It is located in the Southern Dry Zone of Karnataka (Agro Climatic Zone VI) and Region III at 11° 30' to 13° 05' North latitude and 76° 05' to 77° 45' East longitude with an altitude of 695 meters above mean sea level (MSL). The experiment consisted of nine treatments *viz.*, T₁ (100% RDN through FYM), T₂ (50% RDN through FYM + one time application of Jeevamrutha), T₃ (50% RDN through FYM + two time application of Jeevamrutha), T₄ (75% RDN through FYM + one time application of Jeevamrutha), T₅ (75% RDN through FYM + two time application of Jeevamrutha), T₆ (50% RDN through FYM + one time application of Panchagavya), T₇ (50% RDN through FYM + two time application of Panchagavya), T₈ (75% RDN through FYM + one time application of Panchagavya) and T₉ (75% RDN through FYM + two time application of Panchagavya). Panchagavya and Jeevamrutha were applied to soil one at the time of sowing and second application at 30 DAS at the rate of 500 L ha⁻¹. The experiment was laid out in Randomized Block Design with three replications. ~~The experiment was laid out in Randomized Block Design with three replications.~~

Initial soil

Radish variety Arka Nishanth developed at Indian Institute of Horticultural ~~Research, Research~~ Bangalore was used as a test crop. Prior to 15 days of sowing, well decomposed FYM was applied at the rate of 25 t ha⁻¹ to the respective experimental plots as per the treatments and it was incorporated with the top soil after application. Radish seeds were dibbled 2 cm down the ridges at a distance of 15 cm in the soil. Thinning was done at 15 days after sowing by rotation one seedling per hill. Seeds were sown in rows at 30 × 15 cm spacing. All the intercultural operations and plant protection measures recommended for the successful crop growth were followed and timely irrigation was given to maintain the proper moisture in the field for better growth and development of the plants.

Formatted: Highlight

Formatted: Highlight

Randomly five plants from each plot were selected to record the data on plant height (cm), number of leaves per plant, leaf length (cm), leaf area (cm²), leaf area index,), fresh root weight (g plant⁻¹), fresh shoot weight (g plant⁻¹), total fresh weight (g plant⁻¹), dry root weight (g plant⁻¹), dry shoot weight (g plant⁻¹), total dry weight (g plant⁻¹), root length (cm), root diameter (cm), root yield (t ha⁻¹), shoot yield (t ha⁻¹) and total yield (t ha⁻¹).

The obtained data were statistically analyzed as described by Gomez and Gomez (1984). The significance of variation among the treatments was observed by applying ANOVA. The level of significance used in “F” was P = 0.05. Critical difference (CD) values were calculated for the P = 0.05 whenever “F” test was found significant.

3 Results and discussion:

3.1 Results

This section was too shallow

Table should be next to result rather than after discussion.

Formatted: Highlight

There was significant effect of organic manures on all the growth and yield parameters in comparison to the control (Table 1 to 5 and Fig.1). Among the treatments, T₉ (75% RDN through FYM + two time application of Panchagavya) was highly effective for the growth and yield parameters followed by T₅ (75% RDN through FYM + two time application of Jeevamrutha). T₉ (75% RDN through FYM + two time application of Panchagavya) recorded highest growth and yield attributes at 15, 30, 45 DAS and at harvest viz., plant height (8.39, 18.25, 28.16 and 33.06 cm), number of leaves per plant (6.22, 9.11, 16.04 and 18.43), leaf length (17.24, 26.96, 33.90 and 37.23 cm), leaf area (23.91, 136.77, 209.36 and 302.38 cm²), leaf area index (0.08, 0.46, 0.70 and 1.01), fresh root weight (2.99, 61.19, 161.08 and 213.08 g plant⁻¹), fresh shoot weight (8.53, 51.31, 115.29 and 132.43 g plant⁻¹), total fresh weight (11.50, 112.50, 276.37 and 345.51 g plant⁻¹), dry root weight (0.90, 13.13, 26.74 and 43.28 g plant⁻¹), dry shoot weight (1.11, 10.76, 22.01 and 30.49 g plant⁻¹), total dry weight (2.01, 23.89, 48.75 and 73.77 g plant⁻¹), root length (27.91 cm at harvest), root diameter (5.43 cm at harvest), root yield (38.38 t ha⁻¹), shoot yield (30.26 t ha⁻¹) and total yield (68.64 t ha⁻¹). The minimum values of growth and yield attributes were observed with T₂ (50% RDN through FYM + one time application of Jeevamrutha) at all the stages of crop growth.

3.2 Discussion

Increase in growth parameters that could be obtained with application of Panchagavya and Jeevamrutha can be attributed to higher microbial load and growth hormones which might have enhanced the soil biomass thereby sustaining the availability and uptake of applied as well as native soil nutrients which ultimately resulted in better growth and yield (Boraiah *et al.*, 2017). Panchagavya contains kinetin which has a role in enhancing chlorophyll content in plant leaves thus in turn, enhance photosynthetic activity, nitrogen metabolism in plants which ultimately improving the growth parameters (Nileema *et al.*, 2011). Further it may be also due to solubilisation of nutrient in soil, absorption of nutrients and moisture due to soil application of Jeevamrutha (Yogananda *et al.*, 2015). Similar increase in the growth parameters upon application of organic manures was reported in radish by Uddain *et al.* (2010), Pathak *et al.* (2017), Mani *et al.* (2018), Subedi *et al.* (2018), Khatri *et al.* (2019), Gyewali *et al.* (2020) and Basnet *et al.* (2021).

Increase in root length and root diameter might be due to addition of organic manures helps soil micro-organisms to produce polysaccharides and thus lead to better soil structure which is useful for root growth and development (Singh *et al.*, 2016). Decrease in bulk density, better aggregation, increase in porosity and water holding capacity of the soil due to organic manures might have contributed in increasing the root length and root diameter of the plants. Similar results of root length and root diameter were also found in radish by Uddain *et al.* (2010), Kumar *et al.* (2014), Singh *et al.* (2016), Subedi *et al.* (2018), Mani *et al.* (2018) and Gyewali *et al.* (2020).

The maximum root and shoot weight could be attributed to the beneficial effect of organics in improving soil texture, aeration, reducing soil compaction and thus enhances the water and nutrient uptake by increasing the permeability of root cell membrane and stimulating the root growth or it may be attributed to solubilization of plant nutrients by addition of Panchagavya, Jeevamrutha and FYM leading to increase uptake of NPK. Fresh and dry weights of radish root and shoot results corroborate with their results obtained by Bhaktavathsalam and Geetha (2004), Sunandarani and Mallareddy (2007), Kumar *et al.* (2009), Kanaujia *et al.* (2010), Singh *et al.* (2016) and Gyewali *et al.* (2020).

The observed variation in yield of radish with Panchagavya and Jeevamrutha might be attributed to the application of organic liquid manures enhances the availability of native

nutrients to the crops and also improves the soil environment, which stimulated proliferous root system subsequent in better absorption of water and nutrients from the soil resulting in higher uptake and yield (Subramani *et al.*, 2010). The positive response of radish to organic manures with respect to root and shoot yield was evidenced by Kanaujia *et al.* (2010), Jadhav *et al.* (2014), Khalid *et al.* (2015), Ziaf *et al.* (2015), Singh *et al.* (2016) and Gyewali *et al.* (2020) in radish. **Move all result table to result section (put before discussion).**

Formatted: Highlight

Table 1: Effect of liquid organic manures on plant height and number of leaves at different growth stages of radish crop

Treatment	Plant height (cm)				Number of leaves per plant			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	7.28	15.55	20.09	23.95	4.91	7.82	9.88	12.78
T ₂	6.56	14.75	18.38	21.58	4.27	6.71	8.29	10.36
T ₃	6.66	14.81	23.23	27.68	4.29	6.83	12.88	15.03
T ₄	7.73	16.85	20.43	23.73	5.42	8.20	10.32	11.76
T ₅	7.84	16.91	25.85	30.47	5.51	8.34	14.81	17.27
T ₆	7.07	14.87	19.71	22.45	4.51	7.51	9.63	12.03
T ₇	7.11	14.99	24.69	28.40	4.64	7.66	13.63	15.80
T ₈	8.30	18.07	21.97	24.56	6.17	9.04	12.05	14.32
T ₉	8.39	18.25	28.16	33.06	6.22	9.11	16.04	18.43
S.Em±	0.33	0.71	0.96	1.14	0.51	0.35	0.51	0.62
CD@ 5%	0.98	2.13	2.88	3.42	NS	1.05	1.53	1.85

Table 2: Effect of liquid organic manures on leaf length, leaf area and leaf area index at different growth stages of radish crop

Treatment	Leaf length (cm)				Leaf area (cm ²)				Leaf area index			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	13.82	22.02	26.42	30.58	20.65	119.96	145.31	231.40	0.07	0.40	0.48	0.77
T ₂	12.47	20.75	24.91	28.81	19.41	112.93	138.30	225.79	0.06	0.38	0.46	0.75
T ₃	12.56	20.83	26.60	32.06	19.47	113.66	152.50	251.73	0.06	0.38	0.51	0.84
T ₄	15.35	24.57	27.38	30.31	21.63	125.26	156.39	236.63	0.07	0.42	0.52	0.79
T ₅	15.39	24.65	30.03	35.33	21.96	126.62	190.88	290.02	0.07	0.42	0.64	0.97
T ₆	13.17	21.26	26.18	30.07	20.06	116.99	149.03	234.30	0.07	0.39	0.50	0.78
T ₇	13.26	21.33	29.76	32.71	20.32	117.75	166.27	261.91	0.07	0.39	0.55	0.87
T ₈	17.11	26.58	29.04	32.09	23.54	136.14	157.61	257.75	0.08	0.45	0.53	0.86
T ₉	17.24	26.96	33.90	37.23	23.91	136.77	209.36	302.38	0.08	0.46	0.70	1.01

S.Em±	0.63	1.01	1.24	1.42	0.94	5.45	7.01	11.08	0.009	0.02	0.02	0.04
CD@ 5%	1.90	3.04	3.73	4.24	2.82	16.33	21.01	33.22	0.003	0.05	0.07	0.11




Fig 1: Root length and diameter of radish as influenced by application of liquid organic manures at harvest

Table 3: Effect of liquid organic manures on fresh root weight, fresh shoot weight and total fresh weight at different growth stages of radish crop

Treatment	Fresh root weight (g plant ⁻¹)				Fresh shoot weight (g plant ⁻¹)				Total fresh weight (root + shoot) (g plant ⁻¹)			
	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest	15 DAS	30 DAS	45 DAS	At harvest
T ₁	2.79	52.14	123.03	158.69	6.89	44.31	85.91	102.73	9.68	96.45	208.94	261.42
T ₂	2.62	49.36	114.92	141.58	5.06	41.58	82.96	95.39	7.68	90.94	197.88	236.97
T ₃	2.74	49.53	138.55	169.88	5.11	41.87	95.61	112.57	7.85	91.40	234.16	282.45
T ₄	2.81	57.06	126.78	154.44	7.22	48.19	88.09	103.78	10.03	104.58	214.87	258.22
T ₅	2.88	57.98	150.67	189.33	7.47	49.66	105.52	126.60	10.35	106.64	256.19	315.93
T ₆	2.63	51.20	122.42	157.08	6.37	42.15	84.08	99.14	8.99	93.35	206.50	256.22
T ₇	2.69	51.87	140.53	181.48	6.53	42.74	99.02	116.62	9.22	94.61	239.55	298.10
T ₈	2.91	60.77	133.99	166.32	8.52	50.22	90.95	108.37	11.43	110.99	224.94	274.69
T ₉	2.99	61.19	161.08	213.08	8.53	51.31	115.29	132.43	11.50	112.50	276.37	345.51
S.Em ±	0.12	3.17	5.86	7.46	0.33	1.98	4.04	4.97	0.43	4.40	9.97	12.30
CD@ 5%	NS	NS	17.58	22.38	0.98	5.95	12.10	14.91	1.28	13.20	29.90	36.89

Table 4: Effect of liquid organic manures on dry root weight, dry shoot weight and total dry weight at different growth stages of radish crop

Treat ment	Dry root weight (g plant ⁻¹)				Dry shoot weight (g plant ⁻¹)				Total dry weight (root + shoot) (g plant ⁻¹)			
	15 DAS	30 DAS	45 DAS	At harve st	15 DAS	30 DAS	45 DAS	At harve st	15 DAS	30 DAS	45 DAS	At harvest
T ₁	0.75	10.48	19.45	34.67	0.79	8.57	15.78	22.81	1.54	19.05	35.23	57.48
T ₂	0.61	9.63	19.09	33.91	0.66	8.22	14.10	20.32	1.27	17.85	33.19	54.23
T ₃	0.64	9.88	22.17	36.39	0.69	8.33	18.39	24.67	1.33	18.21	40.56	61.06
T ₄	0.83	12.25	20.67	35.18	0.74	9.29	16.80	22.08	1.87	21.54	37.47	57.26
T ₅	0.88	12.86	24.22	41.77	1.05	9.32	20.29	28.49	1.93	22.18	44.51	70.26
T ₆	0.69	10.19	21.07	34.21	0.74	8.39	16.07	22.23	1.43	18.58	37.14	56.44
T ₇	0.73	10.84	23.70	37.47	0.77	8.46	19.65	26.20	1.50	19.30	43.35	63.67
T ₈	0.86	12.74	21.82	35.84	1.06	10.37	17.78	23.73	1.92	23.11	39.60	59.57
T ₉	0.90	13.13	26.74	43.28	1.11	10.76	22.01	30.49	2.01	23.89	48.75	73.77
S.Em ±	0.08	1.04	0.96	1.62	0.24	0.40	0.77	1.07	0.07	0.89	1.73	2.69
CD @ 5%	NS	NS	2.88	4.86	0.71	1.19	2.32	3.22	0.21	2.66	5.19	8.07

Table 5: Effect of liquid organic manures on root, shoot and total yield of radish crop

Treatment	Root yield (t ha ⁻¹)	Shoot yield (t ha ⁻¹)	Total yield (t ha ⁻¹)
T ₁	27.75	22.95	50.70
T ₂	26.19	21.24	47.43
T ₃	31.58	25.32	56.90
T ₄	28.71	22.88	51.59
T ₅	36.08	27.35	63.42
T ₆	27.01	22.11	49.12
T ₇	33.16	26.46	59.62
T ₈	29.66	23.73	53.39
T ₉	38.38	30.26	68.64
S.Em±	1.34	1.08	2.42
CD@ 5%	4.02	3.25	7.27

CONCLUSION:

Formatted: Highlight

The organic manures had statistically significant variation ($p < 0.05$) in all the growth and yield parameters. The performance of T₉ was found superior in all recorded growth and yield parameters followed by T₅. Therefore, 75% RDN through FYM + two times application of Panchagavya and 75% RDN through FYM+ two times application of Jeevamrutha may be suggested to the radish growing farmers to get better results. This experiment should be conducted in a different agro - climatic zones for further validation.

REFERENCE:

- AMALRAJ, E. L. D., KUMAR, G. P., AHMED. M. H. S. K., ABDUL. R. AND KISHORE, N., 2013, Microbiological analysis of panchagavya, vermicompost and FYM and their effect on plant growth promotion of pigeon pea (*Cajanus cajana* L.) in India. *Org. Agric.*, **3**: 23–29.
- BASNET, B., ARYAL, A., NEUPANE, A., BISHAL, K. C., RAI, N. H., ADHIKARI, S., KHANAL, P. AND BASNET, M., 2021, Effect of integrated nutrient management on growth and yield of radish. *J. Agric. Nat. Res.*, **4** (2): 167-174.
- BHAKTAVATHSALAM, R. AND GEETHA, T., 2004, Macronutrient analysis of vermicompost and their effects on the growth of radish plant. *Environ. Ecol.*, **22** (4): 941-947.
- BORAIAH, B., DEVAKUMAR, N., SHUBHA, S. AND PALANNA, K. B., 2017, Effect of panchagavya, jeevamrutha and cow urine on beneficial microorganisms and yield of capsicum (*Capsicum annuum* L. var. grossum). *Int. J. Curr. Microbiol. Appl. Sci.*, **6** (9): 3226-3234.
- GOMEZ, K. A. AND GOMEZ, A. A., 1984, Statistical procedures for agricultural research. 2nd Ed. *John Wiley Sons*, New York.
- GYEWALI, B., MAHARJAN, B., RANA, G., PANDEY, R., PATHAK, R. AND POUDEL, P. R., 2020, Effect of different organic manures on growth, yield, and quality of radish (*Raphanus sativus*). *SAARC J. Agric.*, **18** (2): 101-114.

- HADLEY, P., 1993, Swede, Turnip and Radish, in Handbook of Food Science, Food Technology and Nutrition (R. Macrae, R.K. Robinson, and M.J. Sadler, eds.), Academic Press, London, p.4734.
- JADHAV, P. B., PATEL, D. J., KIREETI, A., PATIL, N. B., DEKHANE, S. S. AND HARAD, N. B., 2014, Effect of different levels of vermicompost on growth and yield of radish cv. Local. *Int. J. Inf. Res. Rev.*, **1** (2): 29-31.
- KANAUIA, S., SINGH, S. P., SINGH, V. B. AND SINGH, A. K., 2010, INM for quality production of radish (*Raphanus sativus* L.) in acid Alfisol. *J. Soils Crops.*, **20** (1): 1-9.
- KHALID, M., YADAV, B. K. AND YADAV, M. P., 2015, Studies on the effect of integrated nutrient management on growth and yield attributes of radish (*Raphanus sativus* L.). *Ann. Hort.*, **8** (1): 81-83.
- KHATRI, K. B., OJHA, R. B., PANDE, K. R., KHANAL, B. R., 2019, Effects of different sources of organic manures in growth and yield of radish (*Raphanus sativus* L.). *Int. J. Appl. Sci. Biotechnol.*, **7** (1): 39-42.
- KUMAR, M., KUMAR, S., RATTAN, P., SHARMA, J. P. AND RAI, G. K., 2009, Response of radish to the use of integrated nutrient management practices. *Veg. Sci.*, **36** (3): 406-408.
- KUMAR, S., SUTANU, M., KUMAR, S. AND SINGH, H. D., 2014, Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese White. *Int. J. Plant Sci.*, **9** (1): 57-60.
- MALI, D. L., SINGH, V., SAROLIA, D. K., TELI, S. K., CHITTORA, A. AND DHAKAR, R., 2018, Effect of organic manures and bio-fertilizers on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese white. *Int. J. Chem. Stud.*, **6** (2): 1095-1098.
- MANI, A. P. AND ANBURANI, A., 2018, Organic nutrient management technique for enhancing growth and physical parameters in radish (*Raphanus sativus* L.). *J. Phytol.*, **10**: 40-42.
- NILEEMA, S. G. AND SREENIVASA, M. N., 2011, Influence of liquid organic manures on growth, nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) in the sterilized soil. *Karnataka J. Agric. Sci.*, **24** (2): 153-157.

- PATHAK, M., TRIPATHY, P., DASH, S. K., SAHU, G. S., AND PATTANAYAK, S. K., 2017, Effect of source of nutrient on growth, yield and quality of radish (*Raphanus sativus* L.) in radish - coriander cropping sequence. *Pharma. Innov. J.*, **6** (12): 496-499.
- RAWAT, S. AND PANT, S. C., 2021, Effect of organic manures, inorganic fertilizers and their combinations on yield of radish (*Raphanus sativus* L.) cv. Japanese White. *Int. J. Recent Adv. Multidisciplinary Top.*, **2** (2): 13-15.
- SADHU, M. K., 1986, Radish crops: Vegetable Crops in India (Bose T.K., and M.G. Som, eds.), Naya Prakash, Calcutta, p. 385.
- SINGH, K. H., NASEERUDDIN AND RANA, D. K., 2016, Effect of organic manures on growth, yield and quality of radish (*Raphanus sativus* L.) cv. Pusa Desi V. *Hort. Flora Res. Spectrum.*, **5** (2): 129-133.
- SUBEDI, A., SRIVASTAVA, M. D., SHARMA AND SHAH, S. C., 2018, Effect of organic and inorganic nutrient sources on growth, yield and quality of radish (*Raphanus sativus* L.) varieties in Chitwan, Nepal. *SAARC J. Agri.*, **16** (1): 61-69.
- SUBRAMANI, A., ANUBURANI, A. AND GAYATHIRI, M., 2010, Response of growth parameters of radish (*Raphanus sativus* L.) to various organic nutrients and biostimulants. *Asian J. Hort.*, **5** (2): 464-466.
- SUNANDARANI, N. AND MALLAREDDY, K., 2007, Effect of different organic manures and inorganic fertilizers on growth, yield and quality of carrot (*Daucus carota* L.). *Karnataka J. Agric. Sci.*, **20** (3): 686-688.
- UDDAIN, J., CHOWDHURY, S. AND RAHMAN, M. J., 2010, Efficacy of different organic manures on growth and productivity of radish (*Raphanus sativus* L.). *Int. J. Agric. Environ. Biotechnol.*, **3** (2): 189-193.
- YOGANANDA, S. B., DEVAKUMAR, N., SHRUTI, M. K. AND NINGARAJU, 2015, Growth and yield of cowpea as influenced by different sources of organic manures, *Nat. Symp. Org. Agric. Sust. Food Sec. Challenges Opp.*, Tamilnadu, India, p. 113.

ZIAF, K., LATIF, U., AMJAD, M. AND SHABIR, M. Z., 2016, Combined use of microbial and synthetic amendments can improve radish (*Raphanus sativus*) yield. *J. Environ. Agric. Sci.*, **6**: 10-15.

UNDER PEER REVIEW