

# ASSESSMENT OF SOIL HEALTH AND SOIL QUALITY OF BOKARO, DISTRICT, JHARKHAND, INDIA

## ABSTRACT

Taking care of soil health is most importance for sustainable crop production. soil quality is an blending of soil processes and provides to a measure change in soil condition as related to following factors such as land use, climate patterns, cropping sequences and farming systems. In the present investigation analysis of different physical and chemical characteristics were made to determine soil health of Bokaro district. Results show that soil bulk density was in normal range, soil pore space % varied from 39.85 to 48.32 g cm<sup>-3</sup>, particle density ranged from 2.24 to 2.69 Mg m<sup>-3</sup>, water holding capacity varied from 34.23 to 43.86 %. soil pH and EC were found to be acidic and non-saline in nature respectively while soil organic carbon and available phosphorus were in medium to high range. Available nitrogen and potassium were found to be low to medium range i.e., 197.0 to 219.0 Kg ha<sup>-1</sup> and 201.96 to 266.01 Kg ha<sup>-1</sup> and the range of phosphorus varied from 16.96 to 26.68 kg ha<sup>-1</sup>. respectively. Deficiency of sulphur followed by zine was found in soils of Bokaro district.

**Keywords:** Soil health, soil quality, pH, EC, sulphur, zinc, soil

## INTRODUCTION

Soil is the biologically energetic, porous medium that has been developed in the uppermost layer of Earth's crust. Soil is one of the critical concepts that substrata of life on the planet, It serving as a reservoir of water and nutrients, as a medium for the filtration and breakdown of dangerous wastes, and as a participant in the biking of carbon and different factors thru the worldwide atmosphere. it's been envolved through weathering techniques driven through organic, climatic, geologic, and topographic impacts.

Soil heterogeneity is the case where soil in a exceedingly small vicinity varies significantly in texture, fertility, topography, moisture content material, drainage etc. Soil heterogeneity is the motives for the numerous nature of cropping and production sample. Soil consists of a solid segment (minerals and organic count) in addition to a porous phase that holds gases and water (Latha and Janaki). Soils provide nutrients, water and area for trees, herbs and different plant life and hence are the basis for existence on the planet. As soil formation is predominantly a vertical system related to climate, parent material, comfort, organisms, time and spatial function and as weathering and humus input is most powerful at the floor and reduces with depth, most soil residences are depth-dependent. The deeper the soil layers are positioned, the decrease is the impact of plant cowl and soil biota and the higher is the impact of weathering, main to specific

depth capabilities for soil houses. for example, soil organic carbon (SOC) decreases with growing soil intensity as natural material enters the soil predominantly from the top after which eventually is incorporated into deeper soil layers.

Soils vary depplyas a function of their position on of the landscape and agricultural management, land use, and overto cultivation intensity. (Ayele *et al.*,2020)

India is the one of the second most populous united states in the global subsequent to China, overall, wherein a huge part of the populace is the based upon the agriculture. There are the stay stock population of India also constitutes 1/6th one of the international populations. To satisfy the food call for of growing human population and forage demand of massive farm animals' populace, herbal forests are not best converted to pasture and the rural fields, however agricultural fields are the additionally put under the excessive strain because of the cOnventional technique of agricultural practices which in the long run results in the deterioration of soil fitness in converted ecosystems. Indian soils not handiest show deficiency of number one nutrients like (Nitrogen, Phosphorous and Potassium) but additionally rest of secondary nutrients (Sulphur, Calcium and Magnesium) and micro nutrients (Boron, Zinc, Copper and Iron and many others.) in maximum important parts of the United States. Instated of these three number one nutrients (N, P, k), deficiency of Sulphur and micro nutrients like Zinc and Boron in many of States, and of Iron, Manganese and Molybdenum in a few States, has been grow to be a restricting component in growing meals productivity.

Jharkhand is the kingdom which is positioned within the eastern part of India. It turned into carved out of the southern member Bihar on 15 November 2000. Jharkhand stocks in its border with the states of Bihar to the north, Uttar Pradesh and Chhattisgarh to the west, Odisha to the south, and West Bengal to the east. Thereby of Ranchi is the country capital and Dumka is the subcapital, even as Jamshedpur is the largest and the biggest commercial town of the country. The nation has been a place of about 79,000 km<sup>2</sup>, one of the smaller states inside the India in phrases of location, and consists of numerous administrative units.

## **MATERIALS AND METHODS**

Bokaro district is lies within the eastern part of Jharkhand nation. It is bounded by the district of Giridih within the north, West Bengal in the south, Dhanbad in the east and Hazaribag in the west. It has a place of 2861 sq. km area and populace of 17,seventy five,961 humans (Census of India, 2001). The district comprises subdivision (Chas and Bermo) and 8 development blocks viz.Chas, Bermo, Gomia, Chandankyari, Jaridih, Kasmar, Peterwar and Nawadih.

Bokaro district is located at Eastern part of Jharkhand state. The climate of the area is moderate to extreme and characterized by hot summer and cold winter. Rainfall in this area is also very moderate... The major cultivated crops in this district are paddy, wheat, maize, pulses, fruits, vegetables etc.

The district has climatic condition slightly distinctive due to less elevation and less forest cover. The summers are hot and winters are bloodless. the highest temperature is recorded as 46degreesCelsius. in wintry weather season temperature come down to three to 4 diploma Celsius. whilst bloodless wind blows, otherwise the temperature tiers among 10 to 15 diploma Celsius. The common rainfall is 1570 mm. Soil samples was collected only from open spaces from depths of 0-15 cm, 15-30 cm and 30-45 cm from different Blocks of Bokaro district, Jharkhand.

### **3. RESULTS AND DISCUSSION**

#### **3.1 Physical Properties of Soil**

##### **3.1.1 Bulk density**

The Table 1 depicted that the maximum the maximum mean bulk density found at (V7) Utasara (1.62 g cm<sup>-3</sup>)and the (V7) was significant higher then (V4) Kamalapur (1.59 g cm<sup>-3</sup>), (V8) Darid (1.55 g cm<sup>-3</sup>), (V6) Gomanjara (1.46 g cm<sup>-3</sup>), (V5) Baraikalan (1.44 g cm<sup>-3</sup>), (V1) Bandhdih (1.41 g cm<sup>-3</sup>). The bulk density decrease with increase in soil depth due to low organic matter and less aggregation. Decrease in bulk density was an indicator for improvement in soil physical properties and congenial environment for crop growth. And the similar finding was reported by **Kumar *et al.* (2010)**

##### **3.1.2 Percent Pore Space**

The Table 1 depicted that the maximum The maximum mean pore space (%) found at (V2) Baradih (46.3 %)and the (V1) Bandhdih (45.8 %)was significant higher then (V3) Bharu(44.9 %), (V6) Gomanjara (44.3%), (V5) Baraikalan (43.8 %), (V7) Utasara (42.4%) then (V9) Keswari (41.6 %), (V8) Darid (41.3%). Decrease in pore space is attributed Increase in compaction in the soil. And the similar finding was reported by **Kumar *et al.* (2010)**


##### **3.1.3 Water holding capacity**

The Table 1 depicted that the maximum Mean Water Holding Capacity (%) found at (V2) Baradih (42.0 %)and the (V1) Bandhdih (41.0%)was significant higher then (V3) Bharu(40.8 %), (V6) Gomanjara


(40.02%),(V5) Baraikalan (39.2 %),(V8) Darid (37.2%) then (V7) Utasara (37.5 %),(V9) Keswari (37.0%). Decrease in pore space is attributed Increase in compaction in the soil. And the similar finding was reported by **Kumar *et al.* (2010)**

**Table 1. Bulk density (Mg m-3), particle density (Mg m-3), pore Space (%) and water holding capacity (%) of soil at different depth**

S.No.	Soil bulk density			Soil partical density			Pore Space (%)			Soil waterholding capacity		
	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 Cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
V <sub>1</sub>	1.40	1.41	1.42	2.27	2.29	2.32	47.42	45.84	44.28	43.86	41.56	40.18
V <sub>2</sub>	1.36	1.41	1.48	2.35	2.36	2.39	48.32	46.75	43.86	44.28	42.39	39.36
V <sub>3</sub>	1.38	1.41	1.44	2.24	2.26	2.30	47.98	44.54	42.26	43.56	40.12	38.95
V <sub>4</sub>	1.56	1.61	1.62	2.65	2.67	2.69	44.35	42.64	41.32	40.95	39.25	38.86
V <sub>5</sub>	1.42	1.44	1.46	2.56	2.57	2.57	45.52	43.86	41.86	40.28	39.86	37.62
V <sub>6</sub>	1.44	1.46	1.50	2.54	2.56	2.61	46.52	44.32	42.16	41.86	40.18	38.78
V <sub>7</sub>	1.57	1.64	1.67	2.51	2.52	2.55	44.32	42.89	40.28	39.62	38.76	34.23
V <sub>8</sub>	1.53	1.54	1.59	2.42	2.44	2.47	42.25	41.89	39.96	38.76	37.56	35.42
V <sub>9</sub>	1.52	1.56	1.58	2.52	2.53	2.53	43.76	41.38	39.85	38.26	37.89	34.89
<b>F-test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S.Em. (±)</b>	0.0099	0.0261	0.0304	0.024	0.035	0.038	0.6365	0.6845	0.6409	0.4950	0.5149	0.0429
<b>C.D@5%</b>	0.0294	0.0778	0.0506	0.073	0.106	0.113	1.8911	2.0338	1.9044	1.4707	1.5298	0.1276



**Fig1: BulkDensity ( $\text{Mg m}^{-3}$ ) and Partical Density( $\text{Mg m}^{-3}$ )**



**Fig2: Pore Space(%) and Water Holding Capacity(%)**

## 3.2 Chemical Properties

### 3.2.1 Soil pH

The Table 2 depicted that the maximum mean soil pH (V9 and V2) of 6.91 was found at Village (Keswari and Baradih) and the minimum pH was found at (V3) 6.47 (Bharu), (V6) 6.34 (Gomanjara), (V5) 5.94 (Barikalan), (V7) 5.74 (Utasara), (V8) 5.68 (Darid), (V1) 5.52 (Bandhidh), (V4) 5.19 (Kamalapur). Hence, the soil acidity minimum and a major crop production constraint in the study area. And the similar finding were reported by **Kumar et al. (2010)**.

### 3.2.2 EC (dS m<sup>-1</sup>)

The Table 2 depicted that the highest mean EC was in (V6)- Gomanjara, (0.06 ds m<sup>-1</sup>) and lowest mean (V1)-Bandhidh (0.02 ds m<sup>-1</sup>) from Hence, all the soils under the study area are safe for all types of crop production with respect to the soluble salt content. And the similar finding was reported by **Kumar et al. (2010)**

### 3.2.3 Organic carbon

Table 2 shows the maximum value of organic carbon mean was in (V6) 0.45% in village (Gomanjara), then in (V5) 0.43% (Barikalan), (V3) 0.41% (Bharu), and (V1 and V2) show 0.40% of village (Bandhidh to Bharu), where the (V2) show 0.39% (Baradih), (V7) 0.37% (Utasara) and (V8-V9) has 0.36% (Darid-Keswari). Soil organic carbon status was found to be medium to high which enables the soil for higher crop production. Medium to higher organic carbon in the study area could be attributed to its presence in the bottom of dense. And the similar finding was reported by **Kumar et al. (2010)**

### 3.2.4 Available nitrogen

The Table 3 depicts that the maximum value of nitrogen is found in soil and the mean value was in (V4)-Kamalapur (213.00 kg ha<sup>-1</sup>), (V2)-Baradih (212.00 kg ha<sup>-1</sup>), (V6)-Gomanjara (211.00 kg ha<sup>-1</sup>), (V3)-Baradih (210.00 kg ha<sup>-1</sup>), (V1-V9)-Bandhidh and Keswari show (209.00 kg ha<sup>-1</sup>), (V5)-Barikalan (208.00 kg ha<sup>-1</sup>), (V7)-Utasara (203.00 kg ha<sup>-1</sup>), (V8)-Darid (200.00 kg ha<sup>-1</sup>). The available nitrogen status in the entire study area was found to be low to medium. The reason may be attributed to the fact that nitrogen content is positively correlated with organic matter content which decreases with depth. And the similar finding was reported by **Kumar et al. (2010)**

### 3.2.5 Available Phosphorus

The Table 3 depicts that the maximum mean was shown in village (V5)-Barikalan (22.99 Kg ha<sup>-1</sup>), then after that (V3)-Bharu show minimum range then other (V5) which was (21.94 Kg ha<sup>-1</sup>), (V3)-Bharu (21.94 kg ha<sup>-1</sup>), (V2) -baradih (21.36 kg ha<sup>-1</sup>), (V1)- Bandhdih (21.35 kg ha<sup>-1</sup>), (V7)-Utasara (20.96 kg ha<sup>-1</sup>), (V8) -Darid (20.13 kg ha<sup>-1</sup>), (V6)- Gomanjara (20.01 kg ha<sup>-1</sup>), (V9)-Keswari (19.03 kg ha<sup>-1</sup>). The available phosphorus status in the entire study area was found to be medium to high. And the similar finding was reported by **Kumaret al. (2010)**

### 3.2.6 Available potassium

The Table 3 depicts the highest range available potassium was found at (V5) -Baraikalan (250.43 kg ha<sup>-1</sup>) and the lowest mean available potassium was found at (V7)-Utasara (211.81 kg ha<sup>-1</sup>). The available potassium status in the entire study area was found to be low to medium. The highest K content observed in the surface horizon and showed more or less decreasing trend with depth. This might be attributed to more intense weathering and release of liable K from organic residues. And the similar finding was reported by **Kumar et al. (2010)**

### 3.2.7 Available Sulphur

The Table 4 depicts that the highest mean available sulphur was in (V3 to V6)-Bharu and Gomanjara (2.59 ppm) and lowest mean available sulphur was found at (V2)-Baradih (0.20 ppm). High deficiency of S in Bokaro soils might be attributed to the upland (Alfisols) leading to leaching of anionic nutrients. Available sulphur was found to be in the range of low to medium and the similar finding was seen by **Kumar et al. (2010)**

### 3.2.7 Available Zinc

The Table 4 depicts that the highest mean available zinc was found at (V6) (0.59 ppm)-Gomanjara and lowest mean (V2 and V7) (0.33 ppm)-Baradih and Utasara. Zn deficiency was observed in all blocks of the Bokaro district. And the similar finding was reported by **Kumar et al. (2010)**

Table 2. pH(w/v), EC (dS m<sup>-1</sup>), organic carbon (%) of soil at different depth

S.No	Soil pH	EC (dS m <sup>-1</sup> )	organic Carbon (%)
------	---------	--------------------------	--------------------

	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
V <sub>1</sub>	5.02	5.60	5.95	0.02	0.02	0.02	0.42	0.41	0.39
V <sub>2</sub>	6.86	6.90	6.99	0.03	0.04	0.05	0.41	0.39	0.38
V <sub>3</sub>	6.18	6.28	6.97	0.04	0.05	0.06	0.43	0.41	0.40
V <sub>4</sub>	4.66	5.01	5.91	0.02	0.04	0.06	0.42	0.41	0.39
V <sub>5</sub>	5.87	5.92	6.03	0.02	0.03	0.04	0.45	0.43	0.42
V <sub>6</sub>	6.97	6.01	6.05	0.05	0.05	0.06	0.47	0.45	0.43
V <sub>7</sub>	5.70	5.14	6.40	0.03	0.03	0.04	0.39	0.37	0.36
V <sub>8</sub>	5.94	6.07	6.07	0.03	0.05	0.06	0.38	0.37	0.35
V <sub>9</sub>	6.85	6.98	6.98	0.03	0.06	0.09	0.37	0.36	0.35
F-test	S	S	S	NS	NS	NS	S	S	S
S.Em. (±)	0.5378	0.5668	0.6618	0.008	0.006	0.009	0.03651	0.036514	0.0365
C.D. @ 5 %	0.1154	0.1354	0.2154	.....	.....	.....		6.9376	6.2073


Table 3. Available Nitrogen (Kg ha<sup>-1</sup>), available phosphorous (Kg ha<sup>-1</sup>), available potassium (Kg ha<sup>-1</sup>) of soil at different depth

S.no	Available Nitrogen (Kg ha <sup>-1</sup> )			Available phosphorus (Kg ha <sup>-1</sup> )			Available potassium (Kg ha <sup>-1</sup> )		
	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm	0-15cm	15-30cm	30-45cm
V1	215.00	209.00	203.00	24.36	20.75	18.96	224.85	212.18	203.56
V2	218.00	213.00	205.00	23.68	21.12	19.28	246.55	231.93	222.36
V3	217.00	209.00	204.00	25.12	22.85	17.85	253.56	242.85	233.72
V4	220.00	213.00	207.00	23.42	21.28	18.54	244.75	236.38	229.63
V5	215.00	208.00	201.00	26.68	22.42	19.87	266.01	249.91	235.37
V6	219.00	212.00	204.00	22.68	19.92	17.45	258.89	246.22	232.65
V7	209.00	203.00	199.00	23.86	20.17	18.86	220.96	211.65	202.84
V8	205.00	200.00	197.00	22.37	20.67	17.37	230.34	220.75	204.12
V9	216.00	210.00	203.00	21.52	18.62	16.96	219.79	205.62	201.96
F-test	S	S	S	S	S	S	S	S	S
S.Em. (±)	2.6239	3.2501	0.4487	0.347482	0.37250	4.2394	3.2857	3.0338	0.2376
C.D. @ 5 %	7.7962	2.7816	1.3348	1.03242	1.10678	1.5961	9.7623	9.0138	0.2376


Table.4 Sulphur(ppm) and Zinc(ppm) of soil of different depth

S. NO	Sulphur(ppm)	Zinc (ppm)
-------	--------------	------------

	0-15cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45cm
V1	3.70	2.77	1.84	0.56	0.37	0.18
V2	0.37	0.18	0.06	0.58	0.33	0.08
V3	2.77	2.59	2.41	0.35	0.18	0.01
V4	3.70	2.22	0.74	0.85	0.40	0.05
V5	2.96	2.40	1.84	0.59	0.37	0.15
V6	2.75	2.59	2.43	0.77	0.59	0.41
V7	3.70	2.59	1.48	0.59	0.33	0.07
V8	2.96	2.40	1.84	0.96	0.40	0.16
V9	3.34	2.22	1.10	0.85	0.57	0.29
F-test	S	S	S	S	S	S
S.Em. (±)	0.0377	0.0447	0.0365	0.0115	0.006	0.032
C.D. @ 5 %	0.0215	0.0055	0.00233	0.0343	0.0186	0.0955




**Fig3:pH**



**Fig4:EC ( $\text{ds m}^{-1}$ )and OC (%)**

UNDER PEER REVIEW



**Fig5: Available Nitrogen, Phosphorous and Potassium (Kg ha<sup>-1</sup>)**



**Fig6: sulphur(ppm) and zinc(ppm)**

## CONCLUSION

It can be concluded that the soils of Bokaro District of Jharkhand state are in good physical and chemical condition. The major crop production constraint in the study area was found to be soil slightly acidity. Due to soil erosion and water logging, most of the soils were found to be deficient in available nitrogen, potassium, zinc and sulphur content. So, application of liming materials along with application of soil test based nitrogenous, potassic, zinc and sulphur containing fertilizers and manures will help to obtain higher crop production as well as sustaining soil health. Hence, it is concluded that to obtain better soil health and soil quality a more attention is needed to pay in the role of macronutrient enhancement in the soil which will lead to optimum economic Yield. And for maintaining soil health and soil quality adoption of organic farming will be an essential step.

## REFERENCES

**A.Kouzani., A. Kaynak., S. Yang Khoo, (2018)** Soil Bulk Density Estimation Methods: A Review. *Pedosphere* 28(4):581-596 DOI:10.1016/S1002-0160(18)60034-7

**A. Balasubramanian,(2017)** CHARACTERISTICS OF SOIL PROFILE, publication/314497793

**Arvind., K.Manas., Denre., R. Prasad, B.K.Agarwal., D.K. Shahi and A.K,(2017).** Benefits of Zinc Fertilization in Hybrid Rice SarkarDepartment of Soil Science and Agricultural Chemistry, Birsa Agricultural University, Ranchi, Jharkhand.

**Ashokkumar., V. Rajani., Behra, R., Hartmann., N.B., Files.(2019)** Soil Quality and Soil Health Department of Agricultural Chemistry & Soil Science College of AgricultureJunagadh Agricultural University Junagadh, Gujarat.RG.2.2.24060.08321

**S.S.Andrews., D.L. Karlen, and C.A. Camberdella, (2004).** Soil Management Assessment Framework: A Quantitative Soil Evaluation Method. *Soil Science Society of America Journal* 68: 1945–1962

**B.N. Ghosh, K. Das., A.K. Sahoo., S. Mukhopadhyay., A. Haldar., D.C. Nayak and S.K. Singh (2019)** Developing land use plan through advanced land resource inventory (LRI) technique in Chhotanagpur plateau region of Jharkhand. ICAR-National Bureau of Soil Survey and Land Use Planning (NBSS&LUP), Regional Centre, Kolkata-700091, West Bengal; ICAR NBSS&LUP, Nagpur-440033, Maharashtra

**Brajendra, Latha PC, Surekha K, Prasad Babu(2014).**, MBB, Gosh TJ, Bhadana VB. Soil Health Testing Manual. Parmar Publishers and Distributors, India.

**FAI (2003 – 2004)** Fertilizer and agriculture statistics, Eastern Region.

**Filip, Z. (2002).** International approach to assessing soil quality by ecologically-related biological parameters. Agriculture, Ecosystems and Environment 88: 169-174.

**Ghosh, B.N., Das, K., Bandyopadhyay, S., Mukhopadhyay, S., Nayak, D.C. and Singh, S.K. (2018)** Impact Assessment of GIS Based Land Resource Inventory Towards Optimizing Agricultural Land Use Plan in Dandakaranya and Eastern ghats Physiographic Confluence of India. J. Indian Soc. Remote Sens., doi.org/10.1007/s12524-017- 0743-1.

**Jackson, M.L. (1958)** Soil Chemical Analysis. pp. 221. niordc.ir/uploads\86\_106\_Binder2.pdf · Soil...

**Koralage. I. S. A., P. Weerasinghe. , N. R. N. Silva and C. S. De Silva(2015)** Phosphorous Assessment\_of\_phosphorus\_availability\_in\_soil/attachment/5ff27a95e35e2b000102cc4d/AS%3A976078315196416%401609726613120/download/The\_Determination\_of\_Available\_Phosphorus\_in\_Soil\_.pdf

**K. Panday., P. K. Thakur., Z. Perween., N. Kumari and R. K. Minz.,(2019)** Status of Soil Nutrients and Fertility of Forest and Nearby Agricultural Land in Bokaro District of Jharkhand) . International Journal of Chemical Studies 2019; SP6: 495-499 E-ISSN: 2321–4902 IJCS 2019; SP6: 495-499

**Kumar., R. Dr. A. T. Jeyaseelan., Shri Ravish Kumar., Dr. P. G. Shanware., Shri K. D. Sood.,(2010)** REPORT ON SOIL RESOURCES OF BOKARO DISTRICT, NORTH CHHOTA NAGPUR COMMISSIONAIRE, JHARKHAND STATE JSAC/TECH-REP/DoIT-GoJ/NRIS/10-11/.

**Key to soil Taxonomy Ninth Edition, 2003**, United States Department of Agriculture (USDA)

**Moebius-Clune, B.N., D.J. Moebius-Clune, B.K. Gugino, O.J. Idowu, R.R. Schindelbeck, A.J. Nortcliff, S. (2002)**. Standardisation of soil quality attributes. Agriculture, Ecosystems and Environment 88: 161-168.

**P.Bastia,,Rakesh Awale,,Stephen Machado,,Rajan Ghimire,, Prakriti Bista (2017)**.soil Health publication/317648217\_Soil.

**Ravinder .S., Tapasya Tiwari Krishna Kumar Singh,.(2023)** Importance of Soil Testing in Sustainable Agriculture Emerging Trends in Agricultural Sciences (pp.173-185)Publisher: P.K. Publishers & Distributors

**Ravinder, K. and Priyanka. (2022)** Viewpoint Soil Quality and Soil Health (e-ISSN: 2582-8223).

**Ramamoorthy. B., Bajaj JC. Soil Fertility Map of India, Fertile News, 1969; 14(8):25.**

**Rakesh Kumar,,kishan .s. Rawat and Brijesh(2012)** Vertical Distribution of Physico-Chemical Properties under Different Topo-sequence in Soils of Jharkhand 1 Department of Soil Science and Agricultural Chemistry, Birsa Agricultural University, Ranchi, 834006, Jharkhand 2 Division of Agricultural Physics, Indian Agricultural Research Institute, New Delhi - 110 012.

**Rakesh, A., Stephen Machado,, Rajan, Ghimire,, Prakrit Bista.(2017)** publication/317648217-Soil health.

**Ristow, H.M. Van Es, J.E. H.A. Shayler, M.B. McBride, D.W. Wolfe, and G.S. Abawi. 2016.** Comprehensive Assessment of Soil Health e Cornell Framework Manual, Edition 3.1. Cornell University: Geneva, NY. <http://www.css.cornell.edu/extension/soil-health/manual.pdf>

**Statistical Handbook of Jharkhand. 2015.** Directorate of Economics and Statistics, GoI, New Delhi

**S. Pinho., Neha, Shreya, Bruno Valentim , Biswajit Paul , Alexandra Guedes, Joana Ribeiro Colin R. Ward., Deolinda Flores(2015)** Multi-technique study of fly ash from the Bokaro and Jharia coalfields (Jharkhand state, India): A contribution to its use as a geoliner). International Journal of Coal Geology, Volume 152, Part B

**Sarkar, D. 2002.** Soil resource information for meeting challenges of land th degradation: A case study in Chotanagpur plateau region. The 4 Dr. R.S. Randhawa Memorial Lecture. J. Indian Soc. Soil Sci.,50(4): 414-437.

**Sarkar, D., Gangopadhyay, S.K. and Velayudhan, M. 2001.** Soil topo sequence relationship and classification in lower outlier of Chhotanagpur plateau. Agropedo., 11: 29-36.

**Sarkar, D., Gangopadhyay, S.K. and Velayudhan, M. 2001.** Soil topo sequence relationship and classification in lower outlier of Chhotanagpur plateau. Agropedo., 11: 29-36.

**Mwendwa.S.1927,** A modication of the Bouyoucos method of soil texture analysis. [https://www.researchgate.net/publication/344561274\\_The\\_Mwendwa\\_Protocol\\_A\\_modification\\_of\\_the\\_Bouyoucos\\_method\\_of\\_soil\\_texture\\_analysis/fulltext/5f80475192851c14bcb93908/The-Mwendwa-Protocol-A-modification-of-the-Bouyoucos-method-of-soil-texture.pdf](https://www.researchgate.net/publication/344561274_The_Mwendwa_Protocol_A_modification_of_the_Bouyoucos_method_of_soil_texture_analysis/fulltext/5f80475192851c14bcb93908/The-Mwendwa-Protocol-A-modification-of-the-Bouyoucos-method-of-soil-texture.pdf)

**OSU. 2009.** Willamette Valley Soil Quality Card Guide EM 8710-E. Oregon State University Extension Service. <https://catalog.extension.oregonstate.edu/em8710>

**Walkley, A. and Black, (1947)** An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. Soil Sci., 37:29-38

**USDA-NRCS. 2001.** Soil Quality Test Kit Guide. USDA-ARS-NRCS Soil Quality Initiative. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_050956.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_050956.pdf)

**USDA-NRCS. 2004. Palouse and Nezperce Prairies Soil Quality Card Guide.**  
[http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/health/assessment/?cid=nrcs142p2\\_05387](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/health/assessment/?cid=nrcs142p2_05387)

UNDER PEER REVIEW