

# A Case report of massive abdomen hemoperitoneum due to uterine fibroid rupture in nulliparous women

## Abstract:

Uterine leiomyoma is benign mass found more commonly in young women at reproductive age and are rarely associated with hemoperitoneum. The direct cause of intraperitoneal bleeding due to fibroids is poorly understood. We report a case of 38 years old women, nulliparous, presented to emergency room with significant abdominal distention/pain and difficulty in breathing. The patient had no obvious history of any medical comorbidities. Incidental findings of multiple large uterine fibroids largest 10x9 cm noted. Patient was hypotensive and tachycardia, underwent emergency laparotomy, intra operative finding of massive intraperitoneal haemorrhage due to rupture of vessels on the surface of a subserous uterine fibroid. Rupture of leiomyoma surface vessel should be considered as one of differential diagnosis, when dealing with a case of intraperitoneal bleeding.

## Keywords:

Uterine fibroid, leiomyomas, intra-abdominal bleeding, massive, hemoperitoneum, rupture, avulsion

## Introduction:

Uterine fibroids grow in the uterus. They are muscular benign tumours that are rarely turn to cancer. They are hypersensitive to oestrogen hormone, avascular, getting its blood supply from the pseudo-capsules of myometrium tissue, and usually surrounded by an abnormal venous plexus. (1) Uterine fibroids are commonly found in 25% of women of reproductive age (1) however, ruptured fibroids are rare and can lead to life-threatening condition due to acute peritoneal bleeding requiring immediate surgical intervention. Rupture Fibroids are more likely to occur on subserous fibroids that have small stalks inside or outside than those intramural or submucous fibroids. As rupture fibroid are rare condition, other common differential diagnosis should be considered in case scenarios of sudden severe pain with shock and hemodynamically un stability, gastrointestinal causes and vascular disorders should be ruled out [3]. Despite advanced medical imaging modalities, the correct diagnosis of rupture fibroid is rarely identified before surgery. (1)

## Case report:

We report a case of 38 years old women, nulliparous, presented to the Nizwa Hospital – Sultanate of Oman–gynaecology emergency room with significant abdominal distention/pain and difficulty in breathing. The patient had no obvious history of any medical comorbidities. Incidental findings of multiple large uterine fibroids largest 10x9 cm noted. On presentation, the patient was hypotensive (blood pressure, 80/40 mm Hg) and tachycardia. The patient had no history of trauma. She had a regular menstrual cycle and she was on day three of her cycle.

Examination revealed a young woman with an average body built, hypotensive, tachycardic, distended abdomen, and tender. Her blood haemoglobin level was 6 g/dl. Quantitative Bhcg was negative. Transabdominal ultrasonography revealed a uterus with thin endometrium, multiple fibroids with distorted anatomy, and significant free fluid in the pouch of Douglas and hepatorenal pouch. The ovaries and adnexa were not visualized because they were obscured by the enlarged, bulky uterus. Emergency abdomen and pelvic CT scan was performed which confirmed moderate abdominopelvic dense free fluid indicating hemoperitoneum, uterus with multiple fibroids some showing central hypodensity, the largest measuring 10x8 cm, small hematoma 4.5x2.6 cm in the pelvis with right ovarian vein suggestive of active bleeding, another large heterogeneous hematoma seen in left lumbar region measuring 11.5x8.5x7.1 cm, ovaries can't be appreciated. ( figure 1 ,2 ) The patient was counselled about the diagnosis of uterine fibroid , the need for emergency laparotomy and written consent obtained.



Figure 1: contrast enhanced coronal reformatted abdomen CT scan showing enlarged uterus with multiple heterogenous fibroids, moderate hemoperitoneum and left lumbar hematoma



Figure 2: contrast enhanced axial pelvis CT scan showing enlarged distorted uterus with multiple heterogenous fibroids, hemoperitoneum and right iliac fossa dense hematoma.

Patient underwent exploratory laparotomy, intraoperatively hemo-peritoneum of 3.5 liters noted, the uterus with multiple numbers of sub-serosal fibroids largest was approximately 10x8 cm at the right side with active bleeding from the vessels seen, cauterized and fibrillar applied with some homeostatic sutures, bleeding was controlled. Both the ovaries and fallopian tubes were normal. The patient received six units of packed red blood cell, four-units of Fresh frozen plasma, and four units of platelets. The patient had an uneventful recovery and was discharged home on day three post operation in a stable condition. Poste

operative patient received GNRH injection monthly for 3 months, advised for laparotomy myomectomy, but she refused surgical intervention as she is nulliparous and worry about the risk of hysterectomy. She opted for uterine artery embolization.

#### Discussion:

Systematic review of cases from 1902 to 2018 reported that intra-abdominal bleeding was commonly due to the rupture of superficial blood vessels over the surface of the fibroid, followed by rupture and avulsion of the fibroid. (1) A presenting complaint was usually of severe abdominal pain with sudden and profound hypovolemic shock. Correct preoperative diagnosis was made in only seven cases on computed tomography imaging. Most commonly performed procedures in such cases were hysterectomy and myomectomy. (1)(11) Mortality was reported in 4 cases which were directly related to complications of uterine fibroids. (1) This systematic review showed that all patients were managed surgically, with no cases managed medically alone or conservatively. (1) Indication for surgery was mainly massive intra-abdominal bleeding with hemodynamic instability (1). The most common surgical approach, was Laparotomy, while laparoscopy successfully performed in only two cases reported and two patients underwent preoperative uterine arterial embolization. (1) Hysterectomy was performed in 58.4% cases, with myomectomy in 36% of cases and ligation of haemorrhagic vessels in 3.2%. (1)

Over half of the identified patients the source of bleeding was a ruptured superficial vessel overlying uterine fibroid. (1)(3) Theory suggest that increased blood flow or mechanical trauma would indirectly cause over- distention and tearing, leading to the onset of bleeding. (5)(6) Venous congestion during menstruation, uterine manipulation (e.g., vaginal examination), increased venous pressure associated with defecation or lifting of heavy objects, and pregnancy all can lead to over distention and tearing of vessels. A few authors have hypothesized that the posterior location of these fibroids predisposes patients to direct-contact injury from the promontory of the sacrum. (1)(3) Trauma such as falls or motor vehicle accidents, has been also linked to clinical presentation. However, these factors were only identified in 26% of the cases that has been reviewed, with 14% related to pregnancy or postpartum period. (1)(3)

Degenerative and sarcomatous changes occur in fibroids, leading to necrosis and perforation can be of the causes of spontaneous rupture of uterine fibroids vessels. Degenerative changes in fibroids are usually due to inadequate blood supply (3) although are present in nearly all tumours, with no significant relationship with the presenting symptoms. (7&8) Degenerative changes were observed in 35% of patients with ruptured fibroids. Environmental factors and hormonal therapy suggest to have some effect, however have not been fully studied and investigated. So far, the full reasons why a fibroid may bleed are not completely understood. (1)(11)

Our patient was admitted with severe abdominal pain and hemodynamically un stable, images showed large subserous fibroids with intraperitoneal bleeding, so as she was un stable decided for laparotomy and proceed. intraoperative finding of bleeding from one of superficial vessels overlying the fibroid. Patient was on day three of her cycle which is going with the theory of venous congestion with mensuration can lead to trauma to those superficial vessels.

Teruo Risai reports a case of a 38-year-old woman who presented with sudden collapse and cardiac arrest (9). An autopsy showed a 3 Liters of hemoperitoneum with a large uterine leiomyoma (weight 2.5 kg). Ruptured splenic aneurysm was found to be the cause of death. (9) This highlights the importance of a multi-disciplinary team of surgeons and gynaecologists while treating such cases (4)

Varras M (10) reported case 55 years old women, post-menopause developed a fever and severe pain in the pelvic area. She had a CT scan of the abdomen and pelvis which show a 12 cm mass growing from the near wall of her uterus. She had Diagnostic laparotomy which confirm bleeding from uterine fibroid vessels. This case illustrates that a ruptured degeneration of uterine leiomyoma can occur in all ages and should be considered as one of the differential diagnoses for all women presenting with abdominal pain and a large fibroid mass regardless of their hormonal status or age. (10)

Treatment of ruptured fibroids first focused on stopping bleeding and ideally saving the uterus, particularly in younger women. Experts suggest that emergency uterine fibroid embolization can play a role in controlling bleeding. In our patient laparotomy was done , cauterization of bleeding vessel done , patient received blood product , myomectomy was not done as our centre is lacking intervention radiology in case indicated and patient was strongly against hysterectomy as she is nulliparous , so further intervention abended as bleeding was controlled and patient was referred later to tertiary centre for Uterine artery embolization .

#### Conclusion:

Uterine fibroids are common benign tumour in women especially at reproductive age, however complications resulting in acute intra peritoneal bleeding are very rare. Degeneration of leiomyomas resulting in rupture has been described in both pregnant and non-pregnant patient. The actual case of those vessels' ruptures remained un clear with many theories This report emphasis that rupture uterine fibroid should be included in the differential diagnosis when dealing with patients with a cystic tumour and massive hemo-peritoneum. This case highlights the need for high clinical suspicion of this rare condition and multidisciplinary approach.

#### Disclosure:

The authors declared no conflicts of interest. A written consent was obtained from the patient

#### References:

1. LIM, W.H., COHEN, S.C. & LAMARO, V.P. Intra-abdominal haemorrhage from uterine fibroids: a systematic review of the literature. BMC Surg, 2020 , volume 20 <https://doi.org/10.1186/s12893-020-00736-5>
- 2- Jenayah AA, Saoudi S, Sferi N, Skander R, Marzouk SB, Cherni A, Sfar E, Chelli D, Boudaya F. Spontaneous subserosal venous rupture overlying a uterine leiomyoma in a

young woman. *Pan Afr Med J.* 2017 Nov 7;28:205. doi: 10.11604/pamj.2017.28.205.12135. PMID: 29610643; PMCID: PMC5878847.

3- Danikas D, Theodorou SJV, Kotrotsios J, Sills C, Cordero PE. Hemoperitoneum from Spontaneous Bleeding of a Uterine Leiomyoma: A Case Report. *The American Surgeon*TM. 1999;65(12):1180-1182

4- Lotterman S. Massive hemoperitoneum resulting from spontaneous rupture of uterine leiomyoma. *Am J Emerg Med.* 2008 Oct;26(8):974.e1-2. doi: 10.1016/j.ajem.2008.02.029. PMID: 18926384

5- Tan YL, Naidu A. Rare postpartum ruptured degenerated fibroid: a case report. *J Obstet Gynaecol Res.* 2014 May;40(5):1423-5. doi: 10.1111/jog.12334. Epub 2014 Apr 2. PMID: 24689652.

6- Goswami N, Alalade A. A rare case of acute abdomen: spontaneous rupture of degenerated fibroid. *BMJ Case Rep.* 2021 Mar 10;14(3):e238010. doi: 10.1136/bcr-2020-238010. PMID: 33692046; PMCID: PMC7949465.

7- Pachy F, Lemerrier D, Dommergues M, Sibony O. Complication rare d'un léiomyome utérin: énucléation spontanée avec hémopéritoine [Unusual complication of uterine leiomyoma: spontaneous avulsion with massive hemoperitoneum]. *J Gynecol Obstet Biol Reprod (Paris).* 2009 May;38(3):239-41. French. doi: 10.1016/j.jgyn.2008.08.008. Epub 2009 Mar 24. PMID: 19179018.

8- Gupta N, Dadhwal V, Misra R, Mittal S, Kiran S. Atypical presentation of a leiomyoma as spontaneous massive haemoperitoneum. *Eur J Obstet Gynecol Reprod Biol.* 2008 May;138(1):120-1. doi: 10.1016/j.ejogrb.2007.01.014. Epub 2007 Mar 26. PMID: 17386968.

9- Dahan MH, Ahmadi R. Spontaneous subserosal venous rupture overlying a uterine leiomyoma. A case report. *The Journal of Reproductive Medicine.* 2002 May;47(5):419-420. PMID: 12063882.

10- Varras M, Antoniou S, Samara Ch, Frakala S, Angelidou-Manika Z, Paissios P. Intraperitoneal haemorrhage secondary to perforation of uterine fibroid after cystic degeneration. Unusual CT findings resembling malignant pelvic tumor: case report. *Eur J Gynaecol Oncol.* 2002;23(6):565-8. PMID: 12556108.

11. Hillary Hu and Joseph Do Woong Choi and Naim Arrage. Unusual presentation of hemoperitoneum secondary to ruptured uterine leiomyoma: a case report. *Gynaecology and Pelvic Medicine journal* .2022 (5). <https://gpm.amegroups.org/article/view/8267>