

Original Research Article

Antimicrobial resistance and ARGs detection in treated final effluent from STPs: An upcoming challenge to the environment

ABSTRACT

Aim: The current study has been designed to analyse the presence of antimicrobial resistance organisms in the hospital effluent.

Study Area and Sampling: Samples for microbiological analysis were collected from two different hospitals in Trivandrum City and carried out further microbiological analysis.

Methodology: In this study, MDRB (Multi Drug Resistant Bacteria) were isolated from the hospital effluent samples collected from two hospitals. Antibiotic susceptibility analysis showed that the 90% screened organisms were resistant to different antibiotics - Tetracycline (30µg), Amikacin (30µg), Gentamycin (10µg), Ciprofloxacin (5µg), Colistin (10µg) and Amoxicillin (30µg). Metagenomic surveillance of effluent helped to assess the efficacy of STPs, at the same time assessing the local clinical antibiotic resistance condition by detecting the presence of antibiotic resistance genes (ARGs) in the hospital effluent.

Results: In this study, a total of 3 antibiotic resistant bacterial strains obtained from hospital STP effluent were identified by 16S rRNA sequence analysis. The sequences of *E.coli*, *Klebsiella* and *Enterobacter* were submitted in Genbank with accession number MT784125, OM978270 and MN437586 respectively. The final effluent from Hospital 1 showed 100% resistance to Tetracycline and 86% resistance to Amoxicillin followed by sensitivity 28%, 22%, 18% and 10% respectively to Ciprofloxacin, Gentamycin, Amikacin and Colistin. The bacterial strains isolated from final effluent of Hospital 2 showed highest resistance to Amikacin and Colistin which is 100% and 86% and 82% resistance to Gentamycin and Ciprofloxacin.. The gene primers used for the respective genes above have been amplified in the sample with a higher efficiency of *16 SrRNA*, *ermB* and *ampC* primers showing a lower Cq value. Thus, these three genes were detected in the samples at high amount which showed the prominent use of the consecutive antibiotics in the clinical field

Keywords: Antibiotic resistance, Ab genes, Final effluent, Metagenomics, Sewage treatment plants

1. INTRODUCTION

An antibiotic is an antimicrobial agent for fighting against bacterial infections, widely used in the treatment and prevention of such infections. They may either kill or inhibit the growth of bacteria and limited number possesses antiprotozoal activity and not at all effective against viruses, that can be treated with some antiviral drugs.

Antimicrobial Resistance occurs when microbes change over time and no longer respond to medicines making infections harder which will make riskier conditions like disease spread, illness etc. The accumulation of resistance features after exposure to various antibiotics and cross-resistance may result in multidrug-resistant (MDR) bacteria. MDR pathogens present a significant threat to global public health and are one of the main clinical concerns [3]. The occurrences and spread of multidrug antibiotic resistance among bacterial pathogens may have serious consequences for human health [4].

Antimicrobial resistance (AMR) is a global threat to public health and environment, the misuse and abuse of antimicrobials in hospital field and human medicine has accelerated the growing worldwide phenomenon of AMR. Hospitals also play a significant role in the

distribution of antibiotic-resistant bacteria into the environment. Antibiotic-resistant bacteria may leave hospitals on colonized patients, but also via the effluent and the waste water from there [1]. Hospital effluent contains a special category of waste that is highly hazardous to public health and environment because of their infectious and toxic characteristics [2]. Hospital effluent contain large amounts of pathogenic bacteria and a variety of substances with antimicrobial activity which will affect in adverse. The effluent from the hospitals consist of pathogens mostly antibiotic resistant microbes mainly from the sources like research, medicinal excretion from the patients etc. Hospital wastewaters are considered generally as hot spots for antibiotic resistance, generating an environment for the exchange of antibiotic resistance genes. After treatment from the STPs water is discharged into environment and on water bodies or surface waters. Certain pathogenic microorganisms can remain in an aquatic environment for longer periods, creating dissemination routes and environmental reservoirs of antibiotic resistance genes (ARGs)[3].

The acquired new resistance mechanisms are the main cause of the emergence and spread leading to antimicrobial resistance continues to threaten our ability to treat common infections. Superbugs are the example for this type of pathogens and also known as pan-resistant bacteria. The persistence of antibiotic-resistant bacteria (ARB) and antibiotic resistant genes (ARGs) in the clinical environment due to the overuse and misuse of humans Antibiotic resistance result in virulence, pathogenicity, disease outbreaks and transmission, leading to prolonged morbidity, hospitalization and even mortality. The control of human infections and diseases becomes problematic once bacterial infections develop resistance to common first-line antibiotics[5].

ARGs are not completely removed by the biological treatments but sometimes remain after the chemical treatments also like the disinfection process etc. This causes the discharge of antibiotic resistance into the environment by the effluent from the treatment plants. Hospital effluent (HEs) are considered as special category because of its hazardous nature. Some studies have reported lower levels of ARB resistance genes (ARGs) in municipal wastewater compared to hospital effluent because of the wide usage of antibiotics [6].

This study is focused on the detection of antimicrobial resistance in wastewater especially from Sewage Treatment plant effluent in hospitals from Trivandrum city and to analyse the presence of ARGs and more broadly antibiotic resistant bacteria.

2. MATERIALS AND METHODS

2.1 Sample Collection

The study was carried out to detect the presence of antibiotic resistant microbes and genes in final effluent of two hospitals in Thiruvananthapuram district, Kerala. Samples for microbiological analysis were collected in sterile glass bottles and placed in icebox and transported to the laboratory for further analysis.

2.2 Isolation and Identification of Antibiotic Resistant bacteria

The presence of coliform bacteria was detected using MPN (Most Probable number) technique. The direct streak plate culture technique was carried out in order to detect the presence of *Pseudomonas* and *Enterobacteriaceae* species like *Staphylococcus*, *Salmonella*, *Shigella* on respective media, *Pseudomonas* Agar, MSA and XLD agar respectively. *E. coli* was further confirmed from positive MPN tubes by culturing on Eosin Methylene Blue (EMB) agar followed by observation for green metallic sheen.

The isolated genomic DNA was checked by agarose gel electrophoresis and was further used for the 16S rDNA amplification. The microbial DNA was amplified using 16srRNA primers. PCR reaction was carried out in SimpliAmp Thermal Cycler, Thermofischer (The Applied Biosystems). A reaction mixture (25 µl reaction volume) was containing 1.5µl of 10µM forward primer, 1.5µl of 10µM reverse primer, 12µl of Takara master mix; 5µl of sterile autoclaved water, and 5µl of template DNA samples. The following universal primers were used;

27F CGGCCAGACTCCTACGGGAGGCAGCA

1492R GCGTGGACTACCAGGGTATCTAATTC

The template DNA was amplified on DNA thermocycler using the PCR conditions 94°C for 4 minutes, 94°C for 30 seconds, 64.5°C for 30 seconds, 72°C for 30 seconds. The total numbers of cycles were 35, with final extension at 72°C for 5 minutes, performed using the programmer. Agarose gel electrophoresis was done for the qualitative analysis of PCR products. Horizontal gel electrophoresis unit was used to run the sample on the gel to determine the size of amplicons. The PCR products were electrophoresed on 2% agarose gel stained with Ethidium Bromide (1mg/ml), run at constant voltage of 50V in 1XTAE buffer. A 100bp DNA ladder was used for the comparative study. The gel documentation was carried

out using Documentation Unit. The remaining PCR product was stored at -20°C for sequencing. PCR products of 16S rRNA of the isolate was obtained through amplification and were purified and sequenced. Sequence results obtained were analyzed using applied bio systems and NCBI-BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi#>). A consensus sequence was generated from forward sequence data using applied bio system software. The consensus sequence was used to perform BLAST against the NCBI Gene Bank database. The first ten sequences were chosen based on their maximum identity score and aligned using the multiple alignment software program. Based on the phylogenetic tree and the pair-wise distance matrix, the closest homolog of each isolate from the NCBI Gene Bank database was identified. The nucleotide sequence of the current isolates was subjected to Blast.

2.3 Antibiotic Susceptibility Test

Antibiotic Susceptibility test for various bacterial isolates was done by antibiotic disks (HiMedia) in accordance with the Clinical and Laboratory Standards Institute (CLSI) by Kirby-Bauer disc diffusion method. For this, the isolates were inoculated into Muller Hinton broth and incubated for 12-24 hrs. The turbidity of the broth for the method was further made equivalent to 0.5 McFarland standards and was used for inoculation over Muller Hinton agar plate. The various antibiotic discs used for the study were Tetracycline (30µg), Amikacin (30µg), Gentamycin (10µg), Ciprofloxacin (5µg), Colistin (10µg) and Amoxicillin (30µg). The discs were placed over the inoculated surface of MHA plates and were incubated for 24 hrs at 37°C to observe the zone of inhibition.

2.4 Metagenomic DNA isolation from water samples:

Total DNA was isolated from the water sample using the soil DNA isolation kit from MagGenome. Briefly, 250ml of the water sample was filtered through an autoclaved nitrocellulose membrane with a pore size of 0.2µm using a sterile syringe filter. The membrane was ground in liquid nitrogen and care was taken to avoid thawing of the sample. The ground powder was later transferred to 2ml centrifuge tubes. Following this, DNA was isolated using the manufacturer's protocol (<http://www.maggenome.com/wp-content/uploads/2022/04/XpressDNA-Soil-kit.pdf>).

The isolated DNA was quantified using nanodrop and further checked for the presence of inhibitions in the PCR amplifications by the amplification of the 16S rRNA gene (27F and 1492R primer pair) using conventional PCR and visualization of gel image. The

concentration of the isolated DNA was found to be 314 µg/ml with an absorbance ratio (A260/A280) of 1.8.

2.5 qRT PCR Amplification of Resistant gene

The isolated DNA was quantified using nanodrop. The concentration of the isolated DNA was found to be 314 µg/ml with an absorbance ratio (A260/A280) of 1.8. These samples were further subjected to qPCR amplifications in the previously obtained dilution of DNA (Sample Concentrations 1:4) using the provided gene primers. The PCR reaction was carried out in 10µl reaction mixture containing 5µl 2X SYBER GREEN qPCR Mix (G bioscience), 1µl of 2µM forward and reverse primer, 1µl of template DNA and 2µl Nuclease free water. The DNA dilution (1:4) was found to have maximum amplification and minimum inhibition. The dilution 1:4 was used in further amplifications using the primer pairs. The primer sequences (Table: 1) used for qPCR using previously published primer sets allowing the amplification of short amplicons (160–420 basepairs).

3. RESULTS

3.1 Isolation and identification of Antibiotic resistant bacteria

In this study, a total of 3 antibiotic resistant bacterial strains obtained from hospital STP effluent were identified by 16S rRNA sequence analysis. The sequences of *E. coli*, *Klebsiella* and *Enterobacter* were submitted in Genbank with accession number MT784125, OM978270 and MN437586 respectively. Here, NCBI BLASTn analysis confirmed the isolates as *E. coli* and *Klebsiella* and *Enterobacter*. The phylogenetic analysis of the isolated strains are represented below (Fig 1&2)




Fig.1.16S rDNA phylogeny of *E. coli* isolate[12, 13]




Fig.2.16S rDNA phylogeny of *Enterobacter* isolate[12, 13]




Fig.3.16S rDNA phylogeny of *Klebsiella* isolate[12, 13]

3.2 Antibiotic Susceptibility Test

Out of the 6 major antibiotics tested, the bacterial strains isolated from final effluent of Hospital 1 exhibited highest resistance to Tetracycline and Amoxicillin. Here the final effluent from Hospital 1 showed 100% resistance to Tetracycline and 86% resistance to Amoxicillin followed by sensitivity 28%, 22%, 18% and 10% respectively to Ciprofloxacin, Gentamycin, Amikacin and Colistin. The bacterial strains isolated from final effluent of Hospital 2 showed highest resistance to Amikacin and Colistin which is 100% and 86% and 82% resistance to Gentamycin and Ciprofloxacin. Antibiotic resistant profile of the selected hospitals is shown below (Figure:4)




Fig.4. Antibiotic resistance pattern of hospital final effluent

3.3 Metagenomic DNA isolation from water samples

For Metagenomic analysis, DNA bands were obtained and the concentration of the isolated DNA was found to be 314 $\mu\text{g/ml}$ with an absorbance ratio of 1.8 (A_{260}/A_{280}). The Gel image of PCR product during DNA isolation is represented in Fig 5




Fig.5. Gel image of PCR product during DNA isolation

The primer sequences used for the metagenomic analysis of the sample is shown below (Table 1).

Table 1: Primer sequences [14]

| Gene | Sequence | Annealing temperature (°C) |
|-------------|------------------------------|----------------------------|
| sul1 | F: CGCACCGGAAACATCGCTGCAC | 68 |
| | R: TGAAGTTCCGCCGCAAGGCTCG | |
| Sul2 | F: TCCGGTGGAGGCCGGTATCTGG | 60 |
| | R: CGGGAATGCCATCTGCCTTGAG | |
| ermB | F: CATTAAACGACGAAACTGGC | 63 |
| | R: GGAACATCTGTGGTATGGCG | |
| ampC/UKSAL1 | F: TTCTATCAAMACTGGCARCC | 55 |
| | R: CCYTTTTATGTACCCAYGA | |
| tet(A) | F: CCTGCGGATCTGGTTCACT | 55 |
| | R: GCCAGCGAGACGAGCAAGA | |
| vanA | F: ATGAATAGATAAAAAGTTGCAATAC | 55 |
| | R: GGAGTAGCTATCCAGCATT | |
| 16S rRNA | F: CCTACGGGAGGCAGCAG | 68 |
| | R: ATTACCGCGGCTGCTGGC | |

The qPCR assay allowed the quantification and detection of Abr genes present in the sample. When testing water samples with this assay, melting curves and amplification curves for the consecutive primers were obtained. Melt curve of the consecutive primers implies the qPCR amplicon in the samples assessed. The melt curves of 16SrRNA sequence & Sul 1, Sul 2 are represented below in Figure 6, Melt curves of amp C & vanA are represented in Figure 7 & Melt curves of ermB & tetA are represented in Figure 8 and further amplification was represented in Figure 9, 10 & 11 respectively.




Fig.6. Melt curve of 16SrRNA sequence& Sul 1, Sul 2 showing the melt profile for PCR amplicons




Fig.7. Melt curve of amp C&vanA sequences showing the melt profile for PCR amplicons.




Fig.8. Melt curve of ermB&tetA sequence showing the melt profile for PCR amplicons




Fig.9. Amplification of 16 S rRNA Sequence, Sul1&Sul2




Fig.10. Amplification of ampC&van A




Fig.11. Amplification of ermB & tet A

.Quantification data of sample replicates and its Quantification cycle mean represents the amount of genes present in the sample (Table 2).

Table 2: Quantification data

| Gene | Sample replicates | Cq Mean |
|---------|-------------------|---------|
| Sul 1 | Hospital 2a | 21.80 |
| Sul 1 | Hospital 2b | 21.41 |
| Sul 1 | Hospital 2c | 20.46 |
| 16SrRNA | Hospital 2a | 19.14 |
| 16SrRNA | Hospital 2b | 15.67 |
| 16SrRNA | Hospital 2c | 15.64 |
| Sul 2 | Hospital 2a | 26.37 |
| Sul 2 | Hospital 2b | 25.82 |
| Sul 2 | Hospital 2c | 25.76 |
| ermB | Hospital 2a | 16.03 |
| ermB | Hospital 2b | 18.05 |
| ermB | Hospital 2c | 17.39 |
| Tet A | Hospital 2a | 21.72 |
| Tet A | Hospital 2b | 22.39 |
| Tet A | Hospital 2c | 22.23 |
| amp C | Hospital 2a | 19.70 |
| amp C | Hospital 2b | 20.19 |
| amp C | Hospital 2c | 19.92 |
| van A | Hospital 2a | 25.48 |
| van A | Hospital 2b | 23.13 |
| van A | Hospital 2c | 23.25 |

Metagenomic DNA isolated from wastewater samples was used to determine ARGs profiles. The gene primers used for the respective genes above have been amplified in the sample with a higher efficiency of *16 SrRNA*, *ermB* and *ampC* primers showing a lower Cq value. Cq value is inversely proportional to the amount of genes in the sample, Cq mean value for *16 SrRNA* was 19.14, 15.67, 15.64 and for *ermB* was 16.03, 18.05, 17.39 and for *ampC* was 19.70, 20.19 and 19.92 respectively, thereby indicating higher relative quantification. Thus, these three genes were detected in the samples at high amount which showed the prominent use of the consecutive antibiotics in the clinical field. While, *Sul 1*, *Sul 2*, *Tet A* and *van A* had higher Cq value of about 21, 25, 22 and 23 respectively indicating lower quantification compared to *16 SrRNA*, *ermB* and *ampC*.

4. DISCUSSION

The evolution of Antimicrobial resistance associated with the STP sector is an emerging threat to health and social well being of humans. Antimicrobial resistance in wastewater and hospital effluent has been noted as one of the increasing microbial threats of this century. Recent studies show the transferability of the resistance genes from effluent to humans through water bodies which is an emerging risk factor for public health [7]. Many studies have spotlighted the role of wastewater as a significant environmental reservoir of AMR as it represents a supreme environment for AMR bacteria (ARB) and antimicrobial resistant genes (ARGs) to persist [8]. Many research have highlighted the role of sewage effluent as a major environmental reservoir of AMR, as it represents an ideal environment for AMR microorganisms and ARGs to persevere. The situation of ARGs is composite, because they are not degradable and can even be spread among microbial communities in the environment through the mechanism of horizontal gene transfer, which is the main resistance mechanism in *Enterobacteriaceae* family.

Ampicillin, Sulfamethoxazole, and Ciprofloxacin, or closely related drugs (Amoxicillin), are among the top 5 antibiotics prescribed for use for adults in the United States [9], and all have been found to occur in WWTPs in varying concentrations and design conditions. After the administration of the antibiotic substances they are released into effluent through patient excreta. Unused drugs are sometimes disposed of down drains. The drugs were released into the effluent where they are not eliminated through the process of sewage treatment method and enter the aquatic environment and eventually reach drinking water. If

the concentrations are high enough, effluent from hospitals, municipal sewage and sewage treatment plants (STPs) may become a reservoir for the selection of resistant bacteria [10].

In the current study, the samples (final effluent) collected from two hospitals were found to have the presence of antibiotic resistant bacteria, *E. Coli* MT784125, *Klebsiella* OM978270 and *Enterobacter sp* MN437586, which were resistant to multiple drugs. The presence of antibiotics in hospitals and treatment plants may help the bacterial genes to persist in the micro biome for a long time. The presence of antibiotics even in small amount in the final effluent after treatment ensures the sustainability of such resistant genes and may dominantly express in the microbial community.

Commonly hospital wastewater contains a high number of multidrug-resistant coliform bacteria and *E. coli*. These strains can transfer multiple resistance genes through the conjugative plasmid and spread them to susceptible bacterial species. Hospital wastewaters frequently contain significant amounts of fecal coliforms, which exhibit resistance or multi-resistance towards various types of antibiotics [11].

Metagenomic analysis benefits the ability to quantify thousands of genes in the sample and the data can be reanalysed if novel genes of interest are identified. Here, in this work, qPCR technique objectively and independently document the abundance of ARG in wastewater effluent and quantified the absolute abundance of target genes in raw wastewater samples. In this study, STP effluent was found as a potential source of AMR by detecting ARGs using a Metagenomics DNA-seq approach. Metagenomic DNA-seq analysis showed that the genes amplified in the sample with a higher efficiency were, *16 SrRNA*, *ermB* and *ampC* primers showing a lower C_q and higher relative quantification values which denotes higher abundance of these 3 genes in the sample. This is an indication of heavy usage of the antibiotics like Erythromycin and Ampicilin in the clinical field.

5. CONCLUSION

The over usage of Antibiotics in hospitals for disease prevention may cause the evolution of antimicrobial resistance. *E. coli* and *Klebsiella pneumoniae* and *Enterobacter* isolated in the study shows high resistance to Tetracycline, Amoxicillin and Colistin which might be the after effect of over exploitation of these antibiotics in clinical field. Also, the resistance genes are suspected to be emanate through the species and its continuous evolution may cause cross resistance to other antibiotics. The metagenomic analysis also confirms the presence of resistant genes in the effluent sample. Hence antibiotic use in hospitals and its final disposal should be

monitored properly to avoid the development of resistance. The study showed a correlation between the antibiotics used in the hospitals and the emergence of antimicrobial resistance among the treated effluent. The results of the study indicate the need for strict control over use of antibiotic in the environment to limit the rapid evolution and spread of antimicrobial resistance to different antibiotics.

References :

[1] Didier Hocquet , Marie Petitjean , Laurence Rohmer , Benoît Valot , Hemantha D Kulasekara, Lucas R Hoffman , Samuel I Miller (2016) “Pyomelanin-producing *Pseudomonas aeruginosa* selected during chronic infections have a large chromosomal deletion which confers resistance to pyocins”, Society for Applied Microbiology and John Wiley & Sons Ltd, Environmental Microbiology Volume 18(10):3482-3493

[2] Sara Rodriguez-Mozaz , Sara Chamorro , Elisabet Marti , Belinda Huerta , Meritxell Gros , Damià Barceló , Jose Luis Balcázar (2015) “Occurrence of antibiotics and antibiotic resistance genes in hospital and urban wastewaters and their impact on the receiving river” Journal of Water , Volume 69:234-242

[3] Nikola Roulová, Petra Mot'ková, Iveta Brožková, Marcela Pejchalová J Water Health (2022) “Antibiotic resistance of *Pseudomonas aeruginosa* isolated from hospital wastewater in the Czech Republic” Journal of Water health, Volume 20 (4): 692–701

[4] Magdalena Pazda , Jolanta Kumirska , Piotr Stepnowski , Ewa Mulkiwicz 2019 Dec “Antibiotic resistance genes identified in wastewater treatment plant systems” Journal of Science of The Total Environment , A review Volume 697: 134023.

[5] Kristina E Rudd , Sarah Charlotte Johnson , Kareha M Agesa , Katya Anne Shackelford , Derrick Tsoi , R Scott Watson , Mohsen Naghavi 2020 “Global, regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study” The Lancet, Volume 395(10219):200-211.

[6] Urska Rozman, Darja Duh; Mojca Cimerman; Sonja Sostar Turk. Journal of Water, Sanitation and Hygiene for Development (2020) “Hospital wastewater effluent: hot spot for

antibiotic resistant bacteria”, *Journal of Water, Sanitation and Hygiene for Development* Volume10 (2): 171–178.

[7]S. Sreejith, ShamnaShajahan, P.R. Prathiush, V.M. Anjana, ArathyViswanathan, VishnuChandran, G.S. Ajith Kumar, R. Jayachandran, JyothisMathew, E.K. Radhakrishnan (2020) “Healthy broilers disseminate antibiotic resistance in response to tetracycline input in feed concentrates”, *Microbial Pathogenesis.*, Volume 149, 104562 149:104562

[8]Nour Fouz , Krisna N. A. Pangesti , Muhammad Yasir , Abdulrahman L. Al-Malki , Esam I. Azhar , Grant A. Hill-Cawthorne and Moataz Abd El Ghany (2020) “The Contribution of Wastewater to the Transmission of Antimicrobial Resistance in the Environment: Implications of Mass Gathering Settings”, *Tropical Medicine and Infectious disease*, 2020, 5(1), 33; <https://doi.org/10.3390/tropicalmed5010033>.

[9]Maitreyee Mukherjee, Edward Laird, Terry J. Gentry, John P. Brooks and Raghupathy Karthikeyan “Increased Antimicrobial and Multidrug Resistance Downstream of Wastewater Treatment Plants in an Urban Watershed”, *Front. Microbiol.*, 24 May 2021 *Sec. Antimicrobials, Resistance and Chemotherapy*, Volume 12 – 2021 <https://doi.org/10.3389/fmicb.2021.657353>.

[10]K. Kummerer, “Significance of antibiotics in the environment”, *Journal of Antimicrobial Chemotherapy*, Volume 52, Issue 1, July 2003, Pages 5–7, <https://doi.org/10.1093/jac/dkg293>.

[11]Tomas Mackul'ak, Klara Cverenkova , Miroslav Feher, Michal Tamas , Miroslav Gal, Monika Naumowicz, Viera Spalkova, and Lucia Birosova “Hospital Wastewater—Source of Specific Micro pollutants, Antibiotic-Resistant Microorganisms, Viruses, and Their Elimination”, *Antibiotics (Basel)*. 2021 Sep; 10(9): 1070, doi: 10.3390/antibiotics10091070.

[12] Kimura M. (1980). “A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences”. *Journal of Molecular Evolution* **16**:111-120.

[13] Tamura K., Stecher G., and Kumar S. (2021). MEGA 11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* <https://doi.org/10.1093/molbev/msab120>.

- [14]C. Stangea, J.P.S. Sidhub, S. Tozec, A. Tiehma (2019) “Comparative removal of antibiotic resistance genes during chlorination, ozonation, and UV treatment” International Journal of Hygiene and Environmental Health 222 (2019) 541–548.

UNDER PEER REVIEW