

Original Research Article
**Prevalence of *Klebsiella pneumoniae* and
Acinetobacter baumannii in urine samples of
pregnant women in South-East, Nigeria.**

ABSTRACT:

*AIMS: To determine the prevalence of *Klebsiella pneumoniae* and *Acinetobacter baumannii* in urine samples of pregnant women attending ante-natal clinics at three health institutions within the study area.*

STUDY DESIGN: This study took a form of a descriptive cross-sectional study.

Place and Duration of Study: Department of Microbiology Federal University Teaching Hospital, Owerri, Teaching Hospital Orlu and General Hospital Okigwe, Imo State between October 2021 and February 2022.

*METHODOLOGY: Urine samples were collected aseptically from 408 pregnant women between 18-45 years and analyzed by culturing unto the surface of freshly prepared MacConkey and Chromagar plates for the isolation of *Klebsiella pneumoniae* and *Acinetobacter baumannii* respectively, and confirmed by appropriate biochemical tests.*

*RESULTS: From the results, there was a prevalence rate of 145/408 (35.54%) for both isolates with *Klebsiella pneumoniae* being more dominant with a prevalence rate of 96/408 (23.53%) as against *Acinetobacter baumannii* that was 49/408 (12.01%). The highest prevalence of the isolates was seen in Owerri zone with a prevalence rate of 37 (9.07%) for *Klebsiella pneumoniae* and 23 (5.64%) for *Acinetobacter baumannii* respectively. The highest age specific prevalence of the isolates within the three zones was observed in the 24-29 year-olds with a prevalence rate of 42(10.29%) in Owerri zone while prevalence of both organisms was higher in pregnant women in their third trimester of pregnancy with a prevalence rate of 72 (17.65%) and multigravida women with a prevalence rate of 217 (53.2%).*

CONCLUSION: The results from this study showed that these isolates are becoming more predominant in causing both symptomatic and asymptomatic urinary tract infection in pregnant women, hence regular check-ups and right treatment of urinary tract infections in pregnant women should be encouraged.

*Keywords: Prevalence, urinary tract infection, *Klebsiella pneumoniae*, *Acinetobacter baumannii*. Urine, pregnant women, Owerri.*

1. INTRODUCTION

Urinary tract infections account for a large percentage of community and health-care acquired infections affecting 2–10% of pregnant women in both underdeveloped, developed and developing countries, and they may be symptomatic or asymptomatic [1]. UTI can be described based on the part of the tract affected; for the

upper tract it is pyelonephritis, while for the lower tract it is cystitis or urethritis [2]. Untreated or undertreated bacteriuria in pregnancy, regardless of symptoms, increases the risk of negative outcomes, including increased maternal and newborn morbidity and death [3]. As a result, screening and treating pregnant women for bacteriuria has become a standard aspect of prenatal treatment.

Urinary tract infection occurs in about 8% of all pregnant women and peaking in about 90% during the period of about 24 weeks of gestation [4]. About 70% of pregnant women have urethral dilatation which remains until delivery in addition to having glucosurea which enhances bacterial growth in urine [5]. Symptoms associated with urinary tract infection includes, pain or burning (discomfort) sensation when urinating, cramps or pain in the lower abdomen, the need to urinate more often than usual, turbid and foul smelling urine, pressure or tenderness in the area of the bladder and when bacteria spread to the kidneys, there can be back pain, chills, fever, nausea and vomiting [1].

Globally, the prevalence of UTIs in pregnant women is between 13% - 33% with symptomatic bacteriuria occurring in 1% - 18% [6], while asymptomatic cases are observed in 2%-45% in different populations [7] among females. The rate of prevalence has varied including epidemiological and cross-sectional researches in developing and under-developed nations. Epidemiological surveillance of outpatient urine culture offers relevant insights into the changing prevalence and even antimicrobial drugs that are susceptible to specific uropathogens [6], particularly in patients presenting nonspecific symptoms. The most significant organisms causing UTI in pregnant women includes, *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacteriaceae* (eskape) [8]. because of their distinct resistance properties, *Acinetobacter baumannii* and *Klebsiella pneumoniae* are becoming more difficult to combat. They are non-lactose-fermenting, catalase-positive, non-motile, non-fastidious, oxidase-negative, and aerobic gram-negative coccobacilli are all members of this genus. Clinically, *A. baumannii* and *K. pneumoniae* are important because they cause UTIs and nosocomial infections and are inherently resistant to a broad range of antimicrobials, with a high risk of acquiring resistance. They are found in roughly 42% of healthcare personnel and patients' hands, according to studies [9]. Their capacity to tolerate desiccation allows them to persist for months in inert things, facilitating their spread in hospitals and the environments.

In many hospitals, infections of the respiratory system, urinary tract, bloodstream (septic), post-surgical (wound) infections, and pneumonia are among the most often reported diseases caused by these bacteria [10]. Although the prevalence of these organisms has been detected in several Nigerian hospitals, there are no mechanisms in place to ameliorate the issue, such as surveillance studies that gives trustworthy data in the state. As a result, investigations to determine the prevalence of this organisms in Imo state is required to close the knowledge gap and provide more foundation for empirical treatment. This research will look at the distribution of *Klebsiella pneumoniae* and *Acinetobacter baumannii*, which was found in urine samples of pregnant women in the state.

2. METHODOLOGY

Study Area

The research was conducted in Imo state Nigeria. The state is divided into three zones, namely; Owerri, Orlu and Okigwe. Hospitals and health centers within these three zones were used as sample collection centers and the samples was analyzed at federal university teaching hospital Owerri, Imo state. Ethical clearance was obtained and informed consent was also obtained from participants. The inclusion criteria included pregnant women within the study area who gave informed consent to participate in the study. Pregnant women without informed consent and who had taken antibiotics within the last 2 weeks of sample collection were excluded.

Sample Collection

Four hundred and eight (408) mid-stream urine samples of apparently healthy pregnant women between the ages of 18-45 years in Imo state were collected in a sterile, dry, leak-proof container and labelled properly.

Using the sampling formula from charan&biswas [11]. The samples were divided within the three geopolitical zones comprising of Okigwe (87), Orlu (121) and Owerri (200). With the aid of a questionnaire, demographic parameters including age, gestational age of pregnancy, and questions related to clinical features of *K. pneumoniae*, *acinetobacter* and urinary tract infection was issued to them. The collected urine samples was placed in a cold box and transported to the microbiology laboratory of federal university teaching hospital Owerri for analysis.

Isolation of Organisms

The Chromagartm(Paris, France) and Macconkey Agar (Titan Biotech Ltd, india) was prepared according to manufacturer's instruction and was used for primary isolation of *Acinetobacter baumannii* and *Klebsiella pneumoniae* respectively by streaking the urine sample unto the surface of the prepared media (12). Colonies from culture plates of both organisms were sub-cultured onto nutrient agar plates, pure cultures were made with repeated streaking and incubated at 37°C for 24 hours.

Identification of Isolates

Identification and confirmation of the organisms from the culture plates of the pure colonies was done by morphological, cultural characteristics and biochemical tests adopted from Cheesbrough, (2006) and the result recorded accordingly. *Klebsiella pneumoniae* appeared as large, mucoid pink colonies without retaining the primary stain and was confirmed to be non-motile, citrate positive, catalase positive and oxidase negative.

Acinetobacter baumannii appeared as round, smooth pink colonies on the chromogenic agar without also retaining the primary stain and was also confirmed to be catalase positive, non-motile, citrate positive and urease negative.

Data Analysis And Interpretation

Statistical analysis was done using SPSS, chi square for categorical data and ANOVA for variable that were more than two.

3. RESULTS AND DISCUSSION

3.1 Total Distribution of the Isolates within the Study Area

Out of the four hundred and eight (408) urine samples of the pregnant women examined, one hundred and forty-five (145) representing 35.57% of the population was infected with either *A. baumannii* or *K. pneumoniae* with a mixed infection of about 1.97%. While two hundred and sixty three (263) representing 64.43% of the population was uninfected.

The overall pattern of distribution of *A. baumannii* and *K. pneumoniae* examined within the study area are shown in the table 1 below. Both organisms were found in all the zones sampled with an overall prevalence of 35.57%. Of the number examined in each zone, owerri zone recorded the highest occurrence of 14.71%, followed by orlu zone; 11.03% while the least percentage occurrence was observed in okigwe zone with 9.80%. The total distribution of *K. pneumoniae* and *A. baumannii* showed that urinary infection was not dependent on demography with $p=0.28$. statistical analysis revealed that there is no correlation between mixed infection as against zones of infection with $P=.05$

Table 1. Total distribution (%) of infections among zones studied.

ZONES	No. of Samples N (%)	No of persons Infected N (%)	No Infected with <i>Klebsiella pneumonia</i> N (%)	No Infected with <i>Acinetobacte r baumannii</i> N (%)	No of Mixed infection <i>Kp/Ab</i> N (%)
Okigwe	87 (21.3)	40 (9.80)	26 (6.37)	14 (3.43)	3 (0.74)
Orlu	121 (29.7)	45 (11.03)	33 (8.09)	12 (2.94)	1 (0.25)
Owerri	200 (49.0)	60 (14.71)	37 (9.07)	23 (5.64)	4 (0.98)
Total	408 (100)	145 (35.54)	96 (23.53)	49 (12.01)	8 (1.97)

* Kp/Ab = *Klebsiella pneumoniae* and *Acinetobacter baumannii*

3.2 Age-related Prevalence of the Isolates within the Study Area

The distribution of the organisms with respect to age of the women examined is shown in Table 2. Results showed that pregnant women between the 24-29 years age bracket had the highest prevalence of the organisms with a prevalence rate of 10.29% while pregnant women between 42 years and above had the least prevalence of 1.72%. From the result below, it was further observed that the prevalence rate of *A. baumannii* was higher in pregnant women between the ages of 36-41 in the three zones with a rate of 3.19% while *K. pneumoniae* occurred more in pregnant women between the 24-29 age brackets amongst the three zones with a rate of 7.35%. further analysis in table 2.revealed that *K. pneumoniae* occurred more between the 24-29 age brackets in the three zones with similar prevalence rate while *A. baumannii* occurred more between the 36-35 age brackets of Owerri zone with a prevalence rate of 34.78% and Orlu zone recorded zero prevalence of *A. baumannii* between the 18-23 age brackets. From the results, it can be deduced that infection and mixed infection was not dependent on both age and zones studied, thus the relationship between the prevalence as against age and zones showed no correlation with a p value of 0.30.

Table 2: Prevalence (%) of the isolates in relation to age of the pregnant women studied.

AGE (YEARS)	No. of Samples N (%)	No of persons Infected N (%)	No Infected with <i>Klebsiella pneumoniae</i> N (%)	No Infected with <i>Acinetobacter baumannii</i> N (%)	No of Mixed infection <i>Kp/Ab</i> N (%)
18-23	63 (15.4)	27 (6.62)	18 (4.41)	9 (2.21)	3 (0.74)
24-29	117 (28.7)	42 (10.29)	30 (7.35)	12 (2.94)	2 (0.49)
30-35	128 (31.4)	34 (8.33)	22 (5.39)	12 (2.94)	2 (0.49)
36-41	74 (18.1)	35 (8.58)	22 (5.39)	13 (3.19)	1 (0.25)
42 and above	26 (6.4)	7 (1.72)	40 (0.98)	3 (0.74)	0
Total	408 (100)	145 (35.54)	96 (23.52)	49 (12.01)	8 (1.97)

Key*: Kp/Ab = *Klebsiella pneumoniae* and *Acinetobacter baumannii*

Table 3: Prevalence (%) of the isolates with respect to age of pregnant within the three zones studied

ZONES AGE (YEARS)	No. of Samples N (%)	No of persons Infected N (%)	No Infected with <i>Klebsiella pneumoniae</i> N (%)	No Infected with <i>Acinetobacter baumannii</i> N (%)	No of Mixed infection <i>Kp/Ab</i> N (%)
Okigwe	N= 87	N=40	N=26	N=14	N=3
18-23	18 (20.7)	12 (30.0)	7 (26.92)	5 (35.71)	1 (50.0)
24-29	22 (25.3)	14 (35.0)	10 (38.46)	4 (28.57)	1 (50.0)
30-35	21 (24.1)	6 (15.0)	5 (19.23)	1 (7.14)	0
36-41	17 (19.5)	6 (15.0)	3 (11.54)	3 (21.42)	1 (50.0)
42 and above	9 (2.2)	2 (5.0)	1 (3.85)	1 (7.14)	0

above					
Orlu	N=121	N=45	N=33	N=12	N=1
18-23	17 (14.0)	4 (8.8)	4 (12.12)	0	0
24-29	28 (23.1)	13 (28.9)	10 (30.30)	3 (25.0)	0
30-35	40 (33.1)	12 (26.7)	9 (27.27)	3 (25.0)	1
36-41	25 (20.7)	14 (31.1)	9 (27.27)	5 (41.67)	0
42 and above	11 (9.1)	2 (4.4)	1 (3.03)	1 (8.33)	0
Owerri	N=200	N=60	N=37	N=23	N=5
18-23	28 (14.0)	11 (18.3)	7 (18.91)	4 (17.39)	2 (40.0)
24-29	67 (33.5)	15 (25.0)	10 (27.02)	5 (21.74)	2 (40.0)
30-35	67 (33.5)	16 (26.7)	8 (21.62)	8 (34.78)	1 (20.0)
36-41	32 (16.0)	15 (25.0)	10 (27.02)	5 (21.74)	0
42 and above	6 (2.0)	3 (5.00)	2 (5.41)	1 (4.35)	0
Total	408 (100)	145 (35.54)	96 (23.52)	49 (12.01)	8 (1.97)

Key* Kp/Ab = *Klebsiella pneumoniae* and *Acinetobacter baumannii*

3.3 PREVALENCE OF THE ISOLATES WITH RELATION TO AGE OF PREGNANCY AND NUMBER OF BIRTHS

From table 4 below, prevalence of both organisms was highest in pregnant women in their third trimester of pregnancy with a rate of 17.65%, followed by those within their second trimester and first trimester with a rate of 13.48% and 4.41% respectively. Prevalence of *K. pneumoniae* was higher within the third trimester of pregnancy (12.75%) while prevalence of *A. baumannii* was higher (5.64%) within the second trimester of pregnancy. Mixed infection occurred at a rate of 1.22% within the second trimester. This shows that infection was not dependent on the trimester of pregnancy with a P value of 0.10.

From table 4 below, prevalence of both organisms was higher in multigravida women with a rate of 14.46% for *K. pneumoniae* and 5.64 % for *A. baumannii*.

Prevalence and mixed occurrence in the overall infection shows that infection was not dependent on the gravidity of pregnancy with a P value of 0.12. Relationship against gravidity in pregnancy showed no correlation between them with a P value of 0.16.

Table 4. Occurrence (%) of the isolates with respect to gestational age of pregnancy

Trimester (gestational age)	No. of Samples N (%)	No of persons Infected N (%)	No Infected with <i>Klebsiella pneumonia</i> N (%)	No Infected with <i>Acinetobacter baumannii</i> N (%)	No of Mixed infection <i>Kp/Ab</i> N (%)
1 st Trimester (0-13 weeks)	68 (16.7)	18 (4.41)	12 (2.94)	6 (1.47)	1 (0.24)
2 nd Trimester (14-27 weeks)	151 (37.0)	55 (13.48)	32 (7.84)	23 (5.64)	5 (1.22)
3 rd Trimester (28-40 weeks)	189 (46.3)	72 (17.65)	52 (12.75)	20 (4.90)	2 (0.49)
Total	408 (100)	145 (35.54)	96 (23.52)	49 (12.01)	8 (1.97)

Key: Kp/Ab = *Klebsiella pneumoniae* and *Acinetobacter baumannii*

Table 5: Prevalence (%) of the isolates with respect to gravidity and parity

PARITY	No. of Samples N (%)	No of persons Infected N (%)	No Infected with <i>Klebsiella pneumonia</i> N (%)	No Infected with <i>Acinetobacter baumannii</i> N (%)	No of Mixed infection <i>Kp/Ab</i> N (%)
Primigravida	118 (28.9)	39 (9.56)	22 (5.39)	17 (4.17)	1 (0.25)
Multigravida	217 (53.2)	82 (20.09)	59 (14.46)	23 (5.64)	6 (1.47)
Grand-multi Gravid	73 (17.9)	24 (5.88)	15 (3.67)	9 (2.21)	1 (0.25)
Total	408 (100)	145(35.54)	96 (23.52)	49 (12.01)	8 (1.97)

Key* Kp/Ab = *Klebsiella pneumoniae* and *Acinetobacter baumannii*

4. DISCUSSION

The criteria for prevalence selection was determined after the isolation and identification of the organisms in the urine samples of pregnant women within the three zones. Pregnant women are at a higher risk of having UTIs due to the physiologic changes in pregnancy. However, in majority of these women, the infections are

asymptomatic. The results from this study revealed that *A. baumannii* and *K. pneumoniae* are endemic in the three zones of Imo state with an overall prevalence of 35.57% (145/408) of the population studied, with *K. pneumoniae* possessing an infection rate of 23.53% and *A. baumannii* possessing an infection rate of 12.01% and a mixed infection of about 1.97%. The level of infection however varied with some zones having higher level of infection than others. This difference in the prevalence rate may be influenced by peculiar ecological characteristics, mode of living and the degree of exposure [13]. This trend of results may not be unconnected to the fact that most of the local governments in these zones studied have rural villages and community arrangements. A close look at the prevalence of these organisms in the three zones of Imo state studied show that Owerri has the highest prevalence of 14.71% out of the 408 sampled; this might be due to the fact that Owerri has the highest number of population sampled due to development, better opportunities and migration of most pregnant women from the rural settings to urban settings for proper ante-natal and counselling, but the high prevalence level was not significant ($p < 0.05$). This prevalence no matter how low could be attributed to factors such as nature of pregnancy, trimester and period of pregnancy, lack of sensitization and awareness, age, ignorance, poor living condition, inadequate sanitation, level of contact with the organism, inappropriate medication, and the indiscriminate use of antibiotics.

The low prevalence of 12.01% of *A. baumannii* infection observed in this study is almost in tandem with the 14.0% reported by Nwadike, Ejide and Kalu, [14] among pregnant women in a university college hospital Ibadan, 10.0% recorded by Bashir, et al. [15] in a study conducted in three different hospitals in Kano state Nigeria alongside 16.0% recorded by Musyoki, et al. [16] in Kenyatta Kenya. The result was in contrast with a higher prevalence of 30.7% recorded by Iregbu, Ogunsola and Odugbemi, [17] in Lagos Nigeria, and at total variance with the 6.0% prevalence rate of *A. baumannii* recorded by Egwu, et al. [18] in Abakaliki, Nigeria, and 8.5% by Odewale, et al. [19].

The infection rate of 23.53% *Klebsiella pneumoniae* observed in this study was in conformity with that of Nwachukwu, Onyebuchi and Michael [20] who observed the prevalence of *klebsiella pneumoniae* at 24.0% as one of the organisms causing urinary tract infections in pregnant women in Onitsha, and Suraiya, et al. [21] who observed a prevalence rate of 23.73% in a tertiary hospital in Bangladesh, but was slightly in contrast with the 19.1% recorded by Damilola, et al. [22] amongst 407 samples in Ondo south West Nigeria. Olamijulo et al. [10] reported a prevalence of 14.6% in a study carried out on 556 pregnant women in Lagos University Teaching Hospital, Nigeria while Afunwa, Nwokediuko and Oklie [23] recorded 20.0% prevalence of *K. pneumoniae* in a study conducted on pregnant women attending ante-natal clinic in Nike district of Enugu. However, the result is also at variance with Umeaku, et al. [24] who recorded a low prevalence of 6% amongst 50 samples in Owerri, Imo state. This finding however confirmed the involvement of both pathogens in causing UTIs in pregnant women.

The distribution of the isolates in relation to maternal age shows a relatively higher frequency of occurrence between the age range of 24–29 with 10.29% and least occurrence in the age range of 42 and above with 1.72%. This is similar to the findings from Anozie, et al. [25] and Okonko et al. [26] in South-Western Nigeria, whose findings showed the highest prevalence rate of 58.6% and 17.9% respectively amongst pregnant women between the ages of 25-30 years but in contrast with the research published by Akobi, et al. [27] and Akerele, et al. [28] who showed high prevalence of *Klebsiella pneumoniae* in pregnant women between 30-35 years. The reason for the prevalence in this age group could be because many women in this age group are likely to be multiparous, ignorant, engaged in regular sexual activity and could be using one form of contraceptive or another.

The distribution of the isolates with respect to gestational age was found to be highest in the third trimester. This agrees with the findings of Okonko, et al. [26] and Afunwa, et al. [23]. It however differs from the findings of Amala and Nwokah [29] who reported higher prevalence of *K. pneumoniae* and *A. baumannii* in the first and second trimesters. These differences may be as a result of fluctuations in urinary progesterone and oestrogens in the different trimesters of pregnancy [29]. The increased prevalence in the third trimester could be due to the increased size of the uterus, urethral dilation, increased bladder volume and urethral tone, which may lead to stasis and encourage bacterial growth in the urine [4].

The distribution of isolates with gravidity in this study shows a higher prevalence among the multigravida participant compared to primagravida and grand-multigravida. This finding agrees with a study conducted by Akpan, et al. [30] but was in contrast to the study showed by Nwachukwu, et al. [20] who recorded an increase prevalence of *K. pneumoniae* and *Acinetobacter spp* in the grand-multi gravida and primagravida respectively. This is probably due to the subsiding of pelvic organs associated with high parity leading to the widening of the urethral orifice and predisposing the woman to micro-organisms which may progress to symptomatic urinary tract infection. The second highest group in this study was the primigravida group, this may be induced by malaria in pregnancy, due to impaired immunity during first pregnancy making the primigravida prone to infection[31].

CONCLUSION

This findings of this study recorded a moderate prevalence of bacteriuria in the study area among the participant which calls for concern due to the possible effect of bacteriuria on the fetus. Also having most of this prevalence being asymptomatic, there is need to educate the pregnant women on personal and environmental hygiene and also need for regular medical check-ups, laboratory screening and treatment with the right antibiotics when infections are identified to prevent birth complications.

CONSENT

As per international standard and university standard, patient(s) written consent was collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard written ethical approval was collected and preserved by the author(s).

REFERENCES

1. Madueke T, Singh N, Manchanda V. Control of multidrug-resistant gram-negative bacteria in low-and middle-income countries, high impact interventions without much resources. *Clinical Microbiology Infection*. 2017; 23:216–218.
2. Akobi OA, Inyinbor HE, Emumwen EG, Ogedengbe SO, Uzoigwe EO, Okorie IE. Incidence of Urinary Tract Infection among Pregnant Women Attending Antenatal Clinic at Federal Medical Centre, Bida, Niger-State, North Central Nigeria. *American Journal of Infect Diseases and Microbiology*. 2015; 2(2):34-38.
3. Okonko IO, Ijandipe LA, Ilusanya OA, Donbraye-Emmanuel OB, Ejembi J, Udeze AO et al. Incidence of urinary tract infection (UTI) among pregnant women in Ibadan, South. Western Nigeria. *African Journal of Biotechnology*. 2009; 8(23):8-10.
4. Delzell J, Michael L. Urinary tract infection during pregnancy. *American journal of Physician*. 2015; 61:713-720.
5. Ayukekbong JA, Ntemgwa M, Atabe AN. The threat of antimicrobial resistance in developing countries: causes and control strategies. *Antimicrobial Resistance Infection Control*. 2017; 6(1):47.

6. Alemu A, Moges F, Tafess K, Kassu A, Anagaw B. (2019). Prevalence of urinary tract infection in pregnant women at University of Gondar Teaching Hospital, North West Ethiopia. BMC Research Notes. 2019; 10:11-16.
7. Olusanya O, Ogunledum A, Fakoya, TA. Asymptomatic significant bacteriuria among pregnant and non-pregnant women in Sagamu, Nigeria. West African Journal of Medicine. 2013; 2(1):27-33.
8. Boucher HW, Talbot G, Bradley, J. Bad bugs, no drugs: no ESKAPE! An update from the infectious diseases society of America. Journal of Clinical Infectious Disease. 2009; 48(1):1–12.
9. Santoso BI, Surya R, Yasmin FA, Irwinda R. The awareness of urinary tract infection management in pregnant women. A qualitative study. Journal of Obstetrics & Gynecology. 2018; 25:92–96.
10. Olamijulo JA, Adewale CO, Olaleye O. Asymptomatic bacteriuria among antenatal women in Lagos, Journal of Obstetrics and Gynaecology. 2016; 36:722-725.
11. Charan J, Biswas, T. How to calculate sample size for different study designs in medical research. Indian journal of psychological medicine. 2013; 35:121-126.
12. Cheesbrough, M. Medical laboratories manual for tropical countries. 2006; 2: 479.
13. Anejo-okopi JA. Bacterial and antibiotic susceptibility pattern of urinary tract infection isolated from asymptomatic and symptomatic diabetic patients attending tertiary hospital in Jos, Nigeria. Trends in Medicine. 2017; 17:1-5.
14. Nwadike VU, Ojide, CK, Kalu EI. Multidrug Resistant *Acinetobacter* Infection and their Antimicrobial Susceptibility Pattern in a Nigerian Tertiary Hospital ICU. African Journal of Infectious Diseases. 2014; 8(1):14-18.
15. Bashir A, Garba I, Aliero AA, Kibiya A, Abubakar MH, Ntulume I. Superbugs-related prolonged admissions in three tertiary hospitals, Kano State, Nigeria. Pan African Medical Journal. 2019;32(166):14-17.
16. Musyoki VM, Masika MM, Mutai W, Gitau W, Kuria A, Muthini C. Antimicrobial susceptibility pattern of *Acinetobacter* isolates from patients in Kenyatta National Hospital, Nairobi, Kenya. Pan African Medical Journal. 2019; 33:146.
17. Iregbu KC, Ogunsola FT, Odugbemi TO. Infections caused by *Acinetobacter* species in pregnant women and their susceptibility to 14 antibiotics in Lagos University Teaching Hospital, Lagos. West African Journal Medicine. 2012;21:226-229.
18. Egwu IH, Iroha IR, Mmaduabuchi EM, Peter IU, Nnabugwu CC, Ali CM et al. Antimicrobial Susceptibility Pattern and Molecular Identification of *Acinetobacter baumannii* in Alex Ekwueme Federal University Teaching Hospital Abakaliki, Nigeria. Journal of Pharmaceutical Research International. 2021; 33(44B):409-419.
19. Odewale G, Adefioye OJ, Ojo J, Adewumi FA, Olowe OA. Multidrug resistance of *Acinetobacter baumannii* in Ladoke Akintola University Teaching Hospital, Osogbo, Nigeria. European Journal of Microbiology and Immunology. 2016;6(3):238-243.
20. Nwachukwu E, Onyebuchi O, Michael O. Prevalence of urinary tract infections in pregnant women in Onitsha, Nigeria. Journal of Bacteriology & Mycology. 2018; 6(5):284-285.

21. Suraiya SJ, Afroz S, Rasheduzzaman MD, Uddin KH, Shamsuzzaman S. Prevalence and Antimicrobial Susceptibility Pattern of Klebsiella Pneumoniae Isolated from Various Clinical Specimens in a Tertiary Care Hospital in Bangladesh. *Medicine Today*. 2020; 32(2):95–99.
22. Damilola EW, Oluduro AO, Abike TO, Chukwudum CV. Phenotypic and molecular characterization of multiple-resistant gram-negative bacteria in urine of pregnant women attending antenatal clinic of Mother and Child hospital, Ondo, Nigeria. *African Journal of microbiology research*. 2021; 15(5):209-216.
23. Afunwa RA, Nwokediuko I, Okolie CE. Asymptomatic Urinary Tract Infection (Bacteriuria) in Pregnant Women attending Antenatal Clinics in a Community Hospital. *International journal of medical science & Technology*. 2017; 7(5):28-34.
24. Umeaku CN, Chimezie CC, Ozoh CN, Nwajiobi FO, Abana CC, Umeaku-Uzor UC. Isolation of Multi-Resistant Bacteria of Patients Suffering from Urinary Tract Infections in Owerri Metropolis, Imo State, Nigeria. *Acta Scientific Microbiology*. 2021; 9:3-16.
25. Anozie OB, Lawani OL, Esike CU, Mamah E, Ajah LO. Prevalence and Common Microbial Isolates of Urinary Tract Infection in Pregnancy; A Four Year Review in a Tertiary Health Institution in Abakaliki, South-East Nigeria. *American Journal of Clinical Medicine Research*. 2019; 4:25-28.
26. Okonko IO, Ijandipe LA, Ilusanya AO, Donbraye-Emmanuel OB, Ejembi J, Udeze AO. Detection of urinary tract infection among pregnant women in Oluyoro Catholic Hospital, Ibadan, South-Western Nigeria. *Malaysian Journal Microbiology*. 2020; 6(1):16-24.
27. Akobi O A, Inyinbor HE, Akobi EC, Emumwen EG, Ogedengbe SO, Uzoigwe E. et al. Incidence of Urinary Tract Infection among Pregnant Women Attending Antenatal Clinic at Federal Medical Centre, Bida, Niger-State, North Central Nigeria. *American Journal of Infect Diseases and Microbiology*. 2015;2(2):34-38.
28. Akerele J, Abhulimen P, Okonofua F. Prevalence of asymptomatic bacteriuria among pregnant women in Benin City, Nigeria. *Journal of Obstetric Gynaecology*. 2021; 21(2):141-144.
29. Amala SE, Nwokah, EG. Prevalence of asymptomatic Bacteriuria among pregnant women attending antenatal clinic in Port Harcourt Township, Nigeria and Antibiogram of isolated Bacteria. *American Journal of Biomedical Science*. 2021;7(2):125-133.
30. Akpan NS, Umoyen AJ, Luka TT, Esua IT, Okon AS, Effiong AU. Bacterial etiologic agents, prevalence and associated risk factors of asymptomatic bacteriuria among pregnant and non-pregnant women in primary health care centers in South-South Nigeria. *International Journal of Medical and Health Research*. 2019; 5:66-76.
31. Aminu KY, Aliyu UU. Asymptomatic Bacteriuria in Pregnant Women in the Antenatal Booking Clinic at Aminu Kano Teaching Hospital, Kano, Nigeria. *Open Journal of Obstetrics and Gynecology*. 2015;5:286-297.