

## Case report

### **PSEUDOMONAS AERUGINOSA MENINGITIS : A CASE REPORT**

#### **Abstract:**

AIM : To present a case of Pseudomonas meningitis with an atypical presentation.

CASE REPORT : A 48 year old male with history of RTA related brain injury and shunt placement , presented with complaints of diarrhoea. On further investigations he was found to have Gram negative meningitis on CSF studies. This case highlights patient's course in hospital.

CONCLUSION : Patient's with prior history of neurosurgical intervention are at higher risk of meningitis due to Gram negative organisms like Pseudomonas aeruginosa and they may present with atypical symptoms.

*KEY WORDS : Chronic diarrhea , Meningitis , Gram negative, Pseudomonas aeruginosa, Ventriculo-peritoneal shunt, Craniectomy, Candidemia ,*

#### **Introduction :**


Chronic meningitis is meningitis lasting more than four weeks, which can be infectious or non-infectious. *Pseudomonas aeruginosa* , a Gram-negative bacillus is a rare cause of meningitis. Recent case series have brought to light a pattern of patients developing meningitis post neurosurgical procedures<sup>(2)</sup> The possibility of a ventriculoperitoneal (VP) shunt infection is most likely to be in the earlier days after shunt insertion. Between 56% and 87% of infections occur within one month of shunt insertion<sup>(1)</sup> The following is a case report discussing course and management of patient with a rather intriguing presentation

#### **Case report :**

A 48 year old male patient, was a known case of Traumatic Brain injury and Type 2 Diabetes mellitus, who presented with loose stools. Previous surgical procedures included Craniectomy done 3 months before admission followed by Ventriculoperitoneal shunt placement and cranioplasty 1 month after previous surgery. He had a prolonged hospital stay of 3 months requiring multiple antibiotics. His presenting complaint was 10-12 episodes of loose stools per day since one month. On initial examination, vitals were stable and Glasgow Coma Scale (GCS) was E4VTM1, generalized rigidity was present and abdomen was soft. Blood investigations showed anemia with neutrophilic leukocytosis , elevated inflammatory markers and dyselectrolytemia. (TABLE 1)

TABLE 1: INITIAL BLOOD TESTS			
ESR	67	UREA	14.2
CRP	55	CREATININE	0.8
HEMOGLOBIN	9	PTINR	0.98
WBC COUNT	12000	APTT	29
PLATELET	673	HBA1C	6.7
POTASSIUM	2.5	ALBUMIN / GLOBULIN	3
MAGNESIUM	1.1	AST / ALT	22 /9.6
PHOSPHORUS	1.8	TOTAL BILRUBIN	1.2
CALCIUM	7.6	DIRECT BILRUBIN	0.7
SODIUM	133	ALP	140

FIGURE 1 : MRI BRAIN WITH CONTRAST – Hydrocephalus involving bilateral lateral ventricles.



In view of background of Traumatic brain injury associated with decreased sensorium, **Magnetic Resonance Imaging (MRI)** brain with contrast was done. (FIGURE 1) The study showed hydrocephalus involving bilateral lateral ventricles and third ventricle with relative sparing of fourth ventricle possibly aqueductal stenosis. Post contrast meningeal enhancement with a thin rim of collection was also noted along left frontoparietal convexity. Patient had two episodes of generalized tonic clonic seizures, each lasting for 2 minutes. **Cerebrospinal Fluid (CSF)** studies revealed elevated total cell count (4,320) with 85% segmented cells and CSF culture grew *Pseudomonas aeruginosa* (TABLE 2). He was started on Antibiotics Cefepime 2g IV Q8H and Amikacin 5mg/kg IV Q8H according to the CSF culture and sensitivity reports. Blood cultures

showed no growth. Few days later, a discharging sinus was noted over the left clavicle. USG of the region showed hypo echoic collection noted around and along **Ventriculoperitoneal (VP)** shunt and fistula between the VP shunt and the skin. Pus swab grew *Pseudomonas aeruginosa*. Neurosurgery was consulted and a joint decision to do VP shunt removal was taken. During procedure, pus was present in the ventricular and abdominal end of the tube. Catheter tip from ventricle and abdominal end were growing *Pseudomonas aeruginosa*. During hospital stay, he developed Lower Respiratory Tract Infection with Mini Bal culture growing *Pseudomonas aeruginosa*. As total cells in CSF were downtrending, patient was taken up for Endoscopic third ventriculostomy. However, patient's sensorium remained status quo. As there was no further rise in the inflammatory markers, antibiotics were de-escalated. As part of the evaluation of recurrent fever spikes, CT Chest was done and showed thrombus in the Right Internal Jugular vein. His blood culture grew *Candida tropicalis* (Candidemia). In spite of the multi-disciplinary approach and aggressive management, he eventually succumbed to his illness.

TABLE 2: CSF REPORTS			
CSF Studies	Total count (Differential Count)	Protein	Sugar
1 <sup>st</sup> CSF	4320 (85% SEGMENTED CELLS , 15% MONONUCLEAR CELLS)	237	26
2 <sup>nd</sup> CSF	60 (2% SEGMENTED , 10% MONONUCLEAR)	60	123

### **Discussion:**

Gram-negative bacterial infections of the central nervous system CNS (commonly *Escherichia Coli*, *Enterobacter* species, *Serratia* species, and *Pseudomonas aeruginosa*) have worse clinical outcomes. The clinical presentation of a ventriculoperitoneal (VP) shunt infection can include the signs and symptoms of meningitis to fever with abdominal pain and peritonitis.<sup>(4)</sup> Previous studies have indicated that patients with GNB CSF shunt infections often appear relatively well at presentation<sup>(5)</sup>. Initial broad spectrum empirical intravenous antimicrobial therapy with appropriate Gram-negative coverage therapy has shown good outcomes with decreased mortality. *Candida tropicalis* Candidemia can also cause mortality especially in long term hospitalized patients

**(6)**

### **Conclusion:**

Patients with neurosurgical intervention are at higher risk of developing GNB meningitis. It is vital to keep an open mind when such patients present with atypical symptoms also. Although there is high mortality associated with Gram negative meningitis, timely

diagnosis and treatment proves to be associated with positive outcomes. A multi-disciplinary approach is crucial in such patients.

**Consent:**All authors declare that informed consent was obtained from the patient (or other approved parties) for publication of this case report.

### References :

1. Ahmad F, Brubaker M, Rajendraprasad SS, Hoeynck B, Clyde BL, Velagapudi M. Challenges in the Management of Gram-Negative Bacterial Infections in Patients With Ventriculoperitoneal Shunt. *Cureus*. 2021 Aug 9;13(8):e17035. doi: 10.7759/cureus.17035. PMID: 34522514; PMCID: PMC8425579.
2. Gallaher C, Norman J, Singh A, Sanderson F. Community-acquired *Pseudomonas aeruginosa* meningitis. *BMJ Case Rep*. 2017 Oct 19;2017:bcr2017221839. doi: 10.1136/bcr-2017-221839. PMID: 29054951; PMCID: PMC5665197.
3. Tunkel AR, Hartman BJ, Kaplan SL, Kaufman BA, Roos KL, Scheld WM, et al. Practice Guidelines for the Management of Bacterial Meningitis. *Clinical Infectious Diseases [Internet]*. 2004 Nov 1;39(9):1267–84.
4. Charalampoudis, P. (2012, January 1). Right Lower Quadrant Abdominal Pain in a Patient with Prior Ventriculoperitoneal Shunting: Consider the Tip! *Case Reports in Medicine*. <https://doi.org/10.1155/2012/253027>. Consulted the .../.../.....
5. Stamos JK, Kaufman BA, Yogev R. Ventriculoperitoneal shunt infections with gram-negative bacteria. *Neurosurgery*. 1993 Nov;33(5):858-62. doi: 10.1227/00006123-199311000-00011. PMID: 8264883.
6. Charalampoudis, P. (2012, January 1). Right Lower Quadrant Abdominal Pain in a Patient with Prior Ventriculoperitoneal Shunting: Consider the Tip! *Case Reports in Medicine*. <https://doi.org/10.1155/2012/253027> . Consulted the... /... / 2....