

Original Research Article

Studies on response of ber (*Ziziphus mauritiana* L.) cv. Umranto fertilizers, micronutrients and bioregulatorson flowering and fruit growth & developments attributes.

ABSTRACT:

Abstract: The investigation entitled “Studies on response of ber (*Ziziphus mauritiana* L.) cv. Umran to fertilizers, micronutrients and bio regulators on “flowering and fruit growth & developments attributes” was carried out during 2019-20 at Fruit Research Station, Imalia, Department of Horticulture JNKVV Jabalpur. The experiment was laid out under randomized complete block design (RCBD) with 3 replications having 12 treatments comprising fertilizers, micronutrients, bioregulators either alone or in different combinations. On the basis of result obtained in present investigation, it is concluded that the application of treatment T₄ [RDF (N:P:K)+Foliar spray of GA₃ + NAA (50 ppm+50 ppm)] was found significantly superior as compared to other treatments for flowering parameter in respect to induce early flower bud formation. The earliest date of flower bud initiation (11.09.2019) and days taken to induce earliest flowering after spray is (22 days), days taken for 50% flowering (8.67).The treatment T₁₁ [RDF (N:P:K)+ NPK -18:18:18 (1.0%) + ZnSO₄ (0.5%) + Borax (0.3%)] proved best for fruit growth and development attributes viz. fruit length (4.53 cm), fruit diameter (3.50 cm), fruit weight (36.52 g), fruit volume (36.56 ml), specific gravity (1.047).The treatment T₁₀ was found next best in respect to almost all the fruit growth and development attributes.

INTRODUCTION:

The Indian jujube, commonly known as ber, is a member of the Rhamnaceae family. It is often referred to as the "King of arid-zone fruits" or the "poor man's apple." The genus *Ziziphus* gets its name from the Arabic word "Zizaif," which is the name of the fruit. The Rhamnaceae family itself comprises approximately 50 genera and more than 600 species that are distributed across tropical and sub-tropical regions in the northern hemisphere (Pareek, 1983). *Ziziphus mauritiana* Lamk. has its origins in India, while *Ziziphus vulgaris* Lamk. is native to China. These two species have been commercially cultivated and cherished in both India and China for centuries and are among the most commercially significant varieties.

The ber tree exhibits rapid growth, characterized by a wide-spreading canopy and a relatively short trunk. Its branches are slender, covered in a downy

texture, and often feature pairs of brown, sharp spines. This tree is exceptionally resilient to drought conditions, owing to several xerophytic adaptations.

The ber fruit stands out as a fruit crop that can yield favourable returns even when grown under rainfed conditions. This is attributed to its remarkable adaptability to a wide range of soils, varying water availability conditions, and diverse climates, although it is sensitive to heavy frosts. It thrives particularly well in arid and semi-arid regions. This adaptability makes it a resilient and economically viable option for cultivation in areas with challenging environmental conditions.

The ber fruit boasts a spongy, sweet, and delicious pulp that is not only flavourful but also packed with essential nutrients. It is known to be an excellent source of vitamins, including vitamin C, vitamin A, and various B vitamins.

Chemical nutrients and bio-regulators, even in small quantities, play a crucial role in promoting the growth and development of plants, ultimately influencing both yield and quality. These substances can bring about changes in plant metabolism by altering the nutritional and hormonal status of the plant (Gadi and Bohra, 2005). 1-Naphthaleneacetic acid (NAA) is an important plant hormone reported to enhance the fruit set, growth, retention, yield and market price of some fruit species also delayed in fruit ripening as well as enhancing fruit formation through cell division and elongation (Dutta and Banik, 2007). Gibberellins are reported to increase fruit set, size, retention and yield as well as improve fruit physico-chemical characteristics and ripening (Rizk-Alla *et al.*, 2011). In addition to plant hormones, micronutrients such as boron (B) and zinc (Zn) have proven to be beneficial in ber fruit cultivation, (Samant *et al.* in 2008) demonstrated that these micronutrients can positively impact fruit set, yield, fruit quality, and the storage life of ber fruits.

MATERIALS AND METHODS

The present investigation was carried out under Kymore plateau and Satpura hill regions during 2019-20 at Fruit Research Station Imalia, Department of Horticulture, College of Agriculture JNKVV Jabalpur, Madhya Pradesh, on Ber cv. Umran and the trees were maintained under a uniform cultural schedule. The experiment was laid out in RBD (Randomized block design) comprising 12 treatment combinations and was replicated thrice. Treatments were given twice i.e., first, before flowering and second, at fruit setting stage. The following

treatment combinations are as follows: T₀: Control, T₁:RDF (N:P:K) (500:250:250) – Soil application, T₂:RDF(N:P:K) + Foliar spray of GA₃ (50 ppm), T₃: RDF(N:P:K) + Foliar spray of NAA (50 ppm), T₄:RDF(N:P:K) + Foliar spray of GA₃ + NAA (50 ppm + 50 ppm), T₅: RDF (N:P:K) + Foliar spray of KNO₃ (1.0 %), T₆: RDF (N:P:K) +Urea (1.5 %) + ZnSO₄ (0.50%), T₇: RDF (N:P:K) + Urea (1.5 %) + ZnSO₄(0.50%) + Borax(0.50%), T₈: RDF (N:P:K)+ Urea (1.5%)+ZnSO₄ (0.5%)+ Borax (0.5%)+CuSO₄ (0.3%), T₉:RDF (N:P:K) + NPK – 18:18:18 (1.0 %), T₁₀: RDF (N:P:K) + NPK -18:18:18 (0.5%) + ZnSO₄ (0.5%) + Borax (0.5%) and T₁₁:RDF (N:P:K) + NPK -18:18:18 (1.0%) + ZnSO₄ (0.5%) + Borax (0.3%).

Preparation of solutions

Solutions of NPK, KNO₃, Urea, ZnSO₄, and Borax were prepared by dissolving them in distilled water to obtain the required concentration as mentioned above. GA₃ and NAA solutions were prepared in distilled water after dissolving them in the required amount of alcohol.

Spraying of chemicals & growth regulators

The first sprayings were done at the time of RDF application before flowering i.e. (20.08.2019). The second spraying was given at the fruit setting stage. The spraying was done with a foot sprayer.

1. Flowering parameters

1.1 Date of flower bud initiation

Five uniform branches around each tree were selected and tagged. The date of flower bud initiation was noted in each tagged branch from the date of foliar application of treatments.

1.2 Days taken for flowering (50%)

Days taken to 50 % flowering in each treatment were counted after the first flowering in the tagged shoots based on visual observation on all four directions tagged branches of each treated plant.

2. Fruit growth & development attributes.

2.1 Fruit length (cm)

The length of each fruit from stalk end to stylar end was measured with the help of vernier calipers at 60, 90, 120, 150 days after the fruit set and the average length of fruit was calculated.

2.2 Fruit diameter (cm)

The diameter of each fruit was measured at the maximum thickness of the fruit by vernier calipers at 60, 90, 120, 150 days after the fruit set, and the average diameter of the fruit was worked out.

2.3 Fruit weight (g)

The average weight of the fruit was calculated after the final picking as per the formula given below:

$$\text{Average fruit weight} = \text{Total weight of fruits (g)} / \text{Number of fruits}$$

2.4 Fruit volume (ml)

Fruits were placed in measuring cylinders of 1-liter capacity filled with water. The replaced water was measured, and the data were recorded as the volume of fruits in ml, then the average value was computed.

2.5 Specific gravity

The specific gravity of the fruit was determined by weighing fruits from each treatment of each replication in top pan balance and then recording displaced water after sinking the fruits in water. The specific gravity was calculated by dividing the weight of the fruit by the volume of the fruit.

RESULTS AND DISCUSSION

The results obtained during the investigation were statistically analyzed and presented under the following heads.

1.0 Flowering parameters

The data pertaining to the Date of flower bud initiation and Days taken for 50% flowering of different treatments are presented in (Table 1). The investigations revealed that the various treatments have influenced this trait greatly. The earliest date of flower bud initiation (11.09.2019) and days taken to induce earliest flowering after spray is (22) and the minimum days to 50% flowering (8.33) were recorded in treatment T₄ [RDF (N:P:K)+Foliar spray of GA₃ + NAA (50 ppm + 50 ppm)] followed by treatment T₁₁ [RDF (N:P:K) + NPK -18:18:18 (1.0%) + ZnSO₄ (0.5%) + Borax (0.3%)], whereas the maximum day taken for flower bud initiation (19.09.2019) and the maximum days (13.67) to 50% flowering was recorded in T₀ (Control).

The early onset of flowering may be attributed to the influence of GA₃, which enhances leaf production and encourages vegetative growth. This boosts photosynthetic rates, leading to increased carbohydrate accumulation and

ultimately the initiation of flower development, as indicated by Ujjwal *et al.* (2018). Additionally, NAA may contribute to the enlargement of cells, further supporting this process. These findings are in line of, Karole and Tiwari, (2016), Parouss *et al.* (2002), Kacha *et al.* (2014).

2.0 Fruit growth & development attributes.

The data about fruit growth & development attributes like fruit length, fruit width, fruit weight, fruit volume, stone length, stone diameter, and stone weight, were observed under various treatments. The investigations revealed that the various treatments have influenced this trait greatly. The maximum fruit length (2.94 cm), (3.59 cm), (3.84 cm), (4.53 cm) at 60,90,120,150, days respectively after the fruit set presented in (Table 2), the maximum fruit diameter (1.85 cm), (2.30 cm), (3.36 cm), (3.50 cm), at 60,90,120,150 days respectively after fruit set presented in (table 3), maximum fruit weight (36.52 g), fruit volume (36.56 ml) and specific gravity (1.047) presented in (table 4) were recorded with the application of treatment T₁₁ [RDF (N:P:K) + NPK -18:18:18 (1.0%) + ZnSO₄ (0.5%) + Borax (0.3%)].

The minimum values of fruit length (2.62 cm), (3.18 cm), (3.52 cm), (3.83 cm) at 60,90,120,150 days respectively after fruit set, fruit diameter (1.56 cm), (1.93 cm), (2.85 cm), (2.99 cm) at 60,90,120,150 days respectively after fruit set, minimum fruit weight (27.50 g), fruit volume (27.22 ml), and specific gravity (0.899 gm/ml) was observed under control T₀, which was at par with treatments T₁₀, T₈, T₇, T₆, T₄, and T₂.

This phenomenon could be attributed to nitrogen, phosphorus, and potash stimulating chlorophyll synthesis and enhancing photosynthetic activity. This, in turn, leads to increased storage of food material in the tissue, ultimately resulting in larger fruit size, as documented by Jat and Kacha (2014). These findings align with Prasad (2005) Dalal *et al.* (2011), Dhayal *et al.* (2011), Mishra *et al.* (2011), and Gill and Singh, (2011).

Boron boosts nitrogen uptake, facilitating photosynthesis and leading to carbohydrate accumulation, ultimately resulting in increased fruit size, as reported by Kamble *et al.* (1994), Sharma *et al.* (2008), Kamble *et al.* (1994), Kumar and Shukla (2010), Singh *et al.* (2016), Nehete *et al.* (2011).

Moreover, Zinc enhances fruit size by stimulating auxin production and improving sugar metabolism. It increases fruit length, diameter, fruit weight and

specific gravity possibly through cell wall regulation, leading to larger and heavier fruits. Zinc maintains elevated auxin levels in fruit parts, and its role in auxin production is well-established. This results in greater fruit width and length due to enhanced water uptake, as reported by Pippal *et al.* (2019). These findings are inline with Joon *et al.*(1984), Kamble *et al.* (1994), Singh and Vashistha (1997), Kamble *et al.* (1994) , Sharma *et al.* (2008), Sharma *et al.* (2011), Pandey *et al.* (1988), Nehete *et al.* (2011), Chandra and Singh (2015), Gami *et al.* (2019).

Table 1. Effect of fertilizers, micronutrients and bio regulators on Date of flower bud initiation, Days to (50%) flowering, initial fruit set percentage and fruit retention percentage.

Treatments	Date of flower bud initiation	Days to flowering 50 %
T0	19.09.2019	13.67
T1	17.09.2019	11.67
T2	14.09.2019	10.67
T3	12.09.2019	9.00
T4	11.09.2019	8.33
T5	15.09.2019	11.67
T6	13.09.2019	10.33
T7	13.09.2019	9.67
T8	12.09.2019	9.33
T9	15.09.2019	11.33
T10	11.09.2019	9.00
T11	11.09.2019	8.67
SEm±	0.79	0.59
C.D. at 5%	2.30	1.71

Table 2. Effect of fertilizers, micronutrients, and bio regulators on fruit length (cm) at different days of growth and development period at (60, 90,120,150 days)

Treatments	Fruit length (cm) at different days			
	60 days	90 days	120 days	150 days
T0	2.62	3.18	3.52	3.83
T1	2.65	3.22	3.55	3.87
T2	2.80	3.37	3.70	4.15
T3	2.78	3.34	3.68	4.12

T4	2.82	3.40	3.72	4.22
T5	2.73	3.32	3.63	4.09
T6	2.84	3.43	3.75	4.25
T7	2.85	3.49	3.78	4.29
T8	2.87	3.53	3.79	4.46
T9	2.70	3.29	3.59	4.05
T10	2.89	3.56	3.81	4.49
T11	2.94	3.59	3.84	4.53
SEm±	0.05	0.077	0.04	0.11
CD at 5%	0.15	0.22	0.14	0.34

Table 3 Effect of fertilizers, micronutrients and bioregulators on fruit Diameter at different days of growth and development period at (60, 90,120,150 days)

Treatments	Fruit Diameter (cm) at different days			
	60	90	120	150
T0	1.56	1.93	2.85	2.99
T1	1.59	1.84	2.98	3.12
T2	1.71	2.13	3.18	3.32
T3	1.68	2.10	3.15	3.19
T4	1.72	2.16	3.20	3.34
T5	1.65	2.05	3.12	3.26
T6	1.74	2.20	3.23	3.37
T7	1.77	2.23	3.25	3.41
T8	1.78	2.25	3.30	3.44
T9	1.62	2.02	3.10	3.24
T10	1.82	2.26	3.34	3.48
T11	1.85	2.30	3.36	3.50

SEm±	0.037	0.049	0.10	0.09
CD at 5%	0.10	0.144	0.29	0.26

Table 4 Effect of fertilizers, micronutrients and bioregulators on fruit weight, fruit volume and specific gravity

Treatments	Fruit weight (gm)	Fruit volume (ml)	Specific gravity(gm/ml)
T0	27.30	27.22	0.899
T1	29.30	28.23	0.923
T2	31.85	32.99	0.965
T3	31.28	32.36	0.966
T4	32.00	34.19	0.935
T5	30.72	31.82	0.965
T6	32.70	35.01	0.934
T7	33.90	35.06	0.966
T8	33.98	35.12	0.967
T9	29.88	30.32	0.929
T10	34.48	35.62	0.998
T11	36.52	36.56	1.047
SEm±	1.05	1.04	0.022
CD at 5%	3.04	3.03	0.065

References :

- Bhatia, S.K., Ahlawat, V.P., Gupta, A.K. & Rana, G.S., (2001). Physicochemical attributes of guava as affected by nitrogen application. *Haryana Journal of Horticultural Sciences*. **30**(1&2): 65-66.
- Chandra, R., & Singh, K.K., (2015). Foliar application of zinc sulphate, magnesium sulphate and copper sulphate on the yield and quality of aonla (*Emblia officinalis* Gaerth L.) cv. "NA- 7" under Garhwal Himalaya. *Himalaya Journal of Medicinal Plants Studies*. **3**(5): 42-45.
- Dalal, R.P.S., Navjot, A.T., & Brar, J.S., (2011). Effect of Foliar Application of Nutrients on Leaf Mineral Composition and Yield of Ber (*Ziziphus mauritiana* Lamk.) under Arid Conditions. *Annals of Arid Zone*. **50**(1): 53-56.
- Dayal, H., Lal, G., Singh, Y.V., Kumar, P., & Singh, D. (2010). Effect of nitrogen, phosphorus and zinc on growth and yield of ber cv. Gola under arid and semi-arid conditions. *Indian Journal of Horticulture*. **67**(2): 277-280.
- Dutta, P., & Banik, A.K., (2005). Effect of foliar feeding of nutrients and plant growth regulators on physico-

- chemical quality of Sardar guava grown in red and lateritic tract of West Bengal. *In International Guava Symposium* **735**: 407- 411
- Gadi, B.R.,& Bohra, S.P.,(2005). Effect of plant growth regulators on photosynthesis and some biochemical parameters in ber cv.Gola. *Indian Journal of Horticulture*. **62**(3):296-297.
- Gami, J., Sonkar, P., Haldar, A.& Patidar, D.K. (2019). Effect of Pre harvest Spray of ZnSO₄, KNO₃ and NAA on Growth, Yield and Quality of Ber (*Zizyphus mauritiana* Lamk.) cv. Seb under Malwa Plateau Conditions. *Int. J. Curr. Microbiol. App. Sci.* **8**(3): 1977-1984.
- Ghosh, S.N., Bera, B., Kundu, A.& Roy, S. (2008). Effect of plant growth regulators on fruit retention, yield and physico-chemical characteristics of fruits in ber 'Banarasi Karka' grown in close spacing. *In International Jujube Symposium*. **840**: 357- 362.\
- Haridayal, P.R., Singh, D.& Lal, G. (2011). Response of ber to nutrient management in relation to yield and physico-chemical composition of fruits under semi-arid condition. *Annals of Arid Zone*. **50**(1): 1-4.
- Iqbal, M., Khan, M.Q., Rehman, K.& Munir, M. (2009). Effect of foliar application of NAA on fruit drop, yield and physico-chemical characteristics of guava (*Psidium guajava* L.) red flesh cultivar. *Journal of Agricultural Research*, **47**(3): 207- 210.
- Jadhao, B.J., Ghawade, S.M., Nandre, D.R., Dalal, S.R.& Jogdande, N.D. (2005). Integrated nutrient management in guava cv. 'Sardar'. *In 1st Guava International Symposium*.
- Jat, G.& Laxmidas, K.H. (2014). Response of guava to foliar application of urea and zinc on fruit set, yield and quality. *Journal of agrisearch*. **1**(2): 157-162.
- Joon, M.S., Singh, R.R.& Daulta, B.S. (1984). Effect of foliar spraying of zinc and urea on yield and physico-chemical composition of ber fruit cv. Gola. *Haryana Journal of Horticultural Sciences*. **13**(3): 110-112.
- Kacha, H.L., Viradia, R.R., Leua, H.N., Giriraj, J.& Tank, A.K. (2012). Effect of NAA, GA₃ and ethrel on yield and quality of phalsa (*Grewia asiatica* L.) under South- Saurashtra condition. *Asian Journal of Horticulture*. **7**(2): 242-245.
- Kamble, A.B., Desai, U.T.& Choudhari, S.M. (1994). Effect of micronutrients on fruit set, fruit retention and yield of ber (*Zizyphus mauritiana* Lamk). *Annals of Arid zone*. **33**(1):53-55.
- Karole, B.& Tiwari, R. (2016). Effect of pre-harvest spray of growth regulators and urea on growth, yield and quality of ber under malwa plateau conditions. *Annals of Plant and Soil Research*. **18**(1): 18-22
- Katiyar, P.N., Yadav, V.& Singh, J.P. (2010). Effect of preharvest spray of NAA, GA₃ and Urea on fruiting, fruit quality and yield of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. *Annals of Horticulture*. **3**(1): 92-94.
- Kumar, S.& Shukla, A.K. (2010). Improvement of old ber cv. Gola orchard through bunding and micro-nutrients management. *Indian Journal of Horticulture*. **67**(3): 322-327.
- Kundu, S.& Mitra, S.K. (1999). Response of guava to foliar spray of copper, boron and zinc. *Indian Agric*. **43**(1): 49-54.
- Madhavi, A.M., Prasad, I., Prabhakar, R. and Girwani, A. (2005). Integrated nutrient management for increased productivity and quality in guava. *1st International Guava Symposium*. Dec. 5- 8. CISH. Lucknow.
- Meena, V.S., Yadav, P.K.& Meena, P.M. (2008). Yields attributes of Ber (*Zizyphus mauritiana* Lamk.) cv. gola as influenced by foliar application of ferrous sulphate and borax. *Agricultural Science Digest*. **28**(3): 219-221.
- Mishra, S., Choudhary, M.R., Yadav, B.L.& Singh, S.P. (2011). Studies on the response of integrated nutrient management on growth and yield of ber. *Indian Journal of Horticulture*. **68**(3): 318-321.
- Nehete, D.S., Padhiar, B.V., Shah, N.I., Bhalerao, P.P., Kolambe, B.N.& Bhalerao, R.R. (2011). Influence of micronutrient spray on flowering, yield, quality and nutrient content in leaf of mango cv. KESAR. *Asian Journal of Horticulture*. **6**(1): 63- 67.
- Pandey, D.K., Pathak, R.A.& Pathak, R.K. (1988). Studies on the Foliar Application of Nutrients and Plant Growth Regulators in Sardar Guava (*Psidium Guajava* L.) Effect on Yield and Fruit Quality. *Indian Journal of Horticulture*, **45**(3): 197-202.
- Pareek, O.P. (1983). *The Jujube*. *Indian Council of Agricultural Research*. New Delhi.
- Paroussi, G., Voyiatzis., D.G., Paroussis, E.& Drogoudi, P.D. (2002). Growth, flowering and yield responses to GA₃ of strawberry grown under different environmental conditions. *Scientia Horticulturae*. **96**(1): 103-113.
- Pippal, R., Lekhi, R., Barholia, A.K., Rana, S.& Rana, P. (2019). Response of guava to foliar spray of zinc, boron and magnesium on growth, development and yield. *J. Pharmac. Phytochem*. **36**(2): 942-946.
- Prasad, R.N. (2005). Effect of N and P on growth, yield and quality of ber grown under rainfed conditions of Indian arid zone. *Indian Journal of Horticulture*. **62**(4): 404-406.
- Rizk-Alla, M.S., El-Wahab, M.A.& Fekry, O.M. (2011). Application of GA₃ and NAA as a means for improving

- yield, fruit quality and storability of Black Monukka grape cv. *Nature and Science*. **9**(1): 1-19.
- Samant, D., Mishra, N.K., Singh, A.K.& Lal, R.L. (2008). Effect of micronutrient sprays on fruit yield and quality during storage in ber cv. Umran under ambient conditions. *Indian Journal of Horticulture*. **65**(4): 399-404.
- Sharma, J.R., Bhatia, S.K., Gupta, R.B., Surender, S.& Rakesh, G. (2008). Influence of nutrients and growth regulators on yield and quality of ber (*Zizyphus mauritiana* Lamk.) fruits cv. Umran. *Haryana Journal of Horticultural Sciences*.**37**(3): 217-219.
- Singh, C.& Bal, J.S. (2006). Effect of nutrients and growth regulators on fruit drop, size and yield of ber (*Zizyphus mauritiana* Lamk.). *Int. J. Agric. Sci.***2**(2): 358-360.
- Singh, V.K., Shanker, K., Tiwari, N.K.& Rao, O.P. (2016). Effect of certain nutrients on fruit set, fruit retention, physical characters and yield of ber fruit (*Zizyphus maurifiana* Lamk). cv. Banarasi Karaka. *International Journal of Bio-Resource and Stress Management*.**7**(1): 648-652.
- Singh, R.S., Vashistha, A. (1997). Effect of foliar spray of nutrient on fruit drop, yield and quality of ber cv. Seb. *Haryana Journal of Horticulture Science*. **26**(1-2): 20-24.
- Ujjwal, V., Singh, M.K., Dev, P., Chaudhary, M., Kumar, A.& Tomar, S. Effect of different levels of GA₃ and NAA on vegetative growth and flowering parameters of tomato (*Solanum lycopersicum* L.).
- Wang, Y., Tao, T., Liao, P.& Deng, Q. (2018). Effects of GA₃, NAA and PCPA on the Fruit Drop Rate and Quality of *Zizyphus jujuba* 'Zhanshanmizao'. In 7th International Conference on Energy and Environmental Protection (ICEEP 2018). *Atlantis Press*.
- Yadav, A.K., Singh, J.K.& Singh, H.K. (2011). Studies on integrated nutrient management in flowering, fruiting, yield and quality of mango cv. Amrapali under high density orcharding. *Indian Journal of Horticulture*. **68**(4): 453-460.
- Yadav, S., Shukla, H.S.& Ram, R.A. (2010). Studies on Foliar Application of NAA, GA, Boric acid, and Ca(NO) on Fruit Retention, Growth, Yield and Quality of Aonla (*Emblica officinalis* Gaertn.) cv. Banarasi. *The Hort, J.*, **Vol23**(2)

UNDER PEER REVIEW