

**Studies on integrated Nutrient Management in Aggregatum Onion (*Allium cepa* var. *aggregatum*)**

**ABSTRACT**

A field experiment was conducted in the Department of Horticulture, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal, U.T. of Puducherry in aggregatum onion [*Allium cepa* var. *aggregatum*] cv 'Perambalur Local', during summer 2023 with the objective of comparing the integrated effects of organics and bioenhancers on growth, bulb yield and quality of aggregatum onion. The experiment was laid out in a Randomized Block Design with two replications involving two factors viz., fertilisers level ( $F_1$  with 6 levels) and bioenhancers ( $F_2$  with 3 levels) forming, eighteen treatments in a factorial way (FRBD). The study material comprised of vermicompost, poultry manure, panchagavya and jeevamirtham, besides recommended FYM and N, P, K fertilisers. Maximum plant height at 30 DAP (31.40 cm), leaf girth at 30 DAP (1.94 cm), number of leaves plant<sup>-1</sup> at 30 DAP (25.80), fresh weight of top (12.26 g), polar (3.17 cm) and equatorial diameter of bulblet (2.94 cm), bulblet weight (9.30 g), number of bulblets plant<sup>-1</sup> (10.00), maximum bulb yield (43.18 g plant<sup>-1</sup> and 6.91 t ha<sup>-1</sup>) and dry matter production of bulbs (788.51 kg ha<sup>-1</sup>) were recorded in the treatment receiving RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure along with the application of jeevamirtham 500 L ha<sup>-1</sup> as soil drench thrice with irrigation viz., at planting, 20 and 45 DAP.

**Keywords:** Vermicompost; Poultry manure; Panchagavya; Jeevamirtham.

**1. INTRODUCTION**

Vegetables are considered essential for providing a balanced diet as they constitute phytonutrients, besides nutraceutical compounds. Being rich sources of carbohydrates,

proteins, vitamins and minerals, vegetables are regarded as protective foods (Singh and Devi 2015). Globally, India ranks second in vegetable production next to China, producing 209.14 m MT of vegetables annually, from an area of 11.37 m ha (Hort. Stat., 2022).

Aggregatum onion (*Allium cepa* var. *aggregatum*), also known as small onion or country onion or multiplier onion is one of the most important commercial bulbous vegetables of the family Alliaceae, widely cultivated in Maharashtra, Madhya Pradesh, Karnataka and Tamil Nadu. Globally, India ranks second in onion production with 31.68 m MT of onion being produced from an estimated area of 1.94 m ha. (Hort. Stat., 2022).

Onion is considered to be the second most important vegetable crop grown in the world after tomato with multiple uses such as vegetable and condiment, hence become an indispensable part of Indian kitchen (Kaur *et al.*, 2015). However, the productivity of onion in India is very low ( $18.64 \text{ t ha}^{-1}$ ) compared to the world's average productivity of  $23.06 \text{ t ha}^{-1}$  and imbalance in use of chemical fertilizers is considered one of the prime cause of reduced productivity and a possible alternative is to use organics as substitutes. Organic farming is a holistic approach capable of improving soil fertility and productivity in a sustained way. Further, organic produces fetch better price in the national and international market due to consumer preference, as organic manures not only sustain the productivity but also improves the quality of the produce, besides reducing the cost of cultivation (Sruthi and Somasundaram, 2019; Kale *et al.* 2022), thereby enhancing the net profit.

Organic substances along with poultry manure, vermicompost and farm yard manure can complement inorganic fertilizers to maintain productivity and natural quality (Tripathy *et al.*, 2017). The present study was hence contemplated to determine the effect of organic sources of nutrients and bioenhancers on growth, yield and quality of aggregatum onion cv. Perambalur Local.

## **2. MATERIALS AND METHODS**

A field experiment entitled "Studies on nutrient management in aggregatum onion (*Allium cepa* var. *aggregatum*)" was carried out in the Department of Horticulture, Pandit Jawaharlal Nehru College of Agriculture and Research Institute, Karaikal, U.T. of Puducherry, India,

during summer 2023. Aggregatum onion type 'Perambalur Local' collected from farmer's field in Padalur village of Perambalur district of Tamil Nadu (India) was used for the study. The treatment materials for the study comprised of vermicompost, poultry manure, panchagavya and jeevamirtham, besides recommended FYM and N, P, K fertilisers.

Panchagavya, an ancient vedic formulation is a blend of five products namely the dung, urine, milk, curd and ghee obtained from indigenous cow. It plays an important role in the overall growth and development of crop plants and increase of yield and profits. It is also eco- friendly (Kumar and Singh 2020). Jeevamrutham is an organic fertilizer capable of enhancing microbial activity in soil and helps in improvement of soil fertility (Kaur, 2020).

In the present experiment, panchagavya was applied as foliar spray at 3 per cent concentration on 15, 30 and 45 days after planting, while freshly prepared jeevamirtham was applied in the soil at planting, 20 and 40 days after planting along with irrigation water.

The experiment was laid out in a Randomized Block Design with two replications with eighteen treatments (Table 1) in a factorial way (FRBD) and the study comprised of 2 factors viz., Fertiliser (Factor 1) and Bioenhancers (Factor 2). There were six levels for Factor 1 and three levels for Factor 2, forming 18 treatment combinations.

## **2.1 Level of factors**

### **2.1.1. Factor 1 – Fertilisers – 6 (F<sub>0</sub> to F<sub>5</sub>)**

F<sub>0</sub> - Absolute control

F<sub>1</sub> - RDF (Recommended Dose of Fertilisers)

F<sub>2</sub> - RDFYF + 75 % N + RDP + RDK + 25 % N through Vermicompost

F<sub>3</sub> - RDFYF + 75 % N + RDP + RDK + 25 % N through Poultry manure

F<sub>4</sub> - RDFYF + 50 % N + RDP + RDK + 50 % N through Vermicompost

F<sub>5</sub> - RDFYF + 50 % N + RDP + RDK + 50 % N through Poultry manure

### **2.1.2. Factor 2 – Bioenhancers – 3 (B<sub>0</sub> to B<sub>2</sub>)**

B<sub>0</sub> - Absolute control

B<sub>1</sub> - 3 per cent Panchagavya foliar spray on 15, 30 and 45<sup>th</sup> DAP

B<sub>2</sub> - Jeevamirtham 500L ha<sup>-1</sup> as soil application, thrice with irrigation viz., at planting, 20 and 45 DAP

UNDER PEER REVIEW

**Table 1. Treatment particulars**

Sl. No.	Treatment	Treatment Particulars
1.	T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	Absolute control
2.	T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
3.	T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
4.	T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	RDF
5.	T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	RDF + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
6.	T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	RDF + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
7.	T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost
8.	T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost + 3% Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
9.	T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Vermicompost + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
10.	T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure
11.	T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
12.	T <sub>12</sub> - F <sub>3</sub> B <sub>2</sub>	RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
13.	T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost
14.	T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost + 3 % Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
15.	T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Vermicompost + Jeevamirtham 500 L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP
16.	T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure
17.	T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure + 3% Panchagavya foliar spray on 15, 30 and 45 <sup>th</sup> DAP
18.	T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	RDFYM + 50 % N + RDP + RDK + 50 % N through Poultry manure + Jeevamirtham 500L ha <sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 <sup>th</sup> and 45 <sup>th</sup> DAP

### 3. RESULT AND DISCUSSION

### **3.1. Effect of organics and bioenhancers on growth and yield of aggregatum onion**

#### **3.1.1. Plant height at 30 DAP (cm)**

The effect of fertiliser and bioenhancers observed on plant height of onion at 30 DAP were found to be significantly influenced while, the interaction effect was found insignificant. The application of RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and Jeevamirtham 500 L ha<sup>-1</sup> as soil application, thrice with irrigation viz., at planting, 20 and 45 DAP (F<sub>3</sub>B<sub>2</sub>) has recorded the highest plant height (31.40 cm) in the experiment (Table 2). The combined application of fertiliser with poultry manure, would have helped in sustained supply of macro and micro nutrients ensuring adequate and progressive release of nitrogen to plant as reported earlier by Datt and Kaur (2023) and Marbong and Swaroop (2018) in bellary onion. Among the bioenhancers used, the soil application of jeevamirtham was found superior to register significant performance with regard to plant height over foliar spraying of 3 per cent panchagavya on 15, 30 and 45 DAP and control and this finding was in conformity to the earlier report of Datt and Kaur (2023).

#### **3.1.2. Leaf girth at 30 DAP (cm)**

The differences observed for the leaf girth of onion at 30 DAP were found to be significantly influenced by fertiliser treatments, bioenhancers and their interaction. The leaf girth (1.94 cm) observed was found to be maximum (Table 2) in F<sub>3</sub> B<sub>2</sub> (RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and jeevamirtham 500 L ha<sup>-1</sup> as soil application thrice with irrigation viz., at planting, 20 and 45 DAP) at all stages of observation and this could be attributed to the better availability and supply of nitrogen in the presence of both poultry manure and jeevamirtham as reported by Chanu *et al.* (2022) in onion.

#### **3.1.3. Number of leaves plant<sup>-1</sup> at 30 DAP**

The differences observed for number of leaves plant<sup>-1</sup> of onion at 30 DAP were found to be significantly influenced by fertiliser treatments and bioenhancers while their interaction effect was found insignificant. The application of RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and jeevamirtham 500 L ha<sup>-1</sup> as soil application thrice with irrigation water viz., at planting, 20 and 45 DAP (F<sub>3</sub>B<sub>2</sub>) was found to give the best result in terms of

number of leaves plant<sup>-1</sup>(25.80)as evidenced from Table 2and such an impact of poultry manure in onion had already been established by Adeyeye *et al.* (2017) and Dapaah *et al.* (2014), while Prajapati and Vekariya (2022) reported significant influence of jeevamirtham on number of leaves in irrigated onion. The enhanced leaf production under combined application of fertilisers with organic manure and bioenhancers is attributable to the improved availability of essential nutrients, particularly nitrogen helping in better synthesis of various protein components required for leaf development, photosynthesis and metabolic processes of plant growth as suggested byDapaah *et al.* (2014) in bellary onion.

#### **3.1.4. Fresh weight of top at harvest (g)**

The fresh weight of top (leaves) estimated at harvest was found to show significant variation among the fertilisers treatments, bioenhancers studied and their interaction in comparison to control.The treatment which received 25 per cent nitrogen through poultry manure with rest of recommended nitrogen, phosphorus and potash as inorganic fertilizers in combination with soil application of jeevamirtham thrice at 500 L ha<sup>-1</sup> while planting, 30 and 45 DAP was found to be the best (12.26 g)in terms of fresh weight of top(Table 2)and such a positive role of integrated application of organic with inorganic sources of nutrients on leaf fresh weight has been observed by Adeyeye *et al.* (2017) and Singh *et al.* (2017). Further, they also suggested that the application of jeevamirtham promoting the beneficial microorganism in the rhizosphere, could have increased phosphorus availabilityleading to better top growth.

#### **3.1.5. Bulb diameter(cm)**

Significant differences in polar as well as equatorial diameter of onion bulbs were observed in the present study among fertiliser treatments, bioenhancers used as well as for their interaction effect. The highest polar as well as theequatorial diameter of bulblet as evident from Table 3 was observed in treatment F<sub>3</sub>B<sub>2</sub> (3.17 cm/2.94 cm)with integrated application of fertilizer and poultry manure as already reported in onion by Anand *et al.* (2017). The application of jeevamirtham has outperformed the foliar application of panchagavya and such a positive influence of jeevamirtham on bulb diameter has also been

reported earlier by Datt and Kaur (2023). Moreover, poultry litter is also considered to be a good source of primary, secondary and micro nutrients, which might have resulted in higher nutrient availability coupled with better uptake of nutrients leading to improved bulb growth as supported by Amanullah *et al.* (2007).

### **3.1.6. Bulblet weight (g)**

Bulb weight is one of the primary yield determinants in onion and it was found to vary significantly with fertiliser treatments given, bioenhancers used and their interaction in the present study. It was observed again that, the application of RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and Jeevamirtham 500L ha<sup>-1</sup> as soil drench, thrice with irrigation *viz.*, at planting, 20 and 45 DAP (F<sub>3</sub>B<sub>2</sub>) had shown maximum bulblet weight of 9.30 g (Table 3). The present findings on the role of poultry manure in improving onion bulblet weight is found to corroborate with the earlier result of Akhil and Singh (2022) and Mahala *et al.* (2018). Among the bioenhancers, the use of jeevamirtham has recorded improved bulblet weight and was found to be in conformity to the earlier findings of Datt and Kaur (2023).

### **3.1.7. Number of bulblets plant<sup>-1</sup>**

The study revealed the presence of significant differences among the various fertiliser and bioenhancers used as well as their interaction on number of bulblets plant<sup>-1</sup> in aggregatum onion. The application of inorganic and organic sources of nutrient in appropriate proportion in the treatment receiving RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and Jeevamirtham 500L ha<sup>-1</sup> as soil application, thrice with irrigation *viz.*, at planting, 20 DAP and 45 DAP (F<sub>3</sub> B<sub>2</sub>) has lead to the production of more number of bulblets plant<sup>-1</sup> (10.00) which is attributable to the highest number of bulblets plant<sup>-1</sup> (Table 3), the improved availability of nutrients from the inorganic fertilisers under the influence of poultry manure besides improvement in physicochemical as well as biological properties of the soil and such a report had been published earlier by Sitaula *et al.* (2020). Among the bioenhancers used jeevamirtham 500 L ha<sup>-1</sup> as soil application thrice with irrigation *viz.*, at planting, 20 and 45 DAP was found superior in producing maximum number of bulblets

plant<sup>-1</sup> and the ability of jeevamirtham improving the availability of nitrogen, phosphorous, potassium and beneficial microbes in the rhizosphere as reported by Manoj *et.al.* (2020) could be attributed for this.

### **3.1.8. Bulb yield plant<sup>-1</sup> (g)**

The existence of significant differences for bulb yield among the fertiliser treatments, bioenhancers used and their interaction was noticed in the present study. The maximum bulb yield of 43.18 g plant<sup>-1</sup> and 6.91 t ha ha<sup>-1</sup> (Table 3) was observed in treatment receiving RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and Jeevamirtham 500L ha<sup>-1</sup> as soil application, thrice with irrigation *viz.*, at planting, 20 and 45 DAP (F<sub>3</sub> B<sub>2</sub>). The presence of poultry manure in treatment F<sub>3</sub> (RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure) acts as good soil amendment by adding nutrients as well as beneficial microorganisms (Richa *et al.*, 2020). Among the two bioenhancer tried, jeevamirtham was found to be superior in terms of bulb yield and this could be the result of improved population of beneficial microbes including bacteria, yeast, fungi, actinomycetes and photosynthetic bacteria, besides enhanced soil enzyme activity contributing to such an yield increase of onion bulbs as suggested by Nitin and Purohit (2021) and Pathak and Shukla (2023). Application of jeevamirtham could have also helped in rapid build up of soil fertility resulting in better growth and yield as proposed by Kaur *et al.* (2020).

### **3.1.9. Dry matter production (kg ha<sup>-1</sup>) of bulbs**

The maximum dry matter production of onion bulb (788.51 kg ha<sup>-1</sup>) observed with the application of inorganic and organic sources of nutrients in right proportion in the treatment receiving RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and Jeevamirtham 500L ha<sup>-1</sup> as soil application, thrice with irrigation *viz.*, at planting, 20 and 45 DAP (F<sub>3</sub> B<sub>2</sub>) is attributable to the increased number of leaves, enhanced photosynthesis and better translocation and storage of assimilates in the bulbs, as well as better soil physiochemical and biological properties as reported earlier by Dapaah *et al.* (2014). The use of jeevamirtham as bioenhancers through soil application was also found to perform

better over the application of panchagavya, which was found to be in line with the report of Nitin and Purohit (2021).

UNDER PEER REVIEW

**Table 2. Effect of organics and bioenhancers on growth of aggregatum onion**

UNDER PEER REVIEW

Treatment	Plant height at 30 DAP (cm)		Leaf girth at 30 DAP (cm)		Number of leaves plant <sup>-1</sup> at 30 DAP		Fresh weight of top at harvest (g)	
T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	21.00		1.18		10.94		1.50	
T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	21.20		1.20		13.00		2.10	
T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	23.20		1.30		13.80		3.28	
T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	27.70		1.50		17.30		4.92	
T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	28.30		1.54		19.10		5.34	
T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	28.40		1.62		19.70		6.06	
T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	28.20		1.52		18.20		5.22	
T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	28.56		1.68		21.70		7.26	
T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	29.06		1.70		23.30		8.02	
T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	28.20		1.54		19.00		5.32	
T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	30.30		1.78		23.50		9.94	
<b>T<sub>12</sub> - F<sub>3</sub>B<sub>2</sub></b>	<b>31.40</b>		<b>1.94</b>		<b>25.80</b>		<b>12.26</b>	
T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	23.68		1.44		14.40		2.22	
T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	25.00		1.46		15.40		3.02	
T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	25.10		1.46		16.30		3.18	
T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	24.86		1.44		15.20		2.50	
T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	26.40		1.48		16.60		4.22	
T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	26.52		1.48		17.00		4.70	
Factor	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)
<b>Fertilisers</b>	0.33	0.70	0.02	0.04	0.73	1.53	0.11	0.23
<b>Bioenhancers</b>	0.23	0.49	0.01	0.03	0.51	1.08	0.08	0.16
<b>Fertilisers x Bioenhancers</b>	0.57	NS	0.04	0.07	1.26	NS	0.19	0.39

Table 3. Effect of organics and bioenhancers on yield parameters of aggregatum onion

Treatment	Polar diameter of bulblet (cm)		Equatorial diameter of bulblet (cm)		Bulblet weight (g)		Number of bulblets plant <sup>-1</sup>		Bulb yield plant <sup>-1</sup> (g)		Bulb yield (t ha <sup>-1</sup> )		Dry matter production of bulbs (kg ha <sup>-1</sup> )	
T <sub>1</sub> - F <sub>0</sub> B <sub>0</sub>	2.02		1.16		3.48		5.00		19.32		1.69		253.73	
T <sub>2</sub> - F <sub>0</sub> B <sub>1</sub>	2.12		1.50		3.72		5.20		19.43		2.88		254.50	
T <sub>3</sub> - F <sub>0</sub> B <sub>2</sub>	2.13		1.63		3.84		5.80		21.85		3.13		256.50	
T <sub>4</sub> - F <sub>1</sub> B <sub>0</sub>	2.54		2.06		6.02		6.60		31.28		4.70		595.01	
T <sub>5</sub> - F <sub>1</sub> B <sub>1</sub>	2.63		2.25		6.22		7.00		35.07		4.87		595.02	
T <sub>6</sub> - F <sub>1</sub> B <sub>2</sub>	2.65		2.34		6.28		7.20		39.27		5.17		595.03	
T <sub>7</sub> - F <sub>2</sub> B <sub>0</sub>	2.56		2.08		6.08		6.80		33.23		4.35		645.20	
T <sub>8</sub> - F <sub>2</sub> B <sub>1</sub>	2.67		2.56		6.50		7.20		39.62		5.56		645.25	
T <sub>9</sub> - F <sub>2</sub> B <sub>2</sub>	2.81		2.55		7.04		7.60		40.36		5.66		646.50	
T <sub>10</sub> - F <sub>3</sub> B <sub>0</sub>	2.57		2.17		6.10		7.00		33.65		4.90		787.00	
T <sub>11</sub> - F <sub>3</sub> B <sub>1</sub>	3.11		2.89		7.36		8.80		41.10		5.79		788.50	
<b>T<sub>12</sub> - F<sub>3</sub>B<sub>2</sub></b>	<b>3.17</b>		<b>2.94</b>		<b>9.30</b>		<b>10.00</b>		<b>43.18</b>		<b>6.91</b>		<b>788.51</b>	
T <sub>13</sub> - F <sub>4</sub> B <sub>0</sub>	2.20		1.79		3.92		6.20		22.53		3.76		422.51	
T <sub>14</sub> - F <sub>4</sub> B <sub>1</sub>	2.34		1.90		4.22		6.40		26.26		4.08		422.66	
T <sub>15</sub> - F <sub>4</sub> B <sub>2</sub>	2.37		1.98		4.44		6.40		27.02		4.12		423.50	
T <sub>16</sub> - F <sub>5</sub> B <sub>0</sub>	2.22		1.79		4.14		6.20		25.87		3.94		474.75	
T <sub>17</sub> - F <sub>5</sub> B <sub>1</sub>	2.40		1.98		4.56		6.40		28.52		4.17		475.71	
T <sub>18</sub> - F <sub>5</sub> B <sub>2</sub>	2.50		2.04		5.94		6.40		29.58		4.18		476.75	
Factor	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)	SEd	CD (p=0.05)
<b>Fertilisers</b>	0.03	0.07	0.03	0.06	0.08	0.17	0.09	0.20	0.32	0.68	0.06	0.13	0.17	0.35
<b>Bioenhancers</b>	0.02	0.05	0.02	0.04	0.06	0.12	0.07	0.14	0.23	0.48	0.04	0.09	0.12	0.25
<b>Fertilisers x Bioenhancers</b>	0.05	0.12	0.05	0.10	0.14	0.30	0.16	0.35	0.56	1.17	0.11	0.22	0.29	0.61

#### 4. CONCLUSION

The results of the experiment clearly revealed the significance of applying RDFYM + 75 % N + RDP + RDK + 25 % N through Poultry manure and jeevamirtham 500 L ha<sup>-1</sup> as soil application, thrice with irrigation viz., at planting, 20 and 45 DAP in terms of growth and yield parameters in aggregatum onion

#### REFERENCES

1. Adeyeye AS, Ishaku MA, Gadu HO, Olalekan KK, Lamid WA. Comparative Effect of Organic and Inorganic Fertilizer Treatments on the Growth and Yield of Onion (*Allium cepa* L.). J Bot Sci. 2017; **6** (2): 8-11.
2. Akhil, Singh A. Effect of organic manure and inorganic fertilizer on growth, yield and quality of garlic (*Allium sativum* L.). The Pharma Innovation J. 2022; **11** (9): 656-658.
3. Amanullah MM, Somasundaram E, Vaiyapuri K, Sathyamoorthi K. Poultry manure to crops – A Review. Agric. Rev. 2007; **28** (3): 216-222.
4. Anand M, Sankari A, Anita B. Influences of Integrated Nutrient Management for Garlic under Nilgiris Condition. Inter J Curr Microbiol & Appl Sci. 2017; **6** (12): 3833-3838.
5. Chanu KI, Devi AKB, Singh UC, Devi NS, Singh SNG. Effect of different levels of nitrogen and spacing on growth and yield of common onion (*Allium cepa* L.) cv. Prema 178 under Manipur condition. The Pharma Innovation J. 2022; **11** (4): 1268-1274.
6. Dapaah HK, Amoh-Koranteng JG, Darkwah K, Borketey-La EB. Influence of Poultry Manure and NPK Fertilization on Growth, Yield and Storability of Onion (*Allium cepa* L.) Growth under Rain-fed Conditions. Am J Exp Agric. 2014; **4** (8): 866-878.
7. Datt R, Kaur H. Impact of organic manures and biofertilizers on growth and yield of onion (*Allium cepa* L.). The Pharma Innovation J. 2023; **12** (6): 4838-4841.

8. Horticulture Statistics. Department of Agriculture and Farmers Welfare. Ministry of Agri. Government of India. 2022.
9. Kale RB, Gadge SS, Rao BV, Mahajan V, Singh M. Good Agriculture Practices in Onion and Garlic Production. ICAR- Directorate of Onion and Garlic Research, Pune, Maharashtra and National Institute of Agricultural Extension Management, Hyderabad, Telangana. 2022; 1-170.
10. Kaur A. JEEVAMRUTHAM: An effective activator of soil microorganisms. Just Agriculture e-Newsletter. 2020; 1(1): 1-5.
11. Kaur J, Brar RK, Sharma R. Effect of organic source of nutrients on yield and quality of onion (*Allium cepa* L.). Indian J Ecol. 2015; 42(1): 266-277.
12. Kaur P, Saini JP, Avnee, Meenakshi. Effect of doses and time of application of Jeevamrit on nutrient uptake and soil health under natural farming system. Inter J Chem Stud. 2020; 8 (6): 2537-2541.
13. Kumar CK, Singh G. Effect of Panchagavya on Growth and Yield: A Review. Inter J Curr Microbiol & Appl Sci. 2020; 9 (12): 617-624.
14. Mahala P, Chaudhary MR, Garhwal OP. Yield and Quality of rabi Onion (*Allium cepa* L.) Influenced by Integrated Nutrient Management. Inter J Curr Microbiol & Appl Sci. 2018; 7 (5): 3313-3321.
15. Manoj KN, Uma V, Kiran SC. Significance of liquid organic manures in sustainable crop production: A review. Inter J Ecol & Environ Sci. 2020; 2 (4): 445-449.
16. Marbong, Swaroop N. Effect of Potassium and Sulphur on the Soil Properties, Growth and Yield of Onion (*Allium cepa* L.) cv. Nasik Red. Inter J Curr Microbiol & Appl Sci. 2018; 7 (6): 1584-1591.
17. Nitin, Purohit HS. Effect of different Jeevamrut based liquid organic formulation on biochemical properties of soil and on plant growth of blackgram [*Vigna mungo* (L.) Hepper] under pot culture conditions. Inter J Chem Stud. 2021; 9 (1): 2280-2283.

18. PathakS, Shukla YR.Differential responses of integration of organic and inorganic manures on growth and yield of garlic (*Allium sativum* L.). *The Pharma. Innovation J.*2023; **12** (2): 640-645.
19. PrajapatiPJ,Vekariya PD.Effect of organics on growth and yield of drip irrigated onion (*Allium cepa* L.) under organic condition.*J Pharmacogn&Phytochem.*2022;**11** (4): 192-194.
20. RichaV, Kumar,Singh J, Sharma N.Poultry Manure and Poultry Waste Management: Review *Inter J Curr Microbiol& Appl Sci.* 2020; **9** (6): 3483-3495.
21. SinghS,Devi MB. Vegetables as a Potential Source of Nutraceuticals and Phytochemicals: A Review. *Inter J Medi & PharmSci.*2015; **5** (2): 1-14.
22. SinghVK,Kumari A,Chaudhar VK, Shree S. Role of Biofertilizers and Chemical Fertilizers for Sustainable Onion (*Allium cepa* L.) Production. *Inter J CurrMicrobiol&Appl Sci.* 2017; **6** (9): 2034-2040.
23. SitaulaHP,Dhakal R,Bhattarai C,Aryal A, Bhandari D. Effect of different combinations of poultry manure and urea on growth, yield and economics of garlic (*Allium sativum* L.) *JAgric NatResour.* 2020; **3** (1): 253-264.
24. SruthiB, Somasundaram E. Influence of organic manures on yield, quality and economics of aggregatum onion (*Allium cepa* L. var. aggregatum) *JPharmacogn&Phytochem.* 2019; **8** (3): 1768-1770.
25. TripathyP,Sahoo BB,Patel D, Das DK.Evaluation of integrated management modules for garlic (*Allium sativum* L.). *J Ecol& Environ.* 2017; **35** (1): 78-83.