

Innovative Reproductive Technology in Animal Breeding: A Review

Formatted: English (United Kingdom)

Abstract

The profitability and the intensity achieved by the genetic improvement is determined by the reproductive performance of the farm animals. The major cause of economic loss is reproductive inefficiency in the livestock sector. In the past few years, several advances ~~has~~ ~~have~~ been made in reproductive technologies (RTs) to increase the efficiency of genetic improvement. These technologies ~~involves-include~~ Artificial Insemination, Sexed semen Technology, Cryopreservation, Multiple Ovulation and Embryo Transfer, In-vitro Embryo Production (IVEP), Transgenesis etc. The problems of infertility of male or female, postpartum infertility in high-yielding animals of high genetic merit individuals and breeding of distant species has been overcome through these technologies. The inclusion of these techniques in animal breeding had made possible to obtain large number of progenies from genetically superior animals. This review, discuss the various innovative reproductive technologies implemented in livestock sector.

Commented [nr1]: Intensity of what? Check minor grammatical errors

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Keywords: Estrus Detection, Artificial Insemination, Sexed semen Technology, Cryopreservation, Multiple Ovulation and Embryo Transfer, In-vitro Embryo Production technology, Transgenesis

Commented [nr2]: The text focuses heavily on the benefits. Briefly mentioning potential drawbacks (e.g., animal welfare concerns, ethical considerations) would provide a more balanced view.

RTs address some infertility issues, they don't overcome all problems. Additionally, breeding distant species is highly complex and ethically debated. If discussing breeding distant species, touch on the ethical considerations.

Introduction

In 2023, the estimated total population in India amounted to approximately 1.43 billion. The improvement in the genetic field of the livestock animals is a major concern over these years. The need for innovative reproduction technologies in livestock animals arises from the desire to improve productivity, health and genetic diversity within livestock populations. By enhancing these technologies, farmers can increase the efficiency of breeding and production, leading to a greater supply of food for the growing human population. RTs can be used to select for and produce livestock with desirable traits, such as high milk production in cows or disease resistance in poultry. This can lead to better quality and quantity of animal products, such as milk, eggs, and meat, which are important source of nutrition for the human population. These technologies can help address challenges such as genetic disorders, low fertility rates, and limited access to superior breeding stock. By using advanced RT, farmers and breeders can more effectively manage genetic traits, improve the health and welfare of animals, and ultimately enhance the overall efficiency and sustainability of livestock production.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr3]: Some sentences could be restructured for improved clarity and flow.

Formatted: English (United Kingdom)

Commented [nr4]: Population of what? Specify. This sentence can be removed as it doesn't directly relate to the topic.

Formatted: English (United Kingdom)

Commented [nr5]: The first two paragraphs discuss the need for innovative technologies in similar ways. Consider combining them for a more concise introduction

Formatted: English (United Kingdom)

Commented [nr6]: Briefly mention the challenges faced in traditional livestock breeding practices

Formatted: English (United Kingdom)

Commented [nr7]: Go beyond basic product quality and quantity. Consider animal welfare improvements and environmental benefits (e.g., reduced methane emissions). Briefly introduce the potential ethical concerns surrounding some RTs

Formatted: English (United Kingdom)

To achieve this, several reproduction technologies are engaged in livestock breeding programme. These technologies play an important role that enables to increase genetic gain. Innovative reproduction technologies in livestock animals have been developed to improve breeding efficiency, genetic selection and overall productivity in livestock industry. The major reproduction technologies that are currently used by livestock breeding sectors are Estrus Detection Artificial Insemination (AI), Sexed semen technology, Cryopreservation, Multiple Ovulation and Embryo Transfer (MOET), In-vitro Embryo transfer Technology and

Transgenesis. These technologies alone or in combination with other techniques are implemented in livestock breeding procedures for efficient production of genetically superior offspring from genetically superior breeds. Additionally, these technologies can play crucial role in conserving rare and endangered breeds, ensuring their survival for future generations.

1. Estrus Detection

The behavioural changes and the physiological signs exhibited by the dairy cows before the ovulation period is referred as Estrus (Mičiaková *et al.* 2018). It is the period during the oestrus cycle when the female animals are ready to mate or sexually receptive for mating to become pregnant (Riaz *et al.* 2023). The behavioural changes occur due to the changes in the estrogen and progesterone levels (Paterson *et al.* 1992). Cows in estrus are more likely to mount other cows or to be mounted by other cows (Roelofs *et al.* 2010). Several secondary signs are being exhibited by the cows in estrus such as increase in movement, reduction in feed and water consumption, swelling of vulva, reddening of vulva, mucus discharge from vulva, sniffing in the genitalia area (Röttgen *et al.* 2018). Different devices are developed eventually for detection of estrus in female animals such as pedometer and accelerometer (Silper *et al.* 2015). Recently, a new device has been emerged for detection of estrus viz infrared thermography (Marquez *et al.* 2021). Infrared thermography cameras are very efficient in detecting minute changes in body temperatures of livestock animals (Nääs *et al.* 2020). To ascertain the appropriate timing for artificial insemination, accuracy and efficient estrus detection is important (Endo 2022).

Table no-1: Studies on application of Infrared Thermography (IRT) to detect estrus in cattle and

Buffalo (Riaz *et al.* 2023)

Sr. no.	References	Livestock animals	Site of Infrared thermography observations
1.	Talukder <i>et al.</i> , 2014	Dairy cows	Vulva and muzzle
2.	Marquez <i>et al.</i> , 2019	Dairy cows	Eye, muzzle, cheek, neck, front right foot, front left foot, rump, flank, vulva area, tail head, and withers
3.	Marquez <i>et al.</i> , 2021	Dairy cows	Vulva
4.	Rajput <i>et al.</i> , 2022	Sahiwal cows	Vulval, eyeball, ear and muzzle
5.	Tiwari <i>et al.</i> , 2022	Sahiwal cows	Muzzle and Vulva
6.	Marquez <i>et al.</i> , 2022	Dairy cows	Vulva

Source: (Riaz *et al.* 2023)

2. Artificial Insemination

Artificial Insemination is the first-generation tool used in reproductive biotechnology (Bharali and Sonowal 2019). AI is a reproduction technology in which semen is manually placed inside the vagina of the female animals (Gelayenew and Asebe 2017). This technology is less complex, less invasive, low cost and therefore the most assuring biotechnology for companion animals (Vishwanath 2003). The first successful insemination was performed by

Commented [nr8]: The phrase "innovative reproduction technologies in livestock animals" is repeated several times. Consider using synonyms like "advanced breeding techniques" or simply "RTs."

Formatted: English (United Kingdom)

Commented [nr9]: Table 1 not connected with text. Link table one with the text.

Formatted: English (United Kingdom)

Formatted: Font color: Red, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Font color: Red, English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr10]: Consider discussing limitations of current estrus detection methods or potential drawbacks of infrared thermography.

Review lacks a critical analysis of the different methods for estrus detection. It doesn't discuss limitations or compare the effectiveness of each method.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr11]: Text not linked to figure 1. Review lacks a critical analysis of AI. It doesn't discuss potential drawbacks or limitations of the technology

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Lord Spallanzani in 1784 in Bitch. For preserving and spreading male genetic material (Souza-Fabjan *et al.* 2023) and to utilize genetic material from male semen donors that have aged or suffered from fertility-limiting injuries, AI is generally used (Fonseca *et al.*, 2019; Souza-Fabjan *et al.*, 2021). To increase the production of local livestock animals, AI is performed by utilizing semen from exotic breeds (Verma *et al.* 2012). The success of Artificial Insemination AI depends on the efficiency of semen collected and preserved and is inclusion of various processes such as collection, Evaluation evaluation, Dilution dilution of semen (Semen semen Extenders extenders), packing (in medium- 0.5 ml and mini straws- 0.25 ml), Semen preservation (cryopreservation), Thawing thawing of semen (Patel *et al.* 2017).




Figure 1: Speculum method for inseminating the cow (Farmers Trend 2014).

3. Sexed Semen Technology

This technology deals with the physical separation of X and Y chromosome bearing spermatozoa in collected semen before inseminating female animals (Joshi *et al.* 2021). Several studies on the semen sexing found that the technology is mostly depend on the DNA content in the X and Y chromosome as X chromosome contains more DNA (3.8%) than Y chromosome (Rahman and Pang 2020; Joshi and Mathur *et al.* 2021).

The significant approaches toward the sperm sexing are the Albumin gradient, the identification of H-Y antigen, the Free-flow electrophoresis, the Detection detection of sex specific proteins, The the Centrifugal centrifugal counter current distribution. Separation of X and Y bearing spermatozoa with the use of albumin gradient was first time reported by Ericsson *et al.* (1973). The success rate reported through this method of sperm sexing was about 75% (Beernink *et al.* 1993; Kumar *et al.* 2017). Separation of X and Y bearing sperm cell in the identification of H-Y antigen depends upon the specific cell surface antigens on X

Commented [nr12]: Check the accuracy of this

Commented [nr13]: Cite

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr14]: Review initially mentions companion animals but primarily focuses on livestock

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr15]: Similar to previous sections, this one lacks a critical analysis. It doesn't discuss limitations or potential drawbacks of sperm sexing

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr16]: Be consistent

Formatted: English (United Kingdom)


Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

and Y sperm cells through magnetic bead or affinity chromatography (Hoppe *et al.* 1984; Hendriksen *et al.* 1996; Blecher *et al.* 1999; Hendriksen 1999 and Kumar *et al.* 2017). Yadav *et al.* (2017) reported that H-Y antigen was present in X- chromosome containing spermatozoa as well as in erythrocytes and premeiotic germ cells and cannot be used for separation of these cells because it does not specifically bind to Y chromosome bearing spermatozoa. Sperm sorting by free flow electrophoresis is done on the basis of the charge (negative by X chromosome and positive by Y chromosome) contained by spermatozoa under electric field conditions (Kaneko *et al.* 1984, Mohri *et al.* 1986 and Kumar *et al.* 2017). Howes and his colleagues in 1997 were the first one to work on the identification of differentially expressed proteins between X and Y spermatozoa but their work was indecisive. In Holstein bulls, the differentially expressed proteins in X and Y sperm cells were profiled by Shen *et al.*; (2021). The study showed that 8 and 23 surface proteins in X spermatozoa were up-regulated and down-regulated. About 81 and 151 proteins were also reported in the study that were exclusively expressed in X and Y spermatozoa respectively. Various immunological approaches for semen sexing can be developed on the basis of the different sperm types and their proteomes differences. Eventually the separation of X and Y chromosome bearing sperm cells can be achieved using the biomarkers that are developed from the these unique or differentially expressed proteins. The other technique used for sperm sorting is Flow Cytometry. The certainty of Flow Cytometry depends on the DNA contents of the particular chromosome bearing spermatozoa i.e., X- bearing sperm contains 3.8% DNA more than Y- bearing sperm (Johnson 2000). The Flow Cytometry technique of sperm sorting Uses Fluorescent Dyes to stain the DNA in sperm (Caroppo 2013; Ribeiro *et al.* 2013). The stain used for staining the DNA is Hoechst 33342 and is non-toxic, initially sperms are being stained with Hoechst 33342 stain and are then exposed to UV laser beam. After exposing it to the UV beam the emitting fluorescent light is detected and analysed (Johnson and Welch 1999).

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)



Formatted: English (United Kingdom)

Figure 2: Flow- cytometry based semen sorting (Naniwa et al., 2018).

4. Cryopreservation

There has been considerable success achieved in preservation of semen, oocytes or embryo in farm animals through ~~Cryopreservation~~ cryopreservation (Huang *et al.* 2018). The mating of female cattle that are not bound to time and places can be possible through cryopreservation of semen with minimized risk of disease transmission (Roca *et al.* 2016). The cryopreservation of gametes and embryo can be achieved by two processes (i) slow freezing (programmed) and (ii) vitrification. In slow freezing, the cells or embryos after being treated with cryoprotectants (CPAs) such as glycerol or dimethyl sulphoxide (DMSO) are subjected for slow freezing at the rate of 1°C per minute by using certain devices such as rate-controlled freezer or a benchtop portable freezing container (Day *et al.* 2007). Vitrification is an ultrarapid cooling process in which gametes and cells are directly placed in CPAs and are immediately driven in liquid nitrogen. Only few minutes are required for the vitrification process if compared with slow freezing process which minimizes the exposure time to sub-physiological conditions (Brambillasca *et al.* 2013). The formation of ice crystal is prevented in vitrification method by rapid cooling and warming rates (Day *et al.* 2007). The cryoprotectants (CPAs) used in cryopreservation are of two types (i) permeating and (ii) non-permeating CPAs. Glycerol and dimethyl sulphoxide are the permeating CPAs, DMSO have high penetrating rate (Saha *et al.* 1996), but can become toxic at higher temperature (Kasai *et al.* 1981). The non-permeating CPAs are Ficoll, Sucrose and Trehalose. ~~The addition of these CPAs increases osmotic pressure and are~~ ~~addition of these CPAs increases osmotic pressure and is~~ also helpful in penetration of Ethylene Glycol (EG) and DMSO. The study also reported that the freezing tolerance of oocytes can be improved by trehalose (Guo *et al.* 2000; Chen *et al.* 2001; Puhlev *et al.* 2001). Frostie, a Hereford-Friesian was the first calf born from frozen thawed embryo in 1973 (Wilmut and Rowson 1973).

Table no-2: Milestone achieved in vitrification in embryo (Gordon 2017)

Year	Species	Researcher
1985	Mouse	Rall and Fahy
1986	Cow	Massip <i>et al.</i>
1989	Rabbit	Smorag <i>et al.</i>
1990	Sheep/Goat	Scieve <i>et al.</i>
1994	Horse	Hochi <i>et al.</i>
1998	Pig	Kobayashi <i>et al.</i>

5. Multiple Ovulation and Embryo Transfer

Walter Heape (1890) was the first to utilize the technique of Multiple Ovulation and Embryo Transfer. The first embryo transfer in bovine was reported in 1949 by Umbaugh. The development of calf for the first time through embryo transfer was done in 1950 (Willet *et al.*

Commented [nr17]: Not linked with text

Formatted: English (United Kingdom)

Commented [nr18]: Similar to previous sections, this one lacks a critical analysis. It doesn't discuss limitations or potential drawbacks of cryopreservation

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr19]: Add a section that analyzes cryopreservation. Discuss limitations like potential damage to cells during freezing/thawing, success rate variations, and the infrastructure required for storage in liquid nitrogen. Briefly mention alternative methods for semen and embryo storage (non-cryogenic)

Formatted: English (United Kingdom)

Commented [nr20]: Not connected with text

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr21]: Incomplete citations

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr22]: Discuss limitations such as the hormonal manipulation required, potential stress on donor animals, and the technical expertise needed for successful implementation. Briefly mention alternative breeding methods for comparison. Briefly explain how the development of cryopreservation and non-surgical techniques revolutionized MOET in the mid-1970s. Eliminate redundancy and rephrase for improved clarity.

Formatted: English (United Kingdom)

1951). Earlier the collection and transfer of the embryos was performed surgically through mid-ventral exposure of uterus and ovaries (Genzebu 2015). The non-surgical recovery and transfer of embryos was made possible in mid 1970s with successful cryopreservation of bovine embryo. The production of the young ones through the process of MOET involves i) Selection of Donor-donor cow; ii) Superovulation of Donor-donor cow; iii) Insemination of Donor cow; iv) Development of Embryo (*in-vivo* or *in-vitro*); v) Embryo recovery (Surgically or Non- surgically); vi) Selection of recipient cow; vii) Synchronization of Receptient cow; viii) Embryo handling, Evaluation and Storage; ix) Embryo Transfer Method (Menta 2023).

5.1 Selection of donor cows

The three important points to consider the selection of donor cow are genetic superiority, reproductive ability and high value of progeny market (Mikkola, 2007; Besenfelder *et al.*, 2020). The body condition score should be appropriate at the time of embryo transfer (Habtie 2019). Rather than selecting donor cows, importance should be given to selection of males as males are normally bred to many females and should be selected more accurately. The selection of genetically superior male is very important as 50% of the genetic material is acquired through males, the high-quality semen is especially used from these bulls (David and Hamilton 2016).

5.2 Superovulation of donor cow

The objective of the superovulation is to induce more ovulations than normal rate and can be induced by administrating either PMSG (pregnant mare serum gonadotropin) commonly known as equine chorionic gonadotropin (eCG) or FSH. PMSG is administered on day 10 of the estrus cycle followed by injections of Prostaglandin within 2-3 days at an interval of 12-24 hr. The eCG stimulates greater super ovulatory response than FSH; although good quality of transferrable embryos is obtained through Follicle Stimulating Hormone (FSH) treatment (Selk 2013). The main disadvantage of PMSG is that it causes antibody formation and ovarian cyst in donor animals (Tekeli 2010). FSH is obtained from horse, sheep and pig (Akyol 2001) and is a pituitary gonadotropin. It is produced in the gonadotroph cells located in the anterior lobe of the pituitary gland (Yilmaz 1999), The FSH should be administered in repeated dose because the half-life of Follicle Stimulating Hormone is nearly 2 hours and given two times a day (Kimura *et al.* 2007). The best time to administer FSH is from 9th to 14th days of the estrus cycle (Kanagawa *et al.* 1995).

It is important to maintained hormonal balance through maintained diet as it is necessary for fertilization and embryo development before entering the uterus, at ampulla-isthmus junction (Nicholas and Smith, 1983; Burnett *et al.*; 2018). The recipient cow should be disease free for embryo reception and to perform number of pregnancies (Selk 2010). For intrauterine embryo development a well-developed corpus luteum played an important role (Mattos *et al.*, 2011; Hansen, 2020). After insemination, the embryos are collected after 7 days from uterus (Habtie 2019).

5.3 Synchronisation of recipient cow

The condition of the reproductive tract of recipient cow should be similar to that of donor cow for the transferring the embryo, for acquiring similar conditions synchronisation of the recipient cow should be done at the same time as donor cow (Galina & Orihuela, 2007). In recipient cow the occurrence of PGF2 α -induced estrus is observed after 60-72 hours,

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr23]: Grammar

Formatted: English (United Kingdom)

Commented [nr24]: Importance of progeny market value is mentioned, however, a deeper analysis of the economic implications of donor selection decisions would enhance the report. Factors such as cost-benefit analysis and return on investment could be explored.

Review predominantly focuses on the importance of genetic superiority and male selection, without addressing potential counterarguments or alternative perspectives. Including a balanced discussion would provide a more nuanced understanding of the topic.

Review suggests prioritizing male selection over donor cow selection, it could benefit from clearer justification or evidence supporting this assertion. This would help understand the rationale behind the proposed shift in focus.

Formatted: English (United Kingdom)

Commented [nr25]: Discuss the effectiveness of superovulation in increasing embryo yield. Briefly mention limitations, such as the potential for overstimulation and reduced embryo quality.

Add a section that analyzes superovulation. Discuss potential health risks for donor animals, the need for precise timing and dosage, and the cost implications of hormone treatments.

Formatted: English (United Kingdom)

therefore the recipient's synchronisation is done 12 hours before the donor cow (Genzebu 2015).




Figure 3: Diagram of the embryo flushing and recovery procedure (Selk, 2010).

6. In-Vitro Embryo Production

The In-vitro Embryo Production process involves three steps- in vitro maturation (IVM), in vitro fertilization (IVF) and in vitro development (IVD) of the resulting embryos (Gelayenew & Asebe 2016). At the time of birth thousands of oocytes are present inside the ovary and are lost due to atresia. The loss of genetic material through atresia can be reduced by harvesting oocytes from ovary followed by IVEP technique (Brakette and Zuelke 1993; Hasler 1998). The first calf produced by in vitro fertilization (IVF) in 1981 and was named "Virgil" (Brakett *et al.* 1982). The main principles of IVEP are selection of donor, selection of recipient animals and the collection of embryos (Kidie 2019). Addition of Nano-Selenium and Nano-Zinc Oxide during oocyte maturation increases maturation rate significantly (Abdel-Halim and Helmi 2017). In vitro fertilisation has also been used to produce the thousands of embryos (Gordon and Lu 1990). IVF through intracytoplasmic sperm injection (ICSI) is prominently used in assisted reproductive technologies (Keskintepe *et al.* 2002). For most cattle farmers this technology is an advantage only for extremely valuable cows that are infertile or fail to respond to superovulation.

6.1 In-vitro Maturation (IVM): Collection of oocytes can be achieved either manually through ultrasound-guided transvaginal follicular aspiration (ovum pick-up) or surgically from the ovaries of slaughtered animals or live animals by mid-ventral exposure of uterus and ovaries. Follicles of 3-5 mm diameter are mostly utilised for oocyte collection. Oocytes with

Commented [nr26]: Rreview only focuses on one aspect of recipient cow preparation. Elaborate on synchronization methods, discussing challenges, conducting comparative analyses, and providing practical implementation tips, would enhance the report's comprehensiveness and practical relevance

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr27]: Link with text

Formatted: English (United Kingdom)

Commented [nr28]: This section lacks a critical perspective. It doesn't discuss limitations such as the specialized equipment, expertise, and cost required for IVEP. Additionally, the success rates of IVEP compared to traditional MOET, and the ethical considerations surrounding large-scale embryo production are not addressed. For most cattle farmers, the focus should be on the applicability and economic feasibility of this technology compared to simpler methods.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

compact cumulus and corona are selected for culture. In-Vitro Maturation of oocytes is conducted for 24 hrs at 38.5°C in presence of 5% CO₂ in a CO₂ incubator.

6.2 Preparation of Spermatozoa: Capacitated sperm is essential for in-vitro fertilization. After capacitation, acrosome reaction in the head of sperm allows the release of certain enzymes which help in penetrating the zona pellucida of the oocyte.

6.3 In-vitro Fertilization (IVF): The capacitated sperm and the collected matured oocytes are co-cultured and are allowed to incubate at body temperature of animal for 6-24 hrs.

6.4 In-vitro embryo development: It is a component of IVF where in resultant embryos (Zygote) are allowed to grow for some time in an artificial medium. Culture is continued for 24 to 36 hrs before a further assessment for embryo cleavage is performed. At this stage, the embryos are normally between two to eight cell (blastomere) stages. Embryo quality is assessed depending on blastomere symmetry and the degree of cytoplasm fragmentation before transfer.




Figure 4: In vitro fertilization and embryonic development (Chowdhury, et al., 2019).

7. Transgenesis

Transgenesis technique involves the manipulation of genes of an organism and deliberately addition of that genome into the genome of organism of same or other species (Shankar and Mehendale, 2014). Palmiter et al., (1982) was the first to use genetic engineering in the improvement of livestock animal. The first success through this technology was obtained in laboratory animal mice by Jaenisch and Mintz in 1974. The first transgenic calf “Rosie” was produced in 1997 (Clark 2002) and was helpful in the production of human protein-enriched milk. It is not possible to make alterations in animal genome through conventional breeding, to fulfil the inadequacies of the conventional breeding programme, transgenesis technique is used. The transgenic animals produced from the process of transgenesis shows great advantages such as Increase in Feed Conversion rates, increase in growth rate of meat animals, increase in muscle mass, improved nutritional quality, Increase in disease resistance

Commented [nr29]: Review advantages and disadvantages of each method, as well as considerations for optimizing culture conditions to enhance oocyte quality and developmental competence

Formatted: English (United Kingdom), Superscript

Formatted: English (United Kingdom)

Formatted: Font: Bold, English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr30]: Too short

Formatted: English (United Kingdom)

Commented [nr31]: Too short

Formatted: English (United Kingdom)

Commented [nr32]: Mention limitations like the subjectivity involved in embryo quality assessment based on blastomere symmetry and fragmentation. Discuss the challenges of maintaining optimal conditions in the artificial culture medium to ensure proper embryo development.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr33]: Mention in text

Formatted: English (United Kingdom)

animals, Xenotransplantation's etc. Various methods are used for the production of transgenic animals such as DNA microinjection, Use of Transposon's, Retrovirus mediated-gene transfer, Embryonic stem cell-mediated (ESC) gene transfer and Lentiviral Transfer of Oocytes and Zygotes (Ahmad *et al.*2018). Transgenic dairy cows are able to produce casein content at higher level (Brophy *et al.*, 2003) and are mastitis resistant Staphylococcus aureus (Wall *et al.*, 2005). Introduction of new genes to the entirely different species is also possible through transgenesis (Magnus and Lali, 2008).

7.1 DNA Microinjection

In DNA microinjection the DNA is directly injected into the pronuclei of the embryos and was first documented technique. Due to the slow reproduction rate of the bovine animals and comparatively low embryo generation due to superovulation, the success through microinjection of DNA is only possible when DNA is microinjected to the blastocyst stage after in-vitro oocyte maturation and fertilisation followed by development till blastocyst stage (Krimpenfort *et al.* 1991).

7.2 Retrovirus-mediated gene transfer

Among the various gene transfer techniques, the retrovirus-mediated gene transfer was highly efficient (Nowrouzi *et al.* 2011). Retroviruses are used as vectors for gene transfer. These vectors efficiently transfer genes due to their affinity and infectivity for specific targeted cell and will result in the successful incorporation of transgene (Koo *et al.* 2014). The genetic material is transferred in the form of RNA into the host cells resulting in the chimera animal, these animals are then subjected to inbred for about 20 generations to produce homozygous transgenic animals which carries desired transgene in each and every cell of these offspring's (Mammohan and Niraj 2010).

7.3 Embryonic stem cell-mediated (ESC) gene transfer

This method involves the insertion of the target DNA into the embryonic stem cells that are cultured through *in-vitro* technique. The nature of the embryonic stem cell is pluripotent and are obtained from the inner cell mass of the blastocyst. The blastocyst stage of the embryo has the potential to differentiate into somatic cells, which lead to creation of complete organism. ESC allows the insertion, removal or modification of DNA sequences.

7.4 Lentivirus transfer of oocyte and embryo

The major drawback of Retrovirus-mediated gene transfer is that the silencing of transgenic locus started during embryonic development or shortly after birth (Chan *et al.* 1998). To overcome this drawback, the vector derived from Lentiviruses are used, the microinjection containing lentiviral vector is injected into the oocyte free from cumulus cells. The further study reported that the oocyte injected with the lentiviral vector produced efficient transgenic cattle and all these animals expressed enhanced green fluorescent protein (eGFP) expression (Hofmann *et al.* 2004). The same study reported that lentiviral gene can be either injected directly through sub zonal injection or indirectly through transduced nuclei into enucleated oocyte. The use of these vectors has some limitations such as integrated vectors sometimes show positional effect, the size of the vector genome restricted to only 8-10 kb (Wu *et al.* 2010), disruption of the endogenous gene through insertion mutagenesis by insertion of vector (Li and Lu 2010; Bednarczyk 2016).

Commented [nr34]: Review ethical considerations and regulatory frameworks surrounding transgenesis, as well as recent advancements and future directions in the field to ensure a balanced and up-to-date perspective

Formatted: English (United Kingdom)

Commented [nr35]: Review limitations beyond the slow reproduction rate of cows

Formatted: English (United Kingdom)

Formatted: Font: Not Italic, English (United Kingdom)

Formatted: English (United Kingdom)

Commented [nr36]: Briefly mention the safety concerns associated with retroviruses, such as the risk of insertional mutagenesis, where viral insertion disrupts essential genes. Briefly acknowledge the ethical concerns surrounding the creation of transgenic animals.

Formatted: English (United Kingdom)

Commented [nr37]: Briefly mention the ethical concerns, technical challenges associated with ESCs, limited applications etc.

Formatted: English (United Kingdom)

Commented [nr38]: Elaborate on the limitations associated with these vectors, such as positional effects and disruption of endogenous genes through insertion mutagenesis. Additionally, include recent studies or advancements in lentiviral vector technology to enhance the completeness of the review

Formatted: English (United Kingdom)

Conclusion

Innovative reproductive technologies have contributed enormously in the field of livestock industry. The implementation of AI technology not only maximizes animals' productivity but also provides opportunity to the individual sires with traits of superior quality for breeding and also reduces the risks of spreading sexually transmitted diseases. With the inclusion of semen sorting technology, selection of offspring of desired sex to increase profit in livestock industry has been made practical. The use of cryopreservation technology made possible the preservation of genetic material, semen or embryo for future use, thereby providing an effective method for the conservation of indigenous livestock, global genetic transport, gene banking, breeding line restoration, and for genetic rescue of endangered species. MOET and IVEP technology help to accelerate the transmission of desirable traits/genetic improvement by increasing offspring of selected males and females and the reduction of the generation interval in livestock populations in a shorter period of time compared to classical approaches. Moreover, with the development of disease-resistant animals and other approaches for enhancing animal production capacity, animal transgenesis has the potential to replace traditional.

References

- Abdel-Halim, B. R. and Helmy, N. A. 2017. Effect of nano-selenium and nano-zinc particles during in vitro maturation on the developmental competence of bovine oocytes. *Animal Production Science*.
- Ahmad, S. R., Mahajan, K., Gupta, T., Gulzar, M. and Yadav. V. 2018. Transgenesis in Animals: Principles and Applications – A Review. *Int. J. Curr. Microbiol. App. Sci.*, 7(10): 3068-3077.
- Akyol N, 2001. Using hormone in cattle embryo transfer. *Lalahan Hay Arařt Enst Derg*, 41: 95-104
- Beermink, F. J., Dmowski, W. P., Ericsson, R. J. 1993. Sex preselection through albumin separation of sperm. *Fertility and Sterility*. 59:382-386.
- Bednarczyk M (2016) Some aspects of poultry biotechnology: a review. *Slovak J. Anim. Sci.* 49(4):157–159
- Besenfelder, U., Brem, G., and Havlicek, V., 2020. Environmental impact on early embryonic development in the bovine species. *Animal*, 14(S1): s103–s112.
- Bharali, K. and Sonowal, J. 2019. Assisted Reproductive Technology (ART) in Animals. *E-zine of Biological Sciences*. Issue -9.

Commented [nr39]: A more balanced conclusion would acknowledge both the advantages potential drawbacks and ethical concerns of these technologies

Formatted: English (United Kingdom)

Commented [nr40]: Not consistent

Formatted: English (United Kingdom)

Commented [nr41]: Consistence

Formatted: English (United Kingdom)

- Blecher, S. R., Howie, R., Li, S., Detmar, J., Blahut, L. 1999. A new approach to immunological sexing of sperm. *Theriogenology*. 52:1309-1321.
- Brambillasca, F., Guglielmo, M. C., Coticchio, G., Mignini-Renzini, M., Dal-Canto, M., Fadini, R. 2013. The current challenges to efficient immature oocyte cryopreservation. *Journal of Assisted Reproduction and Genetics*, 30(12): 1531–153.
- Brackett, B. G., Bousquet, D., Boice, M. L., Donawick, W. J., Evans, J. F., Dressel, M. A. 1982. Normal development following in vitro fertilization in the cow. *Biol. Reprod.* 27:147-58.
- Brackett, B. G. and Zuelke, K. A. 1993. – Analysis of factors involved in the in vitro production of bovine embryos. *Theriogenology*, 39 (1), 43-64.
- Brophy, B., Smolenski, G., Wheeler, T., Wells, D., L'Huillier, P., Laible, G. 2003. Cloned transgenic cattle produce milk with higher levels of β -casein and κ -casein. *Nat. Biotechnol.*, 21:157-62.
- Burnett, T. A., Polsky, L., Kaur, M., and Cerri, R. L. A. 2018. Effect of estrous expression on timing and failure of ovulation of Holstein dairy cows using automated activity monitors. *J. Dairy Sci.*, 101: 11310–11320.
- Caroppo, E. 2013. Sperm sorting for selection of healthy sperm: is it safe and useful? *Fertility and Sterility*. 100(3):695-696.
- Chan, A. W. S., Homan, E. J., Ballou, L. U., Burns, J. C., Bremel, D. R. 1998. Transgenic cattle produced by reverse-transcribed gene transfer in oocytes. *Proc. Natl. Acad. Sci. U. S. A.*; 95:14028–14033
- Chen, S. U., Lien, Y. R., Cheng, Y. Y., Chen, H. F., Ho, H. N. and Yang, Y. S. 2001. Vitrification of mouse oocytes using closed pulled straws (CPS) achieves a high survival and preserves good patterns of meiotic spindles, compared with conventional straws, open pulled straws (OPS) and grids. *Human Reproduction*, 16(11): 2350–2356.
- Chowdhury, M.M.R., X, Lianguang, K, Rami, P, Bun-Young, M, Ayman, J, Myeong-Don, A, Fahmida, I, Jong, L, Hyun-Tae and K, Il-Keun, 2019, Invitro production of sex preselected cattle embryos using a monoclonal antibody raise against bull sperm epitopes. *Animal Reproduction Science*; 205: 156–164

- Clarke, A. R. 2002. Transgenesis techniques: principles and protocols (Vol. 180). *Springer Science & Business Media*.
- David, A. and S. Hamilton, 2016. Hamco cattle co. 18th Annual Angus bull sale. Glenboro, Manitoba Canada.
- Day, J. G. and Stacey, G. N. 2007. Gefriertrocknen. Cryopreservation and freeze-drying protocols. *FEBS Letters*, 2007, 377(2): 281–282.
- Endo, N. 2022. Possible causes and treatment strategies for the estrus and ovulation disorders in dairy cows. *Journal of Reproduction and Development*, Vol. 68, No 2. 85-89.
- Farmers Trend .2014. All there is to know about Artificial Intelligence (AI) in Goats and cows. Medilink Vet. Suppliers.
- Fonseca, J. F., Machado, V. O., Paiva, M. P. S. L., Facó, O., Souza-Fabjan, J. M. G. 2019. Recent advances in goat artificial insemination in Brazil. *Revista Brasileira de Reprodução Animal*. 43, 66–71.
- Galina, C. & Orihuela, A. (2007). The detection of estrus in cattle raised under tropical conditions: What We Know and What We Need To Know Hormones and Behavior, 52, 32-38.
- Galli, C., Lazzari, G. 2003. In vitro production of embryos in farm animals. In: Proceeding of the 19th Scientific Meeting of the European Embryo Transfer Association. Rostock: Germany, 12-13 Sept, p.93- 101.
- Gelayenew, B. and Asebe, G. 2017. Review on Major Assisted Reproductive Technologies. *Global Journal of Science Frontier Research: D Agriculture and Veterinary*. Volume 16 Issue 8.
- Genzebu, D., 2015. A Review of Embryo Transfer Technology in Cattle. *Global Journal of Animal Scientific Research*;3(2): 562-561.
- Gordon, I. 2017. Reproductive technologies in farm Animals. In: Gordon I. In vitro embryo production. 2nd ed. Cambridge: CABI Pub, 100–101
- Gordon, I. and Lu, K. H. 1990. Production of embryos in vitro and its impact on livestock production. *Theriogenology*, 33 (1), 77-87.

- Habtie, A.K., 2019. Review on Growth and Development of Multiple Ovulation and Embryo Transfer Technology in Cattle. Ethiopian Institute of Agricultural Research, Pawe Agricultural Research Center, World Scientific News; 127(3): 191-211.
- Hansen, P. J. 2020. Implications of assisted reproductive technologies for pregnancy outcomes in mammals. *Annu. Rev. Anim. Biosci.*, 8: 395-413.
- Hasler, J. F. 1998. – The current status of oocyte recovery, in vitro embryo production, and embryo transfer in domestic animals, with an emphasis on the bovine. *J. Anim. Sci.*, 76 (Suppl. 3), 52-74.
- Hasler, J. F., 2003. The current status of oocyte recovery, in vitro embryo production, and embryo transfer in domestic animals, with an emphasis on the bovine. *Journal of Animal Science* ;76: 52-74.
- Heape, W. 1890. Preliminary note on the transplantation and growth of mammalian ova within a uterine foster-mother. *Proceedings of the Royal Society of London*, 48: 457-8.
- Hendriksen, P. J. M. 1999. Do X and Y spermatozoa differ in proteins? *Theriogenology*. 52:1295-1307.
- Hendriksen, P. J. M., Welch, G. R., Grootegoed, J. A., Vander-Lende, T., Johnson, L. A. 1996. Comparison of detergent solubilized membrane and soluble proteins from flow cytometrically sorted X- and Y- chromosome bearing porcine spermatozoa by high resolution 2 - D electrophoresis. *Molecular Reproduction and Development*. 45:342-450, 25.
- Hofmann, A., Zakhartchenko, V., Weppert, M., Sebald, H., Wenigerkind, H., Brem, G. and Pfeifer, A. 2004. Generation of transgenic cattle by lentiviral gene transfer into oocytes. *Biology of Reproduction*, 71(2), 405-409.
- Hoppe, P. C. and Koo, G. C. 1984. Reacting mouse sperm with monoclonal H-Y antibodies does not influence sex ratio of eggs fertilized in vitro. *Journal of Reproductive Immunology*. 6:1-9.
- Howes, E. A., Miller, N. G., Dolby, C., Hutchings, A., Butcher, G. W., Jones, R. 1997. A search for sex-specific antigens on bovine spermatozoa using immunological and biochemical techniques to compare the protein profiles of X and Y chromosome-

bearing sperm populations separated by fluorescence-activated cell sorting. *J Reprod Fertil.* 110(2):195-204.

Huang, Z., Gao, L., Hou, Y., Zhu, S., Fu, X. 2018. Cryopreservation of farm animal gametes and embryos: recent updates and progress. *Front. Agr. Sci. Eng.*, 6(1): 42–53.

Jaenisch, R. and Mintz, B. 1974. Simian virus 40 DNA sequences in DNA of healthy adult mice derived from preimplantation blastocysts injected with viral DNA. *Proceedings of the national academy of sciences*, 71(4), 1250-1254.

Johnson, L. A. and Welch, G. R. 1999. Sex preselection: High-speed flow cytometric sorting of X and Y sperm for maximum efficiency. *Theriogenology*. 52(8):1323-1341.

Johnson, L. A. 2000. Sexing mammalian sperm for production of offspring: the state - of -the art. *Animal Reproduction Science*. 60-61: b93-107.

Joshi, H., Mathur, M., Mohanty, A. K., Kumar, S., Kaushik, J. K., Mohanty, T. K., Kumar, D., Singh, S. K., Bhushan, V., Parashar, A., Bhardwaj, P., Yata, V. K. 2021. Semen Sexing in Bovine: Current Status and the Need to Develop Alternative Techniques. *Animal Reproduction Update*. Volume-1, Issue-1.

Kanagawa H, I Shimohira and N Saitoh, 1995. Manual of Bovine Embryo Transfer. National Livestock Breeding Center MAFF, JICA, Japan.

Kaneko, S., Oshiro, S., Kobayashi, T., Itzuka, R., Mohri, H. 1984. Human X, and Y - bearing sperm differ in cell surface sialic acid content. *Biochemical and Biophysical Research Communication*. 124:950-955.

Kasai, M., Niwa, K., Iritani, A. 1981. Effects of various cryoprotective agents on the survival of unfrozen and frozen mouse embryos. *Journal of Reproduction and Fertility*, 63(1): 175–180.

Keskintepe, L., Pacholczyk, G., Machnicka, A., Norris, K., Curuk, M. A., Khan, I. and Brackett B. G. 2002. Bovine blastocyst development from oocytes injected with freeze-dried spermatozoa. *Biol. Reprod.*, 67 (2), 409-415.

Kidie, H.A. 2019. Review on growth and development of multiple ovulation and embryo transfer technology in cattle. *World Scientific News*, 127, 191-211.

- Kimura K, M Hirako, H Iwata, M Aoki, M Kawaguchi and M Seki, 2007. Successful superovulation of cattle by a single administration of FSH in aluminium hydroxide gel. *Theriogenology*, 68: 633-639.
- Koo, B. C., Kwon, M. S. and Kim, T. 2014. Retrovirus-mediated gene transfer. In *Transgenic Animal Technology (Third Edition)* (pp. 167-194).
- Krimpenfort, P., Rademakers, A., Eyestone, W., van der Schans, A., van den Broek, S., Kooiman, P. and de Boer, H. 1991. Generation of transgenic dairy cattle using 'in vitro' embryo production. *Nature Biotechnology*, 9(9), 844.
- Kumar, N., Gebrekidan, B., Gebrewahd, T. T., Hadush, B. 2017. Sexed semen technology in cattle. *Indian Journal of Animal Health*. 56(2):157-168.
- Li, J. J. and Lu, L. Z. 2010. Recent progress on technologies and applications of transgenic poultry. *Afr J Biotechnol* 9(24):3481–3488
- Magnus, P. K. and Lali, F. A. 2008. Transgenic milk. *Veterinary World*, 1(10), 319.
- Manmohan, S. and Niraj, K. 2010. Transgenic animals: production and application. *International Journal of Pharmaceutical Sciences and Research (IJPSR)*, 1(9), 12-22.
- MIČIAKOVÁ MÁRIA, STRAPÁK PETER, SZENCZIOVÁ IVETA, STRAPÁKOVÁ EVA, HANUŠOVSKÝ ONDREJ. 2018. Several Methods of Estrus Detection in Cattle Dams: A Review. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 66(2): 619–625.
- Marquez, H. P., Ambrose, D., Schaefer, A., Cook, N., Bench, C.J. 2019. Infrared thermography and behavioral biometrics associated with estrus indicators and ovulation in estrus-synchronized dairy cows housed in tie stalls. *J. Dairy Sci.*, 102, 4427–4440.
- Marquez, H. P., Schaefer, A., Von-Gaza, H., Ambrose, D., Cook, N., Bench, C. J. 2002. Evaluating automated infrared thermography and vulva exposure tracking as components of an estrus detection platform in a commercial dairy herd. *Animal*, 16, 100585.
- Massip, A., Zwalmen, P. V. D., Scheffen, B., Ectors, F. 1986. Pregnancies following transfer of cattle embryos preserved by vitrification. *Cryo Letters*, 7: 270–273.

- Mattos, M. C., Bastos, M. R., Guardieiro, M. M., Carvalho, J. O., Franco, M. M., Mourão, G. B., and Barros, C. M., 2011. Improvement of embryo production by the replacement of the last two doses of porcine follicle-stimulating hormone with equine chorionic gonadotropin in Sindhi donors. *Anim. Reprod. Sci.*, 125: 119-123.
- Menta, Y. D. (2023). Review on embryo transfer in cattle and its application. *Int. J. Adv. Res. Biol. Sci.* 10(4): 71-87.
- Mikkola, M. 2007. Superovulation and ET in dairy cattle effect of management factors with emphasis on sex-sorted semen. Academic dissertation, pp. 79.
- Mohri, H., Oshio, S., Kaneko, S., Kobayashi, T., Lizuka, R. 1986. Separation and characterization of mammalian X and Y - bearing sperm. *Development growth and differentiation.* 28(1):35-36.
- Nääs, I. A., Garcia, R. G., Caldara, F. R. J. 2020. Infrared thermal image for assessing animal health and welfare. *J. Anim. Behav. Biometeorol.*, 2, 66–72.
- Naniwa, Y., Sakamoto, Y., Toda, S., Uchiyama, K. 2018. Bovine sperm sex-selection technology in Japan. *Reprod Med Biol.*;18(1):17-26.
- Nicholas, F. W., and Smith, C., 1983. Increased rates of genetic change in dairy cattle by ET and splitting. *Anim. Prod.*, 36: 341-353.
- Nowrouzi, A., Glimm, H., Von Kalle, C. and Schmidt, M. (2011). Retroviral vectors: post entry events and genomic alterations. *Viruses*, 3(5), 429-455.
- Palmiter, R. D., Brinster, R. L., Hammer, R. E., Trumbauer, M. E., Rosenfeld, M. G., Birnberg, N. C. and Evans, R. M. 1982. Dramatic growth of mice that develop from eggs microinjected with metallothionein–growth hormone fusion genes. *Nature*, 300(5893), 611.
- Patel, G. K., Haque, N., Madhavatar, M., Chaudhari, A. K., Patel, D. K., Bhalakiya, N., Jammesha, N., Patel P. and Kumar, R. 2017. Artificial insemination: A tool to improve livestock productivity. *Journal of Pharmacognosy and Phytochemistry.* SP1: 307-313.
- Patterson, D. J., Perry, R. C., Kiracofe, G. H., Bellows, R. A., Staigmiller, R. B., Corah, L. R. 1992.

Management Considerations in Heifer Development and Puberty. *J. Anim. Sci.* 70, 4018–

4035.

Perez-Marquez, H. J., Ambrose, D. J., Schaefer, A. L., Cook, N. J., Bench, C. J. 2021. Evaluation of infrared thermography combined with behavioral biometrics for estrus detection in naturally cycling dairy cows. *Animal*, 15, 100205.

Puhlev, I., Guo, N., Brown, D. R., Levine, F. 2001. Desiccation tolerance in human cells. *Cryobiology*, 42(3): 207–217.

Rahman, M. S., Pang, M. G. 2020. New Biological Insights on X and Y Chromosome-Bearing Spermatozoa. *Front Cell Dev Biol.*, 7:388.

Rall, W. F., Fahy, G. M. 1985. Ice-free cryopreservation of mouse embryos at -196 degrees C by vitrification. *Nature*, 313(6003): 573– 575.

Rajput, A. S., Bhakat, M., Mohanty, T. K., Baithalu, R. K., Mir, A. A., Lal, G. S., Singh, M., Rajput, R. K. D., Shah, N. 2022. Identification of estrus using infrared thermography in indigenous dairy animals. *Pharm. Innov. J.*, 1571–1575.

Riaz, U., Idris, M., Ahmed, M., Ali, F., Yang, L. 2023. Infrared Thermography as a Potential Non-Invasive Tool for Estrus Detection in Cattle and Buffaloes. *Animals*, 13, 1425.

Ribeiro, S., Sartorius, G., Geyter, C. 2013. Sorting of spermatozoa with flow cytometry. *Fertility and Sterility.*; 100(3):

Roca, J., Parrilla, I., Gil, M. A., Cuello, C., Martinez, E. A., Rodriguez-Martinez, H. 2016. Non-viable sperm in the ejaculate: lethal escorts for contemporary viable sperm. *Animal Reproduction Science*, 169: 24–31.

Roelofs, J., López-Gatius, F., Hunter, R. H. F., Van-Eerdenburg, F. J. C. M., Hanzen, C. 2010. When Is a Cow in Estrus? Clinical and Practical Aspects. *Theriogenology*, 74, 327–344.

Röttgen, V., Becker, F., Tuchscherer, A., Wrenzycki, C., Düpjan, S., Schön, P.C., Puppe, B. 2018. Vocalization as an Indicator of Estrus Climax in Holstein Heifers during Natural Estrus and Superovulation. *J. Dairy Sci.*, 101, 2383–2394.

- Talukder, S., Thomson, P., Kerrisk, K., Clark, C., Celi, P. J. 2015. Evaluation of infrared thermography body temperature and collar mounted accelerometer and acoustic technology for predicting time of ovulation of cows in a pasture-based system. *Theriogenology*, 83, 739–748.
- Tiwari, S., Singh, Y., Sirohi, R., Yadav, B., Singh, D. N., Gurung, A., Shakya, P. J. T. 2022. Infrared thermographical differentiation of estrus and non-estrus stages of dairy animals. *Pharm. Innov. J.*, 10, 24–28.
- Saha, S., Otoi, T., Takagi, M., Boediono, A., Sumantri, C., Suzuki, T. 1996. Normal calves obtained after direct transfer of vitrified bovine embryos using ethylene glycol, trehalose, and polyvinylpyrrolidone. *Cryobiology*, 33(3): 291–299
- Selk, G., 2013. Embryo Transfer in Cattle. Division of Agriculture and Natural Resource, 3158, 1-4.
- Selk, G., 2013. Embryo Transfer in Cattle DASNR 102 Agriculture Hall, Oklahoma State University Stillwater, OK 74078, ANSI-3158, <http://osufacts.okstate.edu>.
- Shen, D., Zhou, C., Cao, M., Cai, W., Yin, H., Jiang, L., Zhang, S. 2021. Differential Membrane Protein Profile in Bovine X- and Y-Sperm. *J Proteome Res.*, 20(6):3031-3042.
- Silper, B. F., Madureira, A. M. L., Kaur, M., Burnett, T. A., Cerri, R. L. A. 2015. Short Communication: Comparison of Estrus Characteristics in Holstein Heifers by 2 Activity Monitoring Systems. *J. Dairy Sci.*, 98, 3158–3165.
- Souza-Fabjan, J. M. G., Batista, R. I. T. P., Correia, L. F. L., Paramio, M. T., Fonseca, J. F., Freitas, V. J. F., Mermillod, P. 2021. In vitro production of small ruminant embryos: latest improvements and further research. *Reproduction Fertility and Development* 33, 31–54.
- MIČIAKOVÁ MÁRIA, STRAPÁK PETER, SZENCZIOVÁ IVETA, STRAPÁKOVÁ EVA, HANUŠOVSKÝ ONDREJ. 2018. Several Methods of Estrus Detection in Cattle Dams: A Review. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 66(2): 619–625.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

- Spallanzani, L. 1784. Dissertations relative to the natural history of animals and vegetables. Trans. by T. Beddoes in Dissertations Relative to the Natural History of Animals and Vegetables. J. Murray, London, 2:195-199.
- Tekeli T, 2010. Embryo Transfer. In: Alaçam E (Editor): Obstetrics and Infertility in Domestic Animals, 7th edition, Medisan, Ankara, pp: 81-97.
- Umbaugh, R. E., 1951. Superovulation and ovum transfer in cattle. *Fertil Steril.* My-Jun;2(3): 243-52.
- Verma, O. P., Kumar, R., Kumar, A. and Chand, S. 2012. Assisted Reproductive Techniques in Farm Animal - From Artificial Insemination to Nanobiotechnology, *Vet. World.* 5(5):301-310.
- Vishwanath, R. 2003. Artificial insemination: the state of the art. *Theriogenology.* 59: 571-584.
- Willet, E. L., Black, W. G., Casida, L. E., Stone, W. H. and Buckner, P. J. 1951. Successful transplantation of a fertilized bovine ovum. *Science*, 113: 247.
- Wall, R. J., Powell, A. M., Paape, M. J., Kerr, D. E., Bannerman, D. D., Pursel, V. G., Wells, K. D., Talbot, N., Hawk, H. W. 2005. Genetically enhanced cows resist intramammary *Staphylococcus aureus* infection. *Nat Biotechnol.* 23:445-51.
- Wilmot, I., Rowson, L. E. 1973. Experiments on the low-temperature preservation of cow embryos. *Veterinary Record.* 92(26): 686–690 29.
- Wilmot, I., Rowson, L. E. 1973. The successful low-temperature preservation of mouse and cow embryos. *Journal of Reproduction and Fertility.* 33(2): 352–353.
- Wu, Z., Yang, H. and Colosi, P. 2010. Effect of genome size on AAV vector packaging. *Mol Ther* 18:80–86.
- Yadav, S. K., Gangwar, D. K., Singh, J., Tikadar, C. K., Khanna, V. V., Saini, S., Dholpuria, S., Palta, P., Manik, R. S., Singh, M. K., Singla, S. K. 2017. An immunological approach of sperm sexing and different methods for identification of X- and Y-chromosome bearing sperm. *Vet World.* Volume10(5):498-504.
- Yilmaz B, 1999. Hormones and Reproductive Physiology. Feryal Press, Ankara, Turkey.