

# Sero-Prevalence of Human Immunodeficiency Virus Among Students Undergoing Medical Examination in a Tertiary Institution

---

## ABSTRACT

**Aims:** To assess the prevalence of human immunodeficiency virus among students undergoing medical checkup in a tertiary institution

**Study design:** The investigation utilised a cross-sectional study design technique.

**Methodology:** In this present study, 2114 participants were included in the age range 15-49 years. Serological method was used to detect the presence of HIV using a rapid diagnostic technique using determine. Tile method was used to determine the ABO and Rhesus D antigens in the individuals.

**Results:** Out of the 2114 participants sampled, the population studied according to various departments within the university were ME 95 (5%), EM 181 (9%), LIB 101 (5%), SPE 63 (3%), ENG 296 (14%), BIO 423 (20%), CHE 150 (7%), COM 305 (14%), MGT 236 (11%) and ACC 236 (12%). The prevalence of HIV among tertiary institution students is 1%.

**Conclusion:** This research gives an insight into the prevalence of HIV among students in tertiary institution for strategic intervention.

*Keywords: Medical check, HIV, Student population*

## 1. INTRODUCTION

Viruses are microscopic entities that have a core of genetic material, either DNA or RNA. The core is covered with a capsid, a protective coat made of proteins. Around the coat there may be a spiky covering known as the envelope. These spikes are proteins that enable viruses to bind to and enter the host cells (<https://www.medicalnewstoday.com>). The human immunodeficiency virus (HIV) is a chronic viral infectious disease that damages the immune system of the infected individual. It progresses to the development of Acquired Immune Deficiency Syndrome (AIDS) and death without proper medical attention (Jacobs *et al.*, 2023). It is an enveloped lentivirus within the family Retroviridae. Its genetic material has a single nucleic acid structure which is known as an RNA virus (Tomezsko *et al.*, 2020). HIV can be transmitted through direct inoculation into the bloodstream, or after contact and attachment through mucosal surfaces. It can also be transmitted through direct contact with a bodily fluid containing HIV such as blood, semen, vaginal fluid, and breastmilk (Azuo *et al.*, 2012), the transmission could involve anal, vaginal or oral sex, blood transfusion, contaminated hypodermic needles, exchange between mother and child during pregnancy, childbirth or breastfeeding (Talarok and Talaro, 2002; Centre for Disease Control and prevention, 2003). An infection can result when the virus crosses the body epithelial barriers into the fluid compartments, due to the blood, semen, vaginal secretion of infected subjects, harbours high level of free virus and infected leukocytes, which are significant factors in sexual transmission pathway. Its mode of transmission is similar to other kinds of viral infection such as hepatitis, Ebola virus, Lassa fever virus.

HIV is a global health tragedy that has affected over 36.9 million persons in 2019 which translates to 0.5% of global population (Govender *et al.*, 2021). In 2018, 20.6 million people

in the southern and eastern Africa were living HIV and the western and central Africa was 5.0 million (WHO, 2019). Nigeria has the second largest number of people living with HIV/AIDS in Africa accounting for 9.0% of the global burden with prevalence of 3.2%, and burden of 3.1 million people living with HIV (Endalamaw et al., 2020; Govender et al., 2021). This burden impacts negatively on the health system. Report of HIV research in Nigeria showed that Rivers State has the highest HIV prevalence (15.2%) among the states in the country (Awofala et al., 2018). The generalized prevalence of HIV is approximately 3.6% but there are significantly higher rates among most-at-risk-populations (MARPs) (Federal Ministry of Health, 2012). These MARPs include illicit intra-venous drug users, commercial sex workers and men who have sex with men.

Despite the high burden of HIV among residents within Nigeria, the level of testing and management is poor and varies among the different strata of Governments (Dwyer-Lindgren et al., 2019, Cuadros et al., 2017, Coburn et al., 2017). This variation occurs at the local government, state and national levels and could be a result of knowledge gap about transmission in addition to poor management strategies (O'brien-Carelliet al., 2019).

A large HIV population-based survey conducted in Nigeria demonstrated the relevance of geographic disparities across the healthcare settings [Nigeria 2018]. The study revealed a prevalence of 1.3% among individuals aged 15 to 49 years. This prompted the Nigerian government to revise its strategic framework by classifying states into focal point for identification and treatment of HIV/AIDS-infected persons (United states 2019).

In this study, we present the cross-sectional perspective of prevalence, at the grassroot level, of HIV among students undergoing medical examination in a tertiary institution. Majority of these students are between the age of 15 to 49.

## **2. METHODS**

### **2.1 Study Area**

This study was performed at the Ignatius Ajuru University of Education in Port Harcourt of Rivers State using students undergoing medical checkup as subjects.

### **2.2 Study Design**

The study design employed in this research was cross-sectional design. The study sampled 2114 participants using simple random method which included both male and female sex. The sample size of 2114 was determined using the Cochrane formula with the prevalence of HIV at 15%.

### **2.3 Sample Collection**

After pre-test counselling and explanations, venous blood was drawn aseptically from the antecubital fossa of the subject with the use of vacutainer as described by Cheesebrough (2009).

### **2.4 Determination of HIV Prevalence**

The rapid diagnostic technique using **Determine HIV-1/2 set kit (Abbott Diagnostics Medical Co.Limited)** was used to detect the presence of HIV in subjects. The blood was allowed to coagulate thereby allowing the serum to separate **with the aid of a bench centrifuge ( Atlas**

**Medical).** A drop of serum was added to the Determine strip. The presence of double bands indicates positive for HIV while a single band shows negative result.

## 2.5 Determination of ABO and Rh-D Blood Group of Subjects

The serological method was employed to analyse the blood group of the subjects using the tile method.

For ABO and Rh-D blood group, a drop of anti-A, anti-B, anti-AB and anti-D (Atlas Medical), each was placed in the wells on the tile A, B, AB and O. A drop of red cell was added to the part labelled A, B, AB and O and anti A, anti-B, anti-AB and anti-D was dropped in the part labelled A, B, AB and O. This was mixed gently and rocked for about 30sec and observed for agglutination. Presence of agglutination indicated a positive result while absence of agglutination indicates a negative result.

## 2.6 Data Analyses

This study data was presented as percentages. Chi-square analyses were performed where necessary.

## 3. RESULTS

The study sampled 2114 participants. Table 1 shows the age distribution of subjective from different departments. The highest number of persons sampled fall within the 15 to 25 age range group ME 85 (89%), EM 157 (87%), LIB 88 (87%), SPE 43 (68%), ENG 262 (89%), BIO 396 (94%), CHE 132 (88%), COM 262 (86%), MGT 197 (83%), and ACC 250 (95%). This was followed by the age range of 26 to 25 years: ME 9 (10%), EM 14 (8%), LIB 13 (13%), SPE 15 (24%), ENG 33 (11%), BIO 27 (6%), CHE 18 (12%), COM 43 (14%), MGT 38 (16%), and ACC 14 (5%). The least population included in the study was within the age range of 35 to 45 years: ME 1 (1%), EM 10 (6%), LIB 1 (1%), SPE 5 (8%), ENG 1 (0%), BIO 0 (0%), CHE 0 (0%), COM 0 (0%), MGT 1 (1%) and ACC 0 (0%). The population studied according to various departments within the university were ME 95 (5%), EM 181 (9%), LIB 101 (5%), SPE 63 (3%), ENG 296 (14%), BIO 423 (20%), CHE 150 (7%), COM 305 (14%), MGT 236 (11%) and ACC 236 (12%).

**Table 1. Age Distribution of Subjects from Different Departments**

Age Range	ME n(%)	EM n(%)	LIB n(%)	SPE n(%)	ENG n(%)	BIO n(%)	CHE n(%)	COM n(%)	MGT n(%)	ACC n(%)
15-25	85 (89)	157 (87)	88 (87)	43 (68)	262 (89)	396 (94)	132 (88)	262 (86)	197 (83)	250 (95)
26-35	9 (10)	14 (8)	13 (13)	15 (24)	33 (11)	27 (6)	18 (12)	43 (14)	38 (16)	14 (5)
35-45	1 (1)	10 (6)	1 (1)	5 (8)	1 (0)	0 (0)	0 (0)	0 (0)	1 (1)	0 (0)
Total	95	181	101	63	296	423	150	305	236	264

\*ME – Marketing Education, EM – Education Management, LIB – Library and Information Science, SPE -Special Unit, ENG – English, BIO – Biology, COM – Computer, MGT – Management, ACC – Accounting

Table 2 shows the sex distribution of the sampled populations according to males and females are: ME 29 (31%), 66 (69%), EM, 56 (31%), 125 (69%), LIB 27 (27%), 74 (73%), SPE 21(33%), 42 (67%), ENG 37 (13%), 259 (87%), BIO 71 (17%), 352 (83%), CHE 50

(33%), 100 (67%), COM 64 (21%), 241 (79%), MGT 64 (27%), 172 (73%), and ACC 62 (23), 202 (77%).

**Table 2. Sex Distribution of Subjects from Different Departments**

Departments	Male n(%)	Female n(%)	Total n(%)
ME	29 (31)	66 (69)	95 (4)
EM	56 (31)	125 (69)	181 (9)
LIB	27 (27)	74 (73)	101(5)
SPE	21(33)	42 (67)	63 (3)
ENG	37 (13)	259 (87)	296 (14)
BIO	71 (17)	352 (83)	423 (20)
CHE	50 (33)	100 (67)	150 (7)
COM	64 (21)	241 (79)	305 (14)
MGT	64 (27)	172 (73)	236 (11)
ACC	62 (23)	202 (77)	264 (12)

*\*ME – Marketing Education, EM – Education Management, LIB – Library and Information Science, SPE -Special Unit, ENG – English, BIO – Biology, COM – Computer, MGT – Management, ACC – Accounting*

**Table 3. Prevalence of Blood Group Distribution of Subjects from Different Departments**

Departments	O <sup>+</sup> n(%)	O <sup>-</sup> n(%)	A <sup>+</sup> n(%)	A <sup>-</sup> n(%)	B <sup>-</sup> n(%)	B <sup>+</sup> n(%)	AB <sup>+</sup> n(%)	AB <sup>-</sup> n(%)	Total n(%)
ME	60	2	17	2	0	13	1	0	95 (4)
EM	98	6	50	1	0	25	1	0	181 (9)
LIB	53	3	25	0	0	15	4	0	101(5)
SPE	46	2	9	0	1	3	2	0	63 (3)
ENG	158	10	72	2	3	46	5	0	296 (14)
BIO	229	16	108	1	1	62	6	0	423 (20)


CHE	103	3	20	3	0	15	6	0	150 (7)
COM	168	10	68	1	0	54	4	0	305 (14)
MGT	150	7	40	1	1	36	1	0	236 (11)
ACC	144	14	65	3	1	35	2	0	264 (12)
Total	1209	73	474	14	7	304	32	0	

*\*ME – Marketing Education, EM – Education Management, LIB – Library and Information Science, SPE -Special Unit, ENG – English, BIO – Biology, COM – Computer, MGT – Management, ACC – Accounting*

**Table 4. Prevalence of HIV among Subjects across Different Departments**

Departments	Positive n(%)	Negative n(%)	Total n(%)
ME	1 (1)	94 (99)	95 (4)
EM	5 (3)	176 (97)	181 (9)
LIB	2 (2)	99 (98)	101(5)
SPE	0(0)	63 (100)	63 (3)
ENG	1 (0)	295 (100)	296 (14)
BIO	5 (1)	418 (99)	423 (20)
CHE	1 (1)	149 (99)	150 (7)
COM	2 (1)	303 (99)	305 (14)
MGT	3 (1)	233 (99)	236 (11)
ACC	2 (1)	262 (99)	264 (12)
Total	22	2092	

*\*ME – Marketing Education, EM – Education Management, LIB – Library and Information Science, SPE -Special Unit, ENG – English, BIO – Biology, COM – Computer, MGT – Management, ACC – Accounting*



**Figure 1 Prevalence of HIV among subjects across different Departments**

*HIV Prevalence across Blood Groups*

Departments	O <sup>+</sup> n(%)	O <sup>-</sup> n(%)	A <sup>+</sup> n(%)	A <sup>-</sup> n(%)	B <sup>-</sup> n(%)	B <sup>+</sup> n(%)	AB <sup>+</sup> n(%)	AB <sup>-</sup> n(%)	Total n(%)
Prevalence %	0	2	4	5	3	2	2	4	22

#### 4. DISCUSSION

Our study comprehensively provides the prevalence of HIV among students undergoing regular medical examination. Complete reliance on national data does not portray the exact prevalence of HIV infection within every region of the country. Hence, the study assessed the prevalence of HIV and the associated ABO blood grouping for individuals attending a tertiary institution.

The prevalence of HIV in the study is 1%. This is two times higher than the global prevalence as of 2019 (WHO, 2019). Part of the reason for this phenomenon is the low level of awareness within Rivers State of Nigeria. The prevalence of HIV is also lower than the current national prevalence which is 1.8% (Onovo et al., 2023).

The major role of ABO blood grouping is the prevention of transfusion reactions; however, some studies have associated it with the presence of some infectious diseases (Cooling, 2015; Anstee, 2010). The prevalence of HIV was highest in A<sup>-</sup> followed by AB<sup>-</sup>. This

observation opposed the study by Jacobs et al. (2023) which demonstrated higher prevalence in Rhesus positive individual.

The findings of this study highlight the importance of understanding the prevalence of HIV among students in tertiary institutions. The sero-prevalence rate observed underscores the need for targeted interventions, including HIV education, counseling, and testing services, within this population. The identification of associated risk factors can inform the development of comprehensive prevention strategies tailored to the specific needs of students. Limitations of the study, such as sampling bias is evident due to the location. Future research should explore longitudinal studies to monitor trends in HIV prevalence among students and evaluate the effectiveness of interventions aimed at reducing HIV transmission risk.

Finally, this study contributes to our understanding of the sero-prevalence of HIV among students undergoing medical examination in a tertiary institution. The findings underscore the importance of proactive measures to address HIV transmission risk factors and promote the health and well-being of students. Efforts to enhance HIV prevention and support services within tertiary institutions are crucial in mitigating the impact of HIV on young adults and fostering a healthy campus environment.

## **5. CONCLUSION**

The study demonstrates that the prevalence of HIV according to the departments at Ignatius Ajuru University of Education are ME 1 (1%), EM 5 (3%), LIB 2 (2%), SPE 0(0%), ENG 1 (0%), BIO 5 (1%), CHE 1 (1%), COM 2 (1%), MG 3 (1%) and ACC 2 (1%). The average prevalence of the study was 2.2 in 2114 or 0.0.

## **CONSENT**

All authors declare that written informed consent was obtained from the subjects.

## **ETHICAL APPROVAL**

Ethical approval was obtained from the (ethical committee of Health Services Department, Ignatius Ajuru University of Education, Port Harcourt)

## **REFERENCE**

Awofala, A .A. & Ogundele, O.E., (2018).. HIV epidemiology in Nigeria. *Saudi journal of biological sciences*, 25(4), pp.697-703.

Azuonwu, O., Erhabor, O. &Obire, O. (2012). HIV among military personnel in the Niger Delta of Nigeria. *Journal of community health*, 37, 25-31.

Centre for disease control and prevention, (2003). HIV and its transmission.

- Cheesbrough, M. (2009). District Laboratory Practice in Tropical Countries Part 1. 2nd Edition, Cambridge University Press, Cambridge, 195-216.
- Coburn, B.J., Okano, J.T., & Blower, S. (2017). Using geospatial mapping to design HIV elimination strategies for sub-Saharan Africa. *Science Translational Medicine*; 9. pmid:28356504
- Cooling L. (2015). Blood groups in infection and host susceptibility. *Clinical Microbiology Reviews*. pmid:26085552
- Cuadros, D.F., Li, J., Branscum, A.J., Akullian, A., Jia, P., & Mziray, E.N. (2017). Mapping the spatial variability of HIV infection in Sub-Saharan Africa: Effective information for localized HIV prevention and control. *Scientific Reports*; 7:9093. pmid:28831171
- Dwyer-Lindgren, L., Cork, M. A., Sligar, A., Steuben, K.M., Wilson, K.F., Provost, N.R. (2019). Mapping HIV prevalence in sub-Saharan Africa between 2000 and 2017. *Nature* 570:189–193. pmid:31092927
- Endalamaw, A., Mekonnen, M., Geremew, D. (2020). HIV/AIDS treatment failure and associated factors in Ethiopia: meta-analysis. *BMC Public Health* 20, 82 .
- Federal Ministry of Health. (2012). [www.health.gov.ng](http://www.health.gov.ng). Accessed data 12 January 2024.
- Govender, R.D., Hashim, M.J., Khan, M.A. (2021). Global Epidemiology of HIV/AIDS: A Resurgence in North America and Europe. *Journal of Epidemiology and Global Health* 11, 296–301
- Helina, Meri. (2020). United States President's Emergency Plan for AIDS Relief (PEPFAR). Nigeria Country Operational Plan (COP) 2019 Strategic Direction Summary. 2019. [https://www.state.gov/wp-content/uploads/2019/09/Nigeria\\_COP19-Strategic-Directional-Summary\\_public.pdf](https://www.state.gov/wp-content/uploads/2019/09/Nigeria_COP19-Strategic-Directional-Summary_public.pdf) (accessed 28 Mar2020).
- Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. *Clin Exp Allergy*. 2002;32(4):489-98.
- Anstee DJ. (2010). The relationship between blood groups and disease. *Blood*. pmid:20308598
- Jacobs G, Van den Berg K, Vermeulen M, Swanevelder R, Custer B, Murphy EL (2023) Association of ABO and RhD blood groups with the risk of HIV infection. *PLoS ONE* 18(4): e0284975. <https://doi.org/10.1371/journal.pone.0284975>
- McLaren, P.J. & Fellay, J. (2021). HIV-1 and human genetic variation. *Nature Review Genetics* 22, 645–657. <https://doi.org/10.1038/s41576-021-00378-0>
- Nigeria—Nigeria HIV-AIDS Indicator and Impact Survey 2018, NAIS 2018. [https://nigerianstat.gov.ng/nada/index.php/catalog/68/related\\_materials](https://nigerianstat.gov.ng/nada/index.php/catalog/68/related_materials) (accessed 23 Dec2020).
- O'brien-Carelli, C., Steuben, K., Stafford, K.A., Aliogo, R., Alagi, M., Johanns, C.K. (2022). Mapping HIV prevalence in Nigeria using small area estimates to develop a targeted HIV intervention strategy. *PLoS ONE* 17(6): e0268892.
- Onovo, A.A, Adeyemi, A., Onime, D., Kalnoky, M., KagniniwaB., Dessie, K., Lee, L., Parrish, D., Adebobola, B., Ashefor, G., Ogorry, O. and Goldstein, R. (2023). Estimation of HIV prevalence and burden in Nigeria: a Bayesian predictive modelling study. *Lancet*, 62,102098.
- Talarok, p.& Talaro, A. (2002). The RNA Viruses of medical importance. Foundation in microbiology. 4TH (EDT), Milgraw Hill, New York. Pp.750-789.
- Tomezsko, P.J., Corbin, V.D.A., Gupta, P. (2020). Determination of RNA structural diversity and its role in HIV-1 RNA splicing. *Nature* 582, 438–442. <https://doi.org/10.1038/s41586-020-2253-5>
- Weiss, R.A. (1993). How does HIV cause AIDS? *Science* 260, 1273-1279.
- World Health Organization. (2019). Global Health Observatory (GHO) data: Summary of the global HIV epidemic.. <https://www.who.int/gho/hiv/en/>. Accessed 12 Dec 2023.