

Evaluation of strawberry (*Fragaria × ananassa*) runner production in different growing media under semi-automated polyhouse

ABSTRACT :-

The experiment was laid out in RCBD, replicated 3 times with 11 treatments in different combinations i.e., T₀: Soil, T₁: Soil+ Vermicompost (1:1), T₂ : Soil + Vermicompost (1:2), T₃: Soil + Vermicompost(2:1), T₄ : Soil+ Poultry manure (1:1), T₅: Soil + Poultry manure (1:2), T₆: Soil + Poultry manure (2:1), T₇ : Soil +Vermicompost+ Poultry manure (1:1;1), T₈: Soil +Vermicompost+ Poultry manure (2:1;1), T₉: Soil +Vermicompost+ Poultry manure (1:2:1), T₁₀: Soil +Vermicompost+ Poultry manure (1:1:2). Higher growth and yield attributes of strawberry runners, higher quality attribute and also concluded that, treatment T₈ (Soil +Vermicompost+ Poultry manure (2:1:1) performed best in term of runners growth and developed of strawberry under semi automated poly house condition. The highest B:C ratio was also found in the same treatment with 2.15. Among all the treatments, treatment T₈ performed best in terms of runners growth and development of strawberry.

INTRODUCTION:-

Strawberry is one of important fruit crops in temperate region However, it can also be grown in the tropical and subtropical climate. Strawberry (*Fragaria × ananassa* Duct.), a member of the family Rosaceae, is a soft fruited, short-day herbaceous perennial plant that can successfully be grown at optimum day temperatures of 22°C to 25°C and night temperatures of 7°C to 13°C (De and Bhattacharjee, 2012). Commercially grown strawberry (*Fragaria x ananassa* Duct.) is a monoecious octoploid (2n=6x=56) hybrid of two dioecious octoploid species, namely, *Fragaria chiloensis* Duch. and *Fragaria virginiana* Duch. (Bowling 2000) with a basic chromosome number (x) of 7. Botanically, strawberry is an aggregate fruit having seeds on the surface of a red fleshy receptacle (Darnell, 2003). The last decade has witnessed the emergence of strawberry as the leading fruit in the category of soft berries. The area and production under strawberry in the world has increased logarithmically during the last two decades as much of the crop is being grown under protected structures. Strawberries are among the easiest fruits to rise in kitchen garden, decorative pots, hanging basket and are grown in flat or raised beds and protected cultivation. The area under cultivation this crop in subtropical and tropical region is very low and thus the fruit price remains very high (350-500 kg). It is the only fruit crop that starts paying back within five months from fruits and nine months from runner after transplanting of runners it easily propagates itself by the vegetative method of runner production in hilly region. However, it is difficult for the plant to propagate in plains because the mother plant cannot withstand severe hot and drought condition in June and July months and the plant usually dies above 40°C. In plain areas runner production is possible only under 75% shade net and requires proper management of field to safeguard the daughter plants of strawberry during summer.

It is main reasons the runner's production is tropical and subtropical areas are very low. In plain's strawberry growers can not afford due to higher prices per runner and also the farmers seldom go for its cultivation in plain as the planting material are costly to purchase it from the hills. Thus, this experiment was undertaken to evaluate the strawberry runners production in different growing media under semi automated polyhouse.

2. MATERIALS AND METHODS:-

2.1 Geographical location of the experimental site

The experimental site is located at a latitude of 25.41° North and longitude of 81.84 °East, with an altitude of 98 meters above the mean sea level (MSL).

2.2 Experimental details

Treatment Combinations

T₀: Soil, T₁: Soil+ Vermicompost (1:1), T₂ : Soil + Vermicompost (1:2), T₃: Soil + Vermicompost(2:1), T₄ : Soil + Poultry manure (1:1), T₅ : Soil + Poultry manure (1:2), T₆ : Soil + Poultry manure (2:1), T₇ : Soil +Vermicompost+ Poultry manure (1:1:1), T₈ : Soil +Vermicompost+ Poultry manure (2:1:1), T₉ : Soil +Vermicompost+ Poultry manure (1:2:1), T₁₀ : Soil +Vermicompost+ Poultry manure (1:1:2).

3. RESULTS AND DISCUSSION

3.1. Effect of different levels organic Manure on survival percent, numbers of days taken for initiation of runner, chlorophyll content, leaf area, leaf area index of Strawberry runners.

3.1.1. Survival Percent:-

The maximum survival percentage of plants with the treatment T₈ (Soil +Vermicompost+ Poultry manure (2:1:1)) was 83.33 Percent and the minimum survival percentage of germination was found the treatment (control) that application of organic manures covers significantly encouraged the early initiation of germination. So, it is a most important factor which determines early crop production.

Consistency in availability of nutrients through organic manures means might have supplemented the additional nutrient requirement caused due to early runner coupled

with concomitant increase in runners and consecutive fruit development. Enhanced yield was observed in strawberry due to the vegetative growth stimulation by application of organic manures resulted in a positive early in flowering, fruiting stages and increased total yield per plant. (**Madhavi *et al.*, (2021)**).

3.1.2. Number of Days Taken for Initiation of Runner After Planting:-

The organic manures treatments, minimum number of days taken by plants to reach after transplanting were recorded when the plants organic manures with Soil +Vermicompost+ Poultry manure (1:1:2) (T₁₀). It took about 28.12 days after transplanting which was significantly earlier from rest of treatments days after transplanting. Among organic manures treatments maximum number of days taken to runners initiation was observed under control (T₀) (34.56) followed by Soil +Vermicompost (1:2) (T₁) (32.45). Consistency in availability of nutrients through organic manures means might have supplemented the additional nutrient requirement caused due to early runner coupled with concomitant increase in runners and consecutive fruit development. Enhanced yield was observed in strawberry due to the vegetative growth stimulation by application of organic manures resulted in a positive early in flowering, fruiting stages and increased total yield per plant.

3.1.3. Chlorophyll Content SPAD Value:-

The Maximum (42.21 SPAD reading) chlorophyll concentration was recorded in T8 (Soil +Vermicompost+ Poultry manure (2:1:1)). It was followed by T10, Soil +Vermicompost+ Poultry manure (1:1:2) (41.20 SPAD reading), T9 (Soil +Vermicompost+ Poultry manure (1:2:1) (4.81 SPAD reading). Minimum (35.67 SPAD reading) chlorophyll concentration was reported in T0.

3.1.4. Leaf Area (cm²)

The maximum leaf area 97.75 cm² was recorded with T8 (Soil +Vermicompost+ Poultry manure (2:1:1)) followed by 96.74 cm² with T9 (Soil +Vermicompost+ Poultry manure (1:2:1)) and 96.61 cm² with T10 which were statistically at par to each other, whereas the minimum leaf area was recorded 90.26 cm² with T0 (control).

3.1.5. Leaf Area Index:-

The maximum leaf area index 29.82 was recorded with T8 (Soil +Vermicompost+ Poultry manure (2:1:1)) followed by 29.23 cm² with T10 (Soil +Vermicompost+ Poultry manure (1:1:2)) and 28.82 with T9 which were statistically at par to each other, whereas the minimum leaf area was recorded 27.22 with T0 (control). Leaf area index was increased significantly with the utilization of vermicompost and poultry manures at various treatment combinations. Increase in leaf area index might be due to increased growth of plant in the form of height and number of leaves, which accumulated more photosynthetic and thereby increased leaf area. The results are also in confirmation with the findings of reported that combined application of bio-fertilizers, vermicompost with inorganic fertilizers significantly increased the leaf area of strawberry.

Table 1:Effect of different levels organic manure on Survival percent,Numbers of days taken for initiation of runner,Chlorophyll Content, Leaf area, Leaf area index of Strawberry runners

Treatment	Survival Percentage	Numbers of days taken for initiation of runner after planting	Chlorophyll Content (SPAD Value)	Leaf area (cm²)	Leaf area index
T₀	69.67	34.56	35.67	90.26	27.22
T₁	75.33	32.45	36.30	92.52	27.57
T₂	78.67	30.67	36.79	93.64	27.87
T₃	79.33	29.89	37.12	93.92	28.05
T₄	76.33	29.58	37.40	92.56	27.71
T₅	78.33	29.34	38.45	93.59	28.07
T₆	79.33	29.11	39.34	94.94	28.46
T₇	80.00	29.01	39.40	95.74	28.69

T₈	83.33	28.56	42.21	97.75	29.82
T₉	80.00	28.35	40.87	96.74	28.82
T₁₀	82.66	28.12	41.20	96.61	29.23
F test	S	S	S	S	S
SEd±	0.73	0.59	0.63	0.66	1.12
CD at 5%	0.36	0.32	0.35	0.37	1.89
CV	3.23	6.51	5.58	2.36	3.26

3.2 Effect of different levels organic manure, Number of Runners ,Number of leaves,length of runnersof Strawberry runners

3.2.1 Number of Runners:-

The Maximum number of runners per plant with 120 DAS was recorded 34 with T8 (Soil +Vermicompost + Poultry manure) (2:1:1) followed by 32 with T10, (Soil + Vermicompost + Poultry manure) (1:1:2), which was significantly higher as compared to other treatment were as minimum numbers of runners per plant was found 28 with treatment T0 (control).

3.2.2. Number of Leaves:-

The Maximum number of leaves per plant (39) was recorded with T8 Soil +Vermicompost + Poultry manure (2:1:1). Closely followed by T9 Soil + Vermicompost + Poultry manure (1:2:1) (37) at 120 DAS. Minimum Number of leaves per plant was recorded in T0 control (32) at 120 DAS.

3.2.3. Length of Runners(cm):-

The Maximum length of runners (39 cm) was recorded with T8, Soil +Vermicompost+ Poultry manure (2:1:1). Closely followed by T9, Soil +Vermicompost+ Poultry manure (1:2:1) (37 cm) at 120 DAP. Minimum length of runners was recorded in T0 control (32 cm) at 120 DAS.

3.2.4. Spread of Runners Plant (cm):-

The Maximum plant spread at 120 DAP was seen in T8, Soil +Vermicompost+ Poultry manure (2:1:1) , 25.32 cm respectively, closely followed by T9 . Minimum plant spread was recorded in T0 control (21.47cm) respectively.

Table 2:Effect of different levels organic manureNumber of leaves,length of runners,Spread of the runnersof Strawberry runners

Treatment	Numbers of runners	Number of leaves (DAY)	Length of Runners (cm)	Spread of the runner plant (cm)
	120 DAS	120DAS	120 DAS	120 DAS
T ₀	28	32	22.27	21.47
T ₁	30	34	22.95	22.15
T ₂	28	32	23.64	22.84
T ₃	29	34	24.37	23.57
T ₄	32	37	23.27	22.47
T ₅	30	35	24.67	23.87
T ₆	30	36	25.44	24.64
T ₇	31	37	25.66	24.86
T ₈	34	39	26.12	25.32
T ₉	31	36	25.90	25.10
T ₁₀	32	37	25.79	24.99
F test	S	S	S	S
SEd±	0.55	0.68	0.41	0.41
CD at 5%	0.3	0.37	0.22	0.22
CV	6.0	6.37	5.50	5.68

3.3. CONCLUSION

On the basis of present investigation it is concluded that, treatment T₈ (soil+ vermicompost+ poultry manure) (2:1:1) perform best in terms of runners growth and development of strawberry under semi automated polyhouse condition.

References:-

- A.O.A.C.(1990).**Official methods of analysis, Association of Official Analytical Chemists. 5th edition Washington, D.C., USA.
- Abu-Zahra, T.R. and Tahboub, A.A.(2008).** Strawberry (*Fragaria × ananassa* Duch) Growth, Flowering and Yielding as Affected by Different Organic Matter Sources. *Inter. J. of Botany*, **4**: pp. 481-485, 2008.
- Ahmad, D. and Mohammad, J.(2012).** Impact of integrated organic nutrient handling on fruit yields and quality of strawberry. *Journal of Ornamental and Horticultural Plants*, **2**(4): 251-256.
- Ahmadi, E., Honnabyraiah, M.K. Alur, A.S., Adiga, J.D. and Venkat Rao, V. (2017).** Impact of Integrated Nutrient Management on yield and quality parameters of strawberry (*Fragaria × ananassa* Duch.) Cv. "Sabrina" under polyhouse. *Int. J. Curr. Micro. App. Sci.*, **6**(9): 3481-3487.
- Ahmet, E., Hilal, E. Yildiz and Sezai, E. (2010).** Effects of plant growth promoting bacteria (PGPB) on yield, growth and nutrient contents of organically grown strawberry. *Scientia Horticulturae*, **12**(4), 62–66,
- Anonymous(2017).** Horticulture Statistics At Glance 2017, Horticulture Statistics Division, Department of Agriculture, Cooperation & Farmers Welfare Ministry of Agriculture & Farmers Welfare Government of India. p.47.
- Arancon, N.Q., Edwards, C.A., Bierman, P., Welch, C. and Metzger, J.D.(2004).** Influences of vermicomposts on field strawberries: Effect on growth and yields. *Bio-resource Tech.*, **93**(2): 145-153.

- Arancon, N.Q., Edwards, C.A., Lee, S. & Byrne, R. (2006).** Effects of humic acids from vermicompost on plant growth. *European J. Soil Bio.*, 46:65-69.
- Asrey, R. and Singh, R. (2004).** Evaluation of strawberry varieties under semi-arid irrigated region of Punjab. *Indian J. of Hort*, 61(2):122-124.
- Baba, Z. A., Zargar, M. Y. and Mir, S. A. (2010).** Effect of inorganic and biofertilizers on soil physicochemical properties and micronutrient availability in strawberry (*Fragaria × ananassa* Duch.). *Asian Journal of Soil Sciences*, 5(1): 90-93, 2010.
- Bakshi, P., Bhat, D. J., Wali, V. K., Sharma, A. and Iqbal, M. (2014).** Growth, yield and quality of strawberry (*Fragaria × ananassa* Duch.) cv. Chandler as influenced by various mulching materials. *African J. Agri. Res.*, 9(7):701-706.
- Bhat, N.H., (1999).** Response of strawberry cultivars to varied levels of organic manure. M.Sc. Thesis, SK University of Agricultural Sciences and Technology of Kashmir, Srinagar, J&K, India.
- Bilal, A. P., Singh, D. B. and Ahmad, M. F. (2009).** Response of strawberry to biofertilizers under sub-tropical condition of Allahabad. *Prog. Hort.*, 41(1):94-97.
- Dadashpour, A. and Jouki, M. (2012).** Impact of integrated organic nutrient handling on fruit yields and quality of strawberry cv. Kurdistan in Iran. *J. Orntrl. & Hort. Plants*, 2(4): 251-256.
- Dar, G.A., Reshi, T.A., Sheikh, M.A. and Shagoo, P.A. (2010).** Effect of nitrogen, phosphorus and potassium on growth, yield and quality of strawberry (*Fragaria × ananassa* Duch) cv. Sweet Charlie. *Environment and Ecology*, 28(2B):1216-1219.
- El-Hamid, A.S. Aza, Abbou, A.A.; Mansour, S.A.A. and El. Sayed, A.A.A. (2006).** Effect of some bio-fertilizers on the yield, and fruit quality of strawberry. *Anl. Agric. Sci. Moshtohor*, 44(10): 251-264.
- Gupta, A.K. and Tripathi, V.K. (2012).** Efficacy of Azotobacter and vermicompost alone and

in combination on vegetative growth, flowering and yield of strawberry (*Fragaria x ananassa* Duch.) Cv. Chandler. *Progressive Hort.*, **44**(2): 256-261.

Iqbal, U., Wali, V. K., Ravikher and Jamwal, M.(2008). Effect of poultry manure, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. *Haryana J. of Hort. Sci.*, **37**(1/2):28-30.

Iqbal,U.,Wali,V.K.,RavikherandJamwal,M.(2009). Effect of FYM, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. *Notulae Botanicae, Horti Agrobotanici*, **37**(1):139-143.

Jain, N., Bahadur, V. and Mani, A. (2018). Influence of Integrated Nutrient Management practices on physio-chemical attributes in strawberry (*Fragaria x ananassa* Duch.) cv. sweet charlie. *Inno. Farm.*, **3**(1):1-5.

Karma Beer, Kumar S., Gupta, A.K. and Syamal, M.M. (2017). Effect of organic, inorganic and bio-fertilizer on growth, flowering, yield and quality of strawberry (*Fragaria x Ananassa* Duch.) cv. Chandler. *Int. J. Curr. Microbiol. App. Sci.*, **6**(5): 2932-2939.

Khalid, S., Qureshi, K.M., Hafiz, I.A., Khan, K.S. and Qureshi, U.S. (2013). Effect of organic amendments on vegetative growth, fruit and yield quality of strawberry. *Pakistan J. of Agri. Res.*, **26**(2):104-112.

Kirad, K.S., Barche, S. and Singh, D.B. (2009). Response of Integrated Nutrient Management in strawberry (*Fragaria x ananassa* Duch.). *Acta Horti.*, **84**(2):653-656.

Kumar, N., Ram, R.B. and Mishra, P.K. (2015). Effect of vermicompost and *Azotobacter* on quality parameters of strawberry (*Fragaria x ananassa* Duch.) cv. Sweet Charlie. *Inter. J. Agri. Sci. and Res.*, **5**(4): 269-276.

Kumar, R., Collis, J.P., Singh, S. Moharana, D., Rout, S. and Patra, S.S. (2015). Effect of different levels of NPK in combination with FYM on quality of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *Int. J. Agri. and Food Sci.*, **5**(3):92-97.

Lata, R., Dwivedi, D.H., Ram, R.B., Meena, M.L. and Babu, M.

- (2013). Impact of Integrated Nutrient Management on growth parameters of Strawberry Cv. Chandler under Sub-tropical conditions of Lucknow, *I. J. A. B. R.*, Vol. 3(3):418-421, 2013.
- Mishra, A.N. and V.K. Tripathi, V.K. (2011).** Influence of Different Levels of Azotobacter, PSB alone and in combination on vegetative growth, flowering, yield and quality of strawberry cv. Chandler. *Inter. J. Applied Agri. Res.*, 6(3): 203-210.
- Nazir, N. (2005).** Studies on organic farming techniques for production of quality strawberry (*Fragaria × ananassa* Duch.). M Sc. Thesis, SK University of Agricultural Sciences and Technology of Kashmir, Srinagar, Jammu and Kashmir, India.
- Nazir, N., Singh, S.R., Aroosa, K., Masarat, S. and Shabeena, M. (2006).** Yield and growth of strawberry cultivar Senga Sengana as influenced by integrated organic nutrient management system. *Enl. & Eco.*, 24(3):651-654.
- Nazir, N., Singh, S.R., Sharma, M.K. and Banday, F.A. (2012).** Effect of integrated organic nutrient sources on soil nutrient status and microbial population in strawberry field. *Indian J. Hort.*, 69(2): 177-180.
- Nowsheen, N., Singh, S.R., Aroosa, K. and Masarat, S. (2006).** Yield and growth of strawberry cv Senga Segana. *Environ. and Eco.*, 245(3):651-654.
- Pesakovic, M., Karakljic, S., Milenkovic, Z.S. and Mitrovic, O. (2013).** Biofertilizer affecting yield related characteristics of strawberry (*Fragaria × ananassa* Duch.) and soil micro-organisms. *Sci. Hortic.*, 150:238-243, 2013.
- Rajbir, S., Sharma, R. R. and Singh, D. B. (2010).** Effect of vermicompost on plant growth, fruit yield and fruit quality of strawberries in irrigated arid region of northern plains. *Indian J. of Hort.*, 67(3): 318-321.
- Rana, R.K. and Chandel, J.S. (2003).** Effect of bio-fertilizer and nitrogen on growth, yield and fruit quality of strawberry. *Prog. Hort.*, 35(1): 23-30.
- Ranjit, C. and Bandyopadhyay, S. (2014).** Studies on effect of organic, inorganic and biofertilizers

on plant nutrient status and availability of major nutrients in tomato. *Inter. J. of Bio-resource and Stress Management*, **5**(1):93-97, 2014.

Sahoo, S.K. and Singh, D.B. (2005). Effect of different levels of bio-fertilizers on growth, yield and quality of strawberry (*Fragaria x annanasa* Duch.) cv. Sweet Charlie. *Orissa J. Hort.*, **32**(2): 82-85.

Singh, A. and Singh, J.N. (2006). Studies on influence of bio-fertilizers and bioregulators on flowering, yield and fruit quality of strawberry cv. Sweet Charley. *Annals of Agril. Res.*, **27**(3):261-264.

Singh, A. and Singh, J.N. (2009). Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. *Indian J. Hort.*, **66**:220-24.

UNDER PEER REVIEW