

Evaluation of Strawberry (*Fragaria × Ananassa*) Runner Production in Different Growing Media under Semi-automated Polyhouse

ABSTRACT

The experiment was laid out in RCBD, replicated 3 times with 11 treatments in different combinations i.e., T₀: Soil, T₁: Soil+ Vermicompost (1:1), T₂: Soil + Vermicompost (1:2), T₃: Soil + Vermicompost(2:1), T₄: Soil + Poultry manure (1:1), T₅: Soil + Poultry manure (1:2), T₆: Soil + Poultry manure (2:1), T₇: Soil +Vermicompost+ Poultry manure (1:1;1), T₈: Soil +Vermicompost+ Poultry manure (2:1;1), T₉: Soil +Vermicompost+ Poultry manure (1:2:1), T₁₀: Soil +Vermicompost+ Poultry manure (1:1:2). Higher growth and yield attributes of strawberry runners, higher quality attribute and also concluded that, treatment T₈ (Soil +Vermicomposted+ Poultry manure (2:1:1) performed best in term of runners growth and developed of strawberry under semi automated poly house condition. The highest B:C ratio was also found in the same treatment with 2.15. Among all the treatments, treatment T₈ performed best in terms of runners growth and development of strawberry.

Keywords: **Soil, Vermicompost, Poultry manure , plant**

1. INTRODUCTION

“Strawberry is one of important fruit crops in temperate region However, it can also be grown in the tropical and subtropical climate. Strawberry (*Fragaria × ananassa* Duct.), a member of the family Rosaceae, is a soft fruited, short-day herbaceous perennial plant that can successfully be grown at optimum day temperatures of 22°C to 25°C and night temperatures of 7°C to 13°C” (De and Bhattacharjee, 2012). “Commercially grown strawberry (*Fragaria x ananassa* Duct.) is a monoecious octoploid (2n=6x=56) hybrid of two dioecious octoploid species, namely, *Fragaria chiloensis* Duch. and *Fragaria virginiana* Duch. (Bowling 2000) with a basic chromosome number (x) of 7. Botanically, strawberry is an aggregate fruit having seeds on the surface of a red fleshy receptacle” (Darnell, 2003). The last decade has witnessed the emergence of strawberry as the leading fruit in the category of soft berries. The area and production under strawberry in the world has increased logarithmically during the last two decades as much of the crop is being grown under protected structures.

“Strawberries are among the easiest fruits to rise in kitchen garden, decorative pots, hanging basket and are grown in flat or raised beds and protected cultivation. The area under cultivation this crop in subtropical and tropical region is very low and thus the fruit price remains very high (350-500 kg). It is the only fruit crop that starts paying back within five months from fruits and nine months from runner after transplanting of runners it easily propagates itself by the vegetative method of runner production in hilly region” [1,2]. However, it is difficult for the plant to propagate in plains because the mother plant cannot withstand severe hot and drought condition in June and July months and the plant usually dies above 40°C. In plain areas runner production is possible only under 75% shade net and requires proper management of field to safeguard the daughter plants of strawberry during summer.

It is main reasons the runner's production is tropical and subtropical areas are very low. In plain's strawberry growers can not afford due to higher prices per runner and also the farmers seldom go for its cultivation in plain as the planting material are costly to purchase it from the hills. Thus, this experiment was undertaken

to evaluate the strawberry runners production in different growing media under semi automated polyhouse.

2. MATERIALS AND METHODS

2.1 Geographical Location of the Experimental Site

The experimental site is located at a latitude of 25.41° North and longitude of 81.84 °East, with an altitude of 98 meters above the mean sea level (MSL).

2.2 Experimental Details

2.2.1 Treatment combinations

T₀: Soil, T₁: Soil+ Vermicompost (1:1), T₂: Soil + Vermicompost (1:2), T₃: Soil + Vermicompost(2:1), T₄: Soil + Poultry manure (1:1), T₅: Soil + Poultry manure (1:2), T₆: Soil + Poultry manure (2:1), T₇: Soil +Vermicompost+ Poultry manure (1:1;1), T₈: Soil +Vermicompost+ Poultry manure (2:1;1), T₉: Soil +Vermicompost+ Poultry manure (1:2:1), T₁₀: Soil +Vermicompost+ Poultry manure (1:1:2).

3. RESULTS AND DISCUSSION

3.1 Effect of Different Levels Organic Manure on Survival Percent, Numbers of Days taken for Initiation of Runner, Chlorophyll Content, Leaf Area, Leaf Area Index of Strawberry Runners

3.1.1 Survival percent

The maximum survival percentage of plants with the treatment T₈ (Soil +Vermicompost+ Poultry manure (2:1:1)) was 83.33 Percent and the minimum survival percentage of germination was found the treatment (control) that application of organic manures covers significantly encouraged the early initiation of germination. So, it is a most important factor which determines early crop production.

Consistency in availability of nutrients through organic manures means might have supplemented the additional nutrient requirement caused due to early runner coupled with concomitant increase in runners and consecutive fruit development. Enhanced yield was observed in strawberry due to the vegetative growth stimulation by application of organic manures resulted in a positive early in flowering, fruiting

stages and increased total yield per plant. (Madhavi *et al.*, (2021).

3.1.2 Number of days taken for initiation of runner after planting

The organic manures treatments, minimum number of days taken by plants to reach after transplanting were recorded when the plants organic manures with Soil +Vermicompost+ Poultry manure (1:1:2) (T₁₀). It took about 28.12 days after transplanting which was significantly earlier from rest of treatments days after transplanting. Among organic manures treatments maximum number of days taken to runners initiation was observed under control (T₀) (34.56) followed by Soil +Vermicompost (1:2) (T₁) (32.45). Consistency in availability of nutrients through organic manures means might have supplemented the additional nutrient requirement caused due to early runner coupled with concomitant increase in runners and consecutive fruit development. Enhanced yield was observed in strawberry due to the vegetative growth stimulation by application of organic manures resulted in a positive early in flowering, fruiting stages and increased total yield per plant.

3.1.3 Chlorophyll content SPAD value

The Maximum (42.21 SPAD reading) chlorophyll concentration was recorded in T₈ (Soil +Vermicompost+ Poultry manure (2:1:1)). It was followed by T₁₀, Soil +Vermicompost+ Poultry manure (1:1:2) (41.20 SPAD reading), T₉ (Soil +Vermicompost+ Poultry manure (1:2:1)) (4.81 SPAD reading). Minimum (35.67 SPAD reading) chlorophyll concentration was reported in T₀.

3.1.4 Leaf area (cm²)

The maximum leaf area 97.75 cm² was recorded with T₈ (Soil +Vermicompost+ Poultry manure (2:1:1)) followed by 96.74 cm² with T₉ (Soil +Vermicompost+ Poultry manure (1:2:1)) and 96.61 cm² with T₁₀ which were statistically at par to each other, whereas the minimum leaf area was recorded 90.26 cm² with T₀ (control).

3.1.5 Leaf area index

The maximum leaf area index 29.82 was recorded with T₈ (Soil +Vermicompost+ Poultry manure (2:1:1)) followed by 29.23 cm² with T₁₀ (Soil +Vermicompost+ Poultry manure (1:1:2)) and 28.82 with T₉ which were statistically at par to each other, whereas the minimum leaf area was recorded 27.22 with T₀ (control). Leaf area index was increased significantly with the utilization of vermicompost and poultry manures

at various treatment combinations. Increase in leaf area index might be due to increased growth of plant in the form of height and number of leaves, which accumulated more photosynthetic and thereby increased leaf area. The results are also in confirmation with the findings of reported that combined application of bio-fertilizers, vermicompost with inorganic fertilizers significantly increased the leaf area of strawberry.

3.2 Effect of Different Levels Organic Manure, Number of Runners, Number of Leaves, Length of Runners of Strawberry Runners

3.2.1 Number of runners

The Maximum number of runners per plant with 120 DAS was recorded 34 with T8 (Soil +Vermicompost + Poultry manure) (2:1:1) followed by 32 with T10, (Soil + Vermicompost + Poultry manure) (1:1:2), which was significantly

higher as compared to other treatment were as minimum numbers of runners per plant was found 28 with treatment T0 (control).

3.2.2 Number of Leaves

The Maximum number of leaves per plant (39) was recorded with T8 Soil +Vermicompost + Poultry manure (2:1:1). Closely followed by T9 Soil + Vermicompost + Poultry manure (1:2:1) (37) at 120 DAS. Minimum Number of leaves per plant was recorded in T0 control (32) at 120 DAS.

3.2.3 Length of Runners (cm)

The Maximum length of runners (39 cm) was recorded with T8, Soil +Vermicompost+ Poultry manure (2:1:1). Closely followed by T9, Soil +Vermicompost+ Poultry manure (1:2:1) (37 cm) at 120 DAP. Minimum length of runners was recorded in T0 control (32 cm) at 120 DAS.

Table 1. Effect of different levels organic manure on Survival percent, Numbers of days taken for initiation of runner, Chlorophyll Content, Leaf area, Leaf area index of Strawberry runners

| Treatment | Survival Percentage | Numbers of days taken for initiation of runner after planting | Chlorophyll Content (SPAD Value) | Leaf area (cm ²) | Leaf area index |
|-----------------|---------------------|---|----------------------------------|------------------------------|-----------------|
| T ₀ | 69.67 | 34.56 | 35.67 | 90.26 | 27.22 |
| T ₁ | 75.33 | 32.45 | 36.30 | 92.52 | 27.57 |
| T ₂ | 78.67 | 30.67 | 36.79 | 93.64 | 27.87 |
| T ₃ | 79.33 | 29.89 | 37.12 | 93.92 | 28.05 |
| T ₄ | 76.33 | 29.58 | 37.40 | 92.56 | 27.71 |
| T ₅ | 78.33 | 29.34 | 38.45 | 93.59 | 28.07 |
| T ₆ | 79.33 | 29.11 | 39.34 | 94.94 | 28.46 |
| T ₇ | 80.00 | 29.01 | 39.40 | 95.74 | 28.69 |
| T ₈ | 83.33 | 28.56 | 42.21 | 97.75 | 29.82 |
| T ₉ | 80.00 | 28.35 | 40.87 | 96.74 | 28.82 |
| T ₁₀ | 82.66 | 28.12 | 41.20 | 96.61 | 29.23 |
| F test | S | S | S | S | S |
| SEd± | 0.73 | 0.59 | 0.63 | 0.66 | 1.12 |
| CD at 5% | 0.36 | 0.32 | 0.35 | 0.37 | 1.89 |
| CV | 3.23 | 6.51 | 5.58 | 2.36 | 3.26 |

Table 2. Effect of different levels organic manure Number of leaves, length of runners, Spread of the runners of Strawberry runners

| Treatment | Numbers of runners 120 DAS | Number of leaves (DAY) 120DAS | Length of Runners (cm) 120 DAS | Spread of the runner plant (cm) 120 DAS |
|----------------|----------------------------|-------------------------------|--------------------------------|---|
| T ₀ | 28 | 32 | 22.27 | 21.47 |
| T ₁ | 30 | 34 | 22.95 | 22.15 |
| T ₂ | 28 | 32 | 23.64 | 22.84 |
| T ₃ | 29 | 34 | 24.37 | 23.57 |
| T ₄ | 32 | 37 | 23.27 | 22.47 |
| T ₅ | 30 | 35 | 24.67 | 23.87 |
| T ₆ | 30 | 36 | 25.44 | 24.64 |

| | | | | |
|-----------------|------|------|-------|-------|
| T ₇ | 31 | 37 | 25.66 | 24.86 |
| T ₈ | 34 | 39 | 26.12 | 25.32 |
| T ₉ | 31 | 36 | 25.90 | 25.10 |
| T ₁₀ | 32 | 37 | 25.79 | 24.99 |
| F test | S | S | S | S |
| SEd± | 0.55 | 0.68 | 0.41 | 0.41 |
| CD at 5% | 0.3 | 0.37 | 0.22 | 0.22 |
| CV | 6.0 | 6.37 | 5.50 | 5.68 |

3.2.4 Spread of runners plant (cm)

The Maximum plant spread at 120 DAP was seen in T₈, Soil +Vermicompost+ Poultry manure (2:1:1) , 25.32 cm respectively, closely followed by T₉ . Minimum plant spread was recorded in T₀ control (21.47cm) respectively.

4. CONCLUSION

On the basis of present investigation it is concluded that, treatment T₈ (soil+ vermicompost+ poultry manure) (2:1:1) perform best in terms of runners growth and development of strawberry under semi automated polyhouse condition.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Amin B, Qureshi KM, Ghani A, Mahmood A, Shah SA, Khan MF, Khalid S, Iqbal S, Anwar I. Effect of Different Environments And GA3 On Growth And Runners Production Of Strawberry (Frageria Ananasa). Journal of Pharmaceutical Negative Results. 2023;208-18.
- Tarafdar M, Mishra S, Singh RK, Kumar A, Ekka SK. Effect on Vegetative Growth and Development of Strawberry (Frageria ananassa) in Potting Mixture System under Shade Net Conditions cv. Winter Dawn. International Journal of Environment and Climate Change. 2023;13(10):799-808.
- AOAC. Official methods of analysis, Association of Official Analytical Chemists. 5th edition Washington, D.C., USA; 1990.
- Abu-Zahra TR, Tahboub AA. Strawberry (*Frageria x ananassa* Duch) Growth, Flowering and Yielding as Affected by Different Organic Matter Sources. Inter. J. of Botany. 2008;4:481-485.
- Ahmad D, Mohammad J. Impact of integrated organic nutrient handling on fruit yields and quality of strawberry.

- Journal of Ornamental and Horticultural Plants. 2012;2(4):251-256.
- Ahmadi E, Honnabyraiah MK, Alur AS, Adiga JD, Venkat Rao V. Impact of Integrated Nutrient Management on yield and quality parameters of strawberry (*Frageria x ananassa* Duch.) Cv. "Sabrina" under polyhouse. Int. J. Curr. Micro. App. Sci. 2017;6(9):3481-3487.
 - Ahmet E, Hilal E, Yildiz, Sezai E. Effects of plant growth promoting bacteria (PGPB) on yield, growth and nutrient content of organically grown strawberry. Scientia Horticulturae. 2010;12(4):62-66.
 - Anonymous. Horticulture Statistics at Glance, Horticulture Statistics Division, Department of Agriculture, Cooperation and Farmers Welfare Ministry of Agriculture and Farmers Welfare Government of India. 2017;47.
 - Arancon NQ, Edwards CA, Bierman P, Welch C, Metzger JD. Influence of vermicomposts on field strawberries: Effect on growth and yields. Bio-resource Tech. 2004;93(2):145-153.
 - Arancon NQ, Edwards CA, Lee S, Byrne R. Effects of humic acids from vermicompost on plant growth. European J. Soil Bio. 2006;46:65-69.
 - Asrey R, Singh R. Evaluation of strawberry varieties under semi-arid irrigated region of Punjab. Indian J. of Hort. 2004;61(2):122-124.
 - Baba ZA, Zargar MY, Mir SA. Effect of inorganic and biofertilizers on soil physicochemical properties and micronutrient availability in strawberry (*Frageria x ananassa* Duch.). Asian Journal of Soil Sciences. 2010;5(1):90-93.
 - Bakshi P, Bhat DJ, Wali VK, Sharma A, Iqbal M. Growth, yield and quality of strawberry (*Frageria x ananassa* Duch.) cv. Chandler as influenced by various mulching materials. African J. Agri. Res. 2014;9(7):701-706.
 - Bhat NH. Response of strawberry cultivars to varied levels of organic manure. M.Sc. Thesis, SK University of Agricultural

- Sciences and Technology of Kashmir, Srinagar, J and K, India; 1999.
15. Bilal AP, Singh DB, Ahmad MF. Response of strawberry to biofertilizers under subtropical condition of Allahabad. *Prog. Hort.* 2009;41(1):94-97.
 16. Dadashpour A, Jouki M. Impact of integrated organic nutrient handling on fruit yields and quality of strawberry cv. Kurdistan in Iran. *J. Orntl. and Hort. Plants.* 2012;2(4):251-256.
 17. Dar GA, Reshi TA, Sheikh MA, Shagoo PA. Effect of nitrogen, phosphorus and potassium on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch) cv. Sweet Charlie. *Environment and Ecology.* 2010;28(2B):1216-1219.
 18. El-Hamid AS, Aza, Abbou AA, Mansour SAA, Amd El. Sayed AAA. Effect of some bio-fertilizers on the yield, and fruit quality of strawberry. *Anl. Agric. Sci. Moshtohor.* 2006;44(10):251-264.
 19. Gupta AK, Tripathi VK. Efficacy of Azotobacter and vermicompost alone and in combination on vegetative growth, flowering and yield of strawberry (*Fragaria x ananassa* Duch.) Cv. Chandler. *Progressive Hort.* 2012;44(2):256-261.
 20. Iqbal U, Wali VK, Ravikher, Jamwal M. Effect of poultry manure, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. *Haryana J. of Hort. Sci.* 2008;37(1/2):28-30.
 21. Iqbal U, Wali VK, Ravikher, Jamwal M. Effect of FYM, urea and *Azotobacter* on growth, yield and quality of strawberry cv. Chandler. *Notulae Botanicae, Horti Agrobotanici.* 2009;37(1):139-143.
 22. Jain N, Bahadur V, Mani A. Influence of Integrated Nutrient Management practices on physio-chemical attributes in strawberry (*Fragaria x ananassa* Duch.) cv. sweet charlie. *Inno. Farm.* 2018;3(1):1-5.
 23. Karma Beer, Kumar S, Gupta AK, Syamal MM. Effect of organic, inorganic and bio-fertilizer on growth, flowering, yield and quality of strawberry (*Fragaria x Ananassa* Duch.) cv. Chandler. *Int. J. Curr. Microbiol. App. Sci.* 2017;6(5):2932-2939.
 24. Khalid S, Qureshi KM, Hafiz IA, Khan KS, Qureshi US. Effect of organic amendments on vegetative growth, fruit and yield quality of strawberry. *Pakistan J. of Agri. Res.* 2013;26(2):104-112.
 25. Kirad KS, Barche S, Singh DB. Response of Integrated Nutrient Management in strawberry (*Fragaria x ananassa* Duch.). *Acta Horti.* 2009;84(2):653-656.
 26. Kumar N, Ram RB, Mishra PK. Effect of vermicompost and *Azotobacter* on quality parameters of strawberry (*Fragaria x ananassa* Duch.) cv. Sweet Charlie. *Inter. J. Agri. Sci. and Res.* 2015;5(4):269-276.
 27. Kumar R, Collis JP, Singh S, Moharana D, Rout S, Patra SS. Effect of different levels of NPK in combination with FYM on quality of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler. *Int. J. Agri. and Food Sci.* 2015;5(3):92-97.
 28. Lata R, Dwivedi DH, Ram RB, Meena ML, Babu M. Impact of Integrated Nutrient Management on growth parameters of Strawberry Cv. Chandler under Sub-tropical conditions of Lucknow, I. J. A. B. R. 2013;3(3):418-421.
 29. Mishra AN, Tripathi VK. Influence of Different Levels of *Azotobacter*, *PSB* alone and in combination on vegetative growth, flowering, yield and quality of strawberry cv. Chandler. *Inter. J. Applied Agri. Res.* 2011;6(3):203-210.
 30. Nazir N. Studies on organic farming techniques for production of quality strawberry (*Fragaria x ananassa* Duch.). M Sc. Thesis, SK University of Agricultural Sciences and Technology of Kashmir, Srinagar, Jammu and Kashmir, India; 2005.
 31. Nazir N, Singh SR, Aroosa K, Masarat S, Shabeena M. Yield and growth of strawberry cultivar Senga Sengana as influenced by integrated organic nutrient management system. *Enl. and Eco.* 2006;24(3):651-654.
 32. Nazir N, Singh SR, Sharma MK, Banday FA. Effect of integrated organic nutrient sources on soil nutrient status and microbial population in strawberry field. *Indian J. Hort.* 2012;69(2):177-180.
 33. Nowsheen N, Singh SR, Aroosa K, Masarat S. Yield and growth of strawberry cv Senga Segana. *Environ. and Eco.* 2006;245(3):651-654.
 34. Pesakovic M, Karakljajic S, Milenkovic ZS, Mitrovic O. Biofertilizer affecting yield related characteristics of strawberry (*Fragaria x ananassa* Duch.) and soil micro-organisms. *Sci. Hortic.* 2013;150:238-243.
 35. Rajbir S, Sharma RR, Singh DB. Effect of vermicompost on plant growth, fruit yield and fruit quality of strawberries in irrigated arid region of northern plains. *Indian J. of Hort.* 2010;67(3): 318-321.

36. Rana RK, Chandel JS. Effect of bio-fertilizer and nitrogen on growth, yield and fruit quality of strawberry. *Prog. Hort.* 2003;35(1):23-30.
37. Ranjit C, Bandyopadhyay S. Studies on effect of organic, inorganic and bio-fertilizers on plant nutrient status and availability of major nutrients in tomato. *Inter. J. of Bio-resource and Stress Management.* 2014;5(1):93-97.
38. Sahoo SK, Singh DB. Effect of different levels of bio-fertilizers on growth, yield and quality of strawberry (*Fragaria x annanasa* Duch.) cv. Sweet Charlie. *Orissa J. Hort.* 2005;32(2):82-85.
39. Singh A, Singh JN. Studies on influence of bio-fertilizers and bioregulators on flowering, yield and fruit quality of strawberry cv. Sweet Charley. *Annals of Agril. Res.* 2006;27(3):261-264.
40. Singh A, Singh JN. Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. *Indian J. Hort.* 2009;66:220-24.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.