

1 ***In vitro* evaluation of fungicides against anthracnose of betelvine (*Piper betle* L.)**

2
3 **ABSTRACT**

4 **Background:** Betelvine is important commercial crop and the most profitable among all cultivated
5 crops, which plays a vital role in the overall livelihood security of farm families. Diseases are the
6 major yield constraints of crop plants. One of the most serious fungal diseases of dragon fruit is
7 anthracnose caused by *Colletotrichum* species. Since less information available on anthracnose of
8 betel vine, this study was undertaken.

9 **Methods:** The efficacy of non-systemic, systemic and combi fungicides were tested against
10 *Colletotrichum* spp. using poisoned food technique (Vincent (1947) under *in vitro* condition. Six
11 non-systemic fungicides Chlorothalonil 75 % WP, Captan 50 % WP, Mancozeb 75 % WP, Copper
12 oxychloride 50 % WP, Propineb 70 % WP and Copper hydroxide 53.8 % at (250 ppm, 500 ppm and
13 1000 ppm), six systemic fungicides Hexaconazole 5 % EC, Propiconazole 25 % EC, Azoxystrobin
14 25% SC, Tebuconazole 25.9 % EC, Difenconazole 25 % EC and Picoxystrobin 22.5 % SC at
15 (100ppm, 150ppm, 250ppm) and six combi fungicides Propiconazole 13.9 % + Difenconazole
16 13.9 % EC, Tebuconazole 50 % + Trifloxystrobin 25 % WG, Fluopyram 200 g/L + Tebuconazole
17 200 g/L SC, Fluxopyroxad 250 g/l + pyraclostrobin 250g/L, Fluopyram 250 g/L + Trifloxystrobin
18 250 g/L SC, Azoxystrobin 16.7 % + Tricyclazole 33.3 % SC at (150 ppm, 250 ppm, 500 ppm) were
19 evaluated.

20 **Result:** Among six non-systemic fungicides evaluated against *C. gloeosporioides* Copper hydroxide
21 gave 69.90 per cent inhibition which was superior over all other fungicides evaluated and least
22 inhibition was recorded with Mancozeb 40.30 per-cent. Among six different systemic fungicides
23 evaluated against *C. gloeosporioides* Difenconazole, Tebuconazole were found best with inhibition
24 per-centage of 98.28 and 95.17. Out of the six evaluated combination products, propiconazole +
25 difenconazole exhibited the highest inhibition rate at 99.78%. Following closely, Fluopyram 200
26 g/L + Tebuconazole 200 g/L SC and Tebuconazole 50% EC + Trifloxystrobin 25% WG displayed
27 inhibition rates of 89.47% and 87.50% respectively.

28 **Key words:** Betelvine, Anthracnose, *Colletotrichum gloeosporioides*, Fungicide

Comment [11]: Don't use short forms in abstract

Comment [12]: Specify the specie name

Comment [13]: The author cited very old literature. Replaced it

Comment [14]: From where you obtain fungal cultures? Mention source
If you isolate it then mention accession number

Comment [15]: Add "and result revealed that the" between *gloeosporioides* and Copper

Comment [16]: Rewrite this sentence

Comment [17]: Arrange keywords alphabetically

30

INTRODUCTION

31 Betelvine (*Piper betle* L.), widely known as "paan" in the Indian sub-continent, has a long
32 ancient history in India and occupies a significant place in the everyday life of the people as it is
33 used in rituals and as medicine to cure many diseases and disorders. Malaysia is the most probable
34 place of origin of the Betelvine (Chattopadhyay and Maity, 1967). It belongs to the family
35 Piperaceae. Betelvine is a perennial, dioecious, shade-loving, aromatic, evergreen root climber with
36 glossy heart-shaped leaves and white catkin (Hiralal Jana, 2016). It is mainly grown in the tropics
37 and subtropical regions, for its leaves are used as a chewing stimulant. In India, betelvine is grown
38 throughout the country and as an important cash crop in southern parts, mainly in Andhra Pradesh,
39 Karnataka, Kerala, and Tamil Nadu. Betel vine is also cultivated in Assam, Bihar, Madhya Pradesh,
40 Maharashtra, Orissa, Tripura, Uttar Pradesh, and West Bengal with an estimated area of 53,539 ha
41 (Ray, 2008). It is the most important cash crop, and that adequately justifies its nomenclature as the
42 "Green gold of India" (Nutankumar *et al.*, 2014). Betelvine leaves and areca nut are used in many
43 occasions like Hindu religious ceremonies, wedding ceremonies, and pujas. Chewing of pan leaf is
44 an ancient habit that has existed for more than 2000 years (Kumar *et al.*, 2014). The essential
45 diseases challenging betel leaf production are foot rot complex (*Phytophthora* spp, *Sclerotium*
46 *rolfsii*, *Rhizoctonia solani*, *Macrophomina phaseolina*, and *Pythium vexans*), anthracnose
47 (*Colletotrichum* spp.), leaf rot (*Colletotrichum* spp.), Powdery mildew (*Oidium piperis*) and
48 bacterial leaf spot (*Xanthomonas betlicola*). The symptoms of betel vine anthracnose disease on the
49 stem caused by *Colletotrichum* spp. at first appear as tiny, black, circular specks on the green bark
50 of the stem. If conditions are dry, then these specks usually do not increase in size and remain as a
51 black stain on the surface of the stem. The leaf spot disease appears only after the rain and affects
52 only betelvine leaves. The disease infection will not spread to the vine. Environmental factors such
53 as temperature, rainfall, relative humidity, and shade in baroja play vital roles in the disease
54 development. The high relative humidity (92 %) was critical for severe leaf spot disease and led to
55 heavy loss in betel vine crops (Dasgupta and Sen, 1999).

Comment [18]: Don't cite old literature, replace it.

Comment [19]: Reference is missing in reference list.

Comment [110]: Missing reference

Comment [111]: Old literature

56

MATERIAL AND METHODS

57 The poisoned food technique (Nene and Thapliyal 1993) was followed to evaluate the efficacy of
58 non-systemic, systemic fungicides and combi products in inhibiting the mycelial growth of
59 pathogen. The fungus was grown on potato dextrose agar medium for 12 days prior to setting up the

Comment [112]: Cite the literature that published in 20

60 experiment. The potato dextrose agar medium was prepared and melted. The fungicidal suspension
61 was added to the melted medium to obtain the required concentrations on commercial formulation
62 basis of the fungicide. Twenty ml of poisoned medium was poured in each sterilized Petri plates.
63 Suitable check was maintained without addition of fungicide. Mycelial disc of 5 mm was taken
64 from the periphery of 12 days old colony was placed in the center of Petri plates and incubated at 27
65 $\pm 1^{\circ}\text{C}$ for 12 days and three replications were maintained for each treatment. The diameter of the
66 colony was measured in two directions and average was recorded. Per-cent inhibition mycelial
67 growth of the fungus was calculated by using the formula given by Vincent (1947).

Comment [113]: How can you melt the media?

Comment [114]: Add numerical value 20 instead of twenty.

Comment [115]: Media not medium

Comment [116]: Replace the suitable check with "Control treatment".

Comment [117]: Cite recent literature

$$I = \frac{C - T}{C} \times 100$$

70 Where,

71 I = Per-cent inhibition

72 C = Radial growth in control

73 T = Radial growth in treatment (fungicide)

75 RESULTS AND DISCUSSION

76 The assessment of fungicides through *In Vitro* testing proves to be a convenient method
77 for assessing a substantial array of chemicals, gauging their effectiveness in restraining pathogen
78 growth. This approach swiftly furnishes valuable initial insights into the fungicides' potency
79 against the pathogen, offering a concise timeframe for evaluation. These findings then act as a
80 compass for subsequent field trials. In this ongoing study, a total of six contact, six systemic, and
81 six combined fungicide products were examined against *Colletotrichum gloeosporioides*,
82 encompassing three distinct concentrations.

Comment [118]: Italicized In Vitro n text.

Comment [119]: Contact or non systemic?

83 Among six contact fungicides evaluated against *C. gloeosporioides* and Copper hydroxide
84 gave 69.90 per cent inhibition which was superior over all other fungicides evaluated. Which
85 was followed by Copper oxychloride (62.96 %), Chlorothalonil (53.87 %), Propineb (47.85 %),
86 Captan (40.83 %) and least inhibition was recorded with Mancozeb 40.30 per cent. (Figure 1).

Comment [120]: Use symbol of percentage throughout the text.

87 The results were in similarity with work of Parvathy and Girija (2016). Copper based
88 fungicides are effective because it kills the pathogen by denaturing proteins and enzymes in cells of
89 pathogens when they come in contact.

Comment [121]: Similar

90 Among six different systemic fungicides evaluated against *C. gloeosporioides*
91 Difenconazole, Tebuconazole were found best with inhibition percentage of 98.28 and 95.17.
92 These results were similar to the earlier reports made by Prashanth *et al.* (2008), Ahmed *et al.*
93 (2014), Parvathy and Girija (2016) that Difenconazole and Tebuconazole have highest inhibition
94 percentage on the growth of *C. gloeosporioides* (Figure 2).

95 The effectiveness of the Triazole fungicides may be attributed to their interference with the
96 biosynthesis of fungal sterols and inhibit the ergosterol biosynthesis. In many fungi, ergosterol is
97 essential for the structure of cell wall and its absence cause irreparable damage to cell wall leading
98 to death of fungal cell. A similar study was reported for the effectiveness of Triazoles, which inhibit
99 the sterol biosynthesis pathway in fungi (Nene and Thapliyal, 1993).

100 Out of the six evaluated combination products, propiconazole + difenconazole exhibited
101 the highest inhibition rate at 99.78%. Following closely, Fluopyram 200 g/L + Tebuconazole
102 200 g/L SC and Tebuconazole 50% EC + Trifloxystrobin 25% WG displayed inhibition rates
103 of 89.47% and 87.50% respectively. These findings aligned with the research by Prashanth *et*
104 *al.* (2008), Parvathy and Girija (2016), and Pavithra and Benagi (2017) (Figure 3).

Formatted: Font: (Default) Times New Roman, Italic

Formatted: Font: (Default) Times New Roman, Italic

105 The utilization of combination fungicides effectively curbs the development of fungal
106 resistance to systemic fungicides. This is because systemic fungicides disrupt only a single, or
107 occasionally two, functions within the fungal physiology, which can be easily overcome by a
108 singular mutation. On the contrary, non-systemic protectant fungicides impact numerous
109 aspects of fungal physiology, requiring the fungus to undergo multiple changes in order to
110 develop resistance. As a result, the combination of both systemic and non-systemic fungicides
111 yields superior outcomes.

112 CONCLUSION

113 *In-vitro* efficacy of non-systemic, systemic and combi fungicides done to know their
114 efficiency in suppressing the growth of *C. gloeosporioides* revealed the efficacy of copper
115 hydroxide which showed 69.90 per cent inhibition. In case of systemic fungicides maximum
116 per cent inhibition was recorded in Difenconazole (98.28 %) and lowest per cent inhibition

117 by Azoxystrobin (48.18 %). While among combi products Propiconazole 13.9 % EC +
118 Difenconazole 13.9 % EC showed 99.78 per cent followed by Fluopyram 200 g/L +
119 Tebuconazole 200 g/L SC (89.47 %), Tebuconazole 50 % EC + Trifloxystrobin 25 % WG
120 (87.50 %).

Comment [122]: Write one thing throughout the manuscript, symbol or spell of percentage.

Comment [123]:
Write same unit symbols (capital or small) throughout.

121 References

- 122 Ahmed, M. J., Hossain, K. S and Bashar, M. A., (2014). Anthracnose of betel vine and it's *in vitro*
123 management. Dhaka University Journal of Biological Sciences. 23(2): 127-133.
- 124 Chattapdayay, S. P. and Maity, S., (1967). Diseases of betelvine and species. ICAR, New Delhi.
- 125 Dasgupta, B. and Sen, C., (1999). Assessment of *Phytophthora* root rot of betelvine and its
126 management using chemicals. Journal of Mycology and Plant Pathology. 29: 91-95.
- 127 Hiralal Jana, Debabrata Basu and Suchhanda Jana., (2016). Pesticides use pattern of betelvine
128 growers in controlling diseases in Midnapur (east) district of West Bengal. International
129 Journal of Current Advanced Research. 6: 5314-5320.
- 130 Kumar, V., Suryanarayana, M. A. and T., Bindu, H. K., (2014). Status of Betelvine Cultivation in
131 Karnataka, Section of Medicinal Crops, Indian Institute of Horticultural Research,
132 Bangalore, 1- 4.
- 133 Nene. Y. L. and Thapliyal P. N., (1993). Evaluation of fungicides. In: Fungicides in plant disease
134 control (3rd ed.) Oxford, IBM Publishing Co., New Delhi
- 135 Parvathy, R. and Girija, V. K., (2016). In Vitro evaluation of fungicides and organic preparations
136 against *Colletotrichum gloeosporioides* causing anthracnose of black pepper (*Piper nigrum*
137 L.). International Journal of Applied and Pure Science and Agriculture. 2: 2394-2398.
- 138 Pavithra, S. and Benagi, V. I., (2017). In vitro evaluation of fungicides, botanicals and bio-agents
139 against *Colletotrichum gloeosporioides* (Penz.) Penz. and Sacc. the causal agent of
140 anthracnose of pomegranate. Environmental Ecology. 35(2): 671-675.

141 Prashanth, A., Arun, R. S., Naik, M. K., Patil, M. B. and Rajesh, S. P., (2008). Evaluation of
142 fungicides, bio agents and botanicals against pomegranate anthracnose. Indian Journal of
143 Plant Protection. 36(2): 283-287.

144 Vincent, J. M., (1947). Distribution of fungal hyphae in the presence of certain inhibitors. Nature.
145 159: 850.

146

147

148

149

150

151

152

153


154

155

156

157

Fig. 1: Symptoms of anthracnose of dragon fruit



158

159 Fig. 1: *In-vitro* evaluation of non-systemic fungicides against *Colletotrichum gloeosporioides*

160

161 **T1:** Copper oxychloride

162 **T2:** Copper hydroxide

163 **T3:** Propineb

164 **T4:** Chlorothalonil

165 **T5:** Captan

166 **T6:** Mancozeb


167 **C1:** 250 ppm

169 **C2:** 500 ppm

170 **C3:** 1000 ppm

171

172



173

174 Figure 2: *In-vitro* evaluation of systemic fungicides against *Colletotrichum gloeosporioides*

175 **T1:** Difenoconazole

176 **T2:** Tebuconazole

177 **T3:** Propiconazole

178 **T4:** Hexaconazole

179 **T5:** Picoxystrobin

180 **T6:** Azoxystrobin

181

182 **C1:** 250 ppm


183 **C2:** 150 ppm

184 **C3:** 100 ppm

185

186

187



189

190 Figure 3: *In-vitro* evaluation of combi products against *Colletotrichum gloeosporioides*191 **T1:** Propiconazole + Difenconazole192 **T2:** Fluopyram + Tebuconazole193 **T3:** Tebuconazole + Trifloxystrobin194 **T4:** Fluxapyroxad + Pyraclostrobin195 **T5:** Azoxystrobin + Tricyclazole196 **T6:** Fluopyram + Trifloxystrobin

197

198 **C1:** 500 ppm199 **C2:** 250 ppm200 **C3:** 150 ppm

201

202

203

204

205

UNDER PEER REVIEW

