

# Standardization, Physico-Chemical and Organoleptic Evaluation of Value added Ready-To-Serve (RTS) Beverage from Starfruit

## ABSTRACT

The present study was carried out to develop a ready-to-serve (RTS) beverage from starfruit value added with ginger, mint, aloe vera, lemon grass, basil and rosemary with the objective of assessing the physico-chemical and organoleptic properties. The experiment was laid out in randomized block design (RBD) with seven treatments and three replications. Based on the results of assessment, it has been concluded that treatment T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%) performed best in terms of physico-chemical properties viz. pH, acidity, TSS, vitamin C, reducing sugar and total sugar. On the basis of sensory evaluation, treatment T<sub>1</sub> (Starfruit juice + Ginger Juice 2%) was found most acceptable in terms of organoleptic properties viz. colour, taste, flavour and overall acceptability.

**Keywords:** Beverage, Organoleptic properties, Physico-chemical properties, Starfruit,

## INTRODUCTION

“The star fruit of carambola (*Averrhoa carambola* L.) is an attractive fruit of family *Oxalidaceae*, also known as ‘Golden Star’. The fruit, which is mostly consumed fresh or as juice, is rich in vitamins A and C along with iron and fiber. The ripe fruit may be processed into fermented or unfermented drinks, jam and jelly” (Kesavanathet *et al.*, 2015). The physical appearance of the star fruit is green, yellowish green, unripe, semi-ripe, ripe which has sour and fruity smell with perishable properties (Shirsat and Thakor, 2014). “The fruit contains huge variety of bioactive components such as antioxidants, minerals, total phenolics, and dietary fibers. It is also a good source of potassium, copper, as well as folate and panthothenic acid” (Manda *et al.*, 2012). “Yet, the lack of technical knowledge for its further processing has rendered its potentials unutilized which make it the most overlooked fruit for extracting essential therapeutic uses in the entire region” (Mishra, 2000). “Therefore, transportation, handling and, processing within a short span are major concerns in commercial scale processing” (Rayaguru, 2008). “Star fruits are considered to have a number of beneficial health effects. These include antioxidant, hypoglycemic, hypotensive, hypocholesterolemic, anti-inflammatory, anti-infective, antitumor, and immune-boosting effects” (Lakmalet *et al.*, 2021). “It is usually used to treat headaches,

vomiting, coughing, and hangover, as an appetite stimulant, diuretic, antidiarrheal and febrifugal agent” (Cabrini *et al.*, 2011). “Its hypoglycemic effects are considered to be particularly beneficial in patients with diabetes mellitus, and it has been promoted as a traditional remedy for diabetes”(Gunasekara *et al.*, 2011). “However, excessive consumption of star fruit has been associated with the development of oxalate nephropathy in patients with both normal and abnormal baseline renal function” (Neto *et al.*, 2003; Neto *et al.*, 2009)

“Ready to serve drink is popular among every age group because of their refreshing nature and taste. In recent years, there has been a growing interest in using herbal products as dietary adjuncts in the food industry. Functional beverages are one of its kinds which are tapping into consumer interest in health and wellness. People are now more health conscious and they want all good benefits in one drink. The demand for soft drinks always has an increasing trend and there is a great scope for development of value added beverages by utilizing nutritious food with medicinal properties. Therapeutic beverage sector has been reported to be the fastest growing segment in the soft drink industry sector” (Roberts, 2009). “Degenerative diseases such as diabetes, hypertension, atherosclerosis and obesity are the main causes for the reduction in the consumption of synthetic beverages among health conscious people which has led to produce the functional beverages”(Hariharan and Mahendran, 2016). “There is a growing demand for health drinks based on indigenous fruits. Ready-to-serve (RTS) is a type of beverage containing at least 10% fruit juice besides less than 0.3% acid according to the FPO Standard. It is normally not diluted before serving and hence known as RTS beverage” (Behera *et al.*, 2017).

“Ginger is an ancient medicinal as well as spicy plant belonging to Zingiberaceae family and botanically known as *Zingiber officinale* Rosc. Since a very long time ginger is known for its medicinal values as a digestive aid, spiritual beverage, aphrodisiac, antiemetic, anticancer, anti-oxidant, anti-inflammatory and immune stimulating properties” (Malhotra and Singh, 2003). “Mint has been reported to have pharmacological effect such as antimicrobial, anti-inflammatory, antispasmodic, antituesive, anticancer and analgesic” (Raghavan, 2006). “The functional properties and therapeutic benefits of Aloe vera are known worldwide. It is a source of active substances including vitamins, minerals, enzymes, sugar, anthraquinones of phenolic compounds, lignin, saponins, sterols, amino acids and salicylic acid” (Pugh *et al.*, 2001). “Lemongrass is well-known for its antioxidant, anti-microbial, anti-inflammatory, anti-hypertensive, anti-diabetic, anti-

mutagenicity, anxiolytic properties, and for its hypoglycemic and hypolipidemic activities” (Kiani *et al.*, 2022). “Basil has been found to show effectiveness against many fungal, viral, bacterial and protozoal infections. Basil juice is an effective medicine for inflamed eyes and night blindness, which is often caused by vitamin A deficiency” (Boggia *et al.*, 2015; Hosseini-Parvar *et al.*, 2015). “Rosemary, has been used in folk medicine to alleviate several diseases including headache, dysmenorrhea, stomachache, epilepsy, rheumatic pain, spasms, nervous agitation, improvement of memory, hysteria, depression, as well as physical and mental fatigue” (Duke, 2000; Heinrich *et al.*, 2006).

The value added beverages can be prepared from blends of different fruits & vegetables and extracts of herbs having medicinal, nutritional and therapeutic values with acceptable palatability. The development of starfruit RTS beverages blended with ginger, mint, aloe vera, lemon grass, basil, rosemary would provide the opportunities for best use of these perishable raw materials with less post-harvest loss and simultaneously availability of palatable drinks of medicinal values to the consumers. The consumers are becoming health conscious and more careful about their health and fitness subsequently demands for natural beverages with medicinal properties over synthetic one increasing in the market. Considering the aforementioned facts, a ready-to-serve beverage was developed from starfruit, value added with ginger, mint, aloe vera, lemon grass, basil, and rosemary. The objective was to evaluate the physico-chemical properties and sensory acceptability of the starfruit RTS beverage.

## MATERIALS AND METHODS

The experiment was conducted in the Post-harvest Technology Laboratory, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, India, during the year 2022-2024. The experiment was laid out in Randomized Block Design (RBD) with seven treatments and three replications. The treatments were T<sub>0</sub> (Control), T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), T<sub>2</sub> (Starfruit Juice + Mint Juice 2%), T<sub>3</sub> (Starfruit Juice + Aloe vera Juice 2%), T<sub>4</sub> (Starfruit Juice + Lemongrass Juice 2%), T<sub>5</sub> (Starfruit Juice + Basil Juice 2%), T<sub>6</sub> (Starfruit Juice + Rosemary Juice 2%).

### Formulation of Starfruit RTS Beverage

The six different types of starfruit RTS beverages, along with the control, were prepared in a quantity of 1 litre each. Each batch of 1 litre of starfruit RTS beverages was prepared

using 100ml of starfruit juice, 900ml water, 100g sugar, 0.3g citric acid, and 0.1g sodium benzoate. Additionally, for each formulation, 20ml of flavour juice including ginger, mint, aloe vera, lemongrass, basil and rosemary was added for value addition.

#### Methodology for Preparation of Starfruit RTS Beverage

The cleaned starfruits were chopped into small pieces and then processed using a juicer to extract the juice. In a separate container, the RTS beverage was prepared by combining the freshly extracted starfruit juice with water and sugar in the required proportions. Additionally, each sample of the RTS beverage was flavored by adding ginger, mint, aloe vera, lemon grass, basil, and rosemary juice, respectively. The mixture was stirred thoroughly to ensure proper blending of flavours. The prepared RTS beverage was strained through a double layer muslin cloth to obtain the clarified juice devoid of fibre particles.

#### Storage of Starfruit RTS Beverage

Starfruit RTS beverages were filled into sterilized glass bottles and plastic bottles and stored at ambient room temperature.

#### Evaluation of Physico-chemical Properties of Starfruit RTS Beverage

The starfruit RTS beverages were evaluated for various physico-chemical properties like pH, TSS, acidity, vitamin C, reducing sugar and total sugar. The pH content was analyzed by digital pH meter. The TSS content was analyzed by hand refractometer of 0-32 scale. The acidity content was analyzed by titration method. The Vitamin C content was analyzed by 2, 6-dichlorophenol-inndophenol visual titration method. The reducing sugar and total sugar content was analyzed by lane and eynon method. The recorded data of all samples for different parameters were tabulated and statistically analyzed to find out the most suitable treatment combination in terms of physico-chemical properties.

#### Evaluation of Organoleptic Properties of Starfruit RTS Beverage

The starfruit RTS beverages were tested for sensory evaluation by a panel of five judges to determine colour, taste, flavour, and overall acceptability. The samples were evaluated on a 9-point scale using hedonic rating method. Each sample was evaluated and given a score by the panelists on the scale of 1-9 for each parameter, for each sample. The mean scores of all samples of all the five members were tabulated and statistically analyzed to find out which treatment combination is most acceptable in terms of sensory acceptability.

## RESULTS AND DISCUSSION

### Physico-chemical Properties of Starfruit RTS Beverage

The nutritional value of starfruit RTS beverage was evaluated by analyzing its physico-chemical properties like pH, TSS, acidity, vitamin C, reducing sugar and total sugar. The data recorded on physico-chemical properties of starfruit RTS beverage have been presented in Table 1.

#### Effect of different treatments on pH of Starfruit RTS Beverage

Statistical analysis revealed that pH content differed significantly across all treatments. The mean values of pH ranged from 4.23 to 5.14. The maximum pH content 5.14 was recorded in T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), while the minimum pH content 4.23 was recorded in T<sub>6</sub> (Starfruit Juice + Rosemary Juice 2%). Similar results were reported by Sasikumar *et al.* (2015) in Ready-to-Serve (RTS) beverage prepared from blend of *aloe vera* and aonla; Behera *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from residual osmotic solution of starfruit; Hamid *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from mulberry; Thakur *et al.* (2018) in Ready-to-Serve (RTS) beverage prepared from wild aonla. Beverages with lower pH levels exhibited greater stability over time, with reduced rates of spoilage and degradation of sensory attributes. The variations in pH impacted the perception of sweetness, sourness, and overall taste intensity of the beverage.

#### Effect of different treatments on Total Soluble Solids of Starfruit RTS Beverage

Statistical analysis revealed that TSS content differed significantly across all treatments. The mean values of TSS content ranged from 9.42 to 10.12°Brix. The maximum total soluble solids content 10.12°Brix was recorded in T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), while the minimum total soluble solids content 9.42°Brix was recorded in T<sub>5</sub> (Starfruit Juice + Basil Juice 2%). Similar results were reported by Sasikumar *et al.* (2015) in Ready-to-Serve (RTS) prepared from blend of *aloe vera* and aonla; Kharsyntiew *et al.* (2019) in juice prepared from carambola; Das *et al.* (2021) in Ready-to-Serve (RTS) beverage prepared from kinnow and aonla; Deepa and Karetha (2022) in Ready-to-Serve (RTS) beverage prepared from dragon fruit and lime juice. TSS influenced the shelf stability of the RTS beverage. Higher sugar content acted as a preservative by reducing water activity, thereby inhibiting microbial growth and prolonging the beverage shelf life.

#### Effect of different treatments on Acidity of Starfruit RTS Beverage

Statistical analysis revealed that acidity content differed significantly across all treatments. The mean values of acidity content ranged from 0.11 to 0.38%. The maximum acidity content 0.38% was recorded in T<sub>6</sub> (Starfruit Juice + Rosemary Juice 2%), while minimum acidity content 0.11% was recorded in T<sub>0</sub> (Control). Similar results were reported by Harshitha *et*

*al.* (2016) in Ready-to-Serve (RTS) beverage prepared from mango; Sharma *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from mango; Hamid *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from mulberry; Thakur *et al.* (2018) in Ready-to-Serve (RTS) beverage prepared from wild aonla; Bochare *et al.* (2020) in Ready-to-Serve (RTS) beverage prepared from kiwi incorporated with lemongrass. Das *et al.* (2021) in Ready-to-Serve (RTS) beverage prepared from kinnow and aonla; Shagiwal and Deen (2022) in Ready-to-Serve (RTS) beverage prepared from strawberry, ginger and aloe vera blend; Salaria and Reddy (2022) in Ready-to-Serve (RTS) beverage (RTS) prepared from muskmelon. The moderate acidity levels in beverage enhanced flavour perception by adding complexity and brightness to the taste profile. Higher acidity levels helped extend the shelf life of the RTS beverage by acting as a natural preservative.

#### Effect of different treatments on Vitamin C of Starfruit RTS Beverage

Statistical analysis revealed that vitamin C content differed significantly across all treatments. The mean values of vitamin C content ranged from 21.34 to 24.76mg/100ml. The maximum vitamin C content 24.76mg/100ml was recorded in T<sub>1</sub>(Starfruit Juice + Ginger Juice 2%), while the minimum vitamin C content 21.34mg/100ml was recorded in T<sub>0</sub> (Control). Similar results were reported by Behera *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from residual osmotic solution of starfruit; Goyal and Kumar (2017) in Ready-to-Serve (RTS) beverage prepared from apple, guava, carrot and banana; Das *et al.* (2021) in Ready-to-Serve (RTS) beverage prepared from kinnow and aonla. The presence of vitamin C in the RTS beverage offered added nutritional benefits, potentially appealing to health-conscious consumers.

#### Effect of different treatments on Reducing sugar of Starfruit RTS Beverage

Statistical analysis revealed that reducing sugar content differed significantly across all treatments. The mean values of reducing sugar content ranged from 6.90 to 7.31%. The maximum reducing sugar 7.31% was recorded in T<sub>1</sub>(Starfruit Juice + Ginger Juice 2%), while the minimum reducing sugar 6.90% was recorded in T<sub>0</sub> (Control). Similar results were reported by Malav *et al.* (2014) in Ready-to-Serve (RTS) beverage prepared from orange; Hamid *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from mulberry; Bochare *et al.* (2020) in Ready-to-Serve (RTS) beverage prepared from kiwi incorporated with lemongrass; Das *et al.* (2021) in Ready-to-Serve (RTS) beverage prepared from kinnow and aonla. Reducing sugar content correlated positively with perceived sweetness of the beverage. Higher levels of reducing sugars contributed to a thicker consistency and

smoother mouthfeel.

#### Effect of different treatments on Total sugar of Starfruit RTS Beverage

Statistical analysis revealed that total sugar content differed significantly across all treatments.

The mean values of total sugar content ranged from 9.56 to 11.15%. The maximum total sugar 11.15 % was recorded in T<sub>1</sub>(Starfruit Juice + Ginger Juice 2%), while the minimum total sugar 9.56 % was recorded in T<sub>0</sub> (Control). Similar results were reported by Malav *et al.* (2014) in Ready-to-Serve (RTS) beverage prepared from orange; Hamid *et al.* (2017) in Ready-to-Serve (RTS) beverage prepared from mulberry; Kharsyntiew *et al.* (2019) in juice prepared from carambola; Shagiwal and Deen (2022) in Ready-to-Serve (RTS) beverage prepared from strawberry, ginger and aloe vera. The total sugar content played a key role in enhancing the flavour perception and overall sensory appeal of beverage. Total sugar content affected the texture and mouthfeel of the RTS beverage.

#### Organoleptic Properties of Starfruit RTS Beverage

The sensory acceptability of starfruit RTS beverage was evaluated by analyzing its organoleptic properties like colour, taste, flavour and overall acceptability. The data recorded on organoleptic properties of starfruit RTS beverage have been presented in Table 2.

#### Effect of different treatments on Colour of Starfruit RTS Beverage

Statistical analysis revealed that organoleptic score for colour differed significantly across all treatments. The mean organoleptic score for colour ranged from 6.0 to 8.3. The maximum organoleptic score for colour 8.3 was recorded in T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), while the minimum organoleptic score for colour 6.0 was recorded in T<sub>0</sub> (Control).

#### Effect of different treatments on Taste of Starfruit RTS Beverage

Statistical analysis revealed that organoleptic score for taste differed significantly across all treatments. The mean organoleptic score for taste ranged from 5.3 to 8.7. The maximum organoleptic score for taste 8.7 was recorded in T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), while the minimum organoleptic score for taste 5.3 was recorded in T<sub>0</sub> (Control).

#### Effect of different treatments on Flavour of Starfruit RTS Beverage

Statistical analysis revealed that organoleptic score for flavour differed significantly across all treatments. The mean organoleptic score for flavour ranged from 6.0 to 8.3. The maximum organoleptic score for flavour 8.3 was recorded in T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%), while the minimum organoleptic score for flavour 6.0 was recorded in T<sub>0</sub> (Control).

#### Effect of different treatments on Overall acceptability of Starfruit RTS Beverage

Statistical analysis revealed that organoleptic score for overall acceptability differed significantly across all treatments. The mean organoleptic score for overall acceptability ranged from 5.7 to 8.7. The maximum organoleptic score for overall acceptability 8.7 was recorded in T1 (Starfruit Juice + Ginger Juice 2%), while the minimum organoleptic score for overall acceptability 5.7 was recorded in T0 (Control).

Table 1: Effect of different treatments on Physico-chemical properties of Starfruit RTS Beverage

Treatment	pH	TSS (°Brix)	Acidity (%)	Vitamin C (mg/100ml)	Reducing Sugar (%)	Total Sugar (%)
T <sub>0</sub> (Control)	4.53	9.70	0.11	21.34	6.90	9.56
T <sub>1</sub> (Ginger)	5.14	10.12	0.21	24.76	7.31	11.15
T <sub>2</sub> (Mint)	4.62	9.88	0.17	21.20	7.16	10.68
T <sub>3</sub> (Aloe vera)	4.45	10.09	0.29	22.01	7.06	10.04
T <sub>4</sub> (Lemon grass)	4.97	9.78	0.34	23.83	6.86	10.16
T <sub>5</sub> (Basil)	4.71	9.42	0.30	22.57	7.04	10.42
T <sub>6</sub> (Rosemary)	4.23	9.59	0.38	23.37	6.73	10.03
F-test	S	S	S	S	S	S
S.Ed (±)	0.236	0.194	0.062	0.872	0.139	0.341
CV	6.199	2.420	29.598	4.698	2.434	4.060
CD at 5%	0.514	0.422	0.134	1.899	0.304	0.743

Table 2: Effect of different treatments on Organoleptic properties of Starfruit RTS Beverage

Treatment	Colour	Taste	Flavour	Overall acceptability
T <sub>0</sub> (Control)	6.0	5.3	6.0	5.7
T <sub>1</sub> (Ginger)	8.3	8.7	8.3	8.7
T <sub>2</sub> (Mint)	7.7	7.7	8.0	7.7
T <sub>3</sub> (Aloe vera)	6.3	6.3	6.3	6.7
T <sub>4</sub> (Lemon grass)	7.3	7.3	7.3	7.3
T <sub>5</sub> (Basil)	7.0	6.7	6.7	7.0
T <sub>6</sub> (Rosemary)	6.7	7.7	7.3	7.7
F-test	S	S	S	S
S.Ed (±)	0.509	0.642	0.613	0.675
CV	8.849	11.089	10.509	11.414
CD at 5%	1.109	1.400	1.336	1.470

## CONCLUSION

Based on the results of assessment, it has been concluded that T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%) performed best in terms of physico-chemical properties viz pH (5.14), acidity (0.21%), TSS (10.12°Brix), vitamin C (24.76mg/100ml), reducing Sugar (7.31%) and total Sugar (11.15%). On the basis of sensory evaluation, T<sub>1</sub> (Starfruit Juice + Ginger Juice 2%) was found most acceptable with regard to organoleptic properties viz colour (8.3), taste (8.7), flavour (8.3) and overall acceptability (8.7).

The development of starfruit RTS beverages enriched with a variety of herbs presents a promising avenue for value added beverage production. By blending starfruit with extracts of medicinal herbs, these beverages not only offer enhanced palatability but also provide significant health benefits to consumers. Furthermore, this approach facilitates the efficient utilization of perishable raw materials, thereby reducing post-harvest losses.

## REFERENCES

- Behera, G., Panigrahi, S. S. and Rayaguru, K. (2017).** Effect of storage conditions on quality of ready-to-serve beverage from residual osmotic solution of star fruit. *Advances in Food Science and Engineering*, 1(4): 144-151.
- Bochare, S. S., Kshirsagar, R. B. and Bogha, T. T. (2020).** Studies on development of kiwi fruit RTS beverage incorporated with lemongrass. *Journal of Pharmacognosy and*

- Phytochemistry*, 9(4S): 291-296.
- Boggia, R., Zunin, P., Hysenaj, V., Bottino A. and Comite, A. (2015).** Dehydration of basil leaves and impact of processing composition. In: Preedy, V. (ed.) *Processing and Impact on Active Components in Food*. Academic Press, San Diego, California, 645-653.
- Cabrini, D. A., Moresco, H. H., Imazu, P., Silva, C. D. D., Pietrovski, E. F., Mendes, D. A. G. B. and Otuki, M. F. (2011).** Analysis of the potential topical anti-inflammatory activity of *Averrhoa carambola* L. in mice. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Das, A., Bahadur, V., Prasad, V. M. and Mishra, S. (2021).** Standardization, physico-chemical & organoleptic assessment during storage of ready to serve blended kinnow-aonla beverage. *The Pharma Innovation Journal*, 10(12): 700-706.
- Deepa, K., and Karetha, K. M. (2022).** Studies on quality of dragon fruit and lime juice blended RTS beverage. *The Pharma Innovation Journal*, SP-11(8): 750-753.
- Duke, J. A. (2000).** *Handbook of Medicinal Herbs*. Florida, CRC Press, 630-632.
- Goyal, S. K. and Kumar, A. (2017).** Preparation of Ready-To-Serve (RTS) Beverage from multi-fruit juice and storage study at room and refrigeration condition. *Journal of Pharmacognosy and Phytochemistry*, 6(6S): 461-464.
- Gunasekara, L. C., Fernando, P. H. and Sivakanesan, R. (2011).** A preliminary study on the hypoglycaemic effect of *Averrhoa carambola* (Star fruit) in rats. *Sri Lanka: Proceedings of the Peradeniya University Research Sessions*, 16: 83.
- Hamid, T. N., Kumar, P. and Thakur, A. (2017).** Studies on preparation and preservation of ready-to-serve (RTS) beverage from underutilized mulberry (*Morus alba* L.) fruits and its quality evaluation during storage. *International Journal of Current Microbiology and Applied Sciences*, 6(9): 1067-1079.
- Hariharan, G. and Mahendran, T. (2016).** Physico-chemical, sensory and microbial evaluation of ginger-lime ready-to-serve (RTS) functional beverage, sweetened by Palmyra sugar candy. *Imperial Journal of Interdisciplinary Research*, 2(5): 1545-52.
- Harshitha, T., Gehlot, R., Siddiqui, S. and Rekha (2016).** Development and evaluation of mango ready-to-serve drink and squash. *Progressive Research – An International Journal Society for Scientific Development*, 11(3): 349-354.
- Heinrich, M., Kufer, J., Leonti, M. and Pardo-de-Santayana, M. (2006).** Ethnobotany and ethnopharmacology—Interdisciplinary links with the historical sciences. *Journal of ethnopharmacology*, 107(2): 157-160.

- Hosseini-Parvar, S. H., Matia-Merino L. and Golding, M. (2015).** Effect of basil seed gum (BSG) on textural, rheological and microstructural properties of model processed cheese. *Food Hydrocolloids*, 44: 557–567.
- Kesavanath, J., Premakumar, K. and Inthujaa, Y. (2015).** Nutritional and sensory characteristics of star fruit and sweet orange juices blend fruit cordial. *International Journal of Multidisciplinary Studies*, 2(2).
- Kharsyntiew, C. A., Broadway, A. A., Bharti, B. K., Hossain, S. A. and Lal, N. K (2019).** Studies on the preparation of carambola juice. *Journal of Pharmacognosy and Phytochemistry*, 8(6): 1664-1669.
- Kiani, H. S., Ali, A., Zahra, S., Hassan, Z. U., Kubra, K. T., Azam, M. and Zahid, H. F. (2022).** Phytochemical Composition and Pharmacological Potential of Lemongrass (Cymbopogon) and Impact on Gut Microbiota. *Applied Chemistry*, 2: 229-246.
- Lakmal, K., Yasawardene, P., Jayarajah, U. and Seneviratne, S. L. (2021).** Nutritional and medicinal properties of Star fruit (*Averrhoa carambola*): A review. *Food Science & Nutrition*, 9(3): 1810-1823.
- Malav, M., Gupta, R. and Nagar, T. (2014).** Studies on bio-chemical composition of orange based blended ready-to-serve (RTS) beverages. *Bioscience Biotechnology Research Communications*, 7(1): 78-83.
- Malhotra, S. and Singh, A. P. (2003).** Medicinal properties of ginger (*Zingiber officinale*), *Natural Product Radiance*, 2(6): 296-301.
- Manda, H., Vyas, K., Pandya, A. and Singhal, G. (2012).** A complete review on: Averrhoa carambola. *World journal of pharmacy and pharmaceutical sciences*, 1(1): 17-33.
- Mishra, D. K. (2000).** Study on Stone apple (*Aegle marmelos* Correa) Candy, (M. Sc. Thesis submitted to Department of Horticulture, N.D.U.A.T., Faizabad).
- Neto, M. M., da Costa, J. A. C., Garcia- Cairasco, N., Netto, J. C., Nakagawa, B. and Dantas, M. (2003).** Intoxication by star fruit (*Averrhoa carambola*) in 32 uraemic patients: treatment and outcome. *Nephrology Dialysis Transplantation*, 18(1): 120-125.
- Neto, M. M., Silva, G. E., Costa, R. S., Vieira Neto, O. M., Garcia-Cairasco, N., Lopes, N. P. and Dantas, M. (2009).** Star fruit: simultaneous neurotoxic and nephrotoxic effects in people with previously normal renal function. *NDT plus*, 2(6): 485-488.
- Pugh, N., Ross, S. A., ElSohly M. A. and Pasco, D. S. (2001).** Characterization of Aloeride, a new high-molecular-weight polysaccharide from *Aloe vera* with potent

- immunostimulatory activity. *Journal of agricultural and food chemistry*, 49(2): 1030-1034.
- Raghavan, S. (2006).** *Handbook of spices, Seasonings and Flavorings*, Second edition, CPC Press, 133.
- Rayaguru, K., Khan, M. K. and Sahoo, N. R. (2008).** Effect of storage on quality of stone apple ready-to-serve beverage. *Journal of Agricultural Engineering*, 45(1): 62-68.
- Roberts, W. (2009).** Benefiting beverages. *Prepared Foods*, 56: 32-35.
- Salaria, M. and Reddy, A. H. (2022).** Studies on standardization and quality evaluation of RTS beverage from muskmelon (*Cucumis melo* L.) variety Sarda. *The Pharma Innovation Journal*, 11(3): 1952-1956.
- Sasikumar, R. (2015).** Preparation of therapeutic RTS beverage from aloe vera gel and aonla fruit juice and evaluation of storage stability. *Asian Journal of Dairy and Food Research*, 34(2): 151-155.
- Shagiwal, M. and Deen, B. (2022).** Studies on development of ready-to-serve (RTS) beverage from strawberry (*Fragaria ananassa* Duch), ginger (*Zingiber officinale* Rosc) and aloe vera (*Aloe barbadensis* Miller) blend. *The Pharma Innovation Journal*, 11(7): 2308-2317.
- Sharma, D. S., Kaul, R. K., Sood, M. and Gupta, N. (2017).** Studies on stability and quality of jamun-mango blended ready-to-serve beverage. *Indian journal of Horticulture*, 74(02): 299-302.
- Shirsat, B. S. and Thakor, N. J (2014).** Studies on physico-chemical characteristics of Carambola (*Averrhoa carambola* L.) fruit. *Bioved Journal*, 25(2): 121-125.
- Thakur, N. S., Thakur, N., Kumar, P. and Thakur, A. (2018).** Formulation and optimization of vitamin C rich ready-to-serve juice based beverage from wild aonla fruits and its quality evaluation during storage. *Journal of Pharmacognosy and Phytochemistry*, 7(1): 1796-1802.