

Effect of integrated nutrient management on growth of onion (*Allium cepa* L.) cv. Pusa Shobha

ABSTRACT

Aim: The objective of the research was to study the effect of integrated nutrient management on growth of onion (*Allium cepa* L.) cv. Pusa Shobha

Study Design: The experiment was laid out in Randomized Block Design (RBD) with three replications.

Place and Duration of Study: The experiment conducted during *Rabi* season in the years, of 2021-22 and 2022-23 both the year at Horticulture Research Farm, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya Vihar Raebareli Road Lucknow, (U.P).

Methodology: The experiment conducted during *Rabi* season in the years, of 2021-22 and 2022-23 both the year at Horticulture Research Farm, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya Vihar Raebareli Road Lucknow, (U.P). The experiment was laid out in randomized block design with three replications. The treatments consisted of T₀ Control (without fertilizers), T₁- 100% RDF (NPK@150:50:80 Kg/ha): T₂- 100% RDF + FYM (10 t/ha); T₃ -100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₄ - 100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₅ -100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₆ -75% RDF + FYM (10 t/ha); T₇ -75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₈ -75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₉ -75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha). T₁₀ -50% RDF + FYM (10 t/ha); T₁₁- 50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₁₂- 50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₁₃ -50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha).

Results: Revealed that the treatment T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) performed better with respect to growth characters such as maximum values for plant height at 30, 60 & 90 DAT, number of leaves per plant at 30, 60 & 90 DAT, length of leaf at 30, 60 & 90 DAT, fresh and dry weight of onion bulb.

Conclusion: Integrated approach of Vermicompost, FYM and biofertilizer performed better with respect to growth parameters *viz.* plant height, number of leaves, leaf length, fresh and dry weight of onion bulb.

Keywords: INM, RDF, Growth, Biofertilizer, Vermicompost and Farmyard manure.

1. INTRODUCTION

Onion is one of the most important vegetable crops grown in India, having both the food and medicinal values. The onion is a major vegetable and spice crop raised all over the world in temperate, subtropical, and tropical climates. Onion bulb is rich in minerals, especially calcium, potassium and phosphorus besides having fairly good quantities of carbohydrates, proteins and vitamin C and minerals like phosphorus (39 mg), calcium (27 mg), sodium (1.0 mg), iron (0.7 mg) and potassium (1.57 mg) per 100 g of material (Aykroyd, 1963). Although the nutritive value of onion is low, it is greatly valued for its inevitable and extensive usage as a vegetable and medicine as well in the world. Integrated nutrient management (INM) provides excellent opportunities to overcome all the imbalances besides sustaining soil health and enhancing crop production. Generally, vegetables require large quantity of major nutrients like nitrogen, phosphorus and potassium, in addition to secondary nutrients such as zinc, boron, copper, calcium and sulphur for better growth, yield and post-harvest life. Promotion of the use of inorganic fertilizers for supplying these nutrients in the previous years has now become a bone of contention for its detrimental effect on both soil and environment apart from its enormous price hike every year. The use of FYM, vermicompost and biofertilizers in such situation is, therefore, a practically paying proposal. The steady depletion of native soil fertility and the occurrence of multiple nutrient deficiencies in onion fields have led to the identification of nutrient management as a key factor limiting sustainable onion production. All of these issues need field experimentation with alternative options. A gradual shift from using purely organic sources to some proportion of inorganic fertilization is gaining acceptance. This shift has formed the basis for INM, which could involve three nutrient sources: microbial inoculants or biofertilizers including *Azotobacter*, *Azospirillum*, and phosphate solubilising bacteria (PSB); inorganic fertilizers, and organic manures. However, INM further prescribes that selected nutrient inputs be used judiciously to ensure optimum supply of all essential nutrients for sustainable crop production. Onion is a heavy feeder of mineral elements. A crop of 40 t/ha removes approximately 120 kg of N, 50 kg of P₂O₅ and 60 kg of K₂O per ha (Tandon and Tiwari, 2008). Hence, the greater its ability to utilize nutrients for crop production, the greater is the yield potential. Keeping this in the view, the present investigation was undertaken to study the Effect of integrated nutrient management on growth of onion (*Allium cepa* L.) cv. Pusa Shobha

2. MATERIALS AND METHODS

The experiment conducted during *Rabi* season in the years, of 2021-22 and 2022-23 both the year at Horticulture Research Farm, Babasaheb Bhimrao Ambedkar University (A Central University), Vidya Vihar Raebareli Road Lucknow, (U.P). The experiment was laid out in randomized block design with three replications. The treatments consisted of T₀ Control

(without fertilizers), T₁- 100% RDF (NPK@150:50:80 Kg/ha); T₂- 100% RDF + FYM (10 t/ha); T₃ -100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₄ -100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₅ -100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₆ -75% RDF + FYM (10 t/ha); T₇ -75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₈ -75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₉ -75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha). T₁₀ -50% RDF + FYM (10 t/ha); T₁₁- 50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha); T₁₂- 50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha); T₁₃ -50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) were studied in RBD keeping three replications in each treatment plants were transplanted in a plot size 1.8mx1.2m and at the spacing of 15x10cm. Recommended cultural practices was followed to rise a healthy crop observation was recommended on five randomly selected plant for five characters namely as plant height (cm), number of leaf per plant leaf length freshwater bulb (g) dry weight of bulb (g). Data of both the years was statistically analyses as per standard method suggested by Panse and Sukhatme (1985).

3. RESULTS AND DISCUSSION

3.1 Effect of integrated nutrient management on plant height (cm) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Effect of different treatments on plant height at 30,60 & 90 DAT are given in Table-1. As evident from the data, significant difference among the treatments was recorded at 30 DAT (days after transplanting) in both the years. During 2021-22, the maximum plant height at 30 DAT (32.30cm) was recorded with the treatment T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum plant height (16.70 cm) was recorded in case of control T₀. During 2022-23, the plant height was recorded maximum (33.43cm) in case of application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum plant height (15.41cm) was recorded with control T₀.

Effect of different treatments on plant height (cm) at 60 DAT after transplanting is given in Table-1. As evident from the data significant difference among the treatments was recorded during both the years. During 2021-22 the maximum plant height (49.76 cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5

Kg/ha) + PSB (5 Kg/ha) followed by T₁₃. The minimum plant height (32.33 cm) was recorded in case of control T₀. During 2022-23, the maximum plant height (50.76cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by T₁₃ and T₈. The minimum plant height (31.77cm) was recorded in case of control T₀.

Effect of different treatments on plant height (cm) at 90 DAT after transplanting is given in Table-1. As evident from the data significant difference among the treatments was recorded during both the years. During 2021-22, the plant height at 90 DAT after transplanting was maximum (62.34cm) with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with the T₁₃. While the minimum plant height (41.31cm) was recorded in case of control T₀. During 2022-23, the maximum plant height (63.98cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum plant height (42.37cm) was recorded in case of control T₀. The findings of this investigation were in close conformity with those of Kumar *et al.* (2017), Badal *et al.* (2019), Chhabra and Wankhade and Kale (2019), Upadhyay *et al.* (2023).

Table.1. Effect of integrated nutrient management on plant height (cm) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Treatment Details		Plant height (cm)					
		At 30 DAT		At 60 DAT		At 90 DAT	
		2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₀	Control (without fertilizers)	16.70	15.41	32.23	31.77	41.31	42.37
T ₁	100% RDF (NPK@150:50:80 Kg/ha)	22.31	21.82	39.97	40.97	50.37	51.39
T ₂	100% RDF + FYM (10 t/ha)	24.38	24.72	41.21	42.28	52.11	53.18
T ₃	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	26.40	26.81	42.49	43.94	53.4	54.48
T ₄	100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	27.29	28.21	43.3	44.39	54.56	55.96
T ₅	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	28.38	29.37	44.49	45.49	55.76	56.98
T ₆	75% RDF + FYM (10 t/ha)	26.36	27.26	44.87	45.87	56.87	57.89
T ₇	75% RDF + FYM (5 t/ha) + Vermicompost (2	28.40	29.92	45.13	46.83	58.87	59.88

	t/ha)						
T₈	75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	30.68	31.73	47.65	48.69	59.87	60.97
T₉	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	32.30	33.43	49.76	50.76	62.34	63.98
T₁₀	50% RDF + FYM (10 t/ha)	25.48	26.83	43.11	44.14	55.31	56.86
T₁₁	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	27.28	28.25	44.5	45.57	56.37	57.86
T₁₂	50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	26.73	27.83	46.23	47.23	56.13	57.98
T₁₃	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	31.56	32.45	48.12	49.12	61.36	62.54
SE(m) ±		0.336	0.369	0.479	0.462	0.771	0.553
CD (P=0.05)		0.977	1.073	1.395	1.344	2.243	1.608

3.2 Effect of integrated nutrient management on number of leaf per plant of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Data regarding number of leaves per plant have been presented in Table- 2. Effects on number of leaves was significant at plant growth during both the years. During 2021-22, the maximum number of leaves per plant at 30 DAT (8.73) were recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by the T₁₃. The minimum number of leaves per plant (3.12) was recorded in case of control T₀. During 2022-23, the number of leaves per plant was maximum (8.83) in case of application of of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha). The minimum number of leaves per plant (4.15) was recorded in case of control T₀.

Data regarding number of leaves per plant have been presented in Table- 2. Effects on number of leaves was significant at plant growth during both the years. The treatment effects on number of leaves were significant at plant growth during both the years. At 60 DAT, the maximum number of leaves per plant during 2021-22 (10.03) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum number of leaves per plant (4.10) was recorded in case of control T₀. During 2022-23, the maximum number of leaves per plant

(10.13) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum number of leaves per plant (4.15) was recorded in case of control T₀.

As evident from the data, significant effects of different treatments were recorded during both the years of study. During 2021-22, the number of leaves per plant at 90 days after transplanting was the maximum (12.98) with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by the T₁₃, while the minimum number of leaves per plant (7.20) was recorded in case of control T₀. During 2022-23, the maximum number of leaves per plant at 90 days after transplanting (12.99) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by the T₁₃, The minimum number of leaves per plant (7.28) was recorded in case of control T₀. The findings of this investigation were in close conformity with those of Mandalet *al.* (2013), Vachan and Tripathi(2017), Chhabra and Vishwakarma (2019), Singh *et al.* (2019).

Table. 2. Effect of integrated nutrient management on number of leaf per plant of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Treatment Details		Number of leaf per plant					
		At 30 DAT		At 60 DAT		At 90 DAT	
		2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₀	Control (without fertilizers)	3.12	3.15	4.10	4.15	7.20	7.28
T ₁	100% RDF (NPK@150:50:80 Kg/ha)	4.35	4.88	6.31	6.42	9.42	9.51
T ₂	100% RDF + FYM (10 t/ha)	5.06	5.96	6.90	6.97	9.64	9.74
T ₃	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	5.63	5.86	7.30	7.83	10.21	10.31
T ₄	100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	5.80	5.91	7.49	7.59	10.49	10.59
T ₅	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	5.87	5.97	8.86	8.96	11.07	11.17
T ₆	75% RDF + FYM (10 t/ha)	6.81	6.91	8.90	8.99	10.60	10.16
T ₇	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	7.37	7.47	9.22	9.32	11.01	11.11

T₈	75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	7.89	7.99	9.32	9.42	11.92	11.93
T₉	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	8.73	8.83	10.03	10.13	12.98	12.99
T₁₀	50% RDF + FYM (10 t/ha)	6.2	6.32	8.30	8.13	9.85	9.96
T₁₁	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	7.1	7.62	8.95	8.97	10.43	10.53
T₁₂	50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	7.38	7.58	9.10	9.14	11.78	11.88
T₁₃	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	8.12	8.34	9.82	9.93	12.44	12.54
SE(m) ±		0.067	0.079	0.076	0.075	0.088	0.123
CD (P=0.05)		0.197	0.231	0.223	0.218	0.256	0.357

3.3 Effect of integrated nutrient management on length of leaf (cm) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

The data pertaining to length of the leaf has been presented in Table-3. As evident from the significant effects of different treatments on the length of leaf were recorded at 30 days after transplanting of the plant growth during both the years of study, during 2021-22, the maximum length of leaf at 30 DAT (27.10 cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by the T₁₃ which was at par with T₈. The minimum length of leaf (12.98 cm) was recorded in case of control T₀. During 2022-23, the length of leaf was maximum (28.31 cm) in case of application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃ and T₈. The minimum length of leaf (13.99 cm) was recorded in case of control T₀.

As evident from the data significant effects of different treatments on leaf length at 60 DAT, the maximum length of leaf during 2021-22 (47.90 cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which at par with T₈, T₁₂ and T₁₃. while the minimum length of leaf (30.23 cm) was recorded in case of control T₀. During 2022-23, the maximum length of leaf (48.91 cm) was recorded

with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which at par with T₇, T₈, T₁₂ and T₁₃, while the minimum length of leaf (29.31cm) was recorded in case of control T₀.

As evident from the data, significant effects of different treatments was seen on length of leaf at 90 days during both the years of study. During 2021-22, the length of leaf at 90 DAT was maximum (58.36cm) with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with the T₁₃ and T₈. The minimum length of leaf (38.11cm) was recorded in case of control T₀. During 2022-23, the maximum length of leaf (59.36cm) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃ and T₈. The minimum length of leaf (39.19cm) was recorded in case of control T₀. The findings of this investigation were in close conformity with those of Bhatiet *al.* (2018), Vachan and Tripathi(2017), Chhabra and Vishwakarma (2019), Dhakadet *al.* (2019).

Table. 3. Effect of integrated nutrient management on length of leaf (cm) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Treatment Details		Length of leaf (cm)					
		At 30 DAT		At 60 DAT		At 90 DAT	
		2021-22	2022-23	2021-22	2022-23	2021-22	2022-23
T ₀	Control (without fertilizers)	12.98	13.99	30.23	29.31	38.11	39.19
T ₁	100% RDF (NPK@150:50:80 Kg/ha)	20.29	19.86	37.65	38.21	48.76	49.76
T ₂	100% RDF + FYM (10 t/ha)	21.92	22.99	39.23	40.11	49.10	50.10
T ₃	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	22.31	23.42	40.22	41.22	48.99	50.70
T ₄	100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	22.59	25.12	41.20	43.55	50.18	51.18
T ₅	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	23.21	24.32	42.33	43.11	52.10	53.10
T ₆	75% RDF + FYM (10 t/ha)	23.92	24.99	42.78	44.09	52.00	53.11
T ₇	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	24.10	25.23	43.35	45.11	54.33	55.23

T₈	75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	25.43	26.53	45.93	46.99	56.77	57.67
T₉	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	27.10	28.31	47.90	48.91	58.36	59.36
T₁₀	50% RDF + FYM (10 t/ha)	22.10	23.23	42.10	42.31	51.8	52.91
T₁₁	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	23.45	24.56	40.98	41.78	53.89	54.92
T₁₂	50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	24.64	25.75	45.55	46.19	55.88	56.97
T₁₃	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	26.04	27.05	46.28	47.41	57.56	58.56
SE(m) ±		0.285	0.762	1.386	1.396	0.590	0.653
CD (P=0.05)		0.83	2.217	4.031	4.060	1.716	1.900

4.4 Effect of integrated nutrient management on fresh weight of bulb (g) & dry weight of bulb (g) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

The result of present study clearly indicate that data presented in Table-4 on the effect of integrated nutrient management on fresh and dry weight of onion bulb. As evident from the data, significant effects of different treatments was seen on Fresh weight of bulb during both the years of study. During 2021-22, the fresh weight of bulb was maximum (77.78g) with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with the T₁₃. The minimum fresh weight of bulb (54.52g) was recorded in case of control (T₀). During 2022-23, the maximum fresh weight of bulb (78.78g) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₇, T₈, T₁₁, T₁₂ and T₁₃. The minimum fresh weight of bulb (48.98) was recorded in case of control (T₀). Significant effects of different treatments was seen on dry weight of bulb during both the years of study. During 2021-22, the dry weight of bulb was maximum (10.55g) with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) followed by T₁₃ which was at par with the T₈. The minimum dry weight of bulb (4.77g) was recorded in case of control (T₀). During 2022-23, the maximum dry weight of bulb (10.04g) was recorded with application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) which was at par with T₁₃. The minimum dry weight of bulb (4.89) was recorded in case of control (T₀). A better

nutritional environment in the root zone and throughout the plant system may be the cause of the application of rising fertility levels. Since nitrogen serves as the building block for all living things, it is widely known that it is the mineral nutrient that plants need the most of all for growth and development. Due to its dependence on vital substances such as amino acids, protein, nucleic acids, enzymes, co-enzymes, and alkaloids, it also plays a significant part in plant metabolism. Like nitrogen, phosphorus is a nutrient that plants require in quite high concentrations for typical plant growth. Adenosine diphosphate (ADP) and adenosine triphosphate (ATP) are the major P-containing molecules that provide internal energy to plants. The findings of this investigation were in close conformity with those of Kumar *et al.* (2017), Sinha *et al.* (2017), Wankhade and Kale (2019), Kaur and Singh (2019), Chhabra and Vishwakarma (2019) and Gashaw *et al.* (2021), Upadhyay *et al.* (2023) in onion.

Table. 4. Effect of integrated nutrient management on fresh weight of bulb (g) & dry weight of bulb (g) of onion (*Allium cepa* L.) during the year of 2021-22 & 2022-23.

Treatment Details		Fresh weight of bulb (g)		Dry weight of bulb (g)	
		2021-22	2022-23	2021-22	2022-23
T ₀	Control (without fertilizers)	54.52	48.98	4.77	4.89
T ₁	100% RDF (NPK@150:50:80 Kg/ha)	63.88	64.87	5.42	5.55
T ₂	100% RDF + FYM (10 t/ha)	64.75	65.75	6.86	6.92
T ₃	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	65.34	66.34	7.33	7.81
T ₄	100% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	65.23	66.23	5.88	6.09
T ₅	100% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	67.87	68.87	6.22	6.18
T ₆	75% RDF + FYM (10 t/ha)	70.53	71.53	7.97	8.61
T ₇	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	73.76	74.76	8.11	8.01
T ₈	75% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	74.65	75.65	9.12	9.25
T ₉	75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	77.78	78.78	10.55	10.04

T₁₀	50% RDF + FYM (10 t/ha)	69.61	70.61	6.64	6.83
T₁₁	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha)	71.87	72.87	8.57	8.08
T₁₂	50% RDF + FYM (10 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	72.44	73.44	8.12	9.15
T₁₃	50% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha)	75.65	76.65	9.58	9.39
SE(m) ±		1.004	2.281	0.25	0.261
CD (P=0.05)		2.921	6.633	0.72	0.76

4. CONCLUSION

On the basis of results, it could be concluded that the application of T₉- 75% RDF + FYM (5 t/ha) + Vermicompost (2 t/ha) + Azotobacter (5 Kg/ha) + PSB (5 Kg/ha) was found to be the best treatment combination in terms of plant growth parameters of onion. Integrated approach of Vermicompost, FYM and biofertilizer performed better with respect to growth parameters viz. plant height, number of leaves, leaf length, fresh and dry weight of onion bulb.

REFERENCES

- Anil, K., R. B. Ram, M. Sutan, K. Sachin, Y. Rahul and K. R. Meena (2017). Effect of organic manures, biofertilizers and micronutrients on growth, yield and quality of onion (*Allium cepa* L.). *International Journal Agriculture, Science*, 13(2): 236-241
- Aykroyd, U.R. (1963) Indian Council of Medical Research, Special Report Vegetable, National Book Trust India, New Delhi, 42, 188-191.
- Badal, D. S., Dwivedi, A. K., Kumar, V., Singh, S., Prakash, A., Verma, S., & Kumar, J. (2019). Effect of organic manures and inorganic fertilizers on growth, yield and its attributing traits in garlic (*Allium sativum* L.). *Journal of Pharmacognosy and Phytochemistry*, 8(3), 587-590.
- Bagali, A.N., Patil, H.B., Chimmad, V.P., Patil, P.L. And Patil, R.V. (2012). Effect of inorganics and organics on growth and yield of onion (*Allium sativum* L.), *Karnataka Journal of Agricultural Sciences*, 25(1), 112-115.
- Bhati, Vidya, P. K. Yadav, & R. Kumar. (2018). Effect of levels of inorganic fertilizers, organic manure and bio-fertilizer on plant growth attributes of onion (*Allium cepa* L.) cv. N-53 under Hot Arid Region of Western Rajasthan, India. *International Journal Current Microbiology & Applied Science*, 7, 3593-601.
- Chavan, D. L., N. H. Chavan and S. M. Choudhary (2016). Effect of different organic manures on growth and yield of onion (*Allium cepa* L.). *The Bioscan*, 11(4), 2529-2532.

Chhabra, S. and G. Vishwakarma (2019). Effect of integrated nutrient management on growth, yield and quality of onion (*Allium cepa* L.) cv. Palam Lohit. *Journal PharmacognPhytochem.* SP4:73-77.

Dhakad, K. R., Chudasama, V.R., Verma, J., Jalpa, G., Dhaked, K.M., (2019). Effect of integrated nutrient management on growth and yield of onion (*Allium cepa* L.) cv. Palam Lohit under North Gujarat condition. *International Journal of Current Microbiology and Applied Sciences*, 8(4), 1618-1622.

Gashaw, B. (2021). Evaluation of different rates of NPS on growth and yield performances of garlic (*Allium sativum* L.) in Cheha district, Gurage Zone, Ethiopia. *International Journal of Agronomy*: 1-5.

Kaur, A. and S. Singh (2019). Role of various fertilizers and azotobacter (biofertilizer) on the performance of *kharif* onion (*Allium cepa* L.) cv. Agrifound Dark Red. *Journal Pharmacogn. Phytochem.* SP4: 146-151.

Mandal, J., Ghosh, C. and Chattopadhyay, G.N. (2013). Proportional substitution of chemical fertilizers with vermicompost on growth and production potential of onion (*Allium cepa* L.). *International Journal of Bio-resource and Stress Management*, 4(2), 356-357.

Panse VG, Sukhatme PV. (1985). Statistical methods for agriculture worker 4th Edition, ICAR, New Delhi.

Sahoo, B.B., Nayak, B.S., Mohanty, S. K., and Khanda, C. (2022). Significance of biofertilizer incorporation with customized organic manures to reduce inorganic nutrient on growth dynamics, bulb yield and economics of Onion (*Allium cepa* L.) under the western undulating zone of Odisha. *Vegetos*, 1-12.

Singh B.D., Dwivedi A.K., Kumar Vimal, Singh Sopal, Prakash Atul, Verma Somendra and Kumar Jeetendra. (2019). Effect of organic manures and inorganic fertilizers on growth, yield and its attributing traits in garlic (*Allium sativum* L.). *Journal Pharmacognosy Phytochemistry*, 8, 587-590.

Sinha, S. K., R. B. Verma, V. K. Singh, V. B. Patel, G. S. Panwar, D. K. Bharati and R. Kumar (2017). Integrated nutrient management approach for increasing growth, yield and economics of onion (*Allium cepa* L.). *Current Journal Applied Science and Technology* 1-6.

Tandon, H.L.S. and K.N. Tiwari. (2008). Nutrient management in horticultural crops. FDCO publishers, New Delhi.

Upadhyay, A., Singh R. B., Singh P., Lai B, and Singh S. (2023). Effect of integrated nutrient management on yield attributing parameters of *kharif* onion (*Allium cepa* L.). *International Journal of Environment and Climate Change*, 13(10), 4187-4193.

Vachan, Ram and S. M. Tripathi (2017). Study on the effect of bio-fertilizer with chemical fertilizer on plant growth, yield and economics of *rabi* season onion (*Allium cepa* L.) cv. NHRDF Red 2." *Journal Pharmacogn. Phytochem.* 6(5): 1496-1499.

Wankhade S. D. and V. S. Kale (2019). Effect of organic nutrient management on growth and yield of onion (*Allium cepa* L.) cv. Akola Safed. *JournalPharmacogn. Phytochem.* 8(1): 1918-1920.

UNDER PEER REVIEW