

## **Original Research Article**

# **Response of inorganic fertilizer and plant growth regulators on growth and seed yield of radish (*Raphanus sativus* L.)**

### **ABSTRACT**

This study aimed to evaluate the influence of different levels of inorganic fertilizer and plant growth regulator on seed yield of radish (*Raphanus sativus* L.) was formulated and conducted during the rabi season of 2022-23 at vegetable seed production plot, Department of Horticulture (Vegetable & Floriculture), BAC, Sabour Bhagalpur. The experiment was conducted to determine seed production ability at four levels of fertilizers viz, (F<sub>0</sub>-control, F<sub>1</sub>-N40:P30:K40, F<sub>2</sub>-N60:P40:K50, & F<sub>3</sub>-N8P50:K50) as well as seven levels of PGRs viz, P<sub>0</sub> (Control), P<sub>1</sub> (GA<sub>3</sub> @ 50ppm), P<sub>2</sub> (GA<sub>3</sub>@100 ppm), P<sub>3</sub>(GA<sub>3</sub>@150ppm), P<sub>4</sub>(NAA@50ppm), P<sub>5</sub>(NAA@100ppm) and P<sub>6</sub>(NAA@150ppm). Inorganic fertilizers were applied at 20 days and PGRs as foliar spray at 30 and 60 days after seedlings transplanting. The experiment was conducted in RBD (Factorial) and replicated thrice. The combined effect of inorganic fertilizers and Plant growth regulators F<sub>2</sub>P<sub>3</sub> significantly gave maximum diameter of pod (7.64 mm), length of pod (56.61mm), number of seeds/pod (6.18), 1000- seedweight (15.12g) and seed yield (18.23q/ha) and plant height is non significant. The treatment combination F<sub>3</sub>P<sub>6</sub> gave the maximum number of branches (8.06). Hence, F<sub>2</sub>P<sub>3</sub> treatment combination is a best combination for seed yield of radish and enhancing seed yield is our ultimate goal. As we performed the study in a particular location, we recommend multilocation trials in different agro-climatic regions to study and final confirmation of the results.

*Keyword: radish, seed, PGRs, inorganic fertilizer, seedling, transplanting*

### **1. INTRODUCTION**

Among the root vegetables, radish (*Raphanus sativus* L.) is an ancient as well as popular vegetable of tropical and temperate regions and can be cultivated extensively in almost all seasons of the year. Kashi Shweta is a popular cultivated variety suitable for tropical conditions. It is considered to be the native of China and India. It is grown for its pungent taste and flavour and considered as an appetizer. The characteristic pungent flavour of radish is due to the volatile isothiocyanates (trans-4- methyl- thibutenyl - isothiocyanate). The tender leaves are also cooked and eaten as vegetable. It is a nutritious vegetable providing a good source of carbohydrates, protein and vitamins A, B & C (Bakhsh *et al.* 2006). The seed yield of radish depends on variety, cultivation methods, climatic conditions, soil, fertilizer, PGRs as well as edaphic factors etc. The seed quality of radish is also an important factor for higher yield of the crop and seed production of this crop is also

considered a crucial problem for radish improvement and development. Amongst these, good quality seed is the most important and critical input. In most of the vegetable crops, the seed is an insignificant item in the total cost of production but the efficiency of other costlier inputs like fertilizer, irrigation, herbicides, insecticides, fungicides, harvesting and processing revolves around the use of good quality seed. The use of poor quality of seed is the major factor for low production of vegetable crops. Among the major essential nutrients required by the plants for their normal growth and yield. Further it has been observed that farmers usually use urea only as chemical fertilizer without addition of phosphatic and potassic fertilizer, for sustainable cropping system balanced use of nutrients is essential (Mishra & Das, 2014). The role of nitrogen is acceptable as it is a necessary component of protein, nucleic acids, chlorophyll and certain important enzymes (Pervez et al. 2004). The excessive use of nitrogen negatively affects the quality as well yield of crops (Chen et al. 2004). Keeping this point in view, the present investigation was carried out during the rabi season 2022- 23 to study the effect of PGRs and levels of fertilizer on the seed yield of radish cv. Kashi Shweta at Bihar Agricultural University, Sabour, Bhagalpur.

## 2. MATERIAL AND METHODS

Field investigations were undertaken at vegetable seed production plot, department of Horticulture (Vegetable & Floriculture), BAC, Sabour Bhagalpur during the rabi season 2022-23. The variety Kashi Shweta was used for this study. Agronomic practices are followed as per the package of practices. The experiment consisted of four levels of fertilizers *viz*, (F<sub>0</sub>-control, F<sub>1</sub>-N40:P30:K40, F<sub>2</sub>-N60:P40:K50, & F<sub>3</sub>-N8P50:K50) as well as seven levels of PGRs *viz*, P<sub>0</sub> (Control), P<sub>1</sub> (GA<sub>3</sub> @ 50ppm), P<sub>2</sub> (GA<sub>3</sub>@100 ppm), P<sub>3</sub>(GA<sub>3</sub>@150ppm), P<sub>4</sub>(NAA@50ppm), P<sub>5</sub>(NAA@100ppm) and P<sub>6</sub>(NAA@150ppm). Total of 28 treatment combinations were tried in RBD (Factorial) and replicated thrice in a plot size of 2.50 x 1.80 m for each treatment. The treatments were required as per the plan. The observations on plant growth, seed yield and quality components were recorded for the net plot. After the harvest of the crop, recorded on seed yield components like diameter of pod (mm), length of pod (mm), number of seeds/pod, 1000- seed weight (g) and seed yield (q/ha) were recorded. Statistical analysis were performed by following the method of Panse and Sukhatme (1984).

## 3. RESULTS AND DISCUSSION

Plant height is closely related to growth and development of the plant. The maximum plant height (163.42cm) was recorded with the application of P<sub>3</sub> (GA<sub>3</sub> @150 ppm). The increase in plant height might be due to cell elongation and make the plant taller. Similar results were also found by Haokip *et al.* (2016) and Shruthi *et al.* (2016) in carrot. Among different levels of nutrients at maturity stage, the maximum plant height (177.85 cm) was recorded in F<sub>3</sub> (N80: P50: K60) which was found significantly superior to the rest of the nutrients levels. This could be due to sufficient supply of NPK combined with maximum photosynthetic activity, resulting in rapid vegetative growth, physiologically healthier and strong plant morphology. This result was also supported by the earlier findings of Malek *et al.*, (2010) in carrot and Panwar *et al.* (2001) in radish. Whereas interaction effect was found non significant over plant height.

Number of branches/plant was found significantly higher in plants sprayed with NAA as compared to GA<sub>3</sub> and maximum number of branches per plant (6.96) was recorded with the

foliar application of P<sub>6</sub> (NAA @ 150 ppm). The increase in the number of branches due to auxin NAA treatment may be attributed to the activation of cell division and cell elongation in the auxiliary bud, which had a promoting effect on number of branches. The result of the present investigation in terms of number of branches are similar with the result of Meena and Malhotra (2006), Kurmi *et al.* (2019) and Yugandhar *et al.* (2016) in coriander. The highest number of branches (6.81) was found at fertility level F<sub>3</sub> (N80: P50: K60). This might be due to an adequate supply of NPK associated with high photosynthetic activity leading to vigorous vegetative growth, physiologically high stout and healthy plant morphology. Similar results were reported by Haokip *et al.* in carrot. (2016), Sharma and Lal (1994) in radish. The interaction effect of different levels of nutrients and plant growth regulators on number of branches was also found significant. The maximum number of branches (8.06) was recorded in F<sub>3</sub>P<sub>6</sub> (N80: P50: K60+NAA @150 ppm). The number of branches increased due to proper supply of NPK as well as suppression of apical dominance due to spraying of NAA which diverts the polar transport of auxin towards the basal buds that leads to increase in the number of branches/plants.

The effect of different levels of fertilizers on diameter and length of pod was found significant. The maximum diameter (6.58mm) and length(45.24mm) of pod were recorded in F<sub>2</sub>(N60: P40: K50). The reason for increase in pod length and pod diameter is that with high NPK doses plants got more nutrition and other favourable conditions like more water, more air etc. for growth, hence more vigorous plants. These results are in agreement with the findings of Sandhu *et al.* (1965), Mehla *et al.* (1987) in radish. The use of GA<sub>3</sub> could bring significant increment in length and diameter of pods. The maximum length (49.30 mm) and diameter (6.42 mm) of pod were associated with foliar application of GA<sub>3</sub> (P<sub>3</sub>) which proved its superiority over other concentrations of GA<sub>3</sub>. The increase in length and pod diameter may be attributed to increase in the number of cells as well as elongation of cells which is characteristic action of GA<sub>3</sub>. These results are supported by Shruthi *et. al.* (2016) in radish and Alagukanna and Kumar (2003) in fenugreek.

Number of seeds per pod influenced significantly due to the different levels of fertilizer. Among the different levels of fertilizer, maximum number of seeds per pod (5.44) was found in F<sub>2</sub> (N60:P40:50). This could be due to an adequate supply of fertilizer to the root zone which increased the length and diameter of pods. Similar results were observed by Panwar *et. al* (2001). Among the different levels of PGRs, maximum number of seeds per pod (5.60) was recorded in P<sub>3</sub> (GA<sub>3</sub> @ 150 ppm). This might be due to spray of GA<sub>3</sub> results in improved photosynthesis, vegetative growth and mobilization of photosynthates towards reproductive site which finally increase the number of seed per plant. Similar results were observed by Haokip *et al.* (2016).

The different levels of fertilizer have a significant effect on 1000 seed weight. The maximum 1000 seed weight (13.32g). was observed F<sub>2</sub>(N60:P40:K50). The increase in 1000 seed weight due to the application of NPK may be due to the fact that NPK helps in protein synthesis and ultimately increases the plumpness of the grain and hence, the greater supply of NPK must have resulted in bolder seeds having greater 1000seed weight. Similar findings were also found by Rastogi *et al.* (1987),Shukla *et al.* (2012) and Shukla *et al.* (2013) in radish.Maximum 1000 seed weight (13.84g) was observed in P<sub>3</sub> (GA<sub>3</sub> @ 150 ppm). This might be due to enhanced physiological activities like photosynthesis, translocation of fertilizer and photosynthates (Saxena, 1989). Similar results were observed by Malek *et al.* (2011) in carrotand Alagukanna & Kumar (2003) in fenugreek. The interaction effect between nutrients and PGRs was also found significant. The maximum (15.12g) 1000 seed weight was recorded in F<sub>3</sub>P<sub>3</sub>(N80: P50 : K60+ GA<sub>3</sub> @ 150 ppm). This might be due to

enhanced physiological activities like photosynthesis, translocation of fertilizer and photosynthates (Saxena, 1989).

The maximum seed yield (14.01q/ha) was associated with foliar application of GA<sub>3</sub>@150 ppm (P<sub>3</sub>) which proved its superiority over other concentrations of GA<sub>3</sub> and NAA. Similar results were recorded by Malek *et al.* (2011) in carrot, Panwar *et al.* (2001), Mohanta *et al.* (2015). The highest seed yield (q/ha) was recorded at F<sub>2</sub> (N60:P40:K50) and found significantly superior to the rest of the levels. The increase in seed yield (q/ha) may be attributed to increased number of branches per plant along with an increased number of pods, more number of seeds per pod, and maximum weight of pods per plant which favourably resulted in maximum seed yield (q/ha). Similar results were obtained by Shukla *et al.* (2012), Shukla *et al.* (2013).

Table 1: Effect of different levels of chemical fertilizer (F) and PGRs (P) on plant height

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	114.28	118.68	121.85	125.38	116.76	119.81	122.96	119.96
N40:P30:K40 Kg/ha (F <sub>1</sub> )	148.24	153.65	157.78	160.92	150.82	155.36	158.70	155.07
N60:P40:K50 Kg/ha (F <sub>2</sub> )	164.86	171.56	176.70	181.26	168.72	172.65	177.18	173.28
N80:P50:K60 Kg/ha (F <sub>3</sub> )	168.56	175.65	179.48	186.12	173.21	179.08	182.86	177.85
Mean	148.99	154.89	158.95	163.42	152.38	156.73	160.43	
	F	P	FXP					
CD (P = .05)	5.05	6.69	NS					
CV %	5.22							

Table 2: Effect of different levels of chemical fertilizer (F) and PGRs (P) on number of branches/plant.

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	2.52	3.84	4.20	4.68	4.81	5.08	5.48	4.37
N40:P30:K40 Kg/ha (F <sub>1</sub> )	4.22	4.61	5.04	5.62	5.78	6.10	6.58	5.42
N60:P40:K50 Kg/ha (F <sub>2</sub> )	4.93	5.38	5.88	6.56	6.74	7.45	7.68	6.37
N80:P50:K60 Kg/ha (F <sub>3</sub> )	5.49	5.54	6.62	7.52	6.98	7.45	8.06	6.81
Mean	4.29	4.84	5.44	6.09	6.08	6.52	6.95	
	F	P	FXP					
CD (P = .05)	0.18	0.24	0.49					
CV %	5.21							

Table 3: Effect of different levels of chemical fertilizer (F) and PGRs (P) on diameter of pods (mm)

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	4.10	4.32	4.48	4.81	4.08	4.32	4.72	4.40
N40:P30:K40 Kg/ha (F <sub>1</sub> )	4.80	5.75	5.98	6.12	5.65	5.78	5.96	5.72
N60:P40:K50 Kg/ha (F <sub>2</sub> )	5.82	6.28	6.75	7.64	5.96	6.46	7.12	6.58
N80:P50:K60 Kg/ha (F <sub>3</sub> )	5.56	6.12	6.34	7.10	5.88	6.12	4.40	5.93
Mean	5.07	5.62	5.89	6.42	5.39	5.67	5.55	

	F	P	FXP	
CD ( $P = .05$ )	0.24	0.32	0.64	
CV %	7.00			

Table 4: Effect of different levels of chemical fertilizer (F) and PGRs (P) on length of pods (mm)

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	25.40	26.38	35.17	42.56	25.42	30.19	38.32	31.92
N40:P30:K40 Kg/ha (F <sub>1</sub> )	26.49	28.32	40.30	46.65	29.18	32.24	41.36	34.93
N60:P40:K50 Kg/ha (F <sub>2</sub> )	35.26	42.31	46.55	56.61	40.78	44.50	50.64	45.24
N80:P50:K60 Kg/ha (F <sub>3</sub> )	32.30	38.30	44.22	51.36	37.35	41.37	48.45	41.91
Mean	29.86	33.83	41.56	49.30	33.18	37.08	44.69	
	F	P	FXP					
CD ( $P = .05$ )	1.08	1.43	2.87					
CV %	4.55							

Table 5: Effect of different levels of chemical fertilizer (F) and PGRs (P) on number of seeds /pod

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	3.49	3.96	4.02	4.58	3.86	4.08	4.48	4.07
N40:P30:K40 Kg/ha (F <sub>1</sub> )	4.09	4.56	5.12	5.58	4.18	5.21	5.12	4.84
N60:P40:K50 Kg/ha (F <sub>2</sub> )	4.70	5.01	6.02	6.18	5.07	5.32	5.78	5.44
N80:P50:K60 Kg/ha (F <sub>3</sub> )	4.45	5.06	5.58	6.07	4.75	5.16	5.60	5.24
Mean	4.19	4.65	5.19	5.60	4.46	4.94	5.25	
	F	P	FXP					
CD ( $P = .05$ )	0.14	0.18	0.37					
CV %	4.72							

Table 6: Effect of different levels of chemical fertilizer (F) and PGRs (P) on 1000-seed weight

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	10.46	10.84	11.15	11.73	10.75	11.04	11.50	11.07
N40:P30:K40 Kg/ha (F <sub>1</sub> )	10.92	12.05	12.75	13.41	11.70	12.14	13.02	12.28
N60:P40:K50 Kg/ha (F <sub>2</sub> )	11.60	12.49	13.94	15.12	12.38	13.17	14.56	13.32
N80:P50:K60 Kg/ha (F <sub>3</sub> )	11.32	12.35	13.28	15.09	11.78	12.88	13.92	12.95
Mean	11.08	11.93	12.78	13.84	11.65	12.31	13.25	
	F	P	FXP					
CD ( $P = .05$ )	0.34	0.45	0.90					
CV %	4.45							

Table 7: Effect of different levels of chemical fertilizer (F) and PGRs (P) on seed yield (q/ha)

Treatment	Control (P <sub>0</sub> )	GA <sub>3</sub> @50 ppm (P <sub>1</sub> )	GA <sub>3</sub> @100 ppm (P <sub>2</sub> )	GA <sub>3</sub> @150 ppm (P <sub>3</sub> )	NAA@50 ppm (P <sub>4</sub> )	NAA@100 ppm (P <sub>5</sub> )	NAA@150 ppm (P <sub>6</sub> )	Mean
Control	5.12	6.06	7.07	7.68	5.89	6.91	7.44	6.60
N40:P30:K40 Kg/ha (F <sub>1</sub> )	7.21	9.70	11.47	12.29	9.42	11.06	11.90	10.44
N60:P40:K50	10.48	14.62	17.20	18.23	14.13	16.59	17.85	15.59

Kg/ha (F <sub>2</sub> )								
N80:P50:K60 Kg/ha (F <sub>3</sub> )	9.92	14.07	16.63	17.82	13.66	16.07	17.25	15.06
Mean	8.18	11.11	13.09	14.01	10.78	12.66	13.61	
	F	P	FXP					
CD (P = .05)	0.59	0.78	1.57					
CV %	8.07							

#### 4. Conclusion

Therefore, it may be concluded that the combined effect of PGRs and chemical fertilizer gave significant effect on vegetative growth and yield attributing traits for seed production of radish. The application of inorganic fertilizer @ 60:40:50 kg NPK/ha along with foliar spray of GA3 @ 150ppm at 30 & 60 days after seedlings transplanting was found most outstanding for getting higher seed yield, net return with higher benefit-cost ratio and seed yield of radish.

#### REFERENCES

- Alagukannan G, Kumar V. Response of Fenugreek (*Trigonella foenum-graceum* L.) to plant growth substances. South Indian Horticulture. 2003;47 (1-6):367-399.
- Bakhsh KB, Ahmad Z, Hassan S. Estimating indicators of higher yield in radish cultivation. International Journal of Agriculture and Biology.2006;8(6): 783-787.
- Haokip C M, Sharangi AB, Debbarma KAK, Devi R, Karthik CS. Role of plant growth regulators on the growth and yield of coriander (*Coriandrum sativum* L.). Journal of Crop and Weed. 2016; 12(3):33-35.
- Chen BM, Wang ZH, Wang SXLIGX, Song HX, Wang XN (2004) Effects of nitrate supply on plant, nitrate accumulation, metabolic nitrate concentration and nitrate reductase activity in three leafy vegetables. Plant Sciences. 2004;167: 635-643.
- Kurmi AK, Naruka IS, Kushwah SS, Shaktawat RPS. Effect of plant growth regulators on growth, yield and quality of coriander (*Coriandrum sativum* L.). Environment and Ecology2020;38 (3B): 710-714.
- Malek MA, Rahim MA, Begum K. Effects of foliar application of growth regulators on quality seed yield in carrot. Journal of Agroforestry and Environment.2010;p135-138.
- Mehla CP, Lal S, Singh H, Mishra HP. Response to radish (*Raphanus sativus* L.) to varying levels of irrigation and nitrogen. Haryana Journal of Science. 1987;16(3): 287-293.
- Mishra P, Dash D. Rejuvenation of biofertilizer for sustainable agriculture and economic and development. Consilience: The Journal of Sustainable Development, 2014;11(1): 41-61.
- Mohanta HC, Hossain MM, Islam MS, Salam MA, Saha SR. Effect of plant growth regulators on seed yield of carrot. Annals Bangladesh Agriculture.2015;19:23-31.
- Mohanta HC, Hossain MM, Islam MS, Salam MA, Saha SR. Effect of plant growth regulators on seed yield of carrot. Annals Bangladesh Agriculture.2015;19:23-31.
- Panse VGand Sukhatme, PV. Statistical methods for agricultural workers. Third Edition, Indian Council of Agricultural Research, New Delhi.1984

- Panwar AS, Kashyap AS, Baweja HS, Mehta K. Influence of nitrogen and phosphorus on seed yield of radish (*Raphanus sativus* L.) under mid hill conditions of Himalayas. Indian Journal of Hill Farming.2001;14 (1):73-77.
- Pervez MA, Ayub CM, Saleem BA, Virk NA, Mahmood N. Effect of nitrogen levels and spacing on growth and yield of radish (*Raphanus sativus* L.). International Journal of Agriculture and Biology.2004; 06(3): 504- 506.
- Rastogi KB, Sharma PP, Korla BN. Effect of different levels of nitrogen and reductase activity in three leafy vegetables. Plant Sciences.1987;167: 635-643.
- Sandhu KS, Dhesi NS, Kang US. Pod and seed characters of turnip (*Brassica rapa*) as Influenced by N, P, K and Spacing Treatments. Indian Journal of Agronomy.1965;10: 279-82.
- Saxena OP. Role of plant growth regulators in plat productivity studies Proceedings of National Seminar on Strategies in Physiological Regulation of Plant Productivity. Bombay India.1989; 13- 17.
- Sharma SK, Lal G. Effect of nitrogen nutrition, intra-row spacing and steckling size on seed vigour in radish. Horticulture Journal. 1994;7(2): 121-124.
- Shruthi HT, Srinivasa V, Ibaad MH. Influence of spacing growth regulators on growth, flowering, seed yield and quality of radish (*Raphanus sativus* L.) cv. Pusa Chetki. Research in Environment and Life Sciences.2016; 9(3): 288-291
- Shukla YR, Kumar P, Kansal S, Singh M, Kumar S. Effect of integrated nutrient management practices on seed yield and quality of radish,(*Raphanus sativus* L.) cv. Chinese Pink. International Journal of Farm Sciences.2013;3(2):10-18
- Shukla YR, Mehta S Sharma R. Effect of integrated nutrient management on seed yield and quality of radish (*Raphanus sativus* L) cv. Chinese Pink. International Journal of Farm Sciences. 2012;2(1):47-53.
- Yugandhar V, Reddy PSS, Sivaram GT, Ramesh E (2017) Impact of pre-soaking and foliar application of plant growth regulators on growth and seed yield of coriander (*Coriandrum sativum* L.). Journal of Crop and Weed2017;13(1): 100-102.
- Yugandhar V, Reddy PSS, Sivaram GT, Ramesh E (2017) Impact of pre-soaking and foliar application of plant growth regulators on growth and seed yield of coriander (*Coriandrum sativum* L.). Journal of Crop and Weed.2017; 13(1): 100-102.