

Original Research Article

Analgesic, anti-inflammatory and antipyretic activities of ethanol extract from *Ritchieacapparoides* leaf in rodents

Comment [u1]: Methanol or Ethanol?
What did you use?
Ethanol is mentioned only in Title and Abstract Background. But all over document, Methanol is mentioned.

ABSTRACT

Background: In Southern Nigeria, the leaf of *Ritchieacapparoides* is used traditionally in the treatment of pains, fever, malaria and snake bites. Therefore, this study evaluated the analgesic, anti-inflammatory and antipyretic activities ~~from~~ the ethanol extract ~~off~~ from *Ritchieacapparoides* leaves in rodents.

Comment [u2]: Methanol or Ethanol?
What did you use?
Be consistent.

Methods: The Analgesic effect was carried out using acetic acid and tail immersion models in mice, while anti-inflammatory and antipyretic activities were examined using xylene, egg-albumen, brewer's yeast and dinitrophenol models in mice and rats at doses of 125 mg/kg, 250 mg/kg and 500 mg/kg of the methanol extract. Phytochemical screening and oral acute toxicity tests were also carried out on the leaf extract of the plant.

Results: The methanol leaf extract and the standard drug (Aspirin) significantly decreased the number of writhes caused by acetic acid at $P < 0.05$ and $P < 0.01$. There was significant increase in reaction time in standard and extract groups of the tested agent. The extract produced significant $p < 0.05$ and $p < 0.01$ dose related inhibition of oedema which was comparable to aspirin in egg-albumin induced paw oedema model. *Ritchieacapparoides* leaf extract also demonstrated significant $p < 0.05$ and $p < 0.01$ effect in xylene induced mouse ear oedema test compared to dexamethasone. Statistical significant reduction in rectal temperatures were observed in both Brewer's yeast and dinitrophenol induced pyrexia in rats. The methanol leaf extract contain alkaloids, saponins, tannins, flavonoids, terpenoids, steroids and cardiac glycosides. The oral acute toxicity tests was found to be greater than 5000 mg/kg.

Comment [u3]: Methanol or Ethanol?
What did you use?
Be consistent.

Conclusion: The results obtained showed potential analgesic, anti-inflammatory and antipyretic effects of the methanol leaf extract of *R.capparoides* at doses tested which support the claim for the traditional use of the plant in treatment of these disease conditions.

Comment [u4]: Methanol or Ethanol?
What did you use?
Be consistent.

Keywords: *Ritchieacapparoides*; Leaf extract; Analgesic; anti-inflammatory; antipyretic

1. INTRODUCTION

Herbs have proven to be of high importance for prevention and treatment of diseases, and of high value to immunological provocation against much pathologic conditions [1]. Before synthetic drugs were produced, man was completely dependent on medicinal plants for treatment of different diseases [1]. Extracted substances from variety of these plants are commonly and widely used by traditional healers in Nigeria [2]. Constituents in many plants products have exhibited biological and pharmacological activities, which includes antiplasmodial, antidiabetics² anti-inflammatory, antipyretic, antiviral and many other effects [3]. Therefore, searching for novel pain relievers with analgesic, anti-inflammatory and antipyretic activities is a welcome idea [4].

Ritchieacapparoides, Family (Capparidaceae) is an evergreen climber but when alone it is a self-supporting shrub with compound palmate leaves. The leaves can be collected all – year round as the plant can stand dry season. The roots are tuberous and with strong pungent odour when perceived. Like other capparidaceae, this herb is indigenous to the tropics, found mostly in the lowland area of rain forest, especially beside water body and virgin up -lands. As a shrub it grows to a height of few meter(s) and as climber can grow a considerable length of about 5 meters with several branches. The plant is virtually all over the tropical land of Africa and particularly West Africa [5]. In Igboland it is called “Aka-ato or Nti-ato” [5]. The leaf extract exhibited antinociceptive activities in mice [6]. Ethnomedicinally, the root extract of *R. capparoides* are used asanthelmintic [7]. In the South-Western part of Nigeria, decoction of the leaves and root are widely used for the treatment of microbial and plasmodial infections [8, 9]. Its antifungal activity has also been reported [10]. *R. capparoides* leaf preparations has been used inophthalmic condition, conjunctivitis, swellings, wounds and in the treatment of Guinea worm [11].

2. METHODS

2.1 Plant collection

Fresh leaves of *Ritchieacapparoides* were collected from the compound of MaziObasiAgwu in UgoniOkposi-Okwu in Ohaozara LGA of Ebonyi State. This plant was identified and authenticated by a Taxonomist, Dr. E.I. Aigbokhan of the Department of Botany, University of Benin, Ugwowo Campus, Benin City. The voucher number UBH-R443 was deposited in the Herbarium of the Department for future reference.

2.2 Extraction of the leaf

The collected leaves were air-dried for fourteen days at room temperature, and the size was reduced with mortar and pestle. The powdered plant leaf (592 g) was soaked in 1.8 L of methanol for 24 hours. The mixture was sieved using a clean muslin sieve into a conical flask. The filtrate was dried on a water bath at a reduced temperature of 40 °C. The extract yield was stored in a refrigerator before the experiment.

2.3 Experimental animals

Male and female Wistar rats and mice respectively weighing 150-180 g and 20-25 g, were sourced from house of the department of Veterinary Medicine, University of Nigeria, Nsukka. The animals were allowed free access to standard feed and water *ad libitum*. They were kept six each in clean separate cages with saw dust as bedding, which was replaced every two days. The study was conducted according to ethical guidelines on laboratory animal use and care in compliance with Faculty of Basic Clinical Sciences, Nnamdi Azikiwe University Research Policy.

2.4 Phytochemical screening

Phytochemical screening of the extract was carried out using simple chemical tests to detect the presence or absence of secondary constituents as detailed in the literature [12].

2.5 Acute toxicity test

Acute toxicity (LD₅₀) of the extract was determined using Lorke [13] method. This study was performed in two phases and the mice used were deprived of food overnight before extract administration. In phase 1, three groups of three mice per cage were orally administered the

extract in increasing doses of 10 mg/kg, 100 mg/kg and 1000 mg/kg respectively. The mice treated were monitored for the first 4 h and 24 h for signs of toxicity and mortality. With the absence of death after 24 hrs, phase 2 was introduced. Four groups of one mouse were each given the extract orally at doses of 1600 mg/kg, 2900 mg/kg, 5000 mg/kg and 10 mL/kg distilled water. The animals were then observed for signs of toxicity and mortality for 24 h, 48 h and 72 h respectively for late toxicity.

Comment [u5]: Not clear, please clarify

2.6 Analgesic activity

2.6.1 Acetic acid induced writhing in mice

The methanol leaf extract was assessed for analgesic activity using acetic acid-induced writhing method as described by Essien et al., [14]. Albino mice (20-25 g) of both sex randomized into 5 different cages of 6 mice in each cage. Group 1 (drug free) was given normal saline (10 mL/kg), the extract (125 mg/kg, 250 mg/kg and 500 mg/kg) were administered to groups 2-4, accordingly. The positive control received (150 mg/kg) of acetyl salicylic acid (ASA). After thirty minutes, intraperitoneal injection of acetic acid (10 ml/kg, of 0.7%) was administered to each mouse. They were separately placed in a transparent cage for observation. The writhing movements for each mouse was counted for 30 minutes.

2.6.2 Tail immersion test in mice

The method described by Akuodor et al, [15] was used for this study. Thirty randomly selected adult albino mice of both sexes were divided into 5 groups of 6 mice in each cage and were subjected to 24 h fast, but free access to water. Control groups 1 and 5 were treated orally with 10 mL/kg of distilled water and subcutaneously, with 10 mg/kg of morphine, respectively; While groups 2, 3 and 4 received orally, 125, 250 and 500 mg/kg of *R. capparoides* respectively. Thirty minutes after treatment, each mouse was placed in the restrainer cage (Grieve Cooperation, Illinois, U.S.A.), leaving the tail hung out and freely exposed to be dipped in a water bath that was maintained thermo-statistically at $51 \pm 1^\circ\text{C}$. The duration of stay (latency) of the tail in the hot-water bath before the animal withdrew its tail out of the water was recorded. The latency was evaluated at 30, 60, 90 and 120 min.

2.7 Anti-inflammatory activity

2.7.1 Egg-albumin-induced inflammation

Method described by Essien et al [16] was used for this study. Thirty adult Wistar rats of both sex were randomly selected and grouped into 5 with 6 rats per cage. Group 1 received orally, 10 mL/kg of distilled water, while group 2 received 150 mg/kg of aspirin orally. Groups 3, 4 and 5 received orally, 150 mg/kg, 250 mg/kg and 500 mg/kg of *R.capparoides* respectively. After 30 min treatment, the initial ~~volume (size)~~ of each rat's right hind paw was measured and recorded. Thereafter, inflammation was induced in rats by injecting subcutaneously, 0.2 mL of fresh egg-albumin into the sub-plantar of the right hind paw. Assessment of oedema volume using plethysmometer was thereafter measured at 30 min intervals for 120 min.

2.7.2 Xylene induced ear oedema method

This method as described by Akuodor et al., [17] was adopted for the study. Mice ~~recruited~~ were grouped into 5 with 6 animals in each. The animals were orally treated with the leaf extract in graded doses of 125 mg/kg, 250 mg/kg and 500 mg/kg. The negative control group was treated with 10 mL/kg of distilled water, while the positive control was treated with 4 mg/kg of dexamethasone. Oedema was induced in each mouse one hour after treatment with one drop xylene into the inner surface of the right ear. Three hour later, mice were sacrificed and both cut-off to equal size and weighed. The mean difference between the right and left ears were recorded as an indication of inflammation.

Comment [u6]: Find an appropriate term

2.8 Antipyretic activity

2.8.1 Brewer yeast induced pyrexia

The antipyretic activity was evaluated using Brewer yeast induced pyrexia in rats as described [18]. Hyperthermia was subcutaneously induced by administering 20 mL/kg of 20% aqueous suspension of Brewer yeast in distilled water 24 hours before treatment. Thirty Wistar rats of both sex were divided into five groups. Groups 1 and 2 served as negative control (distilled water 10 mL/kg), and positive control (ASA 150 mg/kg), while groups 3, 4, and 5 received 125 mg/kg, 250 mg/kg, and 500 mg/kg of the extract respectively. All drugs were administered orally. Rectal

Comment [u7]: Rephrase the sentence. Method is missing. Sentence is not clear.

temperatures were taken by the use of digital thermometer (Mediklin, China) before yeast injection, 24 h after the injection, and at 1, 2, 3, 4 and 5 hours after drug administration.

2.8.2 D-amphetamine induced pyrexia test

The antipyretic activities of the secondary metabolites of the ethanol leaf extract of *R.capparoides* was screened using D-amphetamine induced pyrexia method [19]. The animals used (wistar rats) of both sex were subjected to 24 hours fasting. The initial temperatures of the selected rats were recorded. The rats were grouped into 5 and 6 rats in each cage. They were each given 5mg/kg of D-amphetamine intraperitoneal in order to induce pyrexia. 24 hours after, the temperatures of the animals were noted for increase and any rat with temperature less than 0.6°C was avoided. Group 1 and 2 were negative (10 mL/kg distilled water) and positive control (ASA 150 mg/kg) respectively. Whereas, group 3, 4 and 4 received orally, 125 mg/kg, 250mg/kg and 500mg/kg of *R.capparoides*methanol leaf extract [respectively](#).

2.9 Statistical analysis

Results are presented as mean±standard error of mean (SEM) and analyzed with statistical package for social sciences (SPSS version 20) using one-way analysis of variance (ANOVA) followed by Dunnett's post hoc test. Difference in the mean $p<0.05$ was statistically considered significant.

3. RESULTS

3.1 Phytochemical analysis

The phytochemical screening of methanol leaf extract of *Ritchieacapparoides* revealed the presence of alkaloids, saponins, tannins, flavonoids, terpenoids, steroids, cardiac glycosides, resins and balsam.

3.2 Acute toxicity studies

There was no observed changes, mortality or signs of toxicity 72 hours after administration of the methanol leaf extract. The animals were all healthy and active throughout the study. Hence, the median lethal dose (LD50) was found to be greater than 5000 mg/kg.

3.2.3 Effect of methanol leaf extract of *R.capparoides* on acetic acid induced writhing in mice

The leaf extract of *R.capparoides* significantly and dose dependently decreased the number of acetic-acid induced writhing in mice at $p < 0.05$ and $p < 0.01$ respectively. The observed effects of the extract at 500 mg/kg was higher than that of 125 mg/kg and 250 mg/kg. This effect was comparable to that of the standard drug (Table 1).

3.2.4 Effect of methanol leaf extract of *R.capparoides* on Tail immersion test in mice

The methanol leaf extract of *R. capparoides* significantly at ($p < 0.05$ and $p < 0.01$) reduced the thermal stimuli in mice. In this test, dose dependent reduction was produced by the extract. However, morphine the standard drug showed stronger protection (Table 2).

3.2.5 Effect of methanol leaf extract of *R. capparoides* on egg-albumen induced paw oedema in rats

The anti-inflammatory activity of *R.capparoides* methanol leaf extract was observed to be dose dependent. There was significant activity at $p < 0.05$ with 125 mg/kg and 250 mg/kg doses of the extract, whereas the highest dose of the extract (500 mg/kg) had $p < 0.01$ significant activity. The reference drug (ASA) had more activity in this study (Table 3). The anti-inflammatory activity began at 1 hour after administration of drug and lasted for 5 hours before completely vanishing.

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

32.6 Effect of methanol leaf extract of *R.capparoides* on xylene induced ear oedema in mice

The anti-inflammatory effect of *R.capparoides* methanol leaf extract against xylene induced ear oedema in mice is shown in Table 4. The extract exhibited significant and dose dependent activity reduction of oedema at ($p < 0.01$) with the highest dose of the extract comparable to the standard drug, dexamethasone.

32.7 Effect of methanol leaf extract of *R. capparoides* on Brewer yeast induced pyrexia in rats

~~Antipyretic studies~~ Table 5 shows the antipyretic effect of the methanol leaf extract of *R. capparoides* determined using Brewer's yeast induced pyrexia in rats. The extract exerted significant and dose dependent antipyretic action at $p < 0.01$, whereas the standard drug showed significant antipyretic activity at $p < 0.01$.

32.8 Effect of the extract on Dinitrophenol – induced pyrexia in rats

The results of the effect of methanol leaf extract of *R.capparoides* against D-amphetamine induced pyrexia is shown in Table 6. There was a progressive dose dependent reduction at $p < 0.05$ in the temperature of rats treated with the methanol extract. The effect of the extract was less than the standard drug, acetylsalicylic acid (ASA)

Table 1: Effect of methanol leaf extract of *R. capparoides* on acetic acid-induced writhing in mice.

Treatment	Dose(mg/kg)	Mean no of writhes	% Inhibition
Distilled water	20 mL/kg	103.67±3.87	-
<i>R. capparoides</i> 125		30.83±2.71	70a
<i>R. capparoides</i> 250		13.33±1.82	87b
<i>R. capparoides</i> 500		5.83±3.80	94b
Aspirin	150	3.50±1.91	97b

Results are mean±SEM; (n=6); ^aP<0.05; ^bP<0.01 compared to control

Table 2: Effect of methanol leaf extract of *R. capparoides* on tail immersion in mice time interval

		Time (min).				
Pre-treatment	After-treatment					
Treatment	Dose mg/kg	0	30	60	90	120
Distilled water	20 mL/kg	10.17 ± 0.45	10.00 ± 0.58	10.67 ± 0.33	10.55 ± 0.22	10.50 ± 0.20
<i>R. capparoides</i>	125	6.50 ± 0.43	8.50 ± 0.50	10.83 ± 0.83	11.67 ± 0.71	13.17 ± 0.60 ^a
<i>R. capparoides</i>	250	7.50 ± 0.76	9.33 ± 0.76	11.83 ± 0.70	13.50 ± 0.56	16.80 ± 0.68 ^b
<i>R. capparoides</i>	500	8.00 ± 0.52	10.17 ± 0.79	12.50 ± 0.88	14.67 ± 0.56	17.50 ± 0.34 ^b
Morphine	10	7.50 ± 0.62	17.50 ± 0.72	19.17 ± 0.60	21.17 ± 0.87	23.33 ± 0.95 ^b

Results are mean ± SEM; (n=6); ^aP < 0.05; ^bP < 0.01 compared to control

Table 3: Effect of methanol leaf extract of *R. capparoides* on xylene-induced ear oedema in mice.

Treatment	Dose (mg/kg)	Weight of right ear	Weight of left ear	Increase in weight	% Inhibition
Distilled water	20mL/kg	0.042±0.01	0.021±0.03	0.021±0.00	-
<i>R. capparoides</i>	125	0.036±0.05	0.019±0.02	0.017±0.00	55 ^a
<i>R. capparoides</i>	250	0.030±0.04	0.017±0.03	0.017±0.00	60 ^b
<i>R. capparoides</i>	5000.	0.029±0.04	0.016±0.03	0.013±0.00	69 ^a
Dexamethasone	4	0.022±0.03	0.012±0.03	0.010±0.00	76 ^b

Results are mean±SEM; (n=6); ^aP<0.05; ^bP<0.01 compared to control

Table 4: The effect of ethanol leaf extract of *R. capparoides* on Egg-albumin-induced paws oedema in rats.

Time(min)	Drug Dose (mg/kg)	0	20	40	60	80	100	120
Distilled water	10mL/kg	1.24±0.02	1.64±0.01	1.72±0.02	1.79±0.02	1.86±0.02	1.93±0.02	2.05±0.02
<i>R. capparoides</i>	125	1.22±0.04	1.60±0.02	1.53±0.03	1.46±0.02	1.38±0.03	1.31±0.02	1.20±0.03a
<i>R. capparoides</i>	250	1.24±0.02	1.63±0.01	1.56±0.02	1.48±0.02	1.30±0.02	1.24±0.02	1.16±0.02b
<i>R. capparoides</i>	500	1.19±0.02	1.57±0.03	1.49±0.03	1.38±0.02	1.30±0.02	1.21±0.03	1.15±0.03b
Aspirin	150	1.17±0.04	1.61±0.02	1.49±0.02	1.39±0.03	1.30±0.03	1.22±0.03	1.12±0.03b

Results are mean ±SEM; (n=6) ^aP< 0.05; ^bP< 0.01 compared to control

Table 5: Effect of methanol leaf extract of *R. capparoides* against yeast induced pyrexia in rats.

Treatment	Dos(mg/kg)	Yeast induction (h)					Drug administration (h)								
		0	24	1	23	4	5	0	24	1	23	4	5		
D. water	20 ml/kg	35.37±0.05	37.52±0.04	37.80±0.02	37.63±0.02	37.42±0.02	37.29±0.02	37.30±0.03							
<i>R. capparoides</i>	150	35.26±0.03	37.25±0.03	36.41±0.01	36.20±0.02	35.62±0.03	35.37±0.03	35.24±0.02a							
<i>R. capparoides</i>	250	35.25±0.02	37.27±0.02	36.50±0.03	36.22±0.01	35.61±0.02	35.36±0.03	35.22±0.03a							
<i>R. capparoides</i>	500	35.23±0.02	37.30±0.02	36.47±0.02	36.15±0.01	35.55±0.03	35.30±0.02	35.10±0.03a							
Aspirin	150	35.22±0.00	36.79±0.02	35.69±0.04	35.43±0.02	35.40±0.01	34.60±0.02	34.30±0.03b							

Results are mean±SEM; (N=6) ^aP<0.05; ^bP<0.01 when compared to control.

Table 6: Effect of methanol leaf extract of *R. capparoides* on de-amphetamine induced pyrexia in rats (hours).

Treatment	Dose (mg/kg)	D-amphetamine induction (h)					Drug administration(h)				
		0	24	1	2	3	4	5			
D. water	20 ml	35.25±0.04	37.39±0.04	37.61±0.04	37.67±0.02	37.46±0.05	37.26±0.03	37.70±0.03			
<i>R. capparoides</i>	150	35.27±0.04	37.29±0.02	36.43±0.03	36.23±0.03	35.51±0.02	35.30±0.02	35.25±0.02	a		
<i>R. capparoides</i>	250	35.20±0.03	37.30±0.02	36.40±0.02	36.23±0.02	35.52±0.02	35.31±0.02	35.23±0.03	a		
<i>R. capparoides</i>	500	35.24±0.02	37.30±0.02	36.33±0.02	36.20±0.03	35.48±0.02	35.26±0.02	35.21±0.01	a		
Aspirin	150	35.25±0.03	37.31±0.02	36.39±0.03	36.22±0.03	35.49±0.01	35.31±0.01	34.17±0.01	b		

Results are mean±SEM; (N=6) ^aP< 0.05; ^bP<0.01 when compared to control

4. DISCUSSION

This study was carried out to establish the potential pharmacological properties of methanol extract of *R. capparoides* based on claims of its use in herbal medicine. The findings of the present study reveal that *R. capparoides* at doses employed exhibited analgesic effects against chemical pains (writhing) induced [20]. Another report revealed the response of mice to acetic acid as a fast and trusted method to test peripheral analgesic effect of herbal agents [14]. In the present work, the extracts of *R. capparoides* and acetylsalicylic acid (aspirin) inhibited acetic acid induced writhing. Results show the extract has peripheral analgesic properties which suggest the action may be directed via inhibition of local peritoneal receptors [21]. However, irrespective of whether this model evaluates peripheral pain relieving action only or otherwise, the results validate the usefulness of this plant as an analgesic in Nigeria. The injection of acetic acid is reported to induce the release of mediators of pain such as prostaglandins and other cyclokinase [22]. This activity suggests that the extract might have acted by inhibiting the actions of cyclooxygenase responsible for producing prostaglandins from arachidonic acids [22]. The analgesic effects produced by *R. capparoides* validate their use in traditional medical practice as analgesics being used by local population in Nigeria.

More so, centrally acting model of analgesia (tail immersion) was carried out to confirm the analgesic activity of this leaf extract. This method of assay used to indicate the involvement of central analgesic mechanism is believed to involve spinal reflex [23, 15]. Reported has it that centrally acting agents like morphine, possess this activity in both types of study, whereas peripherally acting agents such as acetylsalicylic acid has been reported to exert analgesic action only in the writhing test [24]. Most importantly, the action of acetylsalicylic acid in writhing

assay only could be linked to directly inhibit prostaglandin activity or indirectly inhibit prostaglandin secretion by halting cyclo-oxygenase activity [25]. The essential activity in tail immersion test shows involvement of central analgesic mechanism.

Egg-albumin induced paw oedema has proven to be a suitable way of testing anti-inflammatory agents and has widely been used for screening anti-oedematous action of natural products [26]. This process of testing acute inflammation potency is a highly sensitive tool [27]. The development of oedema which depend on the presence of bradykinin and polymorphonuclear leucocytes with proinflammatory factor like prostaglandins [28]. The leaf extract of *R.capparoides* may not have shown activity on the early phase of inflammation, hence may act by inhibiting the release of prostaglandins. Nonsteroidal anti-inflammatory agents like aspirin, may not inhibit the initial phase of edema induced by egg-albumen whereas the second accelerating phase can be antagonized by the drug [29].

Xylene causes irritation in mouse ear leading to accumulation of fluid and oedema and increase in myeloperoxidase enzymatic activity. Suppression of this response may suggest antiphlogistic activity [15]. The methanol extract of *R.capparoides* leaf exerted significant inhibition of ear oedema in mice. This activity suggests the inhibition of phospholipase A2 which has been involved in the pathophysiology of inflammation resulting from xylene [31]. However, dexamethasone used as the reference drug exhibited significant decrease in the ear weight of positive control rats which indicate an inhibition of PLA2.

Antipyretic agents have been shown to antagonize cyclooxygenase activity through increase in prostaglandin E2 by suppressing high temperature [31]. Increase in temperature may result from damaged tissue, infections, and other factors. This process usually give rise to mediators (interleukins and others), which progresses to prostaglandin E2 formation with increase body temperature [32]. The extract under study reduced rats' anal temperature whose effect was similar to that observed in aspirin. The methanol leaf extract of *R.capparoides* was able to bring fever to a control by getting rid of inflammatory symptoms at both peripheral and nervous system thermoregulator zones. This could bring down pyrogenic secreting cytokines while reducing prostaglandin E2 synthesis from cyclooxygenase possibly through the mechanism reputed for paracetamol [33].

The therapeutic potentials of medicinal plants are mostly attributed to the combination of their secondary metabolites. Flavonoids have been reported to target prostaglandins involved in late phase of acute inflammation and pain and they have therefore been associated with analgesic, anti-inflammatory and antipyretic activities [34, 35]. Therefore, it is not surprising to have seen these activities in *R.capparoides* leaf extract. There was no lethality observed during the LD₅₀ test which proves the relative safety of the herbal agent.

Formatted: Subscript

5. CONCLUSION


The results of present study show that the methanolic leaf extract of *R.capparoides* possess analgesic, anti-inflammatory and antipyretic activities, thus confirming the folklore uses of the plant for the treatment of these diseases conditions.

UNDER PEER REVIEW

Ethical approval

The experimental procedures were performed in accordance with Faculty of Basic Clinical Sciences, Nnamdi Azikiwe University, Nnewi Campus Ethics Committee and Guide for Care and Use of Laboratory Animals by The U.S. National Institute of Health (Publication number 85, revised 1996).

UNDER PEER REVIEW



Comment [u8]: Check again your references

REFERENCES

1. Singh A, Malhotra S, Subban R. Anti-inflammatory and analgesic agents from Indian medicinal plants. *Integr Biol* 2008; 3(1):57-72.
2. Akuodor GC, Anyalewechi NA, Udoh FV, Ikoru NC, Akpan JL, Gwotmut MD, Pharmacological evaluation of *Verbena hastata* leaf extract in the relief of fever. *Adv Pharmacol Toxicol* 2011; 12(3):1-8.
3. Hakiman M, Maziah M. Non enzymatic and enzymatic antioxidant activities in aqueous extract of different *Ficus deltoidea* accessions. *Med Plant Res* 2009; 3(3):120-131.
4. Ravelo Y, Molina V, Carbajal D, Fernández L, Fernández JC, Arruzazabala ML, Más R. Evaluation of anti-inflammatory and antinociceptive effects of D-002 (beeswax alcohols) *Nat Med* 2011; 65(2): 330-335.
5. Anowi CF, Utoh-Nedosa UA, Onyegbule AF, Oche G. Antimicrobial activity of methanolic extract of the leaves of *Ritchiea longipedicellata*. *Fam Capparidaceae Inter Pharma* 2012; 2(2):287-293
6. Sofidiya MO, Osuala EC, Onyemaechi CJ, Fageyinbo MS. Evaluation of antinociceptive activity of *Ritchiea longipedicellata* (Capparaceae) leaf extract in mice, *Herbs, Spices Medicinal Plants* 2018; 24:4, 356-367
7. Ajaiyeoba EO, Okogun JI. Anthelmintic activity of roots of *Ritchiea caparoides* var. *longipedicellata*. *Phytotherapy Research* 1996; 10: 436-437

8. Ogbunugafor HA, Okochi VI, Okpuzor J, Emeka P. Tolerance and antiplasmodial screening of

*Ritchealongipedicellata*in *Plasmodium berghei*. *Biochemistry* 2008; 20(1):23–27

9. Taiwo BJ, Akinkunmi EO, Omisore NO. Antimicrobial and Antiplasmodial activities of a Quaternary compound from *Ritcheacapparoides*var*Longipedicullata*. *African Journal of Traditional and Complementary Alternative Medicine* 2013; 10(b) 528-531.

10. Ajaiyeoba EO, Rahman AU, Choudhary IM. Preliminary antifungal and cytotoxicity studies of extracts of *Ritcheacapparoides* var. *longipedicellata*. *Ethnopharmacol* 1998; 62; 243–246.

11. Neuwinger HD. *African traditional medicine: a dictionary of plant use and applications*. Medpharm Scientific, Stuttgart, Germany 2000 589 pp.

12. Evans WC. *Trease and Evans Pharmacognosy*. 15th Edn., Reed Elsevier India Pvt. Ltd., New Delhi, India 2005; 174: 224-535.

13. Lorke D. A new approach to practical acute toxicity testing. *Arch. Toxicol* 1983; 54, 275–287.

14. Essien AD, Edidara Thomas, Essiet GA, Akuodor GC. Anti-inflammatory, antipyretic and anti-nociceptive activities of the ethanol stem bark extract of *Salaciaelembachii*. *British PharmacolToxicol*. 2017; 8(2):9- 16.

15. Akuodor GC, Essien AD, Udia PM, David-Oku E, Chilaka KC, Asika EC, Nwadam SK. Analgesic, Anti-inflammatory and Antipyretic potential of the stem Bark Extract of *Stachytarphetaindica*. *Br J PharmacolToxicol*. 2015;6(1):16-21.

16. Essien AD, Essiet GA, Akuodor GC, Akpan JL, Chilaka KC, Bassey AL. Pharmacological evaluation of the aqueous stem bark extract of *Bombaxbuonopozense* in the relief of pain and fever. *Afr J Pharm Pharmacol*. 2016;10(5):59-65.

17. Akuodor GC, Ohadoma SC, Ofor CC, Megwas AU, Chukwu LC, Ramalan MA, Okorafor DO,

Chilaka KC. Anti-nociceptive, anti-inflammatory and antipyretic activities of the

ethanol root bark extract of *Salacia hembachii* in rats and mice. *International journal of Basic and Clinical Pharmacology* 2021; 10(6):614-620

18. Ohanme EO, Etu KE, Akuodor GC, Nwakelu B, Ofor CC. Evaluation of Anti-Inflammatory, Anti-Nociceptive and Antipyretic Properties of the Ethanol Leaf Extract of *Celosia leptostachya* in Rats and Mice

19. Okokon JE, Nwafor PA. Anti-inflammatory, analgesic and antipyretic activities of ethanolic root extract of *croton zambesicus*. *Pak J Pharm Sci.* 2010;23(4):385-92.

20. Shaa KK, Oguiche S, Watila IM, Ikpa TF. In vitro antimalarial activity of the extracts of *Vernonia amygdalina* commonly used in traditional medicine in Nigeria. *Science World* 2011;6(2):5-9.

21. Mbiantcha M, Kamanyi A, Teponno RB, Taponjou AL, Watcho P, Nguiefack T. Analgesic and Anti-inflammatory Properties of Extracts from the Bulbils of *Dioscorea bulbifera* L. var *Sativa* (Dioscoreaceae) in Mice and Rats. *Evid Based Compl Altern Med.* 2011:912935.

22. Nkeh CNB, Bekwa PCM, Ndebia JE, Kayo M, Mbafor TJ, Iputo EJ. Analgesic and anti-inflammatory properties of *Oxyanthus unilocularis*. *J Med Plants.* 2010;4(10):932-9.

23. Yam MF, Loh YC, Oo CW, Basir R. Overview of Neurological Mechanism of Pain Profile Used for Animal "Pain-Like" Behavioral Study with Proposed Analgesic Pathways. *International Journal of Molecular Sciences.* 2020;21(12):4355.

24. Ezeja M, Ezeigbo I, Madubuike KG. Analgesic activity of the methanolic seed extract of *Buchholziacoriacea*. *Res J Pharm Biol Chem Sci.* 2011;2(1):187- 93.

25. Kumar A, Agarwal K, Kumar MA, Shanker K, Bushra U, Tandon S, et al. Pharmacological and phytochemical evaluation of *Ocimum sanctum* root extract for its anti-inflammatory, analgesic and antipyretic activities. *Pharmacognosy Mag.* 2015;11(42):217-24.

26. Abotsi WKM, Lamptey SB, Afrane S, Boakye-Gyasi E, Umoh RU, Woode E. An evaluation of the anti-inflammatory, antipyretic and analgesic effects of hydroethanol leaf extract of
27. Xu Z, Zhou J, Cai J, Zhu Z, Sun X, Jiang C. Antiinflammation effects of hydrogen saline in LPS activated macrophages and carrageenan induced paw oedema. *Inflammation*. 2012;9:2.
28. Abdulkhaleq LA, Assi MA, Abdullah R, Zamri-Saad M, Taufiq-Yap YH, Hezme MNM. The crucial roles of inflammatory mediators in inflammation: A review. *Veterinary world*. 2018;11(5):627-35.
29. David OE, Akuodor GC, Edet EE, Ogbuji GK, Obiajunwa OJI, Aja DOJ. Antinociceptive, antiinflammatory and antipyretic effects of ethanolic root bark extract of *Icacinasenegalensis* in rodents. *J ApplPharmaceut Sci*. 2016;(02):104-3.
30. Megwasl AU, Akuodor GC, Chukwu LC, Aja DO, Okorie EM, Ogbuagu EC. Analgesic, antiinflammatory and antipyretic activities of ethanol extract of *Annonasenegalensi* leaves in experimental animal models. *Int J Basic ClinPharmacol*. 2020;9(10):1477-84.
31. Ajayi AM, Tanayen JK, Magomere A, Ezeonwumelu JOC. Antinociceptive and anti-inflammatory effects of aqueous extract of *Chenopodiumopulifoliumschrad* leaves. *J Intercult Ethnopharmacol*. 2016;6(1):14-21.
32. Tarkang PA, Okalebo FA, Siminyu JD, Ngugi WN, Mwaura AM, Mugweru J, et al. Pharmacological evidence for the folk use of Nefang: antipyretic, anti-inflammatory and antinociceptive activities of its constituent plants. *BMC complementary and alternative medicine*. 2015;15:174-.
33. Sumanta M, Gouri KD, Suman A. Analgesic, antiinflammatoty and antipyretic studies of *Neolamarckiacadamba* barks. *J Pharma Res*. 2009;1:1133-6.

34. Agoreyo BO, Okoro NC, Choudhary MI. Preliminary phytochemical analysis of two varieties

of *Adenia lobata* and the antioxidant activity their various solvent fractions. *Bayero J Pure and Appl Sci.* 2012;5(1):182-6.

35. Savithamma M, Rao ML, Suhulatha D. Screening of medicinal Plants for secondary metabolites. *MiddleEast J Scient Res.* 2011;8(3):579-84.

UNDER PEER REVIEW