

Modulation of the standing behavior for managing elevated somatic cell count (SCC) in dairy cows

Abstract

Aim: To examine the effect of feed-off time on standing and lying behaviors to manage Somatic Cell Count (SCC) and reduce risk of subclinical mastitis (SCM) in tie stall dairy cows.

Study design: Completely Randomized Design (5 treatments \times 4 replications)

Place and duration of the study: National Cow Research Program, Chitwan, Nepal; from September, 2019 to March, 2020

Methodology: Twenty SCM-free dairy cows were recruited on different feeding strategies (pre-milking, during milking, and post-milking feeding at varying intervals). Behavioral patterns (lying time, bout frequencies and duration, post-milking standing duration/PMSD) were obtained using pedometer and SCC by autoanalyzer fortnightly. The feeding strategy effect on behavioral outcomes and SCC was analyzed for univariate ANOVA by SPSS packages (version 25). The relationship between behavioral patterns and SCC were performed by Pearson's Correlation analysis.

Results: Immediate post-milking feeding (+iPMF) cows showed higher lying time ($P=.003$) and bout duration ($P=.001$) while the number of lying bout was higher ($P=.001$) in pre-milking feeding (-10PMF). The PMSD was logged higher ($P=.001$) in +iPMF than in -10PMF and during-milking feeding cows but was similar with other post-milking feeding groups. Additionally, immediate and 15-minute post-milking feeding (+15PMF) resulted significantly lower SCC at both teat ($P=.001, .005$) and udder levels ($P=.001$) and are consistently associated with longer standing time after milking, suggesting decreased risk of SCM. Furthermore, post-milking standing duration showed significant but negative relationship with SCC of fore teats ($P=.001; r=-0.26$) and at udder level ($P=.001; r=-0.298$) while it was negative and non-significant with hind teats ($r=-0.09$ in LH and $r=-0.06$ in RH).

Conclusion: Post-milking feeding strategies, +iPMF and +15PMF, have the potential to improve dairy cow welfare and udder health by modulating behavioral patterns and reducing SCC. Further research is needed to optimize management practices for maximizing dairy cow welfare and udder health.

Keywords: Post-milking standing time, log₁₀ SCC, feeding strategy, udder health, lying bout

Introduction

The udder health is a prime focus of any dairy husbandry and describes the success of such operation instantly or in long run. Among the udder health issues, subclinical mastitis (SCM) is a common problem faced by dairy farmers and is fairly common at all levels of parity, although the incidence is much higher in multiparous than in primiparous animals [1,2]. Among the diagnostic tools, milk somatic cell count (SCC) is extensively used as a measure to track udder health and the milk quality [2,3] and up to 200,000 SCC/ml milk is the index figure for evaluating the udder health [4]. Leucocytes are the dominating cells during increase in SCC (>200,000/ml milk) and comprises of macrophages, lymphocytes and neutrophils [5]. The increase in SCC is attributed to the excess migration of neutrophils into the milk from bloodstream during inflammation and during this period, more than 90% of the cells present may consist of polymorphonuclear leukocytes. Therefore, a higher SCC indicates a greater risk of intra-mammary infection and potential contamination of milk [6].

That's why SCC is the widely adopted diagnostic biomarker for udder infection, and the specific test for diagnosing SCM [7]. The somatic cells in healthy cow milk consist of macrophages (66-88%) with neutrophils (1-11%), epithelial casts and mononuclear cells but in infected quarter, neutrophil level peaks and reaches up to 90% [8].

SCM is associated with standing and lying behavior of cow and the poor husbandry practices so understanding these behaviors and implementing effective farm management, to cut down odds of IMI is crucial [9,10]. It has long been understood that motivating dairy cows to stand longer after milking (PMSD) reduces the risk of IMI [11]. Research had indicated that providing fresh feed can keep cows on their feet for some period after milking and promotes longer standing time [11–13], which lets sufficient period for involution of teat orifice/canals before the udder contacts the stall surface, thus reducing the chance of bacterial penetration [12,14]. DeVries et al. [15] reported a decreased risk of environmental mastitis in tie-stall cows lying down between 40 to 60 min post-milking compared to those lying within 40 min. In tie-stall housing, cows typically stand for an average of 79 minutes after milking, much longer than in free-stall housing [12,13,16,17]. This change in behavior in free-stall cows may be due to longer time cows spent on pre-milking preparing as well as waiting in queue for milking turn [18]. Further, longer lying time has been found during the night and early morning hours; so shorter PMSD in free-stall housed and parlor milked cows may thus be a consequence of attempting to compensate for lost lying time [19]. In addition, milking three times a day may further disrupt animals' natural behavior patterns and encouraging cows to lie down quickly after milking in free-stall management. However, whether feeding time-based modulation of standing and lying events affects SCC or not in dairy cows is unexplored in Nepalese context. Dairying is facing a continuous challenge to maintain the quality of milk and the large part of this poor-quality milk is associated with the poor udder health, which might be linked to poor behavioral patterns of dairy cows. Therefore, this research was focused to explore the feeding strategy-based modulation of behavioral patterns of dairy cows for managing SCC to reduce the risk of SCM

Materials and methods

Animal selection and research design

A total of 20 SCM free HF and Jersey crossbred cows were randomly selected from the milking herd of National Cow Research Program, National Agriculture Research Council, Rampur, Chitwan, Nepal. Parity, lactation stage, initial SCC and body condition score of the cows were indifferent across treatments (Table 1). SCM free cows (with lower iSCC, 84- 109 × 1000 cells/ml) were selected after screening the milking herd at NCRP with Somatic Cell Kit (Intervet Schering Plough, Whitehouse, NJ). Somatic Cell is an on-farm rapid test and is conducted as detailed by [20]. The experiment was carried out during autumn and winter seasons for the period of 180 days in Completely Randomized Design with 5 treatments each with 4 replications.

-10PMF: Feed offered 10 min before milking

MFD: Feed offered during milking

+iPMF: Feed offered right after milking

+15PMF: Feed offered 15 min after milking

+30PMF: Feed offered 30 min after milking

The study cows, machine or hand-milked at 05.00- 07.00 morning and 16.00- 17.00 evening, were managed under conventional housing system in tail-to-tail fashion with access to free grazing for 3- 4 hours daily at day time and standard housing space was provided according

to guidelines. The study cows were managed on similar husbandry routine, feeding management and feeding materials (dry roughages, green fodders and concentrates). Commercial concentrates on the basis of production level and *ad libitum* green fodders were offered to address the nutritional requirements of the study cows.

Standing and lying behavior

Data Loggers (HOBO Pendant G Data Logger, Onset Computer Corporation, Pocasset, MA, USA) that records leg orientation in 1 min intervals were used for measuring the standing and lying behaviors of dairy cows. Data loggers were placed on the hind limb (1 animal/treatment at each fitting) with bandaging tape for the first 5 days following the milk sampling at a month interval. Five-day logging period is used to improve the accuracy of data, following Ito et al. [21] who demonstrated accurate estimate of lying behavior at herd with 3 days observation period.

Lying times (hr./day), frequency of lying bout (number/day) and lying bout duration (min/bout) were calculated as described by Ledgerwood et al. [22]. The data were downloaded using HOBO Software (Onset Computer Corporation) and worked-out on MS Excel for generating standing and lying patterns. The data logger recorded *g* force across the x, y, and z-axes, ranging from -3.2 to 3.2; a constant of 3.2 was added to all measurements, yielding positive *g* forces (0 to 6.4). Cut-off values were used to stratify logger readings into specific behaviors, including standing, lying, bout frequencies and duration. Data from the y-axis were specifically utilized to assess lying (≥ 2.55) and standing (< 2.55) behaviors, as well as bout characteristics. Finally, the PMSD, the time elapsed between the end of milking and the first instance of lying down thereafter, was worked out.

Table 1. Description of experimental crossbred dairy cow in tie-stall management at NCRP, Rampur, Chitwan

Treatments	# of animals	# of samplings	Mean parity	Mean lactation (mo.)	Mean iSCC ($\times 1000$ cells/ml)	BCS (1 to 5)
-10PMF	4	12	3.25	2.5	87	2.75
MFD	4	12	2.50	3.0	102	2.5
+iPMF	4	12	2.25	1.75	95	2.5
+15PMF	4	12	3.0	2.5	109	3.0
+30PMF	4	12	2.0	2.25	84	2.5

^a BCS evaluated on a 5-point scale (1- emaciated to 5- grossly over fat) [23]

-10PMF: 10 min Pre- milking feeding, MFD: milking during feeding, +iPMF: immediate PM feeding, +15PMF: 15 min PM feeding and +30PMF: 30 min PM feeding, iSCC: initial somatic cell count, BCS: body condition score

Milk SCC analysis

For sampling of milk, teats and udders were cleaned and wiped off with towels soaked in antiseptic solution. Then, approximately 30 ml morning milk were sampled aseptically in sterilized plastic containers from each quarter and each cow separately at fortnightly interval for 180 days. The first 2-3 streaks of foremilk were discarded before collecting milk samples. SCC was performed using Lactoscan SCC (Milkotronic Ltd., Bulgaria; www.milkotronic.com) following the standard procedures; both the absolute ($\times 1000$ cells/ml) and \log_{10} SCC values were determined. The udder level SCC was calculated as a mean of four quarter SCC at each sampling.

Statistical analysis

The effect of feeding strategies on standing and lying behavior as well as SCC at teat and udder level was worked out by univariate analysis of variance with the help of SPSS statistical packages (version 25). The relationship of different events of standing and lying behavior (lying time, number of bouts, bout duration and PMSD) obtained due to the treatment effects and SCC at teat and udder level in dairy cows were performed by Pearson's Correlation analysis. Mean differences were maintained at $p \leq 0.05$.

Results

Standing and lying behavior of dairy cow

The daily standing and lying events (mean and SEM), i. e. lying time (hrs./d), lying bouts (no./d), bout duration (min./d) and post-milking standing duration (min./d) of dairy cows in different feeding strategies is presented in Table 2. The lying time of +iPMF cows was higher ($P = .003$) than other groups and the rest all were statistically similar. Likewise, lying bouts of -10PMF was higher ($P = .001$) compared to +iPMF, +15PMF and +30PMF cows but was similar with MFD, while MFD showed higher ($P = .001$) lying bouts than that of +iPMF and +30PMF. Bout duration was recorded higher ($P = .001$) in +iPMF cows; among the rest feeding strategies, +30PMF was higher ($P = .001$) than -10PMF but was similar with MFD and +15PMF. Lastly, the post-milking standing duration was logged significantly higher ($P = .001$) in +iPMF than -10PMF and MFD but was similar with +15PMF and +30PMF; -10PMF and MFD too were different ($P = .001$).

Table 2. Standing and lying behavior (mean and SEM) of tie stall dairy cow in different feeding strategies at Chitwan, Nepal

Factors	Lying time (hours /d)		Lying bouts (number/d)		Bout duration (minutes/d)		PMSD (minutes/d)	
-10PMF	11.15 ^d	0.22	11.75 ^a	0.32	57.97 ^c	1.70	62.65 ^c	3.18
MFD	11.13 ^b	0.24	11.44 ^{ab}	0.28	58.66 ^{bc}	1.02	91.06 ^b	2.36
+iPMF	11.86 ^a	0.18	10.59 ^c	0.25	68.15 ^a	1.67	105.52 ^a	1.37
+15PMF	11.08 ^b	0.22	10.75 ^{bc}	0.28	62.10 ^b	0.89	101.36 ^a	1.03
+30PMF	10.67 ^b	0.20	10.30 ^c	0.21	62.54 ^b	1.33	100.91 ^a	1.34
F- value	4.14		5.61		8.82		74.32	
P-value	0.003		0.001		0.001		0.001	

Mean within the respective column with dissimilar superscript letters are different at $P < 0.001$ and $P < 0.01$; -10PMF: animals with 10 min Pre-milking feeding, MFD: Animals with milking during feeding, +iPMF: Animals with immediate post-milking feeding, +15PMF: Animals with 15 min post-milking feeding and +30PMF: Animals with 30 min post-milking feeding; PMSD: post-milking standing duration; CI: confidence interval

Milk SCC in dairy cow

Milk SCC at teat- and udder-level (mean \pm SE) in different feeding strategies in dairy cows presented in Table 3. The +iPMF and +15PMF cows experienced lower SCC ($P=.001, .005$) at left quarters, but SCC of +iPMF cows was numerically lower than -10PMF and MFD in left hind teats. In right fore quarter, the +iPMF, +15PMF and +30PMF cows had statistically lower ($P=.001$) SCC but SCC of +iPMF cows was similar to MFD, however a different trend was observed in right hind quarter with significantly lower ($P=.001$) SCC in +iPMF, +15PMF and MFD. Again, SCC of +iPMF cows was comparable to that of -10PMF. At udder level, +iPMF and +15PMF cows evident significantly lower ($P=.001$) SCC. The +iPMF and +15PMF groups consistently showed longer standing time after milking and experienced lower SCC at both teat and udder level (Figure 1).

Table 3. Milk log₁₀ Somatic Cell Count (SCC) at quarter and udder level (mean and SEM) in different treatments in tie stall dairy cow at Chitwan, Nepal

Factors	LF	LH	RF	RH	p SEM	Udder level	SEM
	Log ₁₀ SCC						
-10PMF	5.26 ^a	5.10 ^{ab}	5.36 ^a	5.14 ^{ab}	0.03	5.38 ^a	0.03
MFD	5.22 ^a	5.13 ^{ab}	5.24 ^{ab}	4.67 ^c	0.13	5.40 ^a	0.05
+iPMF	4.74 ^b	4.84 ^{bc}	4.97 ^{bc}	4.93 ^{bc}	0.10	4.97 ^b	0.07
+15PMF	4.51 ^b	4.75 ^c	4.75 ^c	4.78 ^c	0.12	4.96 ^b	0.08
+30PMF	5.31 ^a	5.27 ^a	4.92 ^c	5.33 ^a	0.09	5.37 ^a	0.04
F-value	11.89	3.83	6.36	5.08		15.54	
P-value	0.001	0.005	0.001	0.001		0.001	

Mean log₁₀ SCC within each column with dissimilar superscript letters are significantly different at $P<0.001$ and $P<0.01$; -10PMF: animals with 10 min Pre- milking feeding, MFD: Animals with milking during feeding, +iPMF: Animals with immediate post-milking feeding, +15PMF: Animals with 15 min post-milking feeding and +30PMF: Animals with 30 min post-milking feeding; LF: left fore, LH: left hind, RF: right fore and RH: right hind quarter, Psem: pooled standard error of mean








Figure 1. Relationship of post milking standing duration (PMSD) and mean log₁₀ SCC in LF (a), LH (b), RF (c), RH (d) and at udder level (e) in crossbred dairy cow managed in different feeding time schedule. SCC: Somatic Cell Count, LF: left fore quarter, LH: left hind quarter, RF: right fore quarter and RH: right hind quarter

Correlation of Standing and lying behavior and SCC in dairy cow

The relationship of different events of standing and lying behavior with SCC at teat and udder level in dairy cows is depicted in Table 4. Lying time had significant and positive ($P=.001$). In contrast, lying bouts showed negative and remarkable ($P=.001$) link with bout duration ($r=-0.64$) and post-milking standing duration ($P=.001$; $r=0.30$). Further, bout duration had a significant positive ($P=0.002$) correlation with post-milking standing duration ($r=0.24$), while PMSD showed significant but negative relationship with SCC of fore teats ($P=.001$; $r=-0.26$) and at udder level ($P=.001$; $r=-0.298$). Finally, association of SCC of each teat was significant and positive with udder SCC.

Table 4. Correlation of different events of standing and lying behaviors with teat and udder level somatic cell count in tie stall dairy cow at Chitwan, Nepal

	Lying Time	Lying Bouts	Bout Duration	PMSD	LF SCC	LH SCC	RF SCC	RH SCC
Lying Time								
Lying Bouts	.502 ^{**}							
Bout Duration	.322 ^{**}	-.641 ^{**}						
PMSD	-.045	-.301 ^{**}	.238 ^{**}					
LF SCC	-	-	-	-.263 ^{**}				
LH SCC	-	-	-	-.085	.573 ^{**}			
RF SCC	-	-	-	-.256 ^{**}	.221 ^{**}	.132		
RH SCC	-	-	-	-.056	.161 [*]	.145	.186 [*]	
Udder level SCC	-	-	-	-.298 ^{**}	.725 ^{**}	.629 ^{**}	.520 ^{**}	.458 ^{**}

^{**}Significant at the 0.01 level (2-tailed) and ^{*}at the 0.05 level (2-tailed); LF: left fore, LH: left hind, RF: right fore and RH: right hind quarter, SCC: somatic cell count; PMSD: post-milking standing duration

Discussion

The immediate post-milking feeding group (+iPMF) exhibited higher ($P=.003$) lying times (11.86 hrs./day) compared to other groups. This finding is in agreement with [24] and [25] who reported average lying time in dairy cows to range from 11.4 to 13.7 hrs./day. However, this study showed a longer lying duration than [26,27] who reported 10.5 ± 2.26 and 10.9 ± 2.26 hrs./day. They observed a bit shorter lying time than the Canadian Code of Practice for Dairy Cow [28] which recommended a minimum of 12 h/d lying time for dairy cow. Similarly, previous study [29] described at least 12–13 hrs./day lying time for heifers which is higher than our observations. In contrast, another study [30] described that changes in daily lying patterns of dairy cows coupled with decreased latency to lie down 20 min post-milking but without any alterations in lying duration on feeding green fodders 6 h post-milking. The +iPMF group showed the lowest lying bout frequencies ($P=.001$) than the other groups. This might indicate the more homogenous resting patterns or physiological balance in dairy cow. However, the -10PMF group showed significantly highest number of lying bouts ($P=.001$). The frequency of lying bouts in this study (10.30 to 11.75) was in comparable range to those discussed in earlier works that ranged between 10.7 and 11.9 bouts/day [25,31,32]. In contrast, [26] reported lower number of lying bouts (8.76 ± 3.58) in primiparous Italian dairy cows. Therefore, immediate post-milking feeding promoted a greater comfort with pronounced physiological balance in dairy cows as evidenced by higher mean daily lying duration and expected to have better udder health with lower SCC.

The mean bout duration was recorded to range from 57.97 to 68.15 min./d (Table 2) and the +iPMF group experienced the significantly longer bout duration ($P=.001$) which could be understood as a deeper uninterrupted rest of dairy cows with a healthy udder condition yielding higher production. Moreover, the differences among +iPMF, +15PMF and +30PMF in bout duration, despite all being post-milking feeding strategies, suggest that feeding time alone may not be the only determinant. Both frequencies of lying bouts ($r=0.50$) and their duration ($r=0.32$) showed positive association ($P=.001$) with lying time and were greater in the morning compared to the evening.

The mean lying time recorded in present study was more or less similar at par with [33] who reported an average lying time of 11.4 h/d with a frequency of 9.5 bouts/d but we observed the higher number of lying bouts, might be associated with higher parities of cows. In the same line, another study [26] described the longer lying time linked with higher parity, advanced lactation and higher BCS. Cows in their third or higher parity were observed lying down approximately 0.5 hours more daily compared to primiparous cows. Additionally, cows with a BCS 3.5 rested for about 1 hour more daily than those with a BCS 2.25 [26].

Post-milking feeding strategies led to longer standing durations after milking (PMSD). The +iPMF, +15PMF and +30 PMF groups showed significantly higher PMSD (105.52, 101.36 and 100.91 min.) in comparison to pre- and during milking feeding. The delivery of fresh green fodders had promoted the cows to feed and might have stimulated the longer standing behavior after milking. [34] described that the cows stood on an average for 48 min with free reach to feed after milking while standing duration was only 21 min when they did not have access to feed. Similarly, Shultz [35] reported shorter post-milking standing time in considerably more dairy cows (within 15 min) when feed was scarce after milking compared with when it was abundant. Johansson et al. [36] documented the correlation of time of feeding and lying behavior in dairy cows immediately after milking, who compared lying behavior of cows managed on three feeding strategies, i. e. 1.5 h pre-milking, right at milking, and 1.5 h

post-milking feeding and found significantly lower lying episodes in cows within an hour of milking that were fed during milking. Surprisingly, cows that were fed 1.5 hours after milking spent the more time lying down during the initial hour post-milking. However, it is challenging to directly compare our results with [36] because we partitioned our cows in 5 groups based of feeding time and each group had much less time differences in terms of feed delivery.

The post-milking feeding strategies, especially the cows that were offered green fodders immediately after milking (+iPMF) and 15 min after milking (+15PMF) consistently experienced lower SCC at quarter and udder level in this study. This lower milk SCC is probably associated to the post-milking feeding strategies we adopted in the study which motivated the cows to pay more time for eating and thus resulted in longer standing durations after milking. The earlier studies well described that the presence and delivery of fresh green fodders/forages encourage the cow's feeding behavior and promotes the longer standing time after milking [11,34,35,37–39]. These findings lighting the lower SCC with immediate or 15 min post milking feeding (yielding higher PMSD) is in line with [39] who reported that cows showing higher PMSD associated with post milking feeding experienced lower SCC. In line to the results of this study, [11] reported the association between standing period after milking and incidence of elevated SCC (eSCC) and stated that cows which rested more than 90 minutes after milking had a reduced chance of developing a new eSCC compared to those lying down earlier. Additionally, another study by DeVries et al. [38] found that cows standing for 40 to 60 minutes after milking were less likely to get a new intra-mammary infection. The Lower SCC observed in +iPMF and +15PMF cows in this study could be linked to the phenomenon where teat canals stay open for a period after milking. If the animals stand longer, it allows sufficient time for the teat and associated structures to undergo involution, reducing their degree of contact with the stall surface and significantly lowering the risk of infection [35]. As obvious, the higher SCC in any teats in study animals posed the greater risk of increased SCC and intra-mammary infections in other teats (LH; $r=0.57$, RF; $r=0.13-0.22$, RH; $r=0.15-0.19$) and udder ($r=0.46-0.72$) as a whole. The lower SCC in during milking feeding group (MDF) and 30 min post-milking feeding group (+30PMF) in right hind and fore teats in this study were difficult to describe. Overall, the post-milking feeding strategies, particularly +iPMF and +15PMF consistently yielded in lower SCC, likely attributed to longer standing durations after milking and decreased the risk of intramammary infections.

Conclusion

The cows of immediate and 15 min post-milking feeding group showed significantly higher lying times, lower number of bouts and extended bout durations compared to other groups, indicating that these feeding strategies had promoted greater comfort with more homogenous resting patterns or physiological balance yielding substantial welfare and potentially healthier udder conditions in dairy cows. Post-milking feeding strategies motivated the cows to lie down considerably late after milking for the first time and resulted a longer standing durations after milking compared to pre- and during-milking feeding groups. The said post-milking feeding strategies resulted in consistently lower somatic cell counts (SCC) at both teat and udder levels, indicating a potential reduction in the risk of IMI. Therefore, these findings underscore the importance of post-milking feeding strategies in promoting dairy cow welfare and udder health. However, further research is needed to explore the complex relationships between feeding times, lying behavior, and SCC to optimize dairy cow management.

Ethical approval

The study did not involve approaches that challenge well-being and welfare animals; we, authors, hereby declare that Principles of laboratory animal care (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable.

References

- [1] Ndahetuye JB, Persson Y, Nyman AK, Tukei M, Ongol MP, Båge R. Aetiology and prevalence of subclinical mastitis in dairy herds in peri-urban areas of Kigali in Rwanda. *Trop Anim Health Prod.* 2019;51:2037–44. <https://doi.org/10.1007/s11250-019-01905-2>
- [2] Costa A, de Marchi M, Sagrafoli D, Lanzi H, Amatiste S, Boselli C, et al. Milk somatic cell count and polymorphonuclear cells in healthy quarters of cows that underwent blanket and selective dry therapy: An Italian case study. *Vet Sci.* 2021;8. <https://doi.org/10.3390/vetsci8120298>
- [3] Petzer IM, Karzis J, Donkin EF, Webb EC, Etter EMC. Somatic cell counts thresholds in composite and quarter milk samples as indicator of bovine intramammary infection status. *Onderstepoort J. Vet. Res.* 2017;84:1–10. <https://doi.org/10.4102/ojvr.v84i1.1269>
- [4] Haxhiaj K, Wishart DS, Ametaj BN. Mastitis: What It Is, Current Diagnostics, and the Potential of Metabolomics to Identify New Predictive Biomarkers. *Dairy.* 2022;3:722–46. <https://doi.org/10.3390/dairy3040050>
- [5] Alhussien MN, Dang AK. Milk somatic cells, factors influencing their release, future prospects, and practical utility in dairy animals: An overview. *Vet World.* 2018;11:562. <https://doi.org/10.14202/VETWORLD.2018.562-577>
- [6] Cobirka M, Tancin V, Slama P. Epidemiology and classification of mastitis. *Animals.* 2020;10:1–17. <https://doi.org/10.3390/ani10122212>
- [7] Kitchen BJ. Review of the progress of dairy science: Bovine mastitis: Milk compositional changes and related diagnostic tests. *J. Dairy Res.* 1981;48:167–88. <https://doi.org/10.1017/S0022029900021580>
- [8] Williamson JR, Callaway TR, Rollin E, Ryman VE. Association of pre-treatment somatic cell counts with bacteriological cure following diagnosis of intramammary infection. *Res Vet Sci.* 2022;152:537–45. <https://doi.org/10.1016/j.rvsc.2022.09.015>
- [9] Lovarelli D, Tamburini A, Mattachini G, Zucali M, Riva E, Provolo G, et al. Relating Lying Behavior with Climate, Body Condition Score, and Milk Production in Dairy Cows. *Front Vet Sci.* 2020;7:5–7. <https://doi.org/10.3389/fvets.2020.565415>
- [10] Barkema HW, Schukken YH, Lam TJGM, Beiboer ML, Benedictus G, Brand A. Management practices associated with the incidence rate of clinical mastitis. *J Dairy Sci.* 1999;82:1643–54. [https://doi.org/10.3168/jds.S0022-0302\(99\)75393-2](https://doi.org/10.3168/jds.S0022-0302(99)75393-2)
- [11] Watters MEA, Meijer KMA, Barkema HW, Leslie KE, von Keyserlingk MAG, DeVries TJ. Associations of herd- and cow-level factors, cow lying behavior, and risk of elevated somatic cell count in free-stall housed lactating dairy cows. *Prev Vet Med.* 2013;111:245–55. <https://doi.org/10.1016/j.prevetmed.2013.05.015>

- [12] Tyler JW, Fox LK, Parish SM, Swain J, Johnson DL, Grasseschi HA, et al. Effect of feed availability on post-milking standing time in dairy cows. *Journal of Dairy Research*. 1997;64:617–20. <https://doi.org/10.1017/S0022029997002501>
- [13] DeVries TJ, von Keyserlingk MAG. Time of feed delivery affects the feeding and lying patterns of dairy cows. *J Dairy Sci*. 2005;88:625–31. [https://doi.org/10.3168/jds.S0022-0302\(05\)72726-0](https://doi.org/10.3168/jds.S0022-0302(05)72726-0)
- [14] Johansson B, Uvnäs-Moberg K, Knight CH, Svennersten-Sjaunja K. Effect of feeding before, during and after milking on milk production and the hormones oxytocin, prolactin, gastrin and somatostatin. *J. Dairy Res*. 1999;66:151–63. <https://doi.org/10.1017/S0022029999003532>
- [15] DeVries TJ, Dufour S, Scholl DT. Relationship between feeding strategy, lying behavior patterns, and incidence of intramammary infection in dairy cows. *J Dairy Sci*. 2010;93:1987–97. <https://doi.org/10.3168/JDS.2009-2692>
- [16] DeVries TJ, von Keyserlingk MAG, Beauchemin KA. Frequency of Feed Delivery Affects the Behavior of Lactating Dairy Cows. *J Dairy Sci*. 2005;88:3553–62. [https://doi.org/10.3168/JDS.S0022-0302\(05\)73040-X](https://doi.org/10.3168/JDS.S0022-0302(05)73040-X)
- [17] Fregonesi JA, Tucker CB, Weary DM. Overstocking Reduces Lying Time in Dairy Cows. *J Dairy Sci*. 2007;90:3349–54. <https://doi.org/10.3168/JDS.2006-794>
- [18] Gomez A, Cook NB. Time budgets of lactating dairy cattle in commercial freestall herds. *J Dairy Sci*. 2010;93:5772–81. <https://doi.org/10.3168/jds.2010-3436>
- [19] Mattachini G, Riva E, Provolo G. The lying and standing activity indices of dairy cows in free-stall housing. *Appl Anim Behav Sci*. 2011;129:18–27. <https://doi.org/10.1016/j.applanim.2010.10.003>
- [20] Langoni H, Penachio DS, Nobrega DB, Guimaraes FF, Lucheis SB, Baldini S. Somaticell as a screening method for somatic cell count from bovine milk. *Ciencia Rural*. 2012;42:1095–101.
- [21] Ito K, Weary DM, von Keyserlingk MAG. Lying behavior: Assessing within- and between- herd variation in free-stall-housed dairy cows. *J Dairy Sci*. 2009;92:4412–20. <https://doi.org/10.3168/jds.2009-2235>
- [22] Ledgerwood DN, Winckler C, Tucker CB. Evaluation of data loggers, sampling intervals, and editing techniques for measuring the lying behavior of dairy cattle. *J Dairy Sci*. 2010;93:5129–39. <https://doi.org/10.3168/jds.2009-2945>
- [23] Ohnstad I. Body condition scoring in dairy cattle: monitoring health to improve milk yield and fertility. *Livestock*. 2013;18:70–3. <https://doi.org/10.12968/live.2013.18.3.70>
- [24] Cook NB, Bennett TB, Nordlund K V. Monitoring indices of cow comfort in free-stall-housed dairy herds. *J Dairy Sci*. 2005;88:3876–85. [https://doi.org/10.3168/jds.S0022-0302\(05\)73073-3](https://doi.org/10.3168/jds.S0022-0302(05)73073-3)
- [25] Bewley JM, Boyce RE, Hockin J, Munksgaard L, Eicher SD, Einstein ME, et al. Influence of milk yield, stage of lactation, and body condition on dairy cattle lying behavior measured using an automated activity monitoring sensor. *J. Dairy Res*. 2010;77:1–6. <https://doi.org/10.1017/S0022029909990227>
- [26] Mattachini G, Tamburini A, Zucali M, Bava L, Riva E, Provolo G, et al. Relationships among lying and standing behaviour, body condition score and milk production in

- primiparous cows. *Ital J Anim Sci.* 2020;19:772–82.
<https://doi.org/10.1080/1828051X.2020.1793695>
- [27] Kathambi EK, VanLeeuwen JA, Gitau GK, Kamunde C. Risk factors associated with cows' lying time, stall and cows' own cleanliness in smallholder dairy farms in Kenya. *Vet World.* 2019;12:1085–92. <https://doi.org/10.14202/vetworld.2019.1085-1092>
- [28] NFACC. Code of practice for the care and handling of dairy cattle. vol. 1. 2009.
- [29] Jensen MB, Pedersen LJ, Munksgaard L. The effect of reward duration on demand functions for rest in dairy heifers and lying requirements as measured by demand functions. *Appl Anim Behav Sci.* 2005;90:207–17.
<https://doi.org/10.1016/j.applanim.2004.08.006>
- [30] DeVries TJ, Von Keyserlingk MAG. Time of feed delivery affects the feeding and lying patterns of dairy cows. *J Dairy Sci.* 2005;88:625–31.
[https://doi.org/10.3168/jds.S0022-0302\(05\)72726-0](https://doi.org/10.3168/jds.S0022-0302(05)72726-0)
- [31] Blackie N, Scaife JR, Bleach ECL. Lying behaviour and activity of early lactation Holstein dairy cattle measured using an activity monitor. *Cattle Prac.* 2006;14:139–42.
- [32] Endres MI, Barberg AE. Behavior of dairy cows in an alternative bedded-pack housing system. *J Dairy Sci.* 2007;90:4192–200. <https://doi.org/10.3168/jds.2006-751>
- [33] Westin R, Vaughan A, de Passillé AM, DeVries TJ, Pajor EA, Pellerin D, et al. Lying times of lactating cows on dairy farms with automatic milking systems and the relation to lameness, leg lesions, and body condition score. *J Dairy Sci.* 2016;99:551–61.
<https://doi.org/10.3168/jds.2015-9737>
- [34] Tyler JW, Fox LK, Parish SM, Swain J, Johnson DL, Grasseschi HA, et al. Effect of feed availability on post-milking standing time in dairy cows. *J. Dairy Res.* 1997;64:617–20.
- [35] Shultz TA. Feeding affects post-milking cow activities Abundant feed after milking keeps cows on their feet. *Calif Agric (Berkeley).* 1985;39:29–30.
- [36] Johansson B, Redbo I, Svennersten-Sjaunja K. Effect of feeding before, during and after milking on dairy cow behaviour and the hormone cortisol. *Animal Science.* 1999;68:597–604. <https://doi.org/10.1017/S1357729800050621>
- [37] DeVries TJ, Von Keyserlingk MAG. Time of feed delivery affects the feeding and lying patterns of dairy cows. *J Dairy Sci.* 2005;88:625–31.
[https://doi.org/10.3168/jds.S0022-0302\(05\)72726-0](https://doi.org/10.3168/jds.S0022-0302(05)72726-0)
- [38] DeVries TJ, Dufour S, Scholl DT. Relationship between feeding strategy, lying behavior patterns, and incidence of intramammary infection in dairy cows. *J Dairy Sci.* 2010;93:1987–97. <https://doi.org/10.3168/jds.2009-2692>
- [39] Bharti P, Bhakat C. Association Among Feeding After Milking, Animal Factors and Postmilking Standing Period and Effect of Post-milking Standing Period on Intra-Mammary Infection in Crossbred Cows. *Int J Livest Res.* 2019;9:136–43.
<https://doi.org/10.5455/ijlr.20181027064304>