

Phytochemical and antioxidant investigations of extracts from the leaves of *Macaranga heterophylla* (Müll. Arg.) (Euphorbiaceae), a medicinal species used in Côte d'Ivoire

ABSTRACT

Aims: This work aims to investigate the phytochemical composition and antioxidant potential of the leaves of *Macaranga heterophylla*.

Methodology: For this purpose, phytochemical screening by detection tests and thin layer chromatography (TLC), determination of total phenols content, total flavonoids content and condensed tannins content, and assessment of antioxidant potential by DPPH and reducing power tests were carried out on aqueous crude extracts, ethanolic crude extracts and selective extracts of the leaves of *M. heterophylla*

Results: The percentage yields obtained with ethanol (70%) (28.90 and 24.70% for the ethanolic decoction and ethanolic macerate respectively) are higher than those obtained with water (24.30 and 21.10% for the aqueous decoction and aqueous macerate respectively). The phytochemical screening highlighted the presence of several phytochemical families such as phenolic compounds (coumarins, flavones, tannins), quinones, sterols and polyterpenes, saponosides, glycosides, cardiotoxic glycosides and oligosaccharides. Quantitative analysis of total phenolics, total flavonoids and proanthocyanidols showed that their respective levels in the leaves of *M. heterophylla* varied depending on the solvent and the extraction technique. Concerning total phenolics, the aqueous decoction and ethanolic macerate gave the best total phenol contents (129.04 ± 9.53 and 119.82 ± 2.63 mg EAG/g DM respectively); for total flavonoids, the aqueous decoction gave the best content (33.03 ± 1.61 mg EQ/g DM), while for condensed tannins, the aqueous macerate gave the best content (0.87 ± 0.02 mg ECT/g DM). With regard to DPPH antioxidant activity, the results showed that the ethanolic decoction has more pronounced antioxidant activity than the aqueous decoction, while the aqueous macerate showed better antioxidant activity than the ethanolic macerate. Concerning the reducing power test, the opposite trend was observed.

Conclusion: The present study demonstrated that *M. heterophylla* is a concentrate of secondary metabolites with antioxidant properties, which would explain its use in traditional medicinal practice.

Keywords: *Macaranga heterophylla*, leaves, phytoconstituents, antioxidant activity.

1. INTRODUCTION

Essential natural resources, including medicinal plants, have always been the main source of medicines because of their wealth of biologically active phytochemicals, commonly known as secondary metabolites, which they synthesize [1]. Secondary metabolites are produced as an outcome of primary metabolism [2]. They are not essential for the growth and reproduction of the plant but they play many roles including defense against pathogens, pests and herbivores, response to environmental stresses and mediating organismal interactions [3-5]. As mentioned above, these metabolites also have a wide range of biological activities (antioxidants, antimicrobials, anticancer agents, etc.), making them highly valuable for human health and well-being, hence the use of plants in traditional medicine in

many ancient communities [6-8] Despite this, few plant species have been studied for medicinal applications [9]. To date, 400.000 known plant species have been the subject of chemical and pharmacological studies [10].

Macaranga is the largest genera in the Euphorbiaceae family. It is known to consist of about 300 species from tropical Africa, south-east Asia, Australia and the Pacific region [11].

Previous work on this genus has demonstrated various bioactivities, including antioxidant [12, 13], antimicrobial [14, 15], anti-inflammatory [16, 17], anticancer [18, 19] and other types of biological activity [20-22]. Notwithstanding the great interest in this genus, there is a lack of information about the phytochemical and antioxidant properties of the *M. heterophylla* species we are investigating. It is an ornamental species used for its captivating foliage and flowers and traditionally used to treat snake bites, coughs, certain infectious diseases, as a purgative and to facilitate childbirth [23-27]. This work aims to investigate the phytochemical composition and antioxidant potential of the leaves of *M. heterophylla*.

2. MATERIALS AND METHODS

2.1 Plant Material

The plant material used consists of the leaves of *M. heterophylla*. They were harvested in April 2021 at Petit Yapo (5° 47' 51" N, 4° 8' 21" W), a locality near Azaguié that is a town in the Agnéby-Tiassa region of southern Côte d'Ivoire. After identification and authentication at the Centre National de Floristique (CNF) of the Félix HOUPHOUËT-BOIGNY University (Cocody/Abidjan) in accordance with the existing herbarium (N° UCJ006139), the leaves were cleaned with water and then dried under air conditioning (18° C) for 2 weeks. After drying, they were ground to a powder using an electric grinder (Vorwerk, Thermomix 3300) to improve the contact area with the solvents.

2.2 Methods

2.2.1 Preparation of extracts

Aqueous and ethanolic (70 %) decoctions: 30 g of powder were boiled in 600 ml of distilled water and 600 ml of ethanol (70%) for 10 min, respectively. After cooling to room temperature and filtration through a büchner, the decoctions were concentrated in vacuum using a rotary evaporator (BÜCHI R100), and oven-dried at 45°C until the dry extracts DA (aqueous decoction) and DE (ethanolic decoction) were obtained.

Aqueous and ethanolic (70 %) macerates: 30 g of powder were macerated in 600 ml of distilled water and 600 ml of 70% ethanol for 24 hours, respectively. The solutions obtained were filtered through a büchner. The macerates were concentrated under vacuum using a rotary evaporator (BÜCHI R100), then kept in an oven (45°C) until the dry extracts MA (aqueous macerate) and ME (ethanolic macerate) were obtained.

Selective extracts: an aliquot of each dry DA, DE, MA, ME extract taken up in 100 ml of distilled water was successively exhausted with petroleum ether (3x50 ml) and ethyl acetate (3x50 ml). The different organic fractions obtained constituted the petroleum ether (DA₁, DE₁, MA₁, ME₁) and ethyl acetate (DA₂, DE₂, MA₂, ME₂) extracts.

2.2.2 Qualitative tests

Phytochemical screening to identify the phytochemical families present in the leaves of *M. heterophylla* was carried out on decoctions and macerates, as well as on selective extracts, using either precipitation and color reaction tests [28-32] or thin-layer chromatography (TLC) [29, 33]. These tests focused on phenolic compounds (phenolic acids, coumarins,

flavonoids, tannins), quinones, saponosides, glycosides, sterols and polyterpenes, alkaloids and oligosaccharides.

2.2.3 Quantitative tests

2.2.3.1 Evaluation of total phenolic (TP) contents

Total phenolic contents were determined on decoctions and macerates using the Folin-Ciocalteu colorimetric method, with slight modifications. To 1 ml of each extract, diluted 1/10th with distilled water, were added 1.5 ml of sodium carbonate (Na_2CO_3 (17%, m/v) and 0.5 ml Folin-Ciocalteu reagent (0.5N). The mixture was dark-incubated at room temperature for 30 min. Absorbance was read at 760 nm with a UV-visible spectrophotometer (Spectro AL 800) against a blank without extract. Total phenolics were quantified according to a regression equation ($y = ax + b$) using gallic acid as standard at different concentrations (0.005 to 0.0375 mg/ml) under the same conditions as the sample. Results were expressed in milligrams of gallic acid equivalent per gram of dry matter (mg EAG/g DM) [30, 34].

2.2.3.2 Evaluation of total flavonoid (TF) contents

Total flavonoid contents were determined on decoctions and macerates as previously described [35]. 0.01 g of each crude extract were solubilized in 10 ml of distilled water to give the stock solution, which was diluted 1 :10. To 2 ml of the diluted solution, 2 ml of aluminium chloride (AlCl_3) (2%, m/v) in methanol (MeOH) were added. The mixture was dark-incubated for 15 min. Absorbance was read at 415 nm with a UV-visible spectrophotometer (Spectro AL 800), with distilled water used as the blank. A calibration line was drawn with quercetin at different concentrations (0.0005 to 0.0375 mg/ml). Total flavonoid contents were expressed in milligrams of quercetin equivalent per gram of dry matter (mg EQ/g DM).

2.2.3.3 Evaluation of condenses tannins (CT) or proanthocyanidols contents

Condensed tannins or proanthocyanidols contents were determined on decoctions and macerates as previously described with slight modifications [36]. To 0.2 ml of each sample, 1.5 ml of a methanolic solution of vanillin (4%, m/v) and 0.75 ml of concentrated hydrochloric acid (HCl) were added. The mixture was dark-incubated for 15 min and the absorbance was read at 500 nm with a UV-visible spectrophotometer (Spectro AL 800). Condensed tannins contents were determined from calibration ranges established with catechin at various concentrations (0.009 to 0.15 mg/ml), and expressed in milligrams of catechin equivalent per gram of dry matter (mg EC/g DM).

2.2.4 Evaluation of antioxidant potential

2.2.4.1 DPPH test

The antioxidant potential of decoctions and macerates was assessed using the stable radical 2,2'-diphenyl-1-picrylhydrazyl (DPPH) reduction assay with slight modifications [37]. A concentration range (0.5; 0.25; 0.125; 0.0625; 0.03125; 0.015625 mg/ml) of extracts (decoctions and macerates) and ascorbic acid (vitamine C, used as antioxidant reference) was prepared in ethanol (EtOH). The ethanolic solution of DPPH at concentration of 0.03 mg/ml was also prepared. The reaction mixture used for the test consisted of the extract (1 ml) and the DPPH solution (2.5 ml), introduced into the cuvette of the UV-visible spectrophotometer (Spectro AL 800). The absorbance of the reaction mixture was read at

517 nm every 2 min (from 0 to 30 min) against a blank (1 ml EtOH and 2.5 ml DPPH solution).

2.2.4.2 Reducing power assay

The ability of **decoctions** and macerates to reduce Fe^{3+} to Fe^{2+} was determined according to the methodologies described [38-40] with slight modifications. Extracts and standard agent (0.5 ml) of different concentrations (1 ; 0.5 ; 0.25 ; 0.125 ; 0.0625 and 0.03125 mg/ml) were mixed with 0.5 ml phosphate buffer (0.2 M, pH 6.6) and 0.5 ml potassium ferricyanide ($\text{K}_3[\text{Fe}(\text{CN})_6]$) (1%, m/v), then incubated at 50°C in a water bath for 20 min. The reaction was stopped by adding 0.5 ml of trichloroacetic acid ($\text{CCl}_3\text{CO}_2\text{H}$) solution (10%) and the mixture was centrifuged at 3000 rpm for 10 min. To 0.15 ml of the mixture were added 0.15 ml of distilled water and 0.04 ml of ferric chloride (FeCl_3) (0.1%, m/v). Then, the mixture was incubated for 10 min and the absorbance of the reaction mixture was measured at 700 nm with a multimode microplate reader (VarioskanTM LUX, Thermo Fisher Scientific). A higher absorbance value indicated greater reducing power.

2.3 Statistical analysis

Statistical analysis of the results was carried out using Graph Pad prism version 8.4.2 software to compare total phenols, total flavonoids, total condensed tannins and antioxidant activity. The Student's t-test was used to check whether the means of the variables differed between the samples (the **decoctions** on the one hand and the macerates on the other hand). Differences at $P < 0.05$ were considered statistically significant.

3. RESULTS AND DISCUSSION

3.1 Extraction yields

Extraction yields obtained with 30 g of plant powder are respectively 24.30% for DA, 28.90% for DE, 21.10% for MA and 24.70% for ME (Table 1). We note that the yields obtained with ethanol (70%) are higher than those obtained with water. Indeed, under the same conditions of extraction time and temperature, solvent and sample composition are the most important parameters affecting yield. The use of a hydroalcoholic solvent may therefore facilitate the extraction of phytoconstituents that are soluble in water and/or in the organic solvent [41], which could explain the high yields obtained from ethanolic extractions (70%).

Table 1. Extraction yields

Extracts	Yields (%)
DA	24.30
DE	28.90
MA	21.10
ME	24.70

3.2 Qualitative phytochemical profile

3.2.1 Phytochemical composition using precipitation and color reaction tests

Phytochemical screening revealed the presence of most phytoconstituents found in **decoctions** and macerates, namely phenolic compounds (coumarins, flavones, tannins),

quinones, sterols and polyterpenes, saponosides, glycosides including cardiotonics, **oligosaccharides**; with the exception of flavones for DE, phlobatanins and ketoses. The presence or absence of alkaloids in extracts depends on the type of reagent used (Table 2).

Table 2. Results of phytochemical screening using precipitation or color reaction tests

Phytoconstituents	Tests	DA	DE	MA	ME
Phenolic compounds	FeCl ₃ (2%)	+	+	+	+
Flavonoids	Shinoda (Mg) / HCl	+	-	+	+
	NH ₄ OH (Flavones)	+	+	+	+
Coumarins	NaOH (10%)	+	+	+	+
Tannins	FeCl ₃ (5%)	+	+	+	+
Alkaloids	H ₂ SO ₄	+	+	+	+
	Bürchard	+	+	+	+
	Dragendorff	-	-	-	-
	Mayer	+	-	-	-
	Picric acid	-	-	-	-
Phlobatannins	HCl	-	-	-	-
Sterols and polyterpenes	CH ₃ CO ₃ CH ₃ , H ₂ SO ₄	+	+	+	+
Saponosides	Fi		200		
Glycosides	CHCl ₃ , NH ₄ OH	+	+	+	+
Cardiotonic glycosides	Liebermann-Bürchard	+	+	+	+
Oligosaccharides	Molish	+	+	+	+
	Seliwanoff (ketoses)	-	-	-	-

(+): positive ; (-): negative ; Fi: Foam index

3.2.2 Phytochemical composition by thin layer chromatography (TLC)

Phytochemical screening by precipitation or color reaction tests does not specify the exact type of phytomolecules sought. That is why we also decided to determine the phytochemical composition of extracts by TLC. This analytical planar chromatographic technique enables phytocompounds to be separated according to their migration properties. It was carried out on selective extracts, which are less complex in terms of phytochemical content than matrix extracts (decoctions and macerates). The results are presented below (Table 3). Several secondary metabolites such as sterols, terpenes, triterpenes of the oleanane or ursane type, coumarins, flavonoids and tannins were identified using TLC. The Liebermann-Bürchard reagent revealed sterols as yellow spots under UV/365 nm. This reagent also revealed oleanane or ursane-type triterpenes as red spots under UV/365 nm [42, 43]. Sulfuric vanillin

revealed terpenes as purple spots and sterols as blue spots in visible [43, 44]. Aluminium chloride (AlCl₃, 1%) revealed flavonoids as green, yellow and blue spots under UV/365 nm [28, 43]. Flavonoids were also highlighted by ammonia under UV/365 nm as green spots [28, 45]. Methanolic solution of potassium hydroxide (KOH, 5%) was used to detect coumarins as yellow-orange spots in visible and as yellow, green, blue and blue fluorescent spots under UV/365 nm [28, 46]. In addition, basic lead acetate (C₄H₁₀O₆Pb, 5%) highlighted them as blue and green spots fluorescing under UV/365 nm [47]. Tannins were revealed by FeCl₃ (2%) as gray spots in visible [43, 48]. Examination of the TLC phytochemical profile confirms the results obtained after screening by precipitation and color reaction tests.

Table 3. Results of phytochemical screening using TLC

Selective extracts	Secondary metabolites identified : [Rf], color
Petroleum ether extracts	DA ₁ Sterols ^{e,f} : [0,16] Y ^e ; [0,33] B ^f ; Terpenes ^f : [0,83] O ^f ;
	DE ₁ Terpenes ^f : [0,66] P ^f ; [0,85] P ^f ; [0,96] P ^f ; Sterols ^f : [0,44] B ^f ;
	MA ₁ Terpenes ^f : [0,15] P ^f ; [0,29] P ^f ;
	ME ₁ Sterols ^{e,f} : [0,16] Y ^e ; [0,19] B ^f ; [0,30] Y ^e ; [0,44] Y ^e ; [0,85] Y ^e ; Triterpenes oleanone or ursane ^e : [0,41] R ^e ; [0,55] R ^e ; [0,69] R ^e ;
Ethyl acetate extracts	DA ₂ Coumarins ^{a,h} : [0,14] G ^h ; [0,16] B ^a ; [0,43] Br ^a ; Flavonoids ^b : [0,14] G ^b ; [0,40] G ^b ; Tannins ^c : [0,40] Gy ^c ;
	DE ₂ Coumarins ^{a,h} : [0,05] G ^a ; [0,26] YO ^a ; [0,61] Bf ^h ; [0,65] B ^a -B ^h ; [0,69] B ^h ; [0,75] B ^h ; Flavonoids ^b : [0,05] G ^b ; [0,49] B ^b ; [0,59] B ^b ; [0,65] B ^b ; [0,69] B ^b ; Tannins ^c : [0,26] Gy ^c ; [0,40] Gy ^c ;
	MA ₂ Coumarins ^{a,h} : [0,06] G ^a ; [0,25] YO ^a ; [0,65] Bf ^a ; Flavonoids ^b : [0,06] G ^b ; [0,19] B ^b ; [0,51] B ^b ; Tannins ^c : [0,25] Gy ^c ;
	ME ₂ Coumarins ^{a,h} : [0,16] B ^a -B ^h ; [0,40] YO ^a ; [0,59] Y ^a ; [0,63] YO ^a ; [0,75] Bf ^a -Bf ^h ; Flavonoids ^{b,g} : [0,11] G ^b ; [0,44] G ^g ; [0,61] Bf ^b ; [0,68] G ^g ; [0,69] Bf ^b ; [0,75] Y ^b ; [0,85] Y ^b ; [0,89] G ^g ; Tannins ^c : [0,05] Gy ^c ; [0,44] Gy ^c ;

a : KOH ; b : AlCl₃ ; c : FeCl₃ ; e : Liebermann-Bürchard ; f : sulfuric vanillin ; g : ammoniac ; h : basic lead acetate ; O : orange ; YO : yellow orange , B : blue ; Y : yellow ; G : green ; Bf : blue fluorescent ; P : purple ; R : red ; Gy : gray

3.3 Quantitative phytochemical profile

3.3.1 TP contents

The TP contents (Table 4) were obtained from the linear regression equation ($y = 6.7469x - 0.0353$; $R^2 = 0.994$) established with a concentration ranges of gallic acid. They range from 84.99 to 129.04 mg (EAG/g DM). In the case of **decoctions**, TP contents are significantly higher ($P < 0.05$) in DA (129.04 mg EAG/g DM on average) than in DE (84.99 mg EAG/g DM on average). An opposite trend was observed in previous studies [49, 50]. In the case of

macerates, TP contents are significantly higher ($P < 0.05$) in ME (119.82 mg EAG/g DM on average) than in MA (107.90 mg EAG/g DM on average). A similar trend was observed in work carried out on the species *Helichrysumstoechas* [49], indicating that estimated total phenolic contents are higher in ethanolic maceration (70%) than in aqueous maceration. Previous studies [51-53] have reported that ethanol combined with water provides better extraction of phytophenols. From these results, we deduce that the total phenolic content depends on the type of extraction and the polarity of the solvent used. As a result, aqueous decoction appears to be suitable for better extraction of total phenolics.

Table 4. TP contents

Extracts	TP contents (mg EAG/g DM)
DA	129.04 ± 9.53 a
DE	84.99 ± 4.44 b
MA	107.90 ± 3.15 a
ME	119.82 ± 2.63 b

Results are expressed as mean ± SD (n=3). There is a significant difference between values followed by a different letter ($P < 0.05$) as measured by the Student's t-test

3.3.2TF contents

The TF contents (Table 5) were determined from the linear regression equation ($y = 22.779x - 0.0159$; $R^2 = 0.999$) established with a concentration ranges of quercetin. They range from 13.42 to 33.03 (mg EQ/g DM). They are significantly higher ($P < 0.001$) in DA (33.03 mg EQ/g DM on average) than in DE (13.42 mg EQ/g DM on average). An opposite trend has been reported by other authors [49, 50], mentioning that total flavonoid contents are higher in ethanolic decoction (70%) than in aqueous decoction. For macerates, no significant difference was found between total flavonoid contents ($P > 0.05$). However, according to these authors [54], an ethanol/water mixture (70, 30 ; v/v) would increase the quantity of flavonoids. From the results we have obtained, we deduce that aqueous decoction would be the most suitable method for flavonoid extraction.

Table 5. TF contents

Extracts	TF contents (mg EQ/g DM)
DA	33.03 ± 1.61 a
DE	13.42 ± 1.28 b
MA	23.56 ± 1.33 a
ME	23.42 ± 1.48 a

Results are expressed as mean ± SD (n=3). There is a significant difference between values followed by a different letter ($P < 0.05$) as measured by the Student's t-test

3.3.3CT or proanthocyanidols contents

The CT contents (Table 6) were determined from the linear regression equation ($y = 1.6546x + 0.0086$; $R^2 = 0.998$) established with a concentration ranges of catechin. They range from 0.59 to 0.87 (mg EC/g DM). For **decoctions**, no significant difference was found between CT

contents ($P > 0.05$). Regardless of the solvents used, decoctions extracted practically the same quantities of CT. However, CT contents were significantly higher ($P < 0.05$) in MA (0.87 mg EC/g DM on average) than in ME (0.61 mg EC/g DM on average). A similar trend was observed in the bracts of *Cynarascolumus* L. In fact, CT contents were higher in the aqueous macerate than in the ethanolic macerate. In addition, previous investigations [53, 55] have shown that decoction is more effective at extracting condensed tannins than maceration. However, according to our results, the high content of condensed tannins is obtained by aqueous maceration. This can be explained by the fact that decoction, which consists of immersing the plant in a solvent and boiling for a few minutes, leads to the degradation of heat-sensitive tannins [56].

Table 6. CT contents


Extracts	CT contents(mg EC/g DM)
DA	0,59 ± 0,03 a
DE	0,59 ± 0,03 a
MA	0,87 ± 0,02 a
ME	0,61 ± 0,09 b

Results are expressed as mean ± SD (n=3). There is a significant difference between values followed by a different letter ($P < 0.05$) as measured by the Student's t-test


3.4 Antioxidant profile

3.4.1 DPPH antioxidant activity

Decoctions and macerates showed antioxidant potential with concentration-dependent percentages reduction (PR) of DPPH radical as a function of time. In the first few minutes, a rapid drop in the absorbance of the DPPH radical is observed, followed by a slow step until equilibrium is reached. Concerning the PR of DPPH radical by the **decoctions** (Figures 1), DA showed a maximum DPPH scavenging activity of 86% at 0.5 mg/ml (30 min), whereas for DE was found to be 78.55% at the same concentration and time. Concerning the PR of DPPH radical by the macerates (Figures 2), MA showed a maximum DPPH scavenging activity of 87.27% at 0.5 mg/ml (30 min), whereas for ME was found to be 90.75% at the same concentration and time.



Figures 1: Percentages reduction of DPPH radical by the decoctions (DA and DE) as a function of time



Figures 2: Percentages reduction of DPPH radical by the macerates (MA and ME) as a function of time

A slight drop of PR was observed with decoctions, which could be explained by the loss of heat-sensitive phytoconstituents. However, the PR of DPPH radical by the decoctions and the macerates are lower than that of vitamin C (95.75%) at 0.5 mg/ml (30 min) (Figure 3).




Figure 3: Percentages reduction of DPPH radical by vitamine C as a function of time

Determination of the CR_{50} (median concentration of the sample that reduces 50% of the DPPH) enabled a better assessment of the antioxidant potential of the extracts. The lower its value, the more pronounced the antioxidant activity [57]. The CR_{50} values of decoctions and macerates were determined graphically. (Figure 4).




Figure 4: CR_{50} values of decoctions and macerates as a function of time

Concerning the decoctions, CR_{50} values of DE are lower than those of DA, demonstrating that DE has a more pronounced antioxidant activity than DA. For the macerates, CR_{50} values of MA are lower than those of ME. MA therefore exhibits a better antioxidant activity than ME.

3.4.2 Reducing power

The reducing power assay is often used to assess the ability of compounds to reduce Fe^{3+} to Fe^{2+} [58]. Figure 5 shows the reducing power of decoctions and macerates, compared with ascorbic acid (AA), used as a reference antioxidant.




Figure 5 : Reducing power of decoctions, macerates and ascorbic acid

The reducing power of decoctions and macerates, like that of AA, is concentration-dependent. Indeed, increasing analyte concentrations induce a progressive variation in reducing power. Concerning the decoctions, DA showed a significantly higher reducing power ($P < 0.001$) than DE at 0.5 mg/ml. For the macerates, ME showed a significantly higher reducing power ($P < 0.001$) than MA, suggesting that the high levels of total phenolics in DA and ME are the reason. Phytochemicals have antioxidant properties as hydrogen donors. They are excellent free radical scavengers [59]. These results are in contrast to those of the DPPH test, which shows the antioxidant profile of DE to be better than that of DA, and that of MA to be better than that of ME. The judicious choice of these two methods seems to explain this difference with regard to the type of reduction mechanism [60, 61]. In any case, the presence of antioxidant phytoconstituents in decocted and macerated of the leaves of *M. heterophylla* justifies its traditional use in the treatment of various diseases.

4. CONCLUSION

This work has enabled us to determine the phytochemical and antioxidant profiles of the leaves of *M. heterophylla*. The study of the chemical composition and effects of a plant matrix is based on a rational approach: firstly, the choice of solvent for better extraction of the phytochemicals, and secondly, the choice of extraction method. The percentages of phytoconstituents extracted from the leaves of *M. heterophylla* varies according to solvent and extraction method. The phytochemical screening highlighted the presence of several phytochemical families, including phenolic compounds (coumarins, flavones, tannins), quinones, sterols and polyterpenes, saponosides, glycosides, cardiogenic glycosides and oligosaccharides. Quantitative analysis of total phenolics, total flavonoids and proanthocyanidols showed that their respective levels in the leaves of *M. heterophylla* varied depending on the solvent and the extraction technique. Assessment of the antioxidant power of the decoctions and macerates showed that the plant contains antioxidant phytoactives. These results suggest that the local species could be developed into a phytopreparation for therapeutic use in the form of an herbal tea as water was able to extract phytoconstituents better.

REFERENCES

1. Fouché JG, Marquet A, Hambuckers A. The medicinal plants from plant to medicine. Observatory of the world of plants, Sart-Tilman, Liège(Translated from french). 2000; 1-18.
2. Chandran H, Meena M, Barupal T, Sharma K. Plant tissue culture as a perpetual source for production of industrially important bioactive compounds. Biotechnology reports. 2020; 26:e00450.
3. Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q. Response of plant secondarymetabolites to environmental factors. Molecules. 2018; 23(4):762.
4. Isah T. Stress and defense responses in plant secondarymetabolites production. Biologicalresearch. 2019; 52:39.
5. Guerrieri A, Dong L, Bouwmeester HJ. Role and exploitation of underground chemical signaling in plants. Pest management science. 2019; 75(9):2455-2463.
6. Marchev AS, Yordanova ZP, Georgiev M.I. Green (cell) factories for advanced production of plant secondarymetabolites. Critical reviews in biotechnology. 2020; 40(4):443-458.
7. Wani AK, Akhtar N, Sharma A, El-Zahaby SA. Fighting carcinogenesis with plant metabolites by weakening proliferative signaling and disabling replicative immortality networks of rapidly dividing and invading cancerous cells. Current Drug Delivery. 2023; 20(4):371-386.
8. Builders PF, editor. Plants SecondaryMetabolites: The Key Drivers of the Pharmacological Actions of Medicinal Plants. Herbal Medicine, IntechOpen; 2019.
9. Strzemski M, Wójciak-Kosior M, Sowa I, Załuski D, Verpoorte R. Historical and traditional medical applications of *Carlina acaulis* L.-A critical ethnopharmacological review. Journal of ethnopharmacology. 2019; 239:111842.
10. Balick MJ, Cox PA. Plants, people, and culture: the science of Ethnobotany. 2nd ed. New York :Scientific American Library. 2020.
11. Magadula JJ. Phytochemistry and pharmacology of the genus *Macaranga*: a review. J Med Plants Res. 2014; 8(12):489-503.
12. Ardhanay SD, Mulia DS, Rosawanti P. Antioxidant activity of ethylacetate fraction of *Macaranga triloba* leaves from central Kalimantan. Asian Journal of Pharmaceutical and Clinical Research. 2018; 11(3):40-42.
13. Minarti M, Ariani N, Megawati M, Hidayat A, Hendra M, Primahana G, Darmawan A. Potential Antioxidant Activity Methods DPPH, ABTS, FRAP, Total Phenol and Total Flavonoid Levels of *Macaranga hypoleuca* (Reichb. f. & Zoll.) Leaves Extract and Fractions. In E3S Web of Conferences. EDP Sciences publisher. 2024; 503:07005.
14. Lee JH, Kim YG, Khadke SK., Yamano A, Woo JT, Lee J. Antimicrobial and antibiofilm activities of prenylated flavanones from *Macaranga tanarius*. Phytomedicine. 2019; 63 :153033.
15. Kamso VFK, Fotso CCS, Mbekou IMK, Toussie BT, Lenta BN, Boyom FF, Sewald N, Frese M, Ngadjui BT, Fotso GW. Chemical constituents of *Macaranga occidentalis*, antimicrobial and chemophenetic studies. Molecules. 2022; 27(24): 8820.

16. Asante-Kwatia E, Jibira Y, Mensah AY, Osei-Sarfoh D. *Macaranga barteri* stem bark extract exerts anti-inflammatory and anti-hyperalgesia activity in murine models. *DiscoveryPhytomedicine*. 2019; 6(3):130-137.
17. Remorosa AGB, Tsai PW, De Castro-Cruz KA, Hsueh CC, Chen RY, Chen BY. Deciphering characteristics of *Macaranga tanarius* leaves extract with electron shuttle-associated anti-inflammatory activity via microbial fuel cells, molecular docking, and network pharmacology. *Biochemical Engineering Journal*. 2024; 208:109345.
18. Arung ET, Amirta R, Zhu Q, Amen Y, Shimizu K. Effect of wood, bark and leaf extracts of *Macaranga* trees on cytotoxic activity in some cancer and normal cell lines. *Journal of the Indian Academy of Wood Science*. 2018; 15:115-119.
19. Nizar A, Ravindran R, Palani J, Nripan T, Asha SD, Pynadath MK. Anticancer Effects of Ethanolic Extracts of *Macaranga Peltata* Leaves on Human Oral Squamous Carcinoma Cell Lines: An In Vitro Study. *Journal of Pharmacy and Bioallied Sciences*. 2024; 16(Suppl 2):S1833-S1837.
20. Cahyana AH, Liandi AR, Wicaksono SS, Darmawan A. BIOLOGICAL ACTIVITIES OF EXTRACTS AND SECONDARY METABOLITE OF *Macaranga mappa* BARK AS AN ANTIDIABETIC BY α -GLUCOSIDASE INHIBITOR, ANTIOXIDANT, AND CYTOTOXICITY. *Rasayan Journal of Chemistry*. 2023; 16(1):290-296.
21. Ehile, EH, Kouakou KL, Goze NB, Yapo AP, Ehile EE. Effect of the Aqueous Extract of *Macaranga Barteri* Müll. Arg (Euphorbiaceae) on Acetic Acid Induced Chronic Ulcers in Rats. *African Journal of Biomedical Research*. 2021; 24(3):471-477.
22. Muhaimin M, Yusnaidar Y, Syahri W, Latief M, Putri RD, Utami A, Chaerunisaa AY, Aditama AY, Siregar JE. Antiplasmodial activity of ethanolic extract of *Macaranga gigantea* leaf and its major constituent. *Pharmacognosy Journal*. 2019; 11(6):1181-1188.
23. Bouquet A, Debray M. Medicinal plants of the Ivory Coast. ORSTOM, Paris (Translated from french). 1974; 231.
24. Kerharo J, Bouquet A. Medicinal and toxic plants of the Ivory Coast – Upper Volta : mission to study indigenous pharmacopoeia in A.O.F. Ed Vigot brothers. Paris (Translated from french). 1950; 297.
25. Kerharo J, Adam JG. Second inventory of the medicinal and poisonous plants of Casamance. Ed Masson. Paris (Translated from french). 1963; 773.
26. Kerharo J, Adam JG. The Traditional Senegalese Pharmacopoeia. Medicinal and toxic plants. Ed Vigot brothers. Paris (Translated from french). 1974; 1011.
27. Magassouba FB, Diallo A, Kouyaté M, Mara F, Mara O, Bangoura O, Camara A, Traoré S, Kéita A, Camara MK, Barry R, Kéita S, Oularé K, Barry MS, Donzo M, Camara K, Toté K, Vanden Bherge D, Totté J, Pieters L, Vlietinck AJ, Baldé AM. Ethnobotanical survey and antibacterial activity of some plants used in Guinean traditional medicine. *Journal of ethnopharmacology*. 2007; 114(1):44-53.
28. Ladiguina EY, Safronitch LN, Otriachenkova VE, Balandina IA, Grinkevitch NI. Chemical analysis of medicinal plants. Ed Moskva, Vischaya Chkola. Moscow. (Translated from russia). 1983; 347.
29. Bekro YA, Mamyrbekova-Bekro JA, Boua BB, Tra Bi FH, Ehilé E. Ethnobotanical study and phytochemical screening of *Caesalpinia benthamiana*

- (baill.) herend and zarucchi (caesalpiniaceae). *Science & nature*. 2007; 4(2):217-225.
30. N'Gaman KCC. Phytochemical study and effect of extracts of *Gmelina arborea* Roxb. (Verbenaceae) from Côte d'Ivoire on osmotic stability of erythrocytes. Single PhD thesis. University of Abobo-Adjamé Ivory Coast. (Translated from french). 2013; 152.
 31. Roghini R, Vijayalakshmi K. Phytochemical screening, quantitative analysis of flavonoids and minerals in ethanolic extract of *Citrus paradisi*. *International Journal of Pharmaceutical Sciences and Research*. 2018; 9(11):4859-4864.
 32. Melenteva GA, Antonova LA. *Pharmaceutical chemistry*. Ed Mir. Moscow. 1988; 472.
 33. Brou KG, Mamyrbekova-Békro JA, Dogbo DO, Gogbeu SJ, Békro YA. On the Qualitative Phytochemical Composition of Crude Hydromethanolic extracts of the Leaves of 6 Varieties of *Manihot esculenta* Crantz of Ivory Coast. *European Journal of Scientific Research*. 2010; 45(2):200-211.
 34. Singleton VL, Ortofer R, Lamuela-Raventos RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. In *Methods in enzymology*, Orlando Academic Press. 1999; 299:152-178.
 35. Arvouet GA, Vennat B, Pourrat A, Legret P. Standardization of propolis extract and identification of its main constituents. *Journal de pharmacie de Belgique*. (Translated from french). 1994; 49(6):462-468.
 36. Heilmer D, Vignolini P, Dini MG, Vincieri FF, Romani A. Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. *Food chemistry*. 2006; 99(3):464-469.
 37. Espin JC, Soler-Rivas C, Wichers HJ. Characterization of the total free radical scavenger capacity of vegetable oils and oil fractions using 2, 2-diphenyl-1-picrylhydrazyl radical. *Journal of Agricultural and Food chemistry*. 2000; 48(3):648-656.
 38. Oyaizu M. Studies on products of browning reaction: antioxidative activities of products of browning reaction prepared from glucosamine. *The Japanese journal of nutrition and dietetics*. 1986; 44(6):307-315.
 39. Boonchum W, Peerapornpisal Y, Kanjanapothi D, Pekkoh J, Pumas C, Jamjai U, Amornlerdpison D, Noiraksar T, Vacharapiyasophon P. Antioxidant activity of some seaweed from the Gulf of Thailand. *International Journal of Agriculture and Biology*. 2011; 13(1):95-99.
 40. Luqman S, Srivastava S, Kumar R, Maurya AK, Chanda D. Experimental assessment of *Moringa oleifera* leaf and fruit for its antistress, antioxidant, and scavenging potential using in vitro and in vivo assays. *Evidence-Based Complementary and Alternative Medicine*. 2012; 2012:519084.
 41. Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, Ju YH. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. *Journal of food and drug analysis*. 2014; 22(3):296-302.
 42. Bruneton J. *Pharmacognosy, Phytochemistry, Medicinal plants*. 3rd ed, Lavoisier Tec. & Doc. Paris. (Translated from french). 1999; 1120.
 43. Lagnika L. Phytochemical study and biological activity of natural substances isolated from Beninese plants. PhD thesis. Louis Pasteur University, Starsbourg (France) / Abomey Calavi University (Benin). (Translated from french). 2005; 280.

44. Meliani N, Dib MEA, Allali H, Tabti B. Hypoglycaemic effect of *Berberis vulgaris* L. in normal and streptozotocin-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*. 2011; 1(6):468-471.
45. Ramamurthy V, Schanze KS. *Semiconductor Photochemistry and Photophysics*/Volume ten. 1sted. CRC Press. New-York. 2003; 384.
46. Georgievskii BP, Komissarenko NF, Dmitrou CE. Bioactive compounds in medicinal plants. Ed Naouka. (Translated from french) 1990; 336.
47. Wagner H, Bladt S. *Plant drug analysis: thin layer chromatography Atlas*. 2nd Edition, Springer Science & Business Media. 1996; 384.
48. Ekoumou C. Phytochemical and pharmacological study of 5 traditional recipes used in the treatment of urinary tract infections and cystitis. University of Bamako, (Mali). (Translated from french). 2003; 158.
49. Boubakeur H, Rebbas K, Belhattab R. Antioxidant and antibacterial activity of extracts of *Helichrysum stoechas* (L.) Moench. *Phytothérapie*. (Translated from french). 2018; 16(3):122-132.
50. Lezoul NEH, Belkadi M, Habibi F, Guillén F. Extraction processes with several solvents on total bioactive compounds in different organs of three medicinal plants. *Molecules*. 2020; 25(20):4672.
51. Katalinić V, Možina SS, Skroza D, Generalić I, Abramović H, Miloš M, Ljubenković I, Piskernik S, Pezo I, Terpinč P, Boban M. Polyphenolic profile, antioxidant properties and antimicrobial activity of grape skin extracts of 14 *Vitis vinifera* varieties grown in Dalmatia (Croatia). *Food chemistry*. 2010; 119(2):715-723.
52. Koffi E, Sea T, Dodehe Y, Soro S. Effect of solvent type on extraction of polyphenols from twenty-three Ivorian plants. *Journal of Animal & Plant Sciences*. 2010; 5(3):550-558.
53. Mahmoudi S, Khali M, Mahmoudi N. Study of the extraction of phenolic compounds from different parts of the artichoke flower (*Cynarascolymus* L.). *Nature & Technology*. (Translated from french). 2013; 5(2):35-40.
54. Mulinacci N, Prucher D, Peruzzi M, Romani A, Pinelli P, Giaccherini C, Vincieri FF. Commercial and laboratory extracts from artichoke leaves: estimation of caffeoyl esters and flavonoidic compounds content. *Journal of Pharmaceutical and Biomedical Analysis*. 2004; 34(2):349-357.
55. Benchaachoua A, Bessam HM, Saidi I, Bel-abbes S. Effects of different extraction methods and solvents on the phenolic composition and antioxidant activity of *Silybum marianum* leaves extracts. *International Journal of Medical Science and Clinical Invention*. 2018; 5(3): 3641-3647.
56. Aires A, editor. *Tannins: extraction from plants. Tannins-structural properties, biological properties and current knowledge*. IntechOpen; 2020.
57. Tanoh SK, N'Gaman-Kouassi C, Boa D, Mamyrbekova-Békro JA, Békro YA. Antioxidant activity of hydroethanolic and hydroacetone crude extracts of the organs of four medicinal plants from Côte d'Ivoire. *Nature & Technology*. (Translated from french). 2019; (21):28-34.
58. Bhalodia NR, Nariya PB, Acharya RN, Shukla VJ. In vitro antioxidant activity of hydroalcoholic extract from the fruit pulp of *Cassia fistula* Linn. *AYU (An international quarterly journal of research in Ayurveda)*. 2013; 34(2):209-214.

59. Di Meo F, Lemaury V, Cornil J, Lazzaroni R, Duroux JL, Olivier Y, Trouillas, P. Free radical scavenging by natural polyphenols: Atom versus electron transfer. *The Journal of Physical Chemistry A*. 2013; 117(10), 2082-2092.
60. Mohamed H, Ons M, Yosra ET, Rayda S, Neji G, Moncef N. Chemical composition and antioxidant and radical-scavenging activities of *Periploca laevigata* root bark extracts. *Journal of the Science of Food and Agriculture*. 2009; 89(5):897-905.
61. Irshad M, Zafaryab MD, Singh M, Rizvi M. Comparative analysis of the antioxidant activity of *Cassia fistula* extracts. *International journal of medicinal chemistry*, 2012.

UNDER PEER REVIEW