

EFFECT OF INTERCROPPING WITH LEAFY VEGETABLES ON THE GROWTH, YIELD AND QUALITY OF GLADIOLUS UNDER AGRO-CLIMATIC CONDITIONS OF PRAYAGRAJ

ABSTRACT

Intercropping brings diversity of species in the cropping systems and is considered to make them more resilient against environmental stresses. It can significantly enhance the efficient use of land, water and solar energy besides assuring insurance against failure of one or the other crops. Therefore, an attempt was made to include leafy vegetable crops as intercrops in gladiolus cv. Novalux, for higher productivity and profitability per unit area. The experiment consists of seven treatments with three replications. From the present investigation, it was concluded that the intercrops significantly affected the performance of gladiolus. The intercropping combination T6: Gladiolus + Fenugreek gave the best results in terms of plant height (91.40 cm), number of leaves per plant (18.53), number of shoots per corm (4.93), days to spike initiation (64.33), spike length (73.93 cm), rachis length (49.47 cm), number of florets per spike (12.93), floret size (11.27), vase life (18.53 days), weight of corms (63.67g) and cormlets (6.77g), yield of corms (2385.00kg/ha) and cormlets (225.50kg/ha). The herbage yield per plot was recorded the highest in T3: Fenugreek sole (880.00. g), and intercropping combination of Gladiolus and fenugreek (T6) gave 665.83g. Based on the recent findings, intercropping of fenugreek with gladiolus can be recommended and expanded on a larger scale for additional yield.

Keywords: *Gladiolus grandiflorus*; leafy vegetables; combination crop; interaction effect; yield

INTRODUCTION

Intercropping is growing of two or more crops simultaneously in the same land by utilizing resources such as soil, water, nutrient and solar radiation more effectively [1]. It could be a viable agronomic practice to minimize risk of crop loss, alternate and additional source of income, improve the fertility status of soil, minimize soil erosion and helps to increase the productivity of land per unit area [2]. Intercropping of horticulture and floriculture crops can enhance profit and also ensures better returns. It increases parasitoid survivor-ship, fecundity and retention, and pest suppression in agro-

Comment [Mu1]: Add "the " before it

Comment [Mu2]: on

Comment [Mu3]: nutrients

Comment [Mu4]: sources

ecosystems. According to APEDA, gladiolus has a net production of 174.74 thousand MT in India during 2015-16 as per the data provided by National Horticultural Board [3]. It is regarded as a viable diversification from traditional field crop due to increase per unit return. It is ideal for intercropping and due to its magnificent spikes, its demand as a cut flower in national and international markets is increasing. Whereas, intercropping is a viable agronomic practice for increasing the productivity and profitability from a unit area, vegetable crops have good marketing potential and many types of vegetable crops grown as sole crop are profitable in the region. Therefore, an attempt was made to include the vegetable crops as intercrops in gladiolus for higher productivity and profitability per unit area.

Comment [Mu5]: Add "the " before it

MATERIALS AND METHODS

The experiment was conducted at the Department of Horticulture, Sam Higginbottom University of Agriculture Technology And Sciences, Prayagraj, during September, 2023 to March, 2024. Geographically, Prayagraj is situated at an elevation of 78 meters from mean sea level in the South-Eastern part of Uttar Pradesh. It lies between the parallels of 24° 77' and 25° 47' north latitudes and 81° 19' and 82° 21' east longitudes. The district experiences average maximum temperature ranges between 43°- 47°C which may go as high as 48°C during peak summers (May-June). The minimum average temperature is 2-4°C, which may fall as low as 1°C during peak winter months (December-January) The average rainfall of the district is 960 mm and the monsoon season is spread between July-September. The experiment was laid out in Randomized Block Design with 7 treatments. The treatment combinations include T1: Gladiolus sole, T2: Spinach sole, T3: Fenugreek sole, T4: Dill sole, T5: Gladiolus + Spinach, T6: Gladiolus + Fenugreek, T7: Gladiolus + Dill. The data on growth yield and quality attributing components like plant height (cm), number of leaves per plant, number of shoots per corm, days to spike initiation, spike length (cm), rachis length (cm), number of florets per spike, floret diameter (cm), weight of corms and cormlets (g), yield of corms and cormlets (kg/ha), vase life (days) and herbage yield per plot (g). The data recorded on growth, yield and quality components were subjected to Fisher's method of analysis of variance (ANOVA) as suggested by Panse and Sukhatme[4].

RESULTS AND DISCUSSION

Growth parameters

Significant variation was observed among the intercropping treatments (**Table 1**). The maximum plant height (91.40 cm), number of leaves per plant (18.53) and number of shoots per corm (4.93) were recorded in T6: Gladiolus + Fenugreek. The minimum plant height (81.00 cm), number of leaves per plant (13.83) and number of shoots per corm (2.73) were recorded in T1: Gladiolus sole. Fenugreek is a leguminous crop which fixes nitrogen and also plant stature of fenugreek is smaller compared to other crops grown as intercrop that favours the growth of gladiolus [5, 6]. Similar results were also observed by Singh et al. [7] and Maddukuri [8].

Floral parameters

The flowering parameters such as days to spike initiation, spike length (cm), rachis length (cm), number of florets per spike and floret diameter (cm) also showed significant variation among the different treatments (**Table 1**). The minimum days to initiation of spike (64.33) were observed in T6: Gladiolus + Fenugreek while the maximum days (72.00) were observed in T1: Gladiolus sole. There is translocation of phytohormones to the plants stimulated by nitrogen, which encourages induction of early spike emergence as shown by Marschner et al. [9]. Plants supplied with nitrogen have delayed spike initiation probably due to prolonged vegetative phase in Gladiolus. Complementary findings were reported by Chandana et al. [10] and Lehri et al. [11].

Comment [Mu6]: Add "the " before it

The maximum spike length (73.93 cm) and rachis length (49.47 cm) were recorded in T6: Gladiolus + Fenugreek. The minimum spike length (65.13cm) and rachis length (35.93cm) were recorded in T1: Gladiolus sole. The presence of companion crop, fenugreek, a nitrogen fixing plant might have resulted in increased spike and rachis length due to higher metabolites production along with efficient utilization of nutrients and minerals involved in cell division and cell elongation. Complementary findings were reported by [10,11].

Comment [Mu7]: Hyphen them, please.

Number of florets per spike (12.93) and floret size (11.27) was also recorded as highest in T6: Gladiolus + Fenugreek. Minimum number of florets per spike (10.53) and floret size (9.33) were also recorded in T1: Gladiolus sole. Adequate nitrogen levels can promote healthy vegetative growth and flowering in Gladiolus, potentially resulting in larger florets. Similar finding was observed in gladiolus and marigold paired system which was significantly superior to conventional planting practice in the production of cormels and spike. Intercropping of marigold with gladiolus paired system gave an

Comment [Mu8]: Add "the " before it

additional yield than the pure cropping of gladiolus paired system [12]. Similar result was reported by [7].

Quality parameters

The maximum vase life (days) in normal tap water (**Table 1**) was recorded in T6: Gladiolus + Fenugreek (18.53 days) followed by T7: Gladiolus + Dill (17.67) whereas the minimum vase life was found in T1: Gladiolus sole (13.00). The introduction of fenugreek in association with gladiolus could potentially influence various aspects of gladiolus growth and development, including its vase life. Nitrogen being a vital component of various proteins, therefore supply nitrogen might have resulted in extending the vase life of gladiolus spikes [13,14]. The results are also in conformity with the findings in sandersonia flower [15], in tuberose [16, 17] and in Gladiolus [18,19].

Comment [Mu9]: Add "of " before it

Table 1: Effect of intercropping with leafy vegetables on the growth and quality of gladiolus

Treatment	Plant height (cm)	No. of leaves /plant	No. of shoots /plant	Days to spike initiation	Spike length (cm)	Rachis length (cm)	No. of florets per spike	Floret size (cm)	Vase life(days)
T1: Gladiolus sole	81.00	13.83	2.73	72.00	65.13	35.93	10.53	9.33	13.00
T2: Spinach sole	-	-	-	-	-	-	-	-	-
T3: Fenugreek sole	-	-	-	-	-	-	-	-	-
T4: Dill sole	-	-	-	-	-	-	-	-	-
T5: Gladiolus + Spinach	85.80	15.20	3.67	69.73	67.47	38.47	10.87	10.27	15.60
T6: Gladiolus + Fenugreek	91.40	18.53	4.93	64.33	73.93	49.47	12.93	11.27	18.53
T7: Gladiolus+ Dill	88.80	17.50	4.20	67.40	70.40	41.67	11.87	10.40	17.67
C.d.	3.84	0.42	0.26	2.27	2.20	1.26	0.60	0.35	0.30
Se(m)	1.23	0.14	0.08	0.73	0.71	0.40	0.19	0.11	0.10
Se(d)	1.74	0.19	0.12	1.03	0.99	0.57	0.27	0.16	0.14
C.v.	4.30	2.51	6.47	3.22	3.09	2.95	5.07	3.34	1.79

Corm parameters

Comparison between different treatments revealed that the maximum weight of corms (63.67g) and cormlets (6.77g) were recorded in T6: Gladiolus + Fenugreek while the minimum weight of corms (41.33g) and cormlets (4.18g) were recorded in T1: Gladiolus sole (**Table 2**) and **Fig 1**. Yield of corms (2385.00kg/ha) and cormlets (225.50kg/ha) were also recorded to be highest in T6: Gladiolus + Fenugreek while the lowest corm and cormlet yield (1530.00kg/ha and 152.50kg/ha respectively)

were recorded in T1: Gladiolus sole. Intercropping with fenugreek gives positive results due to phytohormones and secondary metabolites which might have indirect effects on corm formation and yield. The probable cause for enhanced weight with the optimization of nitrogen could be adequate availability of nutrients and minerals, resulting in increase of transfer and accumulation of assimilates and carbohydrates in the daughter corms. The findings are in proximity with the corm characteristics influenced by nitrogen [20,21,22]. Similar finding was recorded in Colocasia grown with Fenugreek in 1:1 ratio with the highest cormlet weight per [23]. The yield of corms and cormlets per square meter with closer spacing was also recorded to be more [24].

Comment [Mu10]: increasing

Comment [Mu11]: Add "the " before it

Table 2: Effect of intercropping with leafy vegetables on the corm yield of gladiolus

Treatment	Weight of corms (g)	Weight of cormlets(g)	Yield of corms (kg/ha)	Yield of cormlets (kg/ha)
T1: Gladiolus sole	41.33	4.18	1,530.00	152.50
T2: Spinach sole	-	-	-	-
T3: Fenugreek sole	-	-	-	-
T4: Dill sole	-	-	-	-
T5: Gladiolus+ Spinach	45.13	5.05	1,692.50	170.08
T6: Gladiolus+ Fenugreek	63.67	6.77	2,385.00	225.50
T7: Gladiolus+ Dill	53.73	5.96	1,972.50	210.25
C.d.	0.98	0.07	48.19	19.40
Se(m)	0.32	0.02	15.47	6.23
Se(d)	0.45	0.03	21.88	8.81
C.v.	1.87	1.18	2.47	9.96




Fig 1: Effect of intercropping on the corm yield of gladiolus

Herbage yield per plot (g)

The intercropping combination T6: Gladiolus+ Fenugreek reported higher yield of 665.83g, while T4: Gladiolus + Spinach (480.00g) gave the minimum herbage yield per plot (Table 3) and Fig 2.. This was due to the synergistic effect between main crop and intercrops whereas, minimum herbage yield was observed with dill and spinach intercropping system due to antagonistic effect between the crops and highly exhaustive nature of intercrops. The highest yield might be due to the nutrients present on the leafy vegetable which helps to increase the availability of additional nutrients to the main crop and also residual effect of nutrient in intercropping system as compared to sole crop of gladiolus. Similar results have been obtained in coleus + mulberry intercropping system [25] and in gladiolus grown alongside marigold [26]. Also, it was revealed that planting of *Coleus forskohlii* and *Phyllanthusamarus* registered higher tuber yield [27]. Similar findings were also reported in [7,8].

Table 3: Effect of intercropping on the herbage yield of spinach, fenugreek and dill

Treatments	Herbage yield per plot (g)		
	Spinach	Fenugreek	Dill
T1: Gladiolus sole	-	-	-
T2: Spinach sole	746.00	-	-
T3: Fenugreek sole	-	880.00	-
T4: Dill sole	-	-	787.00
T5: Gladiolus+ Spinach	480.00	-	-
T6: Gladiolus+ Fenugreek	-	665.83	-
T7: Gladiolus+ Dill	-	-	506.00
C.d.	1.00	1.00	0.88
Se(m)	0.32	0.32	0.30
Se(d)	0.45	0.45	0.39
C.v.	0.32	0.25	0.26




Fig 2: Effect of intercropping on the herbage yield of spinach, fenugreek and dill

Conclusion

From the present investigation, it was observed that sole cropping treatments spinach, fenugreek and dill gave better herbage yield (g). However, it was observed that intercropping combinations *i.e.* T5 (Gladiolus+ Spinach), T6 (Gladiolus+ Fenugreek) and T7 (Gladiolus+ Dill) recorded better performance in the growth, yield and quality of gladiolus in which treatment T6 (Gladiolus+ Fenugreek) gave the best result. Based on the above findings, intercropping of leafy vegetables with gladiolus has found to increase the productivity per unit area. Fenugreek performs well as an intercrop.

Comment [Mu12]: Add "of " before it

Comment [Mu13]: Add "been " before it

Reference

1. Rana MM, Bhuiya MSU, Akhand MIM. Effect of inter cropping of *Sesbania rostrata* with transplant aman rice on yield and profitability. *Int. J. Agriculture and Veterinary Sci.* 2013; 2(1): 10-14.
2. Faroda AS, NL Jishi, Singh R, Saxena A. Resource management for sustainable crop production in arid zone-A review. *Indian Agronomy* 2007; 52(3): 181-193.
3. APEDA. India production of Gladiolus (apeda.gov.in). 2016; Retrieved on 10th July, 2021.
4. Panse VG, Sukhatme, PV. Stastical methods for agricultural workers. Indian Council of Agricultural Research, New Delhi. 1969.
5. Baker EFI. Research on mixed cropping with cereals in Nigeria farming systems: A system for improvement. Proceedings of International Workshop on Farming System. ICRISAT, Hyderabad. 1975; pp. 287-300.
6. Anitha S, Geethakumari VL, Filial GR. Effect of intercrops on nutrient uptake and productivity of chilli-based cropping system. *J. Trop. Agri.* 2001; 39: 60-61.

7. Singh KMP and Singh D. Performance of Coriander, Fenugreek and Soya as Intercrop under Gladiolus based Intercropping system. *Journal of AgriSearch*. 2014; 1(4): 246-250.
8. Maddukuri LD. Development of site-specific integrated nutrient management systems for gladiolus (*Gladiolus hybridus* Hort.) and marigold (*Tagetes patula* L.) using soil test crop response correlation studies. Ph.D. Thesis, ICAR-Indian Agricultural Research Institute, New Delhi. 2017; pp. 58-132.
9. Marschner H, Romheld V. In Vivo Measurement of Root-Induced pH Changes at the Soil-Root Interface: Effect of Plant Species and Nitrogen Source. *Zeitschrift für Pflanzenphysiologie*. 1983;111,249-251.
10. Chandana K, Dorajeerao, AVD. Effect of graded levels of nitrogen and phosphorus on growth and yield of gladiolus (*Gladiolus grandiflorus* L.) cv. White Prosperity in coastal AP, India. *Plant. Arch.* 14(1): 143-150. 2014.
11. Lehri SN, Kurd AA, Rind MA, Bangulzai NA. The response of *Gladiolus tristis* L. to N and P₂O₅ fertilizers. *Sarhad J. Agric.* 2011; 27(2): 185-188.
12. Singh S, Datta SK. Intercropping of French marigold (*Tagetes patula* Linn.) in gladiolus. *Journal of Ornamental Horticulture*, 2006; 9(1): 37-39.
13. Chouhan P, Vidhya Sankar M, Rathor V. Effect of NPK on physicochemical parameters of Gladiolus (*Gladiolus hybridus* Hort.) cv. White Prosperity. *Int. J. Sci. Res. Publ.* 2014; 4(12): 1-5.
14. Kumar K, Singh CN, Beniwal VS, Pinder R, Poonia, RS. Effect of nitrogen fertilizer on different attributes of gladiolus (*Gladiolus grandiflorous*L.) cv. American Beauty. *Int. J. Environ. Agric. Biotech.* 2017; 2(1): 1-3.
15. Clark GE, GK Burge. Effects of planting density, stem pruning treatments, and shade on secondary tuber development in *Sandersonia aurantiaca*. *N. Z. J. Crop Hortic. Sci.* 1997; 25: 73-78..
16. Kumar R, RL Mishra, SK Singh. Post-harvest life of gladiolus cv. Jester Gold as influenced by different doses of nitrogen, phosphorus and potassium. *Indian J. Hortic.* 2008; 65(1): 83-86.
17. Khalaj, M.A., B. Edrisi, and M. Amiri, Effect of nitrogen and plant spacing on nutrient uptake, yield and growth of tuberose (*Polianthes tuberosa* L.). *J. Ornament. Hortic. Plant.* 2012; 2(1): 45-54.
18. Prakash VR, Singh JP, Shankar, VS, Vijai, K. Effect of foliar application of urea and planting geometry on growth and flowering of gladiolus cv. Sapna. *Anna. Hortic.* 2012; 5(2): 192-197.
19. Regar AL, Thumar BV, Mahawer LN, Chawla SL, Meena NK. Effect of spacing and nitrogen on floral and vase life parameter of gladiolus (*Gladiolus grandiflorus* L.) cv. American Beauty. *The BioScan*. 2016; 11(1): 539-542.

20. Baral SP, Baral DR, Gautam DM, Pun UK. Influence of nitrogen on growth performance, cut flower characteristics and corm/cormel production of Gladiolus. *Nepal J. Sci. Tech.* 2012; 13(1): 25-31.
21. Mahadik MK, Chopde N. Influence of nitrogen and potassium on growth and yield of gladiolus corms. *Plant Arch.* 2015; 15(1): 193-196.
22. Singh G, Kumar A, Kant L, Kumar R, Kumar A. Effect of GA3 and urea on growth, and yield of gladiolus (*Gladiolus grandiflorus* L.) cv. Novalux. *International Journal of Chemical Studies*, 2019; 7(5): 3371-3373.
23. Shankar D, Thakur AK, Kanwar RR, Singh DP. Effect of Growing Different Leafy Vegetables as Intercrop on the Yield of Summer Colocasia in Bastar Plateau of Chhattisgarh. *Journal of Root Crops.* 2016; Vol. 42 No. 2, pp. 81-85.
24. Mukhopadhyay TP, Yadav LP. Effect of corm size and spacing on Growth, flowering and corm production in gladiolus. *Haryana J. Hort. Sci.* 1984; 13(3-4): 95-99.
25. Srinivas, NN, Patil GM, Suresh A. Growth and yield of coleus intercropped in Victory-1 mulberry under integrated nutrient management. *Karnataka J. Agric. Sci.* 2008; 21(4): 524-526.
26. Bora, K. Performance of Gladiolus (*Gladiolus grandiflorus* L.) at Different Levels of Fertilizers and Spacing using Marigold (*Tagetes Patula*) as an Inter-Crop. Ph.D Thesis, Assam Agriculture University, Jorhat. pp. 94-95. 2022.
27. Vennila and Jayanthi. Evaluation of productivity indices of *Coleus forskohlii* and *Phyllanthu samarus* intercropping system. *Agric. for Sustainable Development.* 2014; 2 (2): 158-161.