

# Effect of different level of Chitosan and beeswax edible coating on storage life and physio-chemical properties of strawberry (Fragaria × ananassa) cv winter dawn

## ABSTRACT

The present investigation was done to understand the effect of different edible coating on increasing the shelf-life of strawberry at room temperature. Postharvest losses in fruits are a serious problem due to rapid deterioration during storage. Use of edible coatings over fruits is used to improve their quality and enhance the shelf life. The main purpose of this study was to assess the effect of Chitosan and beeswax coating in extending the shelf-life of strawberry including physio -chemical changes at room temperature. The experiment was laid out in completely randomized design (CRD) with 10 treatments and 3 replications. The treatment consisted of coating material with Chitosan (1.5g%), Chitosan +TPP (1.5g+0.5g%), Chitosan +TPP (1.5g+1%), beeswax (5g), beeswax + TPP (5G+1.5%), Chitosan+ beeswax (0.75g+2.5g), Chitosan+ beeswax +TPP(0.75g+2.5g+0.5%), Chitosan+ beeswax +TPP (0.75g+2.5g+1%). Statistical analysis revealed that the treatment chitosan 1.5g+ TPP 1% had the minimum Weight loss 0.08g, while having the maximum amount of Vitamin C 38.45mg/100g, Hardness 8.07 kg/inch<sup>2</sup>, Bulk Density 1.9g/cm<sup>3</sup> and TSS 5.13°Bx. However, Chitosan 1.5g+TPP 0.5% showed maximum Reducing sugar 3.73% and non-reducing sugar 0.92%. While chitosan 0.75g+ beeswax 2.5g+TPP 0.5g have the maximum Total sugar 5.23%.

**Keywords:** Strawberry, Chitosan, Beeswax, Physio-chemical characteristics, Shelf-life

## 1.INTRODUCTION

“Strawberry is one of the most economically important fruits consumed due to its well nutritional value, aroma, attractive and juicy fruits. They are rich in polyphenols and anthocyanins, vitamins and amino acids. The main characteristics related to the quality of the ripe strawberries are their texture, flavour (organic acids and soluble sugars content) and colour”(Campaniello, *et al.*, 2020). “Due to their very active metabolism strawberries are highly perishable and have high physiological post-harvest activities which lead to short ripening and senescence periods that make their marketing a challenge” (Martino, *et al.*, 1998). “Loss of quality in this fruit is connected with its sensitivity to fungal infection and susceptibility to water loss, bruising, mechanical injuries and texture softening due to the lack of protective rind” (Atress, *et al.*, 2010).

Edible films and coatings are promising systems for the improvement of food quality, shelf life, safety, and functionality. The efficiency and functional properties of edible film and coating materials are highly dependent on the inherent characteristics of film-forming

materials, namely biopolymers (such as proteins, carbohydrates, and lipids), plasticizers, and other additives.

“Development of edible coatings has been based upon the use of polysaccharides, proteins, lipids or their combination in different ways. Polysaccharides and proteins are known to form films with good mechanical properties, but with poor permeability, while the lipids form brittle films but with improved permeability. Therefore, the newly developed coatings are focused on combining the polymer matrix with some hydrophobic component” (Navarro-Tarazaga, Sothornvit, & Perez-Gago, 2008; Zúñiga et al., 2012).

Chitosan coatings are gaining significant attention in post-harvest preservation due to their antimicrobial, and antioxidant properties as well as other desirable characteristics including, moisture barrier, gas permeability, biodegradability, film-forming ability, enhanced nutritional value, compatibility with other treatments and compliance with food safety standards. Overall, chitosan coatings have demonstrated promising results in extending the shelf-life of fruits and vegetables while maintaining their quality, making them a significant development in post-harvest preservation that can contribute to both economic and environmental sustainability in the agricultural industry.

Beeswax in biopolymer-based films enhances their barrier and tensile features. The coating is capable of tuning the release of bioactive from the polymer matrices. Beeswax coatings have great potential for reducing food waste, which is a major issue for food security and sustainability. Beeswax is a natural glazing agent that can be used in food to prevent water loss and provide protection during storage. It is often used to prevent water loss and retard shrinkage and spoilage in fruit.

## **2. MATERIAL AND METHOD**

### **2.1 Study location and Treatment combination**

The experiment was conducted at Post Harvest Laboratory, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, and Prayagraj (UP). The experimental design was CRD (Completely Randomized Design) with 10 treatments and 3 replications. The strawberries were procured from Mundera market and then they were washed and kept it on the same day at room temperature. Uniform and good quality strawberry were picked. The coated fruits were kept at room temperature for 9 days and observation on Physiological loss in weight, total soluble solid, vitamin C/ascorbic acid, Total sugar, reducing sugar, non-reducing sugar, bulk density and hardness of the fruit was taken at 3 days interval. The fruits were divided into equal lots. The fruits were washed with water and allowed to dry under fan and then treated with Chitosan (1.5g%), Chitosan +TPP (1.5g+0.5g%), Chitosan +TPP (1.5g+1%), beeswax (5g), beeswax + TPP (5G+1.5%), Chitosan+ beeswax (0.75g+ 2.5g), Chitosan+ beeswax +TPP(0.75g+2.5g+0.5%), Chitosan+ beeswax +TPP (0.75g+2.5g+1%).

## **2.2 PREPARATION AND APPLICATION OF EDIBLE COATING**

Chitosan solution was prepared by dissolving 1.5g, 0.75g chitosan in 100ml of acetic acid with continuous stirring at room temperature. After the complete dissolution of the chitosan, 2g of glycerol and 2g of tween 80 was added and stirred the solution for 30min. The result solution was used for coating. For preparing beeswax 5g of beeswax in 50ml of ethanol heated to 70°C was melted and mixed with 50ml distilled water and the wax solution was emulsified with 25% of tween 80 with respect to the wax content.

The cross -linking of the chitosan coating was carried out with 0.5g, 1g/100g sodium tripolyphosphate, TPP, prepared as aqueous solutions by dissolving the salt in distilled water under vigorous stirring. The cross-linking lasted few minutes and then the excess of the cross-linking solution was bottled with adsorbent paper. Preparation of chitosan and beeswax was done by using 100ml of 0.75g/100g chitosan solution mixed with 5g of wax. The chitosan coating solution was prepared, as described previously then heated to 70°C. The wax was melted in the heated solutions and the whole mixture was homogenized.

The treated strawberry was tested for their physical and quality parameter such as, physiological weight loss, bulk density, hardness, TSS, vitamin C, total sugar, reducing sugar and nonreducing sugar.

## **3. RESULT AND DISCUSSION**

The weight of the fruit was measure at 3 days interval during the conduction of the experiment. This help in the determination of weight loss during the experiment

### **Weight loss (g) of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

Strawberry fruits are highly susceptible to rapid loss of water which results in fruit shrinkage and weakening of the tissue due to their very thin skin. Therefore, the weight loss was followed over the storage period of 9 days to evaluate the effects of the coatings. T<sub>3</sub> chitosan 1.5g +STPP 1% show the lowest amount of weight loss (1.07, 1.20, 0.80). The chitosan + STPP emulsion served as a physical barrier around the fruit which partially closed the stomatal openings and lenticels thereby reducing the rates of transpiration and respiration **Jain et al. (2011).**

### **Hardness (kg/nch<sup>2</sup>) of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

“Strawberries soften considerably during ripening due to degradation of the middle lamella of the cell wall of cortical parenchyma cells”(Perkins-Veazie, 1995). The coating helps in slowing down the softness and prolong the shelf life. Among the different treatments, T<sub>3</sub>

(9.73,8.80,8.63,8.07) retain the maximum hardness after storage closely followed by T2 (9.1, 8.30,7.34,6.78). “The controlled fruits registered the lowest texture (5.13, 3.88, 3.23 and 2.58) at all the days. The decrease in fruit texture with the storage period might be due to breakdown of insoluble pectin to soluble form and also due to cellular disintegration leading to permeability of the cell membrane which ultimately helps in gaseous exchange” (Mahajan *et al.* 2013). “The higher texture in coconut oil and paraffin wax coated fruits might be due to reduced transpiration and respiration along with delayed ethylene production and thus retained more turgidity of the cells of the fruits” (Nanda *et al.* 2001).

### **Bulk density ( $\text{g/cm}^3$ ) of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

Bulk density is a very complex product property and is of great importance for economical and functional reasons. High bulk density is desirable as it would indicate ample abundance of the edible fruit content. By checking the various bulk density data, we were able to identify that the bulk density of the coated strawberry shows significant differences on 3<sup>rd</sup> day, 6<sup>th</sup> day, 9<sup>th</sup> days. The maximum bulk density was found in T3 chitosan 1.5g +STPP 0.5g (3.30, 2.95, 2.63, 1.9).

### **TSS ( $^{\circ}\text{Brix}$ ) of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

It was observed that TSS of the fruits increased from 3<sup>rd</sup> day to 9<sup>th</sup> day of the treatment. The maximum TSS (6.37) at 9 days of storage was recorded in chitosan 1.5g +STPP 1% coating followed by beeswax 5g (5.37) while, the least TSS was recorded in control (4.3). The gradual increase in TSS up to 9<sup>th</sup> day might be due to complete hydrolysis into sugars and the gradual decline after this period is due to no further increase in sugar since it is the primary substrate for respiration (Wills *et al.* 1980). The increase in TSS in wax coated fruits was till the 9<sup>th</sup> day. The decrease of TSS might be due to the fact that wax has the capacity to delay the metabolic activities during ripening and storage of fruits (Fan *et al.* 1999)

**TABLE:1 Physiological weight loss and Hardness (kg/inch<sup>2</sup>) of the strawberry as**

Sl. No.	Treatment combination	No. of days at room temperature											
Sl. No.	Treatment combination	No. of days at room temperature											
		Bulk density (g/cm <sup>3</sup> )					TSS (°Brix)						
		PLW (g)					Hardness (kg/inch <sup>2</sup> )						
		Initial day	3day	6day	9day	Initial day	3day	6day	9day	Initial day	3day	6day	9day
1	T <sub>0</sub> (control)	1.10	1.37	1.97	5.13	3.88	3.27	2.8					
2	T <sub>1</sub> (chitosan)	1.07	1.40	1.97	9.33	8.27	7.73	7.03					
3	T <sub>2</sub> (chitosan + STPP)	1.33	1.40	1.13	9.10	8.30	7.70	7.13					
4	T <sub>3</sub> (chitosan+ STPP)	1.07	1.20	0.80	9.73	8.80	8.63	8.07					
5	T <sub>4</sub> (Beeswax)	1.33	1.60	1.60	9.10	9.3	8.10	6.87					
6	T <sub>5</sub> (beeswax+ SSTPP)	0.93	2.30	1.67	9.53	9	8.40	6.77					
7	T <sub>6</sub> (beeswax+ STPP)	1.27	1.13	1.07	8.43	7.67	7.43	6.33					
8	T <sub>7</sub> (Chitosan+ beeswax)	1.17	1.33	1.40	9.23	7.9	7.67	6.63					
9	T <sub>8</sub> (chitosan+ beeswax+ STPP)	1.13	1.20	1.50	8.2	7.57	7.00	6.20					
10	T <sub>9</sub> (chitosan+ beeswax+ STPP)	1.07	1.27	1.47	9.27	7.67	7.04	6.13					
F test		NS	NS	NS	S	S	S	S					
S.E (d) (±)		0.39	0.40	0.45	0.20	0.20	0.29	0.23					
C.V		41.35	34.42	38.04	2.69	4.29	4.66	4.22					

affected by the different levels of Chitosan and Beeswax solution

1	T <sub>0</sub> (control)	1.97	1.7	1.17	0.77	3.70	4.13	4.2	4.3
2	T <sub>1</sub> (chitosan)	2.13	1.72	1.20	0.91	3.40	3.93	4.5	5.23
3	T <sub>2</sub> (chitosan + STPP)	2.43	2.02	1.7	0.97	5	5.4	5.97	6.37
4	T <sub>3</sub> (chitosan+ STPP)	3.30	2.93	2.63	1.9	3.27	3.83	4.59	5.13
5	T <sub>4</sub> (Beeswax)	2.03	1.78	1.3	0.92	3.47	4	4.67	5.37
6	T <sub>5</sub> (beeswax+ SSTPP)	2.77	2.83	2.38	1.67	3.20	3.77	4.83	5.33
7	T <sub>6</sub> (beeswax+ STPP)	1.95	1.46	1.2	0.95	3.47	4	4.87	4.7
8	T <sub>7</sub> (Chitosan+ beeswax)	2.1	1.79	1.41	1.17	2.97	3.43	4.13	4.7
9	T <sub>8</sub> (chitosan+ beeswax+ STPP)	2.20	1.83	1.34	0.95	4.40	4.7	5.17	5.63
10	T <sub>9</sub> (chitosan+ beeswax+ STPP)	2.01	1.6	1.37	0.96	3.37	3.8	4.47	5.07
F test		S	S	S	S	S	S	S	S
S.E (d) (±)		0.26	0.18	0.19	0.13	0.27	0.28	0.30	0.28
C.V		13.89	11.70	15.18	13.96	9.29	8.31	7.83	6.46

**TABLE:2 Bulk density and TSS of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

### **Total Sugar (%) of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

There was significant on the 3day -6days on the different coatings with respect to the total sugar content of the strawberry fruits. The total sugars of the fruits increased up to 9th day of treatment. Among all the treatments, T7 chitosan0,75g+ beeswax 2.5g coating scored the maximum value with respect to total sugars (5.7%), followed by T5 chitosan+ STPP(5.37%). The increase in sugars might be due to the rapid conservation of polysaccharides into sugars.



1	T <sub>0</sub> (control)	4	4.43	4.8	5	2.57	3.00	3.20	3.2
2	T <sub>1</sub> (chitosan)	4.17	4.8	5	5.2	2.67	2.86	3.20	3.533
3	T <sub>2</sub> (chitosan + STPP)	3.78	4.76	4.77	5.1	2.57	2.99	3.33	3.73
4	T <sub>3</sub> (chitosan+ STPP)	4.02	4.82	4.99	5.2	2.63	2.72	3.27	3.6
5	T <sub>4</sub> (Beeswax)	3.66	4.57	5.01	5.17	2.27	2.72	3.14	3.7
6	T <sub>5</sub> (beeswax+ SSTPP)	3.75	4.53	5.1	5.37	2.34	2.78	3.03	3.63
7	T <sub>6</sub> (beeswax+ STPP)	3.75	4.82	5	5.33	2.30	2.88	3.20	3.63
8	T <sub>7</sub> (Chitosan+ beeswax)	3.96	4.95	5.23	5.7	2.44	2.71	3.06	3.6
9	T <sub>8</sub> (chitosan+ beeswax+ STPP)	3.84	4.39	4.96	5.23	2.27	2.76	3.20	3.67
10	T <sub>9</sub> (chitosan+ beeswax+ STPP)	3.58	4.62	4.93	5.05	2.27	3.23	3.47	3.6
F test		NS	S	S	NS	S	S	NS	NS
S.E (d) (±)		0.22	0.15	0.12	0.13	0.13	0.12	0.17	0.15
C.V		7.10	4.05	3.01	3.09	6.27	5.21	6.30	5

**TABLE:3 Total Sugar and Reducing sugar of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

Sl.	Treatment combination	No. of days at room temperature
-----	-----------------------	---------------------------------

No.		Non reducing sugar%				Vitamin C (mg/ 100g fruit)			
		Initial day	3day	6day	9day	Initial day	3day	6day	9day
1	T <sub>0</sub> (control)	1.17	1	0.8	0.81	41.00	40	39.67	38.07
2	T <sub>1</sub> (chitosan)	1.5	1.17	1	0.85	42.02	44.67	45.63	39.43
3	T <sub>2</sub> (chitosan + STPP)	1.4	1.24	1.1	0.92	41.90	44.67	45.13	39.20
4	T <sub>3</sub> (chitosan+ STPP)	1.53	1.27	1.07	0.83	42.14	43.53	44.96	40
5	T <sub>4</sub> (Beeswax)	1.6	1.4	1.17	0.85	41.00	40.67	45.47	38.43
6	T <sub>5</sub> (beeswax+ SSTPP)	1.5	1.23	1.06	0.73	41.73	43.87	45.17	39.13
7	T <sub>6</sub> (beeswax+ STPP)	1.57	1.267	1.03	0.89	41.57	42.77	43.43	38.93
8	T <sub>7</sub> (Chitosan+ beeswax)	1.57	1.3	1.11	0.89	41.67	41.10	43.56	39
9	T <sub>8</sub> (chitosan+ beeswax+ STPP)	1.37	1.1	0.9	0.63	41.70	42.87	45.73	38.63
10	T <sub>9</sub> (chitosan+ beeswax+ STPP)	1.53	1.4	1.13	0.85	41.07	41.60	43.00	38.97
F test		S	NS	NS	NS	NS	S	S	NS
S.E (d) (±)		0.12	0.12	0.13	0.11	0.77	0.88	1.39	0.69
C.V		9.62	12.26	14.87	16.72	2.27	3.57	3.78	2.17

**TABLE:4 Non reducing sugar and Vitamin C of the strawberry as affected by the different levels of Chitosan and Beeswax solution**

**CONCLUSION:**

The shelf life of the strawberry which was coated with chitosan-based coatings, beeswax, chitosan and beeswax prolonged the storage period of strawberries for nine days at room temperature and slowed down their senescence process compared to uncoated strawberries.

From the present investigation, it was observed that Treatment (T<sub>3</sub>, chitosan 1.5g+ TPP 1%) had the minimum Weight loss of 0.08g, while having the maximum amount of Vitamin C 38.45mg/100g, Hardness 8.07 kg/inch<sup>2</sup>, Bulk Density 1.9g/cm<sup>3</sup> and TSS 5.13°Bx. However, T<sub>2</sub>(Chitosan 1.5g+TPP 0.5%) showed maximum Reducing sugar at 3.73% and non-reducing sugar at 0.92%. While T<sub>7</sub> chitosan (0.75g+ beeswax 2.5g+TPP 0.5g) has the maximum Total sugar of 5.23%. The highest benefit-cost ratio was found in T<sub>2</sub>(Chitosan 1.5g+TPP 0.5%) at 3.80.

## **REFERENCES**

**Aparicio-García, P. F., Ventura-Aguilar, R. I., del Río-García, J. C., Hernández-López, M., Guillén-Sánchez, D., Salazar-Piña, D. A. & Bautista-Baños, S. (2021).** Edible chitosan/propolis coatings and their effect on ripening, development of *Aspergillus flavus*, and sensory quality in fig fruit, during controlled storage. *Plants*, 10(1), 112.

**Alves, V., Costa, N., & Coelho, I. (2010).** Barrier properties of biodegradable composite films based on kappa-carrageenan/pectin blends and mica flakes. *Carbohydrate Polymers*, 79, 269e276.

**Ahmed, W. A. S. S. E. M, Azmat, R., Qayyum, A, Mehmood, A., Masaud, K., Liaquat, M., & Moin, S. S. (2018).** The role of chitosan to prolonged the fresh fruit quality during storage of grapefruit cv. ray ruby. *Pak. J. Bot*, 50(1), 151-159.

**Arnon, H., Zaitsev, Y., Porat, R., & Poverenov, E. (2014).** Effects of carboxymethyl cellulose and chitosan bilayer edible coating on postharvest quality of citrus fruit. *Postharvest Biology and Technology*, 87, 21-26.

**Amin, Usman, (2021):** "Improvement of the performance of chitosan—aloe vera coatings by adding beeswax on postharvest quality of mango fruit." *Foods* 10.10 2240.

**Baswal, A. K., H. S. Dhaliwal, Zora Singh, B. V. C. Mahajan, Anu Kalia, and K. S. Gill.** "Influence of carboxy methylcellulose, chitosan and beeswax coatings on cold storage life and quality of Kinnow mandarin fruit." *Scientia Horticulturae* 260 (2020): 108887.

**Bibi F., Baloch M.K. 2014** "Postharvest quality and shelf life of mango (*Mangifera indica* L.) fruit as affected by various coatings". *J. Food Process. Preserv.* ;38(1):499–550.

**Brishti FH, Misir J, Sarker A. (2013).** Effect of bio preservatives on storage life of papaya (*Carica papaya* L.). *Int J Food Stud.* 2(1).

**Campaniello, D., Bevilacqua, A., Sinigaglia, M., & Corbo, M. R. (2008).** Chitosan: antimicrobial activity and potential applications for preserving minimally processed strawberries. *Food Microbiology*, 25, 992e1000.

**Efendi, Darda, and Heliyana Hermawati.(2010):** "The use of bee wax, chitosan and BAP to prolong shelflife of Mangosteen (*Garcinia mangostana* L.) fruit." *Jurnal Hortikultura Indonesia* 1, no. 1 32-39.

**Eshetu, A., Ibrahim, A. M., Forsido, S. F., & Kuyu, C. G. (2019).** Effect of beeswax and chitosan treatments on quality and shelf life of selected mango (*Mangifera indica* L.) cultivars. *Heliyon*, 5(1).

**Eshun, K. and He, Q. (2004)** Aloe vera: A valuable ingredient for the food, pharmaceutical and cosmetic industries- A review. *Critical reviews in food science and nutrition*, 44(2), 91-93

**Esua, MF and Rauwald, J-W (2006)** Noval bioactive maloyl glucans from aloe vera gel: isolation, structure elucidation and in vitro bioassays. *Carbohydr Res* 341, pp. 355-364

**Fan X, Blankenship S M and Mattheis J P. 1999.** 1-MCP inhibits apple ripening. *Journal of American Society for Horticultural Science* 124: 690–5.

**Foo, S. Y., Nur Hanani, Z. A., Rozzamri, A., Ibadullah, W. W., & Ismail-Fitry, M. R. (2019).** Effect of chitosan–beeswax edible coatings on the shelf-life of sapodilla (*Achras zapota*) fruit. *Journal of Packaging Technology and Research*, 3, 27-34.

**Gol N B and Rao T V R. 2011.** Banana fruit ripening as influenced by edible coatings. *International Journal of Fruit Science* 11: 119–35.

**Hazarik, Lalthanpuui, D Mandal. (2017).** Influence of edible coatings on physico-chemical characteristics and shelf-life of papaya (*Carica papaya*) fruits during ambient storage. *Indian Journal of Agricultural Sciences* 87 (8): 1077–83.

**Ikram EHK, Stanley R, Netzel M, Fanning K, (2005)** Phytochemicals of papaya and its traditional health and culinary uses: A review. *J Food Compos Anal.* 41:201-211.

**Jain, Shashi & Mukherjee, S. (2011).** Enhancing keeping quality of fruits in mango cv. Langra. *Indian Journal of Horticulture.* 68. 142-144.

**Ladaniya M S. 2001.** Response of 'Mosambi' sweet orange (*Citrus sinensis*) to degreening, mechanical waxing, packaging and ambient storage conditions. *Indian Journal of Agricultural*

**Peñarubia, Omar Riego, (2014):** "Beeswax–and candelilla wax–coconut oil edible coatings extend the shelf life of strawberry fruit at refrigeration temperatures." *International Journal of Postharvest Technology and Innovation* 4.2-4 221-234.

**Purwoko, B.S., and Fitriadesi, P. (2000).** Effect of Coating Materials and Storage Temperatures on Quality and Shelf Life of Papaya Fruit. *Buletin Agronomi* 28, 66-72.

**Raju M, Mondal R, Valliath AS, Tejaswi S, Das P. (2023)** Enhancement of quality parameters and shelf-life of papaya fruit (*Carica papaya* L.) by edible coating during storage and transportation. *Plant Science Today.* 10(2): 114–119.

**Ramachandra CT, Rao PS. (2008).** Processing of aloe Vera leaf gel: A review. *Am J Agric Biol Sci.* 3(2):502–510.

**Ramos-García, Margarita,** “Use of chitosan-based edible coatings in combination with other natural compounds, to control *Rhizopus stolonifer* and *Escherichia coli* DH5 $\alpha$  in fresh tomatoes." *Crop Protection* 38 (2012): 1-6.

**Shahnawaz M and Sheikh S A. 2011.** Physicochemical characteristics of jamun fruit. *Journal of Horticulture and Forestry* 3: 301–6.

**Sharmin MR, Islam MN, Alim MA. (2016).** Shelf-life enhancement of papaya with aloe vera gel coating at ambient temperature. *J Bangladesh Agric Univ.* 13(1):131–136.

**Sheoran (2004)** Statistical package for agricultural research workers, CCS, HAU, Hisar

**Sindhu G S, Dhillon W S and Mahajan B V C. 2009.** Effect of waxing and packaging on storage of pear cv. Punjab Beauty. *Indian Journal of Horticulture* 66: 239–44.

**Singh A, Yadav D S, Patel R K, Nath A and Bhuyan M. 2011.** Wax coating and padding materials influence quality and shelf-life of purple passion fruit during storage. *Indian Journal of Horticulture***68**: 246–9.

**Suwanti, R Utami, Kawiji, D Praseptiangga and Khasanah. (2018).**The effect of packaging methods (paper, active paper, and edible coating) on the characteristic of papaya MJ9 in ambient temperature storage. *IOP Conf. Ser.: Earth Environ. Sci.***102** 012087

**Valverde JM, Valero D, Martínez-Romero D, Guillén F, Castillo S, Serrano M. (2005).**Novel edible coating based on aloe vera gel to maintain table grape quality and safety.*J Agric Food Chem.*53(20):7807–7813.

**Velickova, Elena, (2013):** "Impact of chitosan-beeswax edible coatings on the quality of fresh strawberries (*Fragaria ananassa* cv Camarosa) under commercial storage conditions." *LWT-Food Science and Technology* 52.2 80-92.

**Velickova, Elena, (2015)**"Characterization of multilayered and composite edible films from chitosan and beeswax." *Food Science and Technology International* 21.2 (): 83-93.

**Ventura-Aguilar, R. I. (2018):** "Impact of chitosan based edible coatings functionalized with natural compounds on *Colletotrichum fragariae* development and the quality of strawberries." *Food chemistry* 262 142-149.

**Wills R B H, Bembridge P A and Scott K J. 1980.** Use of flesh firmness and other objective tests to determine consumer acceptability of delicious apples. *Australian Journal of Experimental Agriculture and Animal Husbandry***20**: 252–6.

**Yadav M, Kumar N, Singh D B and Singh G K. 2010.** Effect of post-harvest treatments on shelf-life and quality of Kinnow mandarin. *Indian Journal of Horticulture***67**: 243–8.

**Yurisqi Mukdisari, Ketty Suketi, Winarso Drajad Widodo. (2016).**Fruit Coating with Chitosan and Beeswax to Extend Papaya Shelf Life. Fruit Coating with Chitosan and Beeswax to Extend Papaya Shelf Life. *Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University.*

**Zambrano-Zaragoza, Maria L. (2020):** "Effect of nano-edible coating based on beeswax solid lipid nanoparticles on strawberry's preservation." *Coatings* 10.3 253.